



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

High Yolk Testosterone Transfer Is Associated with an Increased Female Metabolic Rate

Tschirren, Barbara; Ziegler, Ann-Kathrin; Canale, Cindy I; Okuliarová, Monika; Zeman, Michal;
Giraudeau, Mathieu

Abstract: Yolk androgens of maternal origin are important mediators of prenatal maternal effects. Although in many species short-term benefits of exposure to high yolk androgen concentrations for the offspring have been observed, females differ substantially in the amount of androgens they transfer to their eggs. It suggests that costs for the offspring or the mother constrain the evolution of maternal hormone transfer. However, to date, the nature of these costs remains poorly understood. Unlike most previous work that focused on potential costs for the offspring, we here investigated whether high yolk testosterone transfer is associated with metabolic costs (i.e., a higher metabolic rate) for the mother. We show that Japanese quail (*Coturnix japonica*) females that deposit higher testosterone concentrations into their eggs have a higher resting metabolic rate. Because a higher metabolic rate is often associated with a shorter life span, this relationship may explain the negative association between yolk testosterone transfer and female longevity observed in the wild. Our results suggest that metabolic costs for the mother can balance the short-term benefits of yolk testosterone exposure for the offspring, thereby contributing to the maintenance of variation in maternal yolk hormone transfer in natural populations.

DOI: <https://doi.org/10.1086/687571>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-126759>

Accepted Version

Originally published at:

Tschirren, Barbara; Ziegler, Ann-Kathrin; Canale, Cindy I; Okuliarová, Monika; Zeman, Michal; Giraudeau, Mathieu (2016). High Yolk Testosterone Transfer Is Associated with an Increased Female Metabolic Rate. *Physiological and Biochemical Zoology*, 89(5):448-452.

DOI: <https://doi.org/10.1086/687571>

1 *Brief Communication*

2 *Physiol Biochem Zool.* 2016 Sep-Oct;89(5):448-52. doi: 10.1086/687571

3 **High yolk testosterone transfer is associated with an increased female metabolic rate**

4

5 Barbara Tschirren^{a*}, Ann-Kathrin Ziegler^a, Cindy I. Canale^a, Monika Okuliarová^b,

6 Michal Zeman^b & Mathieu Giraudeau^{ac}

7 ^aDepartment of Evolutionary Biology and Environmental Studies, University of Zurich,

8 Winterthurerstrasse 190, 8057 Zurich, Switzerland

9 ^bDepartment of Animal Physiology and Ethology, Faculty of Natural Sciences, Comenius

10 University, Bratislava, Slovak Republic

11 ^cCentre for Ecology and Conservation, University of Exeter, Penryn, TR10 9FE

12 Cornwall, UK

13

14 *Correspondence: Barbara Tschirren, Department of Evolutionary Biology and

15 Environmental Studies, University of Zurich, Winterthurerstrasse 190, 8057 Zurich,

16 Switzerland; email: barbara.tschirren@ieu.uzh.ch, phone: +41 44 635 47 77, Fax: +41 44

17 635 47 80

18

19 Running title: Metabolic costs of yolk testosterone transfer

20

21

22 **What Is Already Known:** Yolk hormones of maternal origin are important mediators or

23 prenatal maternal effects in oviparous species. Within populations, there is large variation

24 in yolk hormone transfer observed among females, but the mechanisms that contribute to
25 the maintenance of this variation are poorly understood.

26

27 **What This Study Adds:** Unlike most previous work that has focused on potential costs
28 for the offspring, we here show that yolk hormone transfer is associated with metabolic
29 costs for the mother. These costs for the mother can balance beneficial effects on
30 offspring and thereby contribute to the maintenance of variation in maternal yolk
31 hormone transfer in animal populations.

32

33 **Abstract**

34 Yolk androgens of maternal origin are important mediators of prenatal maternal effects.
35 Although in many species short-term benefits of exposure to high yolk androgen
36 concentrations for the offspring have been observed, females differ substantially in the
37 amount of androgens they transfer to their eggs. It suggests that costs for the offspring or
38 the mother constrain the evolution of maternal hormone transfer. However, to date the
39 nature of these costs remains poorly understood. Unlike most previous work that focused
40 on potential costs for the offspring, we here investigated if high yolk testosterone transfer
41 is associated with metabolic costs (i.e. a higher metabolic rate) for the mother. We show
42 that Japanese quail (*Coturnix japonica*) females that deposit higher testosterone
43 concentrations into their eggs have a higher resting metabolic rate (RMR). Because a
44 higher metabolic rate is often associated with a shorter lifespan, this relationship may
45 explain the negative association between yolk testosterone transfer and female longevity
46 observed in the wild. Our results suggest that metabolic costs for the mother can balance
47 the short-term benefits of yolk testosterone exposure for the offspring, thereby
48 contributing to the maintenance of variation in maternal yolk hormone transfer in natural
49 populations.

50

51 **Keywords:** maternal effects; reproductive physiology; metabolic rate; trade-off;
52 maintenance of variation; metabolic costs; yolk hormone transfer

53 **Introduction**

54 Maternally transferred yolk androgens are important mediators of prenatal maternal
55 effects in birds and other oviparous species (Schwabl 1993). Many experimental studies
56 have shown that exposure to high androgen concentrations before birth can have
57 beneficial short-term effects on offspring, including an enhanced begging capacity or
58 boosted growth (reviewed in Groothuis et al. 2005b; Gil 2008). This has raised the
59 question why not all females deposit high androgen concentrations into their eggs, and
60 how the large between-female variation in yolk androgen deposition is maintained in
61 natural populations.

62 In an attempt to answer this question, researchers have mostly focused on
63 potential costs of prenatal exposure to high androgen concentrations for the offspring,
64 including immunosuppression (Groothuis et al. 2005a; Tschirren et al. 2005), metabolic
65 costs (Eising et al. 2003; Tobler et al. 2007; Nilsson et al. 2011) or sexual antagonism
66 (Ruuskanen et al. 2012; Tschirren 2015). The results of these studies are mixed. Costs
67 associated with yolk androgen transfer for the mother, on the other hand, have received
68 less attention (but see Pilz et al. 2003; Gil et al. 2006), even though the recent finding that
69 female collared flycatchers (*Ficedula albicollis*) that deposit high testosterone
70 concentrations into their eggs live shorter (Tschirren et al. 2014) indicates that direct
71 costs for the mother may be substantial.

72 A negative association between resting metabolic rate (RMR) and lifespan is the
73 cornerstone of the rate of living hypothesis (Pearl 1928), and evidence for such a
74 relationship has been found among, as well as within species (Ruggiero et al. 2008; but
75 see e.g. Duarte and Speakman 2014). If testosterone transfer to the eggs is associated with

76 metabolic costs for the mother, this could provide a proximate explanation for why
77 females that deposit high testosterone concentrations into their eggs live shorter. Here,
78 we tested if high maternal testosterone transfer is associated with an increased female
79 metabolic rate in a precocial bird, the Japanese quail (*Coturnix japonica*) to explore if
80 metabolic costs for the mother may constrain the evolution of maternal yolk hormone
81 deposition.

82

83 **Material and Methods**

84 *Study population*

85 The study was conducted in a population of Japanese quail (*Coturnix japonica*)
86 maintained at the University of Zurich, Switzerland. Males and females were housed in
87 separate outdoor aviaries (7 x 5.5 m each). For reproduction, male-female pairs (N = 40
88 pairs; age: 154-184 days) were transferred to cages (122 x 50 x 50 cm) within our
89 breeding facility. Cages contained *ad libitum* food, water, grit, a source of calcium, a
90 shelter and a sand bath. The bottom of the cages was lined with sawdust. The breeding
91 facility was kept on a 16 : 8 h light : dark cycle at $20 \pm 3^{\circ}\text{C}$ (see Pick et al. 2016 for
92 details).

93

94 *Yolk testosterone analysis*

95 We collected the 5th egg of each female on the day it was laid (natural clutch size: 7-14
96 eggs (Hoffmann 1988)), separated yolk and albumen, weighed ($\pm 0.01\text{g}$) and
97 homogenised the yolk, and froze it at -20°C (N = 40). Yolk testosterone (yolk T)
98 extraction and radioimmunoassay were performed following previously published

99 protocols (Okuliarová et al. 2011). In short, 100-110 mg of yolk were spiked with
100 approximately 2500 dpm of [³H]-testosterone (PerkinElmer, USA) and extracted twice
101 with a mixture of diethyl and petroleum ether. Yolk testosterone concentrations (pg / mg
102 yolk) were quantified in 10µl aliquots using [1,2,6,7-³H]-testosterone (PerkinElmer,
103 USA, specific activity 63.47 Ci / mmol) and a specific antibody generated in rabbits
104 against testosterone-3-(carboxy-methyl) oxime bovine serum albumin conjugate (Zeman
105 et al. 1986). The sensitivity of the assay was 1.62 ± 0.17 pg per tube. The mean recovery
106 rate \pm SD was $79.3 \pm 6.4\%$. The samples were analyzed in two assays. The intra- and
107 inter-assay coefficients of variation were 4.7% and 6.5%, respectively. Measured yolk
108 testosterone concentrations were log transformed before analysis.

109

110 *Metabolic rate*

111 Resting metabolic rate (RMR) of females (N = 40) was measured over six consecutive
112 nights (seven birds per night) in sealed plastic metabolic chambers (3.9 l, 234 x 165 x
113 165mm, Lock & Lock, Hanacobi Co. Ltd., Korea) during the birds' rest phase (6pm –
114 8:30am) in a dark room at 25-28°C, which is within the species' thermoneutral zone
115 (Ben-Hamo et al. 2010). Food was withdrawn from the cages for two hours before
116 measurements started to ensure a post-absorptive state. We measured the fractional
117 content of oxygen and carbon dioxide in the air using an eight-channel open flow
118 respirometry system (Sable Systems, Las Vegas, USA). Before each trial, the CO₂
119 analyzer was zeroed using CO₂-free air (dry nitrogen, 99.99% N₂, PanGas, Switzerland)
120 and spanned using a 1.002 % mol CO₂ mixture (balance N₂, PanGas, Switzerland). The
121 O₂ analyzer was spanned to 20.95% by flushing dry air through the system. During a

122 trial, external air was pumped into the chambers (seven containing a bird and one empty
123 control chamber) at a flow rate of 1650-1700 ml / min (Dual channel bench field pump,
124 3.5lpm maximal flow per channel, PP-2-1; Flow Bar Mass Flow Meter FB-8-1, Sable
125 System, USA). All gas flow connections passed through ultra-low permeability Tygon
126 tube (internal diameter: 8 mm).

127 Each recording sequence lasted 45 minutes and consisted of one round of measurements
128 of O₂, CO₂, flow rate and temperature for each of the seven chambers containing a bird.
129 The empty control chamber was measured at the beginning and the end of each sequence.
130 Excurrent air from the chambers was pushed through a RM-8-2 respirometry multiplexer
131 (Sable Systems International, Las Vegas, USA) programmed to serially divert individual
132 gas streams every five minutes. A subsample of each gas stream (250 ml / min) was
133 pulled through a dessicant column (magnesium perchlorate, Sigma-Aldrich, USA) before
134 being analyzed every second over a five minute period by a fuel cell O₂ analyzer and a
135 dual wavelength infrared bench CO₂ analyzer (Foxbox, Sable System, USA). To avoid
136 taking measurements during the drift period (i.e. warming up) of the Foxbox, it was
137 turned on 4 hours before the first recording started. Because of residual air in the system,
138 we excluded the first 100 seconds of each 5 min measurement. In total, we obtained 18-
139 20 200 sec measurement sequences per bird during the course of the night.

140 Oxygen consumption rates (VO₂, ml / min) were determined by comparing the oxygen
141 content of the metabolic chamber containing birds (F_e) and the empty control chamber
142 (F_i, baseline). Baseline O₂ and CO₂ were determined by regressing all control chamber
143 readings against time for each 45 minute sequence. VO₂ was calculated using the
144 following equation, which corrects for flow rate (FR) and CO₂ concentration: $VO_2 = FR$

145 * $((F_{iO_2} - F_{eO_2}) - F_{eO_2} * (F_{eCO_2} - F_{iCO_2})) / (1 - F_{eO_2})$. We determined the mean of 60
146 consecutive seconds of lowest VO_2 within a range of variation of < 0.015 ml / min (i.e.,
147 three-fold the standard deviation of the control chamber). RMR (W or J / sec) was
148 estimated from VO_2 and the respiratory exchange ratio (RER; VCO_2 / VO_2) using the
149 thermal equivalence data in Withers (1992). The average RER was 0.70, indicating a
150 predominance of lipid metabolism. The birds were weighed (± 0.1 g) after the RMR
151 measurement. Metabolic rate was measured after egg collection (see above). All females
152 were in breeding condition (i.e. egg laying) and in the same stage of the breeding cycle
153 when RMR was measured.

154 All experiments conform to the relevant regulatory standards and were conducted
155 under licenses provided by the Veterinary Office of the Canton of Zurich, Switzerland
156 (195/2010; 14/2014; 156).

157

158 *Statistical analysis*

159 We tested for an association between yolk T concentration and female RMR using a
160 linear mixed model that included female RMR and body mass as fixed effects. In addition,
161 we ran the same model but with the residuals of a linear regression of RMR on body mass
162 (residual RMR) as fixed effect. Because some of the females used in this study were
163 sisters, we included Family ID (N = 29 families) as a random effect in the models.
164 Residuals of the models were normally distributed. *P* values were obtained by comparing
165 two nested models, with and without the variable of interest, using likelihood ratio tests.
166 All analyses were performed in R (R Development Core Team 2011).

167

168 **Results**

169 We observed a significant positive association between the amount of testosterone a
170 female transfers to her eggs and her RMR ($\chi^2 = 5.496$, $P = 0.019$; Fig. 1A). Similarly,
171 there was a positive association between yolk T and a female's residual RMR ($\chi^2 = 5.243$,
172 $P = 0.022$; Fig. 1B). Although there was a strong positive relationship between RMR and
173 body mass ($\chi^2 = 18.688$, $P < 0.001$), no significant association between a female's body
174 mass and the T concentration in her eggs was observed ($\chi^2 = 0.423$, $P = 0.516$; Fig. 1C).
175 The results were similar when analyzing yolk T content instead of yolk T concentration
176 (yolk T content – mother RMR: $\chi^2 = 6.781$, $P = 0.009$; yolk T content – residual mother
177 RMR: $\chi^2 = 6.566$, $P = 0.010$).

178

179 **Discussion**

180 Our study shows that the transfer of high T concentrations to the eggs is associated with
181 an increased female metabolic rate. If the increased RMR associated with high yolk T
182 transfer affects a female's daily energy budget, which is likely (Buchanan et al. 2001), the
183 resulting allocation trade-off may play a key role in balancing costs and benefits of
184 differential maternal egg provisioning, and contribute to the maintenance of variation in
185 maternal yolk hormone transfer among females.

186 Currently, we can only speculate about the proximate mechanisms that underlie
187 the observed relationship between yolk T concentrations and female metabolic rate.
188 Components involved in the endocrine control of reproductive physiology could either
189 directly or indirectly explain the association. For example, the production of testosterone
190 in the theca and granulosa cells of the follicular wall (Hackl et al. 2003) or the transfer of

191 testosterone from the site of production to the developing yolk might be energetically
192 demanding. Alternatively, the observed association could be indirect, mediated, for
193 example, by circulating T levels in the mother. Indeed, experimental studies have shown
194 that high plasma T levels can lead to an increased metabolic rate in both birds (Buchanan
195 et al. 2001) and men (Welle et al. 1992). However, this scenario is unlikely given that in
196 quail maternal plasma T levels and T concentrations in the eggs appear to be unrelated
197 (Hackl et al. 2003; Okuliarová et al. 2011). Finally, behavioral differences among
198 females may indirectly affect both yolk testosterone transfer and metabolism (van Oers et
199 al. 2011; Bouwhuis et al. 2014).

200 Irrespective of the exact physiological or behavioural mechanisms that link testosterone
201 transfer to the egg and female metabolic rate, costs associated with a high metabolic rate
202 (e.g. Ruggiero et al. 2008; Dowling and Simmons 2009), may (at least partly) explain
203 why females that transfer high yolk T concentrations to their eggs have a shorter lifespan
204 (Tschirren et al. 2014). Ultimately, our study highlights that to understand the evolution
205 of maternal effects as well as their mediators, it is crucial to consider the costs and
206 benefits for all players involved.

207

208 **Acknowledgements**

209 We thank the quail team for help with animal husbandry and data collection and three
210 reviewers for comments on the manuscript. The study was supported by the Swiss
211 National Science Foundation (PP00P3_128386 and PP00P3_157455) and the Fonds zur
212 Förderung des akademischen Nachwuchses (FAN).

213 **Literature Cited**

- 214 Ben-Hamo M., B. Pinshow, M.D. McCue, S.R. McWilliams, and U. Bauchinger. 2010.
215 Fasting triggers hypothermia, and ambient temperature modulates its depth in
216 Japanese quail *Coturnix japonica*. *Comp Biochem Physiol A* 156:84-91.
- 217 Bouwhuis S., J.L. Quinn, B.C. Sheldon, and S. Verhulst. 2014. Personality and basal
218 metabolic rate in a wild bird population. *Oikos* 123:56-62.
- 219 Buchanan K.L., M.R. Evans, A.R. Goldsmith, D.M. Bryant, and L.V. Rowe. 2001.
220 Testosterone influences basal metabolic rate in male house sparrows: a new cost
221 of dominance signalling? *Proc R Soc B* 268:1337-1344.
- 222 Dowling D.K. and L.W. Simmons. 2009. Reactive oxygen species as universal
223 constraints in life-history evolution. *Proc R Soc B* 276:1737-1745.
- 224 Duarte L.C. and J.R. Speakman. 2014. Low resting metabolic rate is associated with
225 greater lifespan because of a confounding effect of body fatness. *Age*
226 36:doi:10.1007/s11357-11014-19731-11353.
- 227 Eising C.M., G.H. Visser, W. Müller, and T.G.G. Groothuis. 2003. Steroids for free? No
228 metabolic costs of elevated maternal androgen levels in the black-headed gull. *J*
229 *Exp Biol* 206:3211-3218.
- 230 Gil D. 2008. Hormones in avian eggs: Physiology, ecology and behavior. *Adv Study*
231 *Behav* 38:337-398.
- 232 Gil D., A. Marzal, F. de Lope, M. Puerta, and A.P. Møller. 2006. Female house martins
233 (*Delichon urbica*) reduce egg androgen deposition in response to a challenge of
234 their immune system. *Behav Ecol Sociobiol* 60:96-100.

235 Groothuis T.G.G., C.M. Eising, C. Dijkstra, and W. Müller. 2005a. Balancing between
236 costs and benefits of maternal hormone deposition in avian eggs. *Biol Lett* 1:78-
237 81.

238 Groothuis T.G.G., W. Müller, N. von Engelhardt, C. Carere, and C. Eising. 2005b.
239 Maternal hormones as a tool to adjust offspring phenotype in avian species.
240 *Neurosci Biobehav Rev* 29:329-352.

241 Hackl R., V. Bromundt, J. Daisley, K. Kotrschal, and E. Möstl. 2003. Distribution and
242 origin of steroid hormones in the yolk of Japanese quail eggs (*Coturnix coturnix*
243 *japonica*). *J Comp Physiol B* 173:327-331.

244 Hoffmann E. 1988. *Coturnix quail*. Yi Hsien Publishing Co., Taipei.

245 Nilsson J.F., M. Tobler, J.A. Nilsson, and M.I. Sandell. 2011. Long-lasting consequences
246 of elevated yolk testosterone for metabolism in the Zebra finch. *Physiol Biochem*
247 *Zool* 84:287-291.

248 Okuliarová M., T.G.G. Groothuis, P. Skrobanek, and M. Zeman. 2011. Experimental
249 evidence for genetic heritability of maternal hormone transfer to offspring. *Am*
250 *Nat* 177:824-834.

251 Pearl R. 1928. *The rate of living*. University of London Press, London.

252 Pick J.L., P. Hutter, and B. Tschirren. 2016. In search of genetic constraints limiting the
253 evolution of egg size: direct and correlated responses to artificial selection on a
254 prenatal maternal effector. *Heredity*. doi: 10.1038/hdy.2016.16.

255 Pilz K.M., H.G. Smith, M.I. Sandell, and H. Schwabl. 2003. Interfemale variation in egg
256 yolk androgen allocation in the European starling: do high-quality females invest
257 more? *Anim Behav* 65:841-850.

258 R Development Core Team. 2011. R: A language and environment for statistical
259 computing. <http://www.R-project.org>.

260 Ruggiero C., E.J. Metter, V. Melenovsky, A. Cherubini, S.S. Najjar, A. Ble, U. Senin,
261 D.L. Longo, and L. Ferrucci. 2008. High basal metabolic rate is a risk factor for
262 mortality: The Baltimore Longitudinal Study of Aging. *J. Gerontol. A* 63:698-706.

263 Ruuskanen S., B. Doligez, N. Pitala, L. Gustafsson, and T. Laaksonen. 2012. Long-term
264 fitness consequences of high yolk androgen levels: sons pay the costs. *Funct Ecol*
265 26:884-894.

266 Schwabl H. 1993. Yolk is a source of maternal testosterone for developing birds. *Proc*
267 *Natl Acad Sci U S A* 90:11446-11450.

268 Tobler M., J.A. Nilsson, and J.F. Nilsson. 2007. Costly steroids: egg testosterone
269 modulates nestling metabolic rate in the zebra finch. *Biol Lett* 3:408-410.

270 Tschirren B. 2015. Differential effects of maternal yolk androgens on male and female
271 offspring: a role for sex-specific selection? *PLoS ONE* 10:e0133673.
272 doi:0133610.0131371/journal.pone.0133673.

273 Tschirren B., E. Postma, L. Gustafsson, T.G.G. Groothuis, and B. Doligez. 2014. Natural
274 selection acts in opposite ways on correlated hormonal mediators of prenatal
275 maternal effects in a wild bird population. *Ecol Lett* 17:1310-1315.

276 Tschirren B., V. Saladin, P.S. Fitze, H. Schwabl, and H. Richner. 2005. Maternal yolk
277 testosterone does not modulate parasite susceptibility or immune function in great
278 tit nestlings. *J Anim Ecol* 74:675-682.

279 van Oers K., K.L. Buchanan, T.E. Thomas, and P.J. Drent. 2011. Correlated response to
280 selection of testosterone levels and immunocompetence in lines selected for avian
281 personality. *Anim Behav* 81:1055-1061.

282 Welle S., R. Jozefowicz, G. Forbes, and R.C. Griggs. 1992. Effect of testosterone on
283 metabolic rate and body composition in normal men and men with muscular
284 dystrophy. *J Clin Endocrinol Metab* 74:332-335.

285 Withers P.C. 1992. *Comparative animal physiology*. Saunders College Pub., Fort Worth.


286 Zeman M., J. Kosutzky, and E. Bobakova. 1986. Testosterone concentration in the
287 seminal plasma of cocks. *Br Poult Sci* 27:261-266.

288

289 **Figure Legend**

290

291 Fig. 1. Relationship between testosterone concentrations in the eggs and A) a female's
292 resting metabolic rate (RMR), B) the residuals of a regression of female RMR on body
293 mass (residual RMR), and C) a female's body mass (N = 40 females).



294