



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
Main Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2017

---

**Wild type agr-negative livestock-associated MRSA exhibits high adhesive  
capacity to human and porcine cells**

Bünter, Julia P; Seth-Smith, Helena M B; Rüegg, Simon; Heikinheimo, Annamari; Borel, Nicole; Jöhler, Sophia

DOI: <https://doi.org/10.1016/j.resmic.2016.09.006>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-127229>

Journal Article

Accepted Version

Originally published at:

Bünter, Julia P; Seth-Smith, Helena M B; Rüegg, Simon; Heikinheimo, Annamari; Borel, Nicole; Jöhler, Sophia (2017). Wild type agr-negative livestock-associated MRSA exhibits high adhesive capacity to human and porcine cells. *Research in Microbiology*, 168(2):130-138.

DOI: <https://doi.org/10.1016/j.resmic.2016.09.006>

1 **Wild type *agr*-negative livestock-associated MRSA exhibits high**  
2 **adhesive capacity to human and porcine cells**

3

4 Julia P. Bünter<sup>1</sup>, Helena M.B. Seth-Smith<sup>1</sup>, Simon Rüegg<sup>2</sup>, Annamari Heikinheimo<sup>4</sup>, Nicole  
5 Borel<sup>1\*</sup>†, Sophia Johler<sup>3</sup>†

6

7

8 <sup>1</sup> Institute for Veterinary Pathology, Vetsuisse Faculty, University of Zurich,  
9 Winterthurerstrasse 268, CH-8057 Zurich, Switzerland

10

11 <sup>2</sup> Section of Epidemiology, Vetsuisse Faculty, University of Zürich, Winterthurerstrasse 270,  
12 CH-8057 Zurich, Switzerland

13

14 <sup>3</sup> Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich,  
15 Winterthurerstrasse 272, CH-8057 Zurich, Switzerland

16

17 <sup>4</sup> Department of Food Hygiene and Environmental Health, Faculty of Veterinary Medicine,  
18 University of Helsinki, Agnes Sjöberginkatu 2, P.O. Box 66, FI-00014 Helsinki, Finland

19

20 \* Corresponding author

21 Email: n.borel@access.uzh.ch (NB)

22 †These authors contributed equally to this work.

## 23 Abstract

24

25 Methicillin-resistant *Staphylococcus aureus* (MRSA) are one of the leading causes of  
26 nosocomial infections and a major public health concern worldwide. During the last decade,  
27 MRSA of CC398 have emerged as important colonizers of livestock. These strains also  
28 represent an increasing cause of human infections. A recent study reporting a new dominant  
29 *spa* type among MRSA from Finish fattening pigs (CC398/t2741) identified a strain lacking  
30 both the global virulence regulator gene locus *agr* and the adhesion gene *fnbB*. The aim of  
31 this study was to characterize this *agr/fnbB*-negative livestock-associated MRSA strain in  
32 terms of growth, hemolysis, and adhesive capacity, and to provide data on its genomic  
33 background. To this end, growth curves and hemolysis patterns were generated and adhesion  
34 assays on human keratinocyte and porcine nasal mucosa cell lines were performed. Whole  
35 genome sequencing was used to determine the nature and extent of the relevant deletions in  
36 the livestock strains. For comparison, an *agr*-positive, *fnbB*-negative CC398/t2741 strain  
37 from the same pig herd, an *agr/fnbB*- positive CC398/t034 strain from another pig herd, and  
38 one human MRSA strain and its isogenic  $\Delta$ *agr* knockout mutant were used. The *agr*-negative  
39 strains adhered significantly better to human and porcine host cells than the *agr*-positive  
40 control strains. For the *agr*-positive porcine MRSA strains, cytotoxic effects on porcine  
41 mucosal cells were observed. The strong adhesive capacity of the naturally *agr*-negative  
42 livestock-associated MRSA in combination with diminished cytotoxic effects is likely  
43 favorable for inducing persistent colonization in pigs. Independent of the host cell type,  
44 similar adhesive capacities of the naturally *agr*-negative livestock-associated MRSA and the  
45 human MRSA strain were shown. Our results indicate that loss of *agr* in the livestock-

46 associated MRSA strain investigated in this study may have increased its potential to be  
47 transmitted to and amongst humans.

## 48 **Introduction**

49

50 Livestock-associated MRSA (LA-MRSA) strains have emerged worldwide. The strains  
51 typically belong to CC398 and are detected in particularly high rates among pigs [1-4]. LA-  
52 MRSA CC398 are thought to have originated from human methicillin-susceptible  
53 *Staphylococcus aureus* and acquired methicillin resistance during the host jump, at the same  
54 time losing genes important for colonization and infection in humans [5]. Even though LA-  
55 MRSA of CC398 lack important virulence factors, they have been suggested to exhibit an  
56 enhanced ability to acquire virulence factors through mobile genetic elements [6].

57 Transmission of CC398 from animals to humans has been described and occupational  
58 exposure to pigs was shown to increase nasal carriage rates (pig farmers: 20-86%;  
59 veterinarians: 4.6-45%) compared to the general population (0.8-3%) [2, 4, 7-14]. In addition,  
60 in recent years, there have been an increasing number of reports of MRSA CC398 strains  
61 causing invasive infections in humans without contact to livestock [15-18]. Therefore, the  
62 transmissibility and the virulence potential of LA-MRSA of CC398 need to be closely  
63 monitored.

64 Heikinheimo et al. [19] recently identified a new dominant *spa* type among LA-MRSA  
65 strains colonizing Finnish fattening pigs (CC398/t2741). Interestingly, for one CC398/t2741  
66 strain isolated in this study, no genes within the accessory gene regulator locus (*agr*) or the  
67 fibronectin-binding protein B (*fnbB*) were detected by DNA microarray. The quorum sensing  
68 *agr* system is a major virulence gene regulator. Activation of *agr* has been shown to inhibit  
69 expression of certain cell-wall associated proteins and to increase exoprotein expression  
70 during the post-exponential phase of growth [20, 21]. While *agr* plays a major role in acute

71 infections, *agr*-defective mutants can frequently be isolated from bacteraemic patients and  
72 were linked to persistent infections [22] and increased mortality [23]. The *fnbB* gene encodes  
73 the fibronectin binding protein B (FnBPB), which belongs to the group of microbial surface  
74 components recognizing adhesive matrix molecules (MSCRAMMs). It facilitates adhesion of  
75 *S. aureus* to the components of the host cells and has a role in invasion [24]. Combined  
76 FnBPA and FnBPB function has been shown to be essential to induce severe infection [25].  
77 The aim of this study was to characterize the naturally *agr/fnbB*-negative CC398/t2741 LA-  
78 MRSA strain detected in Finland with regard to growth, hemolysis, and adhesive capacity,  
79 and to describe the genomic background of the strain.

80

## 81 **Methods**

### 82 **Bacterial strains**

83 An overview of the strains used in this study is presented in Table 1. The naturally  
84 *agr/fnbB*-negative LA-MRSA strain investigated in this study (Fin47\_H17) was isolated from  
85 the carcass of a fattening pig in Finland [19]. Two porcine LA-MRSA strains isolated in the  
86 same study were used as controls: Fin46\_H17, an *agr*-positive, *fnbB*-negative isolate from the  
87 same herd (herd 17), and Fin48\_H18, an *agr/fnbB*-positive isolate from another herd (herd  
88 18). As additional controls, a MRSA strain that had caused recurring skin infections in  
89 humans (MN10) [26] and its isogenic *agr* knockout mutant (MN10\_  $\Delta$ *agr*) were used.  
90 MN10\_  $\Delta$ *agr* was obtained by transduction of the *agr* knockout from RN6911 to MN10 using  
91 phage 80 $\alpha$  and protocols previously described [27]. Correct deletion of *agr* in putative  
92 mutants grown on tetracycline selective plates was confirmed by PCR.

93

94 **Table 1: Strains used in this study.** Presence/absence of *agr* and *fnbB* is indicated as  
 95 previously determined by DNA microarray [19].

Strain ID	Characteristics	Country	Year	Source [reference]
<b>Fin47_H17</b>	CC398/t2741, <i>agr</i> <sup>-</sup> , <i>fnbB</i> <sup>-</sup>	FI	2015	Pig carcass, herd 17 [19]
<b>Fin46_H17</b>	CC398/t2741, <i>agr</i> <sup>+</sup> , <i>fnbB</i> <sup>-</sup>	FI	2015	Pig nares, herd 17 [19]
<b>Fin48_H18</b>	CC398/t034, <i>agr</i> <sup>+</sup> , <i>fnbB</i> <sup>+</sup>	FI	2015	Pig nares, herd 18 [19]
<b>MN10</b>	CC5/t002, <i>agr</i> <sup>+</sup> , <i>fnbB</i> <sup>+</sup>	CH	2013	Human skin infection [26]
<b>MN10_Δ<i>agr</i></b>	CC5/t002, NM10 <i>agr</i> :: <i>tetM</i> , <i>agr</i> <sup>-</sup> , <i>fnbB</i> <sup>+</sup>	CH	2013	This study
<b>RN27</b>	80alpha lysogen	-	-	Brigitte Berger-Bächi [28]
<b>RN6911</b>	<i>agr</i> operon deleted ( <i>agr</i> :: <i>tetM</i> )	-	-	Brigitte Berger-Bächi [28]

96

## 97 **Growth parameters**

98           Single colonies were transferred from 5% sheep blood agar to 50 mL of Luria Bertani  
99 (LB) broth (Becton Dickinson, Allschwil, Switzerland) and grown for 24 h at 37°C (225  
100 rpm). Growth of all strains in LB was determined by viable cell counts after 3 h, 6 h, 9 h, and  
101 24 h using 10-fold dilution series and plate count agar (Sigma-Aldrich, Stockholm, Sweden),  
102 with incubation at 37°C for 18-24 h. Growth parameters such as exponential phase growth  
103 rate and maximum cell density were determined using DMFit 3.0 [29].

104

## 105 **Hemolysis**

106           Screening for hemolytic activity was performed as previously described [30]. Briefly,  
107 alpha- and delta-hemolytic activity was determined by perpendicular streaking to the beta-  
108 hemolysin producing *S. aureus* reference strain RN4220 on 5% sheep blood agar and  
109 incubation over night at 37°C. In this assay, beta hemolysis results in a turbid zone. Alpha  
110 hemolytic activity of the test strain is inhibited by beta hemolysis of RN4220 where the  
111 strains intersect. Delta hemolysis of the test strain is synergistic with beta hemolysin of  
112 RN4220, resulting in an amplified zone of clearing where the strains intersect [31]. Gamma  
113 hemolytic activity cannot be detected in this assay, as it is inhibited by agar [32]. Subsequent  
114 cold shock (exposure to 4°C for 12h) was used to determine beta-hemolysis activity.

115

## 116 **Adhesion assay**

117           Two different cell lines were used for adhesion assays: human cell line HaCaT  
118 (Human adult low Calcium high Temperature keratinocytes, CLS Cell Lines Service GmbH,

119 Eppelheim, Germany; [33]) and porcine cell line PT-K75 (porcine nasal turbinate/mucosa,  
120 CRL-2528, ATCC, Manassas, USA). The medium for the adhesion assay consisted of  
121 DMEM supplemented with 4.5 g/L glucose, 584 mg/L L-glutamine and 10% FCS (CLS Cell  
122 Lines Service GmbH) for the HaCaT cells and DMEM supplemented with 4.5 g/L glucose,  
123 580mg/L L-glutamine, 110mg/L sodium pyruvate (GIBCO, Thermo Fisher Scientific,  
124 Invitrogen, Carlsbad, CA, USA) and 10% FCS (BioConcept, Allschwil, Switzerland) for the  
125 PT-K75 cells.

126 After at least one passage in a 75cm<sup>2</sup> canted neck tissue culture flask with vented cap  
127 (Corning, Sigma-Aldrich), HaCaT-cells were seeded at a concentration of  $2 \times 10^5$  cells/well  
128 and PT-K75-cells at a concentration of  $1.25 \times 10^5$  cells/well in a 24-well flat-bottom cell  
129 culture plate with low-evaporation lid (Techno Plastic Products AG, TPP, Trasadingen,  
130 Switzerland). The culture plates were incubated for 24 hours at 37°C and 5% CO<sub>2</sub> to reach a  
131 confluency of at least 80%.

132 Bacterial strains were prepared by growth on 5% sheep blood agar overnight at 37°C.  
133 Overnight cultures were prepared by inoculation of 5 mL LB broth with a single colony and  
134 incubation at 37°C (225 rpm shaking) for 15 hours. Cultures were subsequently adjusted to  
135  $OD_{590} = 0.40$  ( $\sim 10^8$  CFU/ml) for each strain. The bacteria were used at a multiplicity of  
136 infection (MOI) of 10 to infect both cell lines (HaCaT and PT-K75) in 24-well plates.  
137 Uninfected cells were used as negative controls. After 15 min of incubation at 37°C, cells  
138 were imaged, before monolayers were washed eight times with Dulbecco's phosphate  
139 buffered saline (DPBS, GIBCO) and cells were harvested after five minutes of incubation  
140 with Trypsin-EDTA (0.25%, GIBCO) by scraping with a cell scraper (TPP). Bacterial cell  
141 counts were determined by 10-fold dilution series and plate count agar. Three independent

142 biological experiments were performed, each using three replicates per strain. For imaging  
143 purposes, the Nikon Eclipse Ti-U inverted microscope was used at 200-fold magnification  
144 with software NIS-Elements AR Analysis 4.3 (Nikon AG, Egg, Switzerland).

145

## 146 **Statistical analysis**

147 All statistical analyses with the exception of growth parameter analyses were  
148 performed using R, 3.2. Results were compared using a gaussian linear model of the form  
149  $\text{count} \sim \text{strain} + \text{host.cellline}$ . Growth parameters were compared between strains using SPSS  
150 Statistics 22 (SPSS Inc.Chicago, Illinois). Growth parameters such as exponential phase  
151 growth rate and maximum cell density were determined using DMFit 3.0 (Baranyi &  
152 Roberts, 1994) and compared using one-way ANOVA. Differences were considered  
153 statistically significant if  $p < 0.05$ .

154

## 155 **Whole genome sequencing**

156 DNA was extracted from overnight cultures grown on 5% sheep blood agar plates  
157 using the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) according to the  
158 manufacturer's instructions. The porcine LA-MRSA strains Fin47\_H17, Fin46\_H17, and  
159 Fin48\_H18 were sequenced on the Illumina Miseq platform with 250bp paired end reads  
160 following NEBNextUltra library creation. Raw sequencing reads were adaptor and quality  
161 trimmed and filtered using Trimmomatic version 0.32 [34]. Mapping against the CC398  
162 reference strain S0385 (EMBL Accession AM990992) [6] using BWA [35] indicated  
163 chromosome coverage of each genome to be: Fin47\_H17 =  $132.8x \pm 76.2$ , Fin46\_H17 =

164 140.6 x  $\pm$ 71.2, Fin48\_H18 = 115.1x  $\pm$  53.7. *De novo* assembly used SPAdes v 3.1.0 in multi-  
165 cell mode [36]. Scaffolds were rearranged in ACT [37] against S0385 using files generated in  
166 WebACT (<http://www.webact.org/>). Resulting assemblies comprise 28, 25 and 27 scaffolds  
167 respectively, with scaffolds under 1kb, which did not match the reference, excluded. Target  
168 coding sequences (CDSs) involved in adhesion were identified and analysed in all genomes  
169 using ACT and, where necessary, single CDSs were aligned using ClustalW. The phylogeny  
170 was generated in <http://wgsastaging.pathogensurveillance.net/> using available assembled  
171 CC398 genomes for context [6, 38-42]. Antibiotic resistance phenotypes were also predicted  
172 using this online tool.

173 All read data and assemblies have been deposited with ENA under project number  
174 (PRJEB14187).

175

## 176 **Results**

### 177 **Growth parameters**

178 No statistically significant differences were observed between growth of the five  
179 strains in LB broth with regard to exponential phase growth rate or maximum cell density  
180 (Fig. 1). However, the naturally *agr*-negative LA-MRSA Fin47\_H17 exhibited the lowest  
181 exponential phase growth rate ( $\mu_{\max} = 0.50$ ) compared to all other strains ( $\mu_{\max} = 0.51$ - $0.57$ ).  
182 Fin47\_H17 also reached the lowest maximum cell density (8.97 log CfU/mL) compared to all  
183 other strains tested (9.70-9.91 log CfU/mL).

184

185 **Fig. 1: Growth of all tested strains in LB over 24h.** The tested strains exhibited no  
186 statistically significant differences in exponential phase growth rate or maximum cell density  
187 reached (error bars: 95% CI).

188

### 189 **Haemolysis pattern**

190 Hemolysis patterns after incubation at 37°C overnight and after subsequent cold  
191 exposure are shown in Fig. 2. No hemolysis was visible for both the naturally *agr*-negative  
192 Fin47\_H17 strain and *agr* knockout mutant MN10\_  $\Delta agr$ . Parental strain MN10 exhibited  
193 alpha and delta hemolysis. Fin46\_H17 exhibited alpha and beta hemolysis, and Fin48\_H18  
194 exhibited beta and delta hemolysis.

195

196 **Fig. 2: Hemolysis.** The *agr*-negative strains exhibited no hemolysis (Fin47\_H17 and MN\_  
197  $\Delta agr$ ). MN10 exhibited alpha and delta hemolysis, Fin46\_H17 exhibited alpha and beta  
198 hemolysis, and Fin48\_H18 exhibited beta and delta hemolysis.

199

## 200 **Adhesion properties**

201 While no morphological changes were observed in HaCaT cells after bacterial  
202 infection, infection of PT-K75 cells with the *agr*-positive porcine LA-MRSA strains  
203 (Fin46\_H17 and Fin48\_H18) resulted in a cytotoxic effect including shrinking, rounding and  
204 detachment of the cells (Fig. 3). To minimize cytotoxicity, a trial experiment was performed  
205 on PT-K75 cells with an MOI of 1: cytotoxic effects were still observed (data not shown). In  
206 contrast, neither the naturally *agr*-negative porcine LA-MRSA strain Fin47\_H17 nor the  
207 human-associated MRSA strain MN10 and its isogenic *agr* knockout mutant induced any  
208 morphological changes in the PT-K75 cells (Fig. 3).

209 Counts of adherent bacteria revealed that the *agr*-negative strains Fin47\_H17 and  
210 MN10\_ $\Delta agr$  displayed increased adhesive capacity in both human HaCaT and porcine PT-  
211 K75 cells (Fig. 4) compared to the *agr*-positive LA-MRSA and the human-associated  
212 parental MRSA strain MN10, respectively. Median colony counts for Fin47\_H17 were 3,600  
213 for HaCaT and 14,000 for PT-K75 cells per culture well. These results show significantly  
214 higher adhesive capacity of this strain compared to the *agr*-positive strains Fin46\_H17 (790  
215 to HaCaT,  $p = 1.3e-06$ ), Fin48\_H18 (500 to HaCaT,  $p = 2.0e-07$ ) or MN10 (1,200 to HaCaT,  
216  $p = 4.3e-13$ ; 7,200 to PT-K75,  $p = 0.001$ ). The median colony counts for the MN10\_ $\Delta agr$   
217 mutant strain (3,190 to HaCaT and 35,000 to PT-K75) were significantly higher than those of  
218 MN10 (1,200 to HaCaT,  $p = 0.003$ ; 7,200 to PT-K75,  $p = 4.5e-05$ ). Thus, the naturally *agr*-

219 negative LA-MRSA and MN10\_Δagr exhibited significantly higher adhesive capacity than  
220 the agr-positive LA-MRSA strains and parental strain MN10, respectively. Overall, the tested  
221 porcine and human MRSA strains adhered significantly better to the porcine than to the  
222 human cells ( $p = 1.3e-06$ ).

223

224 **Fig. 3: Cytotoxic effects on porcine cells.** Representative images of the two cell lines before  
225 ( $T_0$ ) and shortly after ( $T_{15 \text{ min}}$ ) infection. Bacterial strains used were the porcine LA-MRSA  
226 strains Fin47\_H17, Fin46\_H17, and Fin48\_H18, as well as the MRSA strain linked to human  
227 infections (MN10) and its isogenic knockout mutant MN10\_Δagr. Cytotoxic effects of the  
228 agr-positive porcine LA-MRSA strains Fin46\_H17 and Fin48\_H18 on PT-K75 cells can be  
229 identified at  $T_{15 \text{ min}}$ .

230

231 **Fig. 4: Adhesion to A) human cells and B) porcine cells: colony counts for different**  
232 **strains.** The plots indicate the median (horizontal line), the range from the first to the third  
233 quartile (25–75%; box), and the extreme values (whiskers) of the colony counts for each  
234 strain after adhesion to A) human cells (HaCaT) and B) porcine cells (PT-K75). The data  
235 resulted from three independent biological experiments with three replicates for each strain  
236 ( $n=9$ ).

237

## 238 **Genomic analysis**

239 Whole genome sequencing and phylogenetic analysis of strains within CC398  
240 confirmed that Fin47\_H17 is closely related to Fin46\_H17, with Fin48\_H18 more distantly  
241 related (S1 Fig).

242 In the naturally *agr*-negative LA-MRSA Fin47\_H17, a deletion of 3,789 bp covering *hld*,  
243 *agrB* and *agrC* (Fig. 5A) was identified. In both this strain and the closely related strain  
244 Fin46\_H17, a further deletion completely removing *fnbB* was detected that leaves the  
245 adjacent *fnbA* intact (Fig. 5B). However, as *fnbB* in both Fin48\_H18 and reference strain  
246 S0385 is truncated after 349 amino acids with a single nucleotide mutation causing a  
247 premature stop codon, the functional relevance of this is unclear.

248 Genes associated with adhesion were investigated within the genomes (S2 Table). Such genes  
249 often have varying repeat lengths, and this was identified in these strains in the genes: *spa*,  
250 *sdrC* and the hypothetical protein encoding CDS *SAPIG1791*; with the further genes *coa*,  
251 *sdrD*, *clfA*, *clfB*, *ebhA* and *cna* almost identical between the three porcine LA-MRSA strains,  
252 but variable in comparison to homologues in S0385. The *vwb* gene, encoding a secreted von  
253 Willebrand factor-binding protein, was found to be absent at the equivalent chromosomal  
254 locus in Fin48\_H18, with a 13.7 kb deletion covering this region. All other genes investigated  
255 were found to be identical or almost identical to those in S0385 (*eno*, *isdB*, *isdA*, *ecb*, *fib*,  
256 *ebpS*, *map*, *sdrH*, *agrA*, *isaB*).

257 The strains carry alternative complements of plasmids and phage (S2 Table) and the  
258 predicted antibiotic resistance profiles (Fin47\_H17 and Fin46\_H17: PEN<sup>R</sup>, TET<sup>R</sup>, ERY<sup>R</sup>,

259 MET<sup>R</sup>; Fin 48\_H18: PEN<sup>R</sup>, TET<sup>R</sup>, MET<sup>R</sup>, TRI<sup>R</sup>) largely correspond to those phenotypically  
260 determined previously, with the exception of clindamycin [19].

261

262 **Fig. 5: Comparison of genome loci around *agr* and *fnbB*.** Each line refers to a genomic  
263 section in the strain indicated. Arrows indicate CDSs. Grey bars show homology between  
264 loci according to the scale. **A.** *agr* locus showing the extent of the deletion in Fin47\_H18,  
265 covering *SAPIG2071-2075*. *hld* is shown in yellow and the *agr* locus (*agrBCA*) in green.  
266 Only *argA* remains in Fin47\_H17. **B.** *fnbB* locus showing the deletion of *fnbB* (*SAPIG2551*)  
267 in both Fin47\_H17 and Fin46\_H17. *fnbB* in both S0385 and Fin48\_H18 is truncated by a  
268 stop codon and is shown in brown for a pseudogene. *fnbA* (yellow) displays some homology  
269 to *fnbB*. Figure was drawn in Easyfig [43].

270

## 271 Discussion

272 Heikinheimo et al. [19] investigated LA-MRSA isolated from Finnish fattening pigs  
273 at slaughter. They found that CC398/t2741 strains were predominant among the isolated LA-  
274 MRSA. Interestingly, one of the isolated strains (Fin47\_H17) did not harbor genes from the  
275 *agr* locus. In this study, we determined the genomic basis of this loss of *agr* and its effect on  
276 growth, hemolytic activity, and adhesive capacity to human and porcine cell lines.

277 Loss of *agr* did not result in altered growth parameters, but had a strong impact on  
278 hemolytic activity consistent with previous studies suggesting that expression of alpha and  
279 delta hemolysins is strongly induced by *agr* [30, 31]. Hemolysins are widely recognized as

280 important virulence factors in *S. aureus* infections. However, selective survival of *agr*-  
281 defective non-hemolytic *S. aureus* in wound and abscess models has been demonstrated [44].

282 In this study, both *agr*-negative strains tested exhibited increased adhesive capacity to  
283 both human and porcine cells. These results are consistent with previous findings suggesting  
284 *agr* to inhibit the expression of adhesion proteins [45-48]. Loss of *agr* was shown to  
285 significantly increase adhesive capacity to human endothelial [49] and mesothelial cells [50].  
286 In a bovine mammary epithelial cell line (MAC-T), two- to threefold higher numbers of  
287 internalized viable bacteria of an *agr* mutant were recovered compared to the wild type [51].

288 LA-MRSA CC398 was reported to cause more cell damage within epithelial cells  
289 than community-associated and hospital-acquired MRSA [52]. In this study, pronounced  
290 cytotoxic effects of the *agr*-positive porcine LA-MRSA strains on porcine cells were  
291 identified. In contrast, the closely related *agr*-negative porcine LA-MRSA Fin47\_H17 did not  
292 induce cytotoxic effects. This may be due to reduced expression levels of *agr*-dependent  
293 cytotoxic proteins such as alpha hemolysin [53]. It is also consistent with a previous study  
294 reporting cytotoxic effects in MAC-T cells upon infection with RN6390, but no cytotoxic  
295 effects in *agr* mutant strain RN6911 [51]. The finding that porcine *agr*-positive LA-MRSA  
296 only led to cytotoxicity in the porcine cell line could be due to host-specific factors  
297 influencing susceptibility to alpha hemolysin [54].

298 Whole genome sequencing revealed that the *fnbB* gene in Fin48\_H18 is truncated, potentially  
299 impairing functionality of FnBPB, whereas it is fully deleted from the genomes of strains  
300 Fin47\_17 and Fin46\_17. We observed no significant differences concerning adherence to  
301 porcine cells of Fin48\_H18 and Fin46\_17. For strain Newman, it has been suggested that  
302 truncation of *fnbB* was transferred to *fnbA*, resulting in complete secretion FnBPs to the

303 culture medium and loss of cell wall anchor function [55]. Consistent with these findings,  
304 reduced adhesive capacity of a *fnbAB* mutant to HaCaT cells has been described [56].

305 We observed no significant differences in adhesive capacity between porcine and  
306 human MRSA apart from the described influence of *agr*. However, irrespective of strain  
307 origin (LA-MRSA or human MRSA), adherence to porcine cells significantly exceeded  
308 adherence to human cells. LA-MRSA CC398 do not only colonize livestock, but can also be  
309 transmitted to humans with direct livestock exposure or even between humans. Although  
310 decreased adherence of LA-MRSA compared to community/hospital-associated MRSA to  
311 human endothelial and epithelial cells was observed [52], LA-MRSA strains may rapidly  
312 adapt to the human host. Our results suggest that LA-MRSA are capable of adherence to both  
313 porcine and human cells with the same efficiency as a community-acquired MRSA control  
314 strain linked to recurring infections within a family [26].

315

## 316 **Conclusion**

317 We were able to show that the recently reported naturally *agr*-negative and *fnbB*-negative  
318 LA-MRSA strain Fin47\_H17 exhibits high adhesive capacity to porcine and human cells.  
319 Our results suggest that loss of *agr* leads to increased adhesive capacity and that porcine LA-  
320 MRSA can exhibit pronounced *agr*-dependent cytotoxic effects on porcine cells. Our findings  
321 provide further evidence for the importance of LA-MRSA as an emerging public health  
322 concern.

323

324

## 325 **Acknowledgments**

326           The authors are grateful to Prof. Roger Stephan, Prof. Angelika Lehner, Dr. Taurai  
327 Tasara, Dr. Claudia Guldemann and Dr. Henna-Maria Sihto from the Institute for Food Safety  
328 and Hygiene, University of Zurich and Dr. Cory Ann Leonard, Dr. Hanna Marti, med. vet.  
329 Sabrina Wanninger and med. vet. Jasmin Kuratli from the Institute for Veterinary Pathology,  
330 University of Zurich, Vetsuisse Faculty, for helpful advice, discussion and laboratory support.  
331 We also thank the Functional Genomics Centre Zurich for technical assistance.

332

## 333 **Conflict of interest statement**

334           The authors declare that they have no conflict of interest with respect to the research,  
335 authorship, and/or publication of this article.

336

## 337 **References**

338 Alt K, Fetsch A, Schroeter A, Guerra B, Hammerl JA, Hertwig S, et al. Factors associated  
339 with the occurrence of MRSA CC398 in herds of fattening pigs in Germany. *BMC Vet Res.*  
340 2011;7:69. doi: 10.1186/1746-6148-7-69.

341

342 Smith TC, Male MJ, Harper AL, Kroeger JS, Tinkler GP, Moritz ED, et al. Methicillin-  
343 resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine  
344 and swine workers. *PLoS One.* 2009;4(1):e4258. doi: 10.1371/journal.pone.0004258.

345  
346 de Neeling AJ, van den Broek MJ, Spalburg EC, van Santen-Verheувel MG, Dam-Deisz  
347 WD, Boshuizen HC, et al. High prevalence of methicillin resistant *Staphylococcus aureus* in  
348 pigs. Vet Microbiol. 2007;122(3-4):366-72  
349  
350 Khanna T, Friendship R, Dewey C, Weese JS. Methicillin resistant *Staphylococcus aureus*  
351 colonization in pigs and pig farmers. Vet Microbiol. 2008;128(3-4):298-303.  
352  
353 Price LB1, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. *Staphylococcus*  
354 *aureus* CC398: host adaptation and emergence of methicillin resistance in livestock. MBio.  
355 2012;3(1). pii: e00305-11. doi: 10.1128/mBio.00305-11. Print 2012.  
356  
357 Schijffelen MJ, Boel CH, van Strijp JA, Fluit AC 2010. Whole genome analysis of a  
358 livestock-associated methicillin-resistant *Staphylococcus aureus* ST398 isolate from a case of  
359 human endocarditis. BMC Genomics 2010;11: 376. doi: 10.1186/1471-2164-11-376  
360  
361 Ellis MW, Hospenthal DR, Dooley DP, Gray PJ, Murray CK. Natural history of community-  
362 acquired methicillin-resistant *Staphylococcus aureus* colonization and infection in soldiers.  
363 Clin Infect Dis. 2004;39(7):971-9.  
364 Kuehnert MJ, Kruszon-Moran D, Hill HA, McQuillan G, McAllister SK, Fosheim G, et al.  
365 Prevalence of *Staphylococcus aureus* nasal colonization in the United States, 2001-2002. J  
366 Infect Dis. 2006;193(2):172-9.  
367

368 Wulf M, van Nes A, Eikelenboom-Boskamp A, de Vries J, Melchers W, Klaassen C, et al.  
369 Methicillin-resistant *Staphylococcus aureus* in veterinary doctors and students, the  
370 Netherlands. Emerg Infect Dis. 2006;12(12):1939-41.  
371  
372 Wulf MW, Sørnum M, van Nes A, Skov R, Melchers WJ, Klaassen CH, et al. Prevalence of  
373 methicillin-resistant *Staphylococcus aureus* among veterinarians: an international study. Clin  
374 Microbiol Infect. 2008;14(1):29-34.  
375  
376 van Belkum A, Melles DC, Peeters JK, van Leeuwen WB, van Duijkeren E, Huijsdens XW,  
377 et al. Methicillin-resistant and -susceptible *Staphylococcus aureus* sequence type 398 in pigs  
378 and humans. Emerg Infect Dis. 2008;14(3):479-83. doi: 10.3201/eid1403.070760.  
379  
380 Cuny C, Nathaus R, Layer F, Strommenger B, Altmann D, Witte W. Nasal colonization of  
381 humans with methicillin-resistant *Staphylococcus aureus* (MRSA) CC398 with and without  
382 exposure to pigs. PLoS One. 2009;4(8):e6800. doi: 10.1371  
383  
384 Denis O, Suetens C, Hallin M, Catry B, Ramboer I, Dispas M, et al. Methicillin-resistant  
385 *Staphylococcus aureus* ST398 in swine farm personnel, Belgium. Emerg Infect Dis.  
386 2009;15(7):1098-101. doi: 10.3201/eid1507.080652.  
387  
388 Huber H, Giezendanner N, Stephan R, Zweifel C. Genotypes, Antibiotic Resistance Profiles  
389 and Microarray-Based Characterization of Methicillin-Resistant *Staphylococcus aureus*  
390 Strains Isolated from Livestock and Veterinarians in Switzerland. Zoonoses Public Health.  
391 2011;58(5):343-9. doi: 10.1111/j.1863-2378.2010.01353.x.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

392  
393 Wulf MW, Markestein A, van der Linden FT, Voss A, Klaassen C, Verduin CM. First  
394 outbreak of methicillin-resistant *Staphylococcus aureus* ST398 in a Dutch hospital, June  
395 2007. Euro Surveill. 2008;13(9). pii: 8051.  
396  
397 Valentin-Domelier AS, Girard M, Bertrand X, Violette J, François P, Donnio P-Y, et al.  
398 Methicillin-susceptible ST398 *Staphylococcus aureus* responsible for bloodstream infections:  
399 an emerging human-adapted subclone? PLoS One. 2011;6:e28369.  
400 10.1371/journal.pone.0028369  
401  
402 Uhlemann AC, Porcella SF, Trivedi S, Sullivan SB, Hafer C, Kennedy AD, et al.  
403 Identification of a highly transmissible animal-independent *Staphylococcus aureus* ST398  
404 clone with distinct genomic and cell adhesion properties. MBio. 2012;3(2). pii: e00027-12.  
405 doi: 10.1128/mBio.00027-12.  
406  
407 Verkade E, Bergmans AM, Budding AE, van Belkum A, Savelkoul P, Buiting AG, et al.  
408 Recent emergence of *Staphylococcus aureus* clonal complex 398 in human blood culture.  
409 PLoS ONE 2012;7:e41855.  
410  
411 Heikinheimo A, Johler S, Karvonen L, Julmi J, Fredriksson-Ahomaa M, Stephan R. New  
412 dominant spa type t2741 in livestock-associated MRSA (CC398-MRSA-V) in Finnish  
413 fattening pigs at slaughter. Antimicrob Resist Infect Control. 2016;5:6. doi: 10.1186/s13756-  
414 016-0105-8.  
415

416 Bronner S, Monteil H, Prevost G. Regulation of virulence determinants in *Staphylococcus*  
417 *aureus*: complexity and applications. FEMS Microbiol. Rev. 2004;28:183–200.  
418

419 Tseng CW, Stewart GC. Rot repression of enterotoxin b expression in *Staphylococcus aureus*.  
420 J. bacteriol., 2005 July 21. ed. 187:5301–5309.  
421

422 Fowler VGJr, Sakoulas G, McIntyre LM, Meka VG, Arbeit RD, Cabell CH, et al. Persistent  
423 bacteremia due to methicillin-resistant *Staphylococcus aureus* infection is associated with agr  
424 dysfunction and low-level in vitro resistance to thrombin-induced platelet microbicidal  
425 protein. J Infect Dis. 2004;190:1140–9.  
426

427 Schweizer ML, Furuno JP, Sakoulas G, Johnson JK, Harris AD, Shardell MD, et al.  
428 Increased mortality with accessory gene regulator (agr) dysfunction in *Staphylococcus aureus*  
429 among bacteremic patients. Antimicrob Agents Chemother. 2011;55:1082–7.  
430

431 Foster TJ, Geoghegan JA, Ganesh VK, Höök M. Adhesion, invasion and evasion: the many  
432 functions of the surface proteins of *Staphylococcus aureus*. Nat Rev Microbiol. 2013;12:49–  
433 62.  
434

435 Shinji H, Yosizawa Y, Tajima A, Iwase T, Sugimoto S, Seki K, et al. Role of fibronectin-  
436 binding proteins A and B in in vitro cellular infections and in vivo septic infections by  
437 *Staphylococcus aureus*. Infect Immun. 2011;79:2215–23.  
438

439 Nüesch-Inderbinnen MT, Stalder U, Johler S, Hächler H, Stephan R, Nüesch H-J. Intrafamilial  
440 spread of a Panton-Valentine leukocidin-positive community-acquired methicillin-resistant  
441 *Staphylococcus aureus* belonging to the paediatric clone ST5 SSCmecIV. JMM Case Reports  
442 2014 Sept 12. doi:10.1099/jmmcr.0.001859.

443  
444 Charpentier E, Anton AI, Barry P, Alfonso B, Fang Y, Novick RP. Novel cassette-based  
445 shuttle vector system for gram-positive bacteria. Appl Environ Microb. 2004;70:6076–85.

446  
447 Novick RP. Analysis by transduction of mutations affecting penicillinase formation in  
448 *Staphylococcus aureus*. J Gen Microbiol. 1963;33:121–36.

449  
450 Baranyi J, Roberts TA. A dynamic approach to predicting bacterial growth in food. Int J Food  
451 Microbiol. 1994;23(3-4):277-94.

452  
453 Herbert S, Ziebandt AK, Ohlsen K, Schäfer T, Hecker M, Albrecht D, et al. Repair of global  
454 regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical  
455 isolates. Infect Immun. 2010;78(6):2877-89. doi: 10.1128/IAI.00088-10.

456  
457 Adhikari RP, Arvidson S, Novick RP. A nonsense mutation in agrA accounts for the defect in  
458 agr expression and the avirulence of *Staphylococcus aureus* 8325-4 traP::kan. Infect. Immun.  
459 2007;75:4534–4540.

460  
461 Wiseman, GM. The hemolysins of *Staphylococcus aureus*. Bacteriol. Rev. 1975;39:317–344.

462

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

463 Boukamp P, Petrussevska RT, Breitkreutz D, Hornung J, Markham A, Fusenig NE. Normal  
464 keratinization in a spontaneously immortalized aneuploid human keratinocyte cell line. *J Cell*  
465 *Biol.* 1988;106(3):761-71.  
466

467 Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence  
468 data. *Bioinformatics* 2014;30:2114-2120. doi: 10.1093/bioinformatics/btu170  
469

470 Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler Transform.  
471 *Bioinformatics* 2009;25:1754-1760.  
472

473 Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a  
474 new genome assembly algorithm and its applications to single-cell sequencing. *J Comput*  
475 *Biol* 2012;19: 455-477. doi: 10.1089/cmb.2012.0021.  
476

477 Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. ACT: the  
478 Artemis Comparison Tool. *Bioinformatics* 2005;21:3422-3423. doi:  
479 10.1093/bioinformatics/bti553  
480

481 Uhlemann AC, Porcella SF, Trivedi S, Sullivan SB, Hafer C, Kennedy AD, et al.  
482 Identification of a highly transmissible animal-independent *Staphylococcus aureus* ST398  
483 clone with distinct genomic and cell adhesion properties. *MBio.* 2012 Feb 28;3(2). pii:  
484 e00027-12. doi: 10.1128/mBio.00027-12. Print 2012.  
485

486 van der Mee-Marquet N, Hernandez D, Bertrand X, Quentin R, Corvaglia AR, François P.  
487 Whole-Genome Sequence of the Ancestral Animal-Borne ST398 *Staphylococcus aureus*  
488 Strain S123. *Genome Announc.* 2013 Aug 29;1(5). pii: e00692-13. doi:  
489 10.1128/genomeA.00692-13.  
490  
491 Hernandez D, van der Mee-Marquet N, Kluytmans J, Donnio PY, Quentin R, Corvaglia AR,  
492 et al. Whole-Genome Sequences of *Staphylococcus aureus* ST398 Strains of Animal Origin.  
493 *Genome Announc.* 2013 Sep 26;1(5). pii: e00689-13. doi: 10.1128/genomeA.00689-13.  
494  
495 Corvaglia AR1, François P, Bertrand X, Quentin R, Hernandez D, van der Mee-Marquet N.  
496 Whole-Genome Sequences of Two *Staphylococcus aureus* ST398 Strains of Human Origin,  
497 S94 and S100. *Genome Announc.* 2013 Aug 29;1(5). pii: e00691-13. doi:  
498 10.1128/genomeA.00691-13.  
499  
500 Golding GR, Bryden L, Levett PN, McDonald RR, Wong A, Graham MR, et al. Whole-  
501 genome sequence of livestock-associated st398 methicillin-resistant *staphylococcus aureus*  
502 Isolated from Humans in Canada. *J Bacteriol.* 2012 Dec;194(23):6627-8. doi:  
503 10.1128/JB.01680-12.  
504  
505 Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.  
506 *Bioinformatics.* 2011 Apr 1;27(7):1009-10. doi: 10.1093/bioinformatics/btr039.  
507

508 Schwan WR, Langhorne MH, Ritchie HD, Stover CK. Loss of hemolysin expression in  
509 *Staphylococcus aureus* agr mutants correlates with selective survival during mixed infections  
510 in murina abscesses and wounds. FEMS Immunol Med Microbiol. 2003;38(1):23-8.  
511

512 Patel AH, Kornblum J, Kreiswirth B, Novick R, Foster TJ. Regulation of the protein A-  
513 encoding gene in *Staphylococcus aureus*. Gene. 1992;114:25–34  
514

515 Saravia-Otten P, Müller HP, Arvidson S. Transcription of *Staphylococcus aureus* fibronectin  
516 binding protein genes is negatively regulated by agr and an agr-independent mechanism. J.  
517 Bacteriol. 1997;179:5259–5263.  
518

519 Projan SJ, Novick RP. The molecular basis of pathogenicity. In: Crossley KB, Archer GL,  
520 editors. The staphylococci in human disease. Churchill Livingstone, New York, N.Y; 1997.  
521 pp 55–82.  
522

523 Wolz C, Pöhlmann-Dietze P, Steinhuber A, Chien YT, Manna A, van Wamel W, et al. Agr-  
524 independent regulation of fibronectin-binding protein(s) by the regulatory locus sar in  
525 *Staphylococcus aureus*. Mol Microbiol. 2000;36(1): 230-243.  
526

527 Pöhlmann-Dietze P, Ulrich M, Kiser KB, Döring G, Lee JC, Fournier JM, et al. Adherence of  
528 *Staphylococcus aureus* to endothelial cells: influence of capsular polysaccharide, global  
529 regulator agr, and bacterial growth phase. Infect Immun. 2000;68(9):4865-71.  
530

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

531 Poston SM, Glancey GR, Wyatt JE, Hogan T, Foster TJ. Co-elimination of mec and spa  
532 genes in *Staphylococcus aureus* and the effect of agr and protein A production on bacterial  
533 adherence to cell monolayers. J Med Microbiol. 1993;39(6):422-8.

Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert

534  
535 Wesson CA, Liou LE, Todd KM, Bohach GA, Trumble WR, Bayles KW. *Staphylococcus*  
536 *aureus* Agr and Sar global regulators influence internalization and induction of apoptosis.  
537 Infect Immun. 1998;66(11):5238-43.

538  
539 Ballhausen B, Jung P, Kriegeskorte A, Makgotlho PE, Ruffing U, von Müller L, et al. LA-  
540 MRSA CC398 differ from classical community acquired-MRSA and hospital acquired-  
541 MRSA lineages: functional analysis of infection and colonization processes. Int J Med  
542 Microbiol. 2014;304(7):777-86. doi: 10.1016/j.ijmm.2014.06.006.

Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert

543  
544 Ezechuk YV, Leung DY, Middleton MH, Bina P, Reiser R, Norris DA. Staphylococcal  
545 toxins and protein A differentially induce cytotoxicity and release of tumor necrosis factor-  
546 alpha from human keratinocytes. J Invest Dermatol. 1996;107(4):603-9.

547  
548 Popov LM, Marceau CD, Starkl PM, Lumb JH, Shah J, Guerrero D, et al. The adherens  
549 junctions control susceptibility to *Staphylococcus aureus*  $\alpha$ -toxin. Proc Natl Acad Sci U S A.  
550 2015;112(46):14337-42. doi: 10.1073/pnas.1510265112.

Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert  
Feldfunktion geändert

551  
552 Grundmeier M, Hussain M, Becker P, Heilmann C, Peters G, Sinha B. Truncation of  
553 fibronectin-binding proteins in *Staphylococcus aureus* strain Newman leads to deficient

554 adherence and host cell invasion due to loss of the cell wall anchor function. Infect Immun.  
555 2004;72(12):7155-63.

556

557 Bur S, Preissner KT, Herrmann M, Bischoff M. The *Staphylococcus aureus* extracellular  
558 adherence protein promotes bacterial internalization by keratinocytes independent of  
559 fibronectin-binding proteins. J Invest Dermatol. 2013;133(8):2004-12. doi:  
560 0.1038/jid.2013.87.

561

562

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

## 563 **Supporting information**

564 **S1 Fig. Phylogenetic tree.** Phylogenetic tree of whole genome sequenced isolates in the  
565 context of other sequenced CC398 strains. Scale bar shows the number of substitutions within  
566 the expected core genome (1,799,838 bp).

567

568 **S2 Table. Variable elements.** Table providing an overview of variable elements within the  
569 genomes of porcine LA-MRSA strains Fin47\_H17, Fin46\_17, and Fin48\_H18 compared to  
570 reference strain S0385.

571

572

573

574

575

576

577

578

579

580

581

582 **Supplementary Table 1: Genomic analysis of adhesion associated genes.** S0385 data from Uhlemann 2012, with additions. Abbreviations:  
583 V/S = variable/same (compared to strain indicated in superscript), A = absent at this location, I = insertion, X = aa substitution,  $\psi$  = pseudogenes,  
584 wt = wild type, aa =amino acid,  $\Delta$  = deletion, SNP = single nucleotide polymorphism.

Gene	S0385 locus tag <sup>1</sup> SAPIG	S085	NM01	Fin47_H17	Fin46_H17	Fin48_H18	Comment
<i>spa</i>	0122	8 aa $\Delta$ at aa390	2 X at aa390	wt	wt	I (8 aa at aa390)	Variable repeat length
<i>coa</i>	0240	81 bp $\Delta$	wt	wt	wt	wt	Variable repeat length
<i>vwb</i>	0483	wt	A	V <sup>S0385</sup>	V <sup>S0385</sup>	A (at this location)	Variable sequence/presence
<i>sdrC</i>	0636	174 bp $\Delta$	wt	1 SNP	S <sup>Fin47_H17</sup>	I (24aa at aa737)	Variable repeat length
<i>sdrD</i>	0637	42 bp $\Delta$ and 4 X	$\Delta$ 54 bp	wt	wt	wt	Variable repeat length

							and SNPs
<i>eno</i>	0855			S <sup>S0385</sup>	S <sup>S0385</sup>	1 SNP, non synonymous	
<i>clfA</i>	0866-7	ψ truncated, 2 frameshifts	wt	V <sup>S0385</sup>	S <sup>Fin47_H17</sup>	V <sup>S0385</sup>	Variable repeat length
<i>isdB</i>	1125	SNP, 9 bp Δ	wt	S <sup>S0385</sup>	S <sup>S0385</sup>	I <sup>S0385</sup>	
<i>isdA</i>	1126			S <sup>S0385</sup>	S <sup>S0385</sup>	1 SNP, synonymous	
<i>ecb</i>	1150	wt	wt	S <sup>S0385</sup>	S <sup>S0385</sup>	S <sup>S0385</sup>	
<i>fib</i>	1154	wt	wt	S <sup>S0385</sup>	S <sup>S0385</sup>	S <sup>S0385</sup>	
<i>ebhA</i>	1434	53 aa Δ at aa9851 and 1 X	wt	1 aa X	1 X	2 X	Variable repeat length
<i>ebpS</i>	1480	wt	wt	S <sup>S0385</sup>	S <sup>S0385</sup>	S <sup>S0385</sup>	
hypothetical	1791	174 bp Δ	wt	1 aa change	S <sup>Fin47_H17</sup>	wt	Variable repeat length

				(aa39)			
<i>map</i>	1981			S <sup>S0385</sup>	S <sup>S0385</sup>	S <sup>S0385</sup>	
<i>sdrH</i>	2069	wt	Δ 39 bp	S <sup>S0385</sup>	S <sup>S0385</sup>	S <sup>S0385</sup>	
<i>hld</i>	2072	wt	wt	ψ A	wt	wt	Variably present
<i>agrB</i>	2073	wt	wt	A	wt	wt	Variably present
<i>agrA</i>	2076			S <sup>S0385</sup>	S <sup>S0385</sup>	S <sup>S0385</sup>	
<i>fnbB</i>	2550-1	ψ truncated, stop codon	wt	Deleted	Deleted	ψ truncated, stop codon	Often defunct
<i>fnbA</i>	2553	wt	wt	Δ9 bp in repeat region, 12 aa changes	S <sup>Fin47_H17</sup>	wt	Variable
<i>clfB</i>	2679	ψ truncated	Δ42aa at aa819	wt	wt	wt	Variable repeats / truncated
<i>isaB</i>	2688			S <sup>S0385</sup>	S <sup>S0385</sup>	1 SNP,	

						synonymous	
<i>cna</i>	2740	wt	$\Delta$ B domain	$\Delta$ 187aas, 4 X	As Fin47	$\Delta$ 187aas, 4 X	Variable in this clade

585