



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2011

More on transfusion and adverse outcome: it's time to change

Spahn, D R ; Shander, A ; Hofmann, A ; Berman, M F

DOI: <https://doi.org/10.1097/ALN.0b013e3182054d3e>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-53769>

Journal Article

Published Version

Originally published at:

Spahn, D R; Shander, A; Hofmann, A; Berman, M F (2011). More on transfusion and adverse outcome: it's time to change. *Anesthesiology*, 114(2):234-236.

DOI: <https://doi.org/10.1097/ALN.0b013e3182054d3e>

More on Transfusion and Adverse Outcome

It's Time to Change

PREOPERATIVE anemia and perioperative allogeneic erythrocyte (RBC) transfusion have both been shown to be independently associated with adverse outcome, such as increased mortality, postoperative infections, lung and kidney injury, myocardial infarction, stroke, and increased length of hospital stay.^{1–6} The relative contribution of each of these conditions, and their interactions, is difficult to specify. The study by Glance *et al.* in this issue of ANESTHESIOLOGY is important, because the authors studied more than 10,000 patients with preoperative anemia who underwent elective noncardiac surgery and received up to 2 units of RBCs intraoperatively. The study used records from the American College of Surgeons National Surgical Quality Improvement Program (ACS NSQIP) database.⁷

In an attempt to reduce confounders, the investigators included only patients within a preoperative hemoglobin range of 6–10 g/dl. In this hemoglobin range, according to American Society of Anesthesiologists guidelines, RBC transfusions may be administered on a clinical basis.⁸ Thus, these RBC transfusions had likely occurred after limited blood loss. Observed outcomes thus may be attributed more clearly to RBC transfusion than if patients had started with normal preoperative hemoglobin and, as a consequence, had to have lost large amounts of blood during ill-controlled surgery before being transfused. In the latter case, it would be difficult to say whether the adverse outcomes were due to hemodynamic instability and other effects of acute blood loss or due to the RBC transfusion itself. Since in the study by Glance *et al.*⁷ all patients were anemic preoperatively and may not have experienced massive blood loss before the index transfusion, the observed outcomes may be attributed quite clearly to the RBC transfusion administered. Also, the negative impact on outcome was substantial: Transfusion of only 1 or 2 units of RBCs resulted in more septic (odds ratio [OR]: 1.43), pulmonary (OR: 1.79), thromboembolic (OR: 1.77), and wound (OR: 1.87) complications and increased mortality (OR: 1.29).

The key strengths of this study lie in the number of the patients and the detailed data collection; the shortcomings are shared with many other transfusion cohort studies. Despite various efforts, it is always difficult to call transfusion as the cause of an adverse outcome, as opposed to some other associated patient comorbidity. A brief look at table 1 of the

Glance *et al.* article indicates that the transfused cohort was generally in worse shape at baseline, compared with the not-transfused group (lower hematocrit, older age, more dependent, more comorbidities, etc.). Statistical methods are used to remedy this commonly seen disparity. Multivariate analysis can be used to adjust for confounders (the method used in the current study), or transfused patients can be matched against nontransfused patients (with varying degrees of complexity). Although these approaches often do marvels, they are not “cures,” because they are ultimately limited by the available variables and their data. Do we have all the potential confounders captured in our dataset? How certain are we of the primary reasons behind giving transfusion to every single case? What was the tipping point that made the physician decide to give blood to this patient, but not the other patient? These questions are often impossible to answer accurately in cohort studies, and they undermine the validity of the results.

Although the multivariate adjustment used in this study is a valid method, more data on the models would have been helpful (*e.g.*, strength of association as indicated by pseudo-R-squared or the reasoning behind creating the specific hematocrit categories). In addition, use of a matching approach may have allowed for more transparent interpretation of the results (at expense of reduced sample size). Given the large sample size, it would have been interesting to see whether the transfused and not-transfused patients could have been effectively matched to each other based on the risk factors for receiving transfusions (and, possibly, known risk factors of each outcome), thus allowing a more direct comparison.

In addition, large database analyses do have limitations of their own. Problems with miscoding of procedures and diagnoses are well recognized.⁹ This study by Glance *et al.*, however, relies on the NSQIP database, which is more accurate than other state or federal administrative datasets. Data are collected by trained surgical chart reviewers, and participating hospitals undergo interrater reliability audits.¹⁰ Still, for any particular study, a generalized database such as NSQIP will not provide all the information that one might wish for.

In the case of the present study, we do not know the extent of pre- or postoperative transfusion in the study patients; because of NSQIP data definitions, patients could have received up to 4 units preoperatively and 4 units postoperatively. Transfusion before or after surgery, but not during,

Accepted for publication October 25, 2010. The authors are not supported by, nor maintain any financial interest in, any commercial activity that may be associated with the topic of this article.

|| American College of Surgeons. Participation requirements. In: National Surgical Quality Improvement Program. Available at: https://acsnsqip.org/main/get_started_requirements.asp. Accessed July 20, 2010.

Copyright © 2011, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins. Anesthesiology 2011; 114: 234–6

◆ This Editorial View accompanies the following article: Glance LG, Dick AW, Mukamel DB, Fleming FJ, Zollo RA, Wissler R, Salloum R, Meredith W, Osler TM: Association between intraoperative blood transfusion and mortality and morbidity in patients undergoing noncardiac surgery. ANESTHESIOLOGY 2011; 114:283–92.

might decrease the detected effect of intraoperative transfusion; alternatively, those patients requiring pre- or postoperative transfusion might be more likely to receive blood intraoperatively. NSQIP does not record intraoperative nadir hematocrit or immediate postoperative hematocrit. Thus, we do not know if lower intraoperative hematocrit was associated with worse outcome. Because NSQIP data do not include a hospital identifier, the final results may have been confounded by hospital-based differences in practice, including differences in local transfusion triggers.¹⁰

Results of the present study are in line with previous studies in general,³ cardiac,¹ orthopedic,⁴ and trauma surgery.^{5,6} Also, in a recent cohort data analysis in 239,286 elderly Veterans Administration patients with and without preoperative anemia RBC transfusions resulted in increased 30-day mortality (adjusted OR: 1.37 [1.27–1.48]).¹¹ Only in a 1-min subgroup of patients with a preoperative hematocrit of less than or equal to 24% (1.9% of the study population) did RBC transfusion result in a reduced mortality, whereas in patients with a preoperative hematocrit of greater than 30% (more than 80% of the study population), RBC transfusions actually increased mortality significantly (ORs: 1.37–1.59). Interestingly, in patients who eventually died, late postoperative complications, such as failure to wean from the ventilator and unplanned intubation indicative of lung injury, renal failure, and infectious complications, were very similar to the ones found in the article by Glance *et al.*,⁷ and they were more frequent than in nontransfused patients.¹¹

Another aspect of the study by Glance *et al.*⁷ deserves mentioning, namely the fact that nonelective cases were excluded. Therefore, it should have been possible to treat pre-existing anemia before surgery. This is important, because preoperative anemia is frequent,^{2,11–13} easy to detect, and—in many situations—treatable at relatively low costs.¹⁴ At least, in orthopedic surgery, there is strong evidence that preoperative treatment of anemia results not only in reduced RBC transfusion, but also in improved outcome.¹⁵

Where do we go from here? The call for prospective, randomized studies on the benefit or detriment of allogeneic RBC transfusions is the usual conclusion after studies of the likes of Glance *et al.* are published. This conclusion is certainly politically correct; however, if Glance *et al.* and the slew of others are correct—which is a valid possibility—can we take the responsibility to continue today's widespread transfusion practice, an approach that is putting patients at risk?

Contrarians to the above still overly rely on banked blood products, despite mounting evidence that patient blood management is achievable, avoids exposure to allogeneic blood products, improves patient outcome, and saves resources.^{15,16} The ease of ignoring preoperative anemia as well as the simplicity of ordering and transfusing RBCs and thereby completely ignoring the burgeoning evidence of adverse transfusion outcome are not in the best interest of patients. Maintaining the clinical status quo under such circumstances would

not be accepted in any other field of medicine in the context of current safety and quality standards.

Working on studies regarding benefit or detriment of allogeneic RBC transfusions is important, but more so is acting already today according to patient blood management principles; it is time for a change toward better patient care.

Donat R. Spahn, M.D., F.R.C.A.,* Aryeh Shander, M.D., F.C.C.P., F.C.C.M.,† Axel Hofmann, M.E.,‡ Mitchell F. Berman, M.D., M.P.H.§ *Institute of Anesthesiology and Head Medical, Section Anesthesiology, Intensive Care Medicine and OR-Management, University Hospital Zurich, Zurich, Switzerland. donat.spahn@usz.ch. †Department of Anesthesiology and Critical Care and Hyperbaric Medicine and Institute for Patient Blood Management and Bloodless Medicine and Surgery, Englewood Hospital and Medical Center, Englewood, New Jersey. ‡Medical Society for Blood Management, Laxenburg, Austria, and Center for Population Health Research, Curtin University, Western Australia. §Department of Anesthesiology, Columbia University College of Physicians and Surgeons, New York, New York.

References

1. Karkouti K, Wijeyesundera DN, Beattie WS: Risk associated with preoperative anemia in cardiac surgery: A multicenter cohort study. *Circulation* 2008; 117:478–84
2. Beattie WS, Karkouti K, Wijeyesundera DN, Tait G: Risk associated with preoperative anemia in noncardiac surgery: A single-center cohort study. *ANESTHESIOLOGY* 2009; 110:574–81
3. Bernard AC, Davenport DL, Chang PK, Vaughan TB, Zwischenberger JB: Intraoperative transfusion of 1 U to 2 U packed red blood cells is associated with increased 30-day mortality, surgical-site infection, pneumonia, and sepsis in general surgery patients. *J Am Coll Surg* 2009; 208:931–7
4. Pedersen AB, Mehnert F, Overgaard S, Johnsen SP: Allogeneic blood transfusion and prognosis following total hip replacement: A population-based follow up study. *BMC Musculoskelet Disord* 2009; 10:167
5. Charles A, Shaikh AA, Walters M, Huehl S, Pomerantz R: Blood transfusion is an independent predictor of mortality after blunt trauma. *Am Surg* 2007; 73:1–5
6. Chaiwat O, Lang JD, Vavilala MS, Wang J, MacKenzie EJ, Jurkovich GJ, Rivara FP: Early packed red blood cell transfusion and acute respiratory distress syndrome after trauma. *ANESTHESIOLOGY* 2009; 110:351–60
7. Glance LG, Dick AW, Mukamel DB, Fleming FJ, Zollo RA, Wissler R, Salloum R, Meredith W, Osler TM: Association between intraoperative blood transfusion and mortality and morbidity in patients undergoing noncardiac surgery. *ANESTHESIOLOGY* 2011; 114:283–92.
8. Practice guidelines for perioperative blood transfusion and adjuvant therapies: An updated report by the American Society of Anesthesiologists Task Force on Perioperative Blood Transfusion and Adjuvant Therapies. *ANESTHESIOLOGY* 2006; 105:198–208
9. Woodworth GF, Baird CJ, Garces-Ambrossi G, Tonascia J, Tamargo RJ: Inaccuracy of the administrative database: Comparative analysis of two databases for the diagnosis and treatment of intracranial aneurysms. *Neurosurgery* 2009; 65:251–6
10. Gombotz H, Rehak PH, Shander A, Hofmann A: Blood use in elective surgery: The Austrian benchmark study. *Transfusion* 2007; 47:1468–80
11. Wu WC, Smith TS, Henderson WG, Eaton CB, Poses RM, Uttley G, Mor V, Sharma SC, Vezeridis M, Khuri SF, Friedmann PD: Opera-

- tive blood loss, blood transfusion, and 30-day mortality in older patients after major noncardiac surgery. *Ann Surg* 2010; 252:11-7
12. Shander A, Knight K, Thurer R, Adamson J, Spence R: Prevalence and outcomes of anemia in surgery: A systematic review of the literature. *Am J Med* 2004; 116(Suppl 7A):58-69S
 13. Goodnough LT, Vizmeg K, Sobecks R, Schwarz A, Soegiarso W: Prevalence and classification of anemia in elective orthopedic surgery patients: Implications for blood conservation programs. *Vox Sang* 1992; 63:90-5
 14. Goodnough LT, Shander A, Spivak JL, Waters JH, Friedman AJ, Carson JL, Keating EM, Maddox T, Spence R: Detection, evaluation, and management of anemia in the elective surgical patient. *Anesth Analg* 2005; 101:1858-61
 15. Spahn DR: Anemia and patient blood management in hip and knee surgery: A systematic review of the literature. *ANESTHESIOLOGY* 2010; 113:482-95
 16. Moskowitz DM, Klein JJ, Shander A, Cousineau KM, Goldweit RS, Bodian C, Perelman SI, Kang H, Fink DA, Rothman HC, Ergin MA: Predictors of transfusion requirements for cardiac surgical procedures at a blood conservation center. *Ann Thorac Surg* 2004; 77:626-34