



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2013

---

**Environmental determinants of *Ixodes ricinus* ticks and the incidence of *Borrelia burgdorferi sensu lato*, the agent of Lyme borreliosis, in Scotland**

James, M C ; Bowman, A S ; Forbes, K J ; Lewis, F ; McLeod, J E ; Gilbert, L

DOI: <https://doi.org/10.1017/S003118201200145X>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-71415>

Journal Article

Accepted Version

Originally published at:

James, M C; Bowman, A S; Forbes, K J; Lewis, F; McLeod, J E; Gilbert, L (2013). Environmental determinants of *Ixodes ricinus* ticks and the incidence of *Borrelia burgdorferi sensu lato*, the agent of Lyme borreliosis, in Scotland. *Parasitology*, 140(2):237-246.

DOI: <https://doi.org/10.1017/S003118201200145X>

# Environmental determinants of *Ixodes ricinus* ticks and the incidence of *Borrelia burgdorferi* sensu lato, the agent of Lyme borreliosis, in Scotland

M. C. JAMES<sup>1,2,4</sup>, A. S. BOWMAN<sup>1</sup>, K. J. FORBES<sup>2</sup>, F. LEWIS<sup>3</sup>, J. E. MCLEOD<sup>4</sup>  
and L. GILBERT<sup>4\*</sup>

<sup>1</sup>*Institute of Biological and Environmental Sciences, Zoology Building, University of Aberdeen, Tillydrone Avenue, Aberdeen AB24 2TZ, UK*

<sup>2</sup>*Section of Immunology and Infection, Medical School, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK*

<sup>3</sup>*Section of Epidemiology, Vetsuisse, University of Zürich, 8057 Zürich, Switzerland*

<sup>4</sup>*James Hutton Institute, Macaulay Drive, Craigiebuckler, Aberdeen, AB15 8QH, UK*

(Received 6 March 2012; revised 14 June and 6 August 2012; accepted 6 August 2012)

## SUMMARY

Lyme borreliosis (LB) is the most common arthropod-borne disease of humans in the Northern hemisphere. In Europe, the causative agent, *Borrelia burgdorferi* sensu lato complex, is principally vectored by *Ixodes ricinus* ticks. The aim of this study was to identify environmental factors influencing questing *I. ricinus* nymph abundance and *B. burgdorferi* s.l. infection in questing nymphs using a large-scale survey across Scotland. Ticks, host dung and vegetation were surveyed at 25 woodland sites, and climatic variables from a Geographical Information System (GIS) were extracted for each site. A total of 2397 10 m<sup>2</sup> transect surveys were conducted and 13 250 *I. ricinus* nymphs counted. Questing nymphs were assayed for *B. burgdorferi* s.l. and the average infection prevalence was 5.6% (range 0.8–13.9%). More questing nymphs and higher incidence of *B. burgdorferi* s.l. infection were found in areas with higher deer abundance and in mixed/deciduous compared to coniferous forests, as well as weaker correlations with season, altitude, rainfall and ground vegetation. No correlation was found between nymph abundance and infection prevalence within the ranges encountered. An understanding of the environmental conditions associated with tick abundance and pathogen prevalence may be used to reduce risk of exposure and to predict future pathogen prevalence and distributions under environmental changes.

Key words: woodland, nymphs, hosts, PCR, GIS, negative binomial hurdle model, disease.

## INTRODUCTION

Lyme Borreliosis (LB) is the most common arthropod-borne medical infection in the Northern Hemisphere (Kurtenbach *et al.* 2006) and reported cases are increasing rapidly in some areas. For example, there has been an approximately 11-fold increase in reported cases in the past decade in Scotland (Health Protection Scotland, 2011). The causative agent of LB is the spirochaete bacterium *Borrelia burgdorferi* sensu lato complex that is vectored by *Ixodes* ticks. Currently no vaccine is available against this pathogen, so reducing disease risk is best achieved by minimizing the risk of exposure to infected tick bites, combined with the swift removal of attached ticks and early recognition and treatment of symptoms. One means of reducing the risk of exposure to infected tick bites is to have an understanding of the environmental conditions associated with high infection risk. Knowledge of the environmental determinants of *Ixodes* tick and *B. burgdorferi*

s.l. abundance is also useful to estimate future pathogen distribution and prevalence under scenarios of environmental changes such as changes in climate, land use, habitat and hosts.

The transmission cycle of *B. burgdorferi* s.l. is complex and is dependent on the tick vector and the *B. burgdorferi* s.l. transmission competence of the reservoir hosts. In northern and western Europe, the primary vector of *B. burgdorferi* s.l. is *I. ricinus*, the sheep tick. *Ixodes ricinus* ticks spend the vast majority of their lives in the undergrowth (either questing for a passing host, in development or in diapause). However, the completion of their 3-stage life cycle (larva, nymph and adult) depends on obtaining a bloodmeal from a host for each of the life stages. The most important hosts for feeding larval and nymphal stages are birds and small mammals, while large ungulates such as sheep and deer are more important for feeding adults and are therefore considered 'tick reproduction hosts' (Gray, 1998; Humair *et al.* 1999). *Ixodes ricinus* abundance may also be influenced by climatic variables (including micro-climate induced by local vegetation characteristics). For example, adult and nymph *I. ricinus* ticks require temperatures above 7 °C to

\* Corresponding author: James Hutton Institute, Macaulay Drive, Craigiebuckler, Aberdeen AB15 8QH, UK. Tel: +44 (0)1224 395187. Fax: +44 (0) 844 928 5429. E-mail: Lucy.Gilbert@hutton.ac.uk

start questing (e.g. Randolph, 2004 and references therein) and can quest for longer without desiccation at higher humidities (Randolph *et al.* 2000; Eisen *et al.* 2002; Piesman and Gern, 2004).

*Borrelia burgdorferi* s.l. is a complex of *Borrelia* species, each varying in their main transmission hosts, LB symptoms and spatial distributions. In Europe, small mammals transmit *B. afzelii* while many bird species can competently transmit *B. garinii* or *B. valaisiana*. *B. burgdorferi* sensu stricto can be transmitted by mammalian, avian and reptilian hosts (e.g. Hanincová *et al.* 2003a,b). While deer are extremely important in maintaining tick populations, they are incompetent reservoirs for *B. burgdorferi* s.l. (Telford *et al.* 1988; Jaenson and Tälleklint, 1992). In contrast to studies on *I. ricinus*, fewer studies have tested for associations between environmental observations and *B. burgdorferi* s.l. prevalence in ticks. In theory, we would predict *B. burgdorferi* s.l. prevalence to be influenced by the density of competent transmission hosts (Ostfeld and Keesing, 2000). However, in practice, abundance estimates for small mammals and birds (the main competent transmission hosts in Europe) are much more difficult to acquire on a landscape scale than are other environmental variables such as climate, habitat and large mammals (the main *I. ricinus* reproduction hosts). Therefore, proxies for small mammals and birds are often used, so instead, for example, it could be predicted, that *B. burgdorferi* s.l. prevalence correlates with the amount of habitat preferred by transmission hosts (see Prusinski *et al.* 2006). This approach is, of course, not perfect, since the main small mammal species (wood mice *Apodemus sylvaticus*, bank voles *Myodes glareolus* and common shrews *Sorex araneus*) may differ in their transmission efficiencies of *B. burgdorferi* s.l. (Kurtenbach *et al.* 1994). However, they are all considered to be competent transmission hosts and have broadly similar habitat preferences in that semi-natural mixed or deciduous woodlands that have understory vegetation are likely to harbour much larger densities of these hosts than are, say, coniferous plantations (e.g. Flowerdew, 1993).

#### Objectives and predictions

The maintenance of *B. burgdorferi* s.l. requires *B. burgdorferi* s.l. transmission hosts, tick reproduction hosts and ticks to occur in close spatial proximity in environmental conditions suitable for all three (as per Killilea *et al.* 2008 for *B. burgdorferi* s.s.). The primary aim of this study was to identify the environmental factors influencing both *I. ricinus* abundance and the presence of *B. burgdorferi* s.l. in ticks. Such information is important for informing disease mitigation strategies and predictions of future risk under scenarios of environmental change. Our approach combines data from field surveys of ticks,

vegetation and large mammal dung counts with climate variables from a GIS. We conducted the study in Scotland, where *I. ricinus* are increasing in abundance (Kirby *et al.* 2004), Lyme borreliosis cases have increased dramatically (11-fold) over the last decade (Health Protection Scotland, 2011), and no large-scale surveys of multiple sites have yet been conducted. We predicted that *I. ricinus* nymph abundance would be greater in areas that have higher large mammal abundance (tick reproduction hosts), in areas with a warmer and wetter climate (conducive to tick survival) and in habitats that both produce a micro-climate favourable to tick survival and favourable to hosts of immature ticks such as small mammals and birds (i.e. semi-natural mixed or deciduous woodland rather than commercial conifer forest). We predicted that questing *I. ricinus* nymphs would be more likely to be infected with *B. burgdorferi* s.l. in conditions favourable for competent transmission hosts (small mammals and birds), such as in mixed or deciduous (rather than coniferous) woods, or depending on ground vegetation type (see Prusinski *et al.* 2006).

#### MATERIALS AND METHODS

##### Field surveys

Studies were conducted in woodlands because this was the habitat type associated with the most Lyme borreliosis cases in Scotland (James, 2010). Surveys were carried out in 2007 and 2008 at 25 woodland sites known to be endemic for LB across Scotland (56° 02' N–57° 51' N and 2° 30' W–6° 12' W), chosen for geographical spread and to provide a variety of environmental characteristics. Woodlands were categorized broadly as either semi-natural mixed/deciduous ( $n=13$ ), typically *Quercus* and *Betula* species, or coniferous ( $n=12$ ), typically *Picea* and *Pinus* species, and they also differed in altitude (1–503 metres above sea level), and deer communities (Table 1).

Surveys were conducted from April to October to cover the main questing season of *I. ricinus* in Scotland. A blanket drag method was used in which questing ticks were collected by dragging a 1 m<sup>2</sup> white blanket across the ground vegetation for 10 m before inspection, counting and collection (Gray and Lohan, 1982; Gilbert, 2010). This technique provides an estimated 'index of relative abundance' of ticks that are questing at the time of sampling between different areas, rather than an absolute density or abundance of ticks in an area.

The location of each blanket drag was recorded on a global positioning system (GPS) device (Garmin eTrex H, Southampton, UK), with latitude and longitude recorded for use in statistical analysis, along with the time and date (noted as Julian day for statistical analysis). Relative humidity and

Table 1. Woodland type, altitude, deer species and years surveyed of the 25 sites across Scotland

Site	Woodland type	Mean altitude (masl)	Deer species present	Years surveyed
AN	Coniferous	280	Roe	2007, 2008
AP	Coniferous	50	0	2008
BM	Coniferous	360	Red + Roe	2008
CB	Coniferous	25	Roe	2007, 2008
WB	Coniferous	212	Roe	2007, 2008
GC	Coniferous	50	Red	2007
CM	Coniferous	220	0	2007
LV	Coniferous	188	Roe	2007
QC	Coniferous	423	Red	2007, 2008
GM	Coniferous	83	0	2008
IN	Coniferous	297	Red + Roe	2007
LA	Coniferous	50	0	2008
LR	Deciduous	230	0	2008
MA	Deciduous	93	Roe	2007, 2008
MW	Deciduous	27	Roe	2007
QD	Deciduous	421	Red	2007
QW	Deciduous	62	Roe	2007
SH	Deciduous	82	0	2008
TB	Deciduous	70	Red + Roe	2007, 2008
CR	Deciduous	234	Red	2007, 2008
AB	Deciduous	86	Roe	2007
DR	Deciduous	85	Roe	2007, 2008
DV	Deciduous	40	Red	2008
FZ	Deciduous	170	0	2008
GD	Deciduous	22	0	2007, 2008

temperature were recorded at ground vegetation height using a temperature-humidity probe (RS212-540, RS Components, Northants, UK). For statistical analysis, values of relative humidity were averaged (per site and per am/pm). The ground vegetation height was measured at 3 points along the 10 m drag using a sward stick and the mean value per blanket drag was used for analysis (Gilbert, 2010). The dominant ground vegetation was categorized into 4 classes: (1) grasses and herbaceous species, (2) ericaceous and *Vaccinium* species, (3) moss species and (4) bracken and ferns.

Dung of red deer *Cervus elephus* and roe deer *Capreolus capreolus* were identified (Sergeant and Morris, 2003) and counts were conducted over each 1 m × 10 m blanket drag area. For statistical analysis we calculated an ‘index of relative abundance’ of roe and red deer by averaging each blanket drag’s dung counts for each site (see Gilbert, 2010).

Between 50 and 200 drags were performed at each site in total, each at least 50 m apart, in a semi-random way so as to sample each site in a representative manner. Sites were visited between 1 and 3 times per year over 1 or 2 years. Dragging was not performed during or just after rain when the ground vegetation was wet enough to soak the blanket, nor when the temperature was below 7 °C as ticks may not begin questing until the mean weekly max temp is above 7 °C (Randolph, 2004 and references therein).

Questing nymphs are the most important stage for transmitting *B. burgdorferi* s.l. to humans, as questing

larvae do not transmit the pathogen (they are unlikely to be infected as they have not yet taken a bloodmeal and transovarial transmission is thought to be absent or rare (Hubálek and Halouzka, 1998) and adults are far less numerous than nymphs. In addition, due to their smaller size, nymphs are less likely than adult ticks to be noticed and removed before *B. burgdorferi* s.l. is transmitted from the biting tick (Falco *et al.* 1999). Robertson *et al.* (2000) estimated that 82% of human tick bites from a forested area in England were from nymphs. This study therefore focuses on the abundance of, and the *B. burgdorferi* s.l. infection in, questing *I. ricinus* nymphs. Nymphs were placed in plastic vials containing 70% ethanol for later analysis of *B. burgdorferi* s.l. To confirm that the ticks collected were *I. ricinus*, 2000 collected nymphs were randomly selected and identified to species level using a light dissecting microscope and species key (Hillyard, 1996).

#### *Borrelia burgdorferi* s.l. detection

*Borrelia burgdorferi* s.l. detection assays were conducted on individual nymphs for those collected in 2007. Using knowledge gained from 2007 and to save time and resources, nymphs were pooled in groups of 5 for those collected in 2008 (see Supplementary Online material for further explanation and details). For analysis, 1 nymph (if collected in 2007) or a pool of 5 nymphs (if collected in 2008) were randomly selected for the *B. burgdorferi* s.l. assay from each of 1218 blanket drags.

DNA extraction was performed by mechanical destruction of the nymphs and ammonia extraction (Guy and Stanek, 1991; James *et al.* 2011).

To identify which nymphs were infected with *B. burgdorferi* s.l. we used a nested polymerase chain reaction (PCR) to detect the intergenic spacer region between the 5S and 23S rRNA genes (5S-23S rRNA IGS), as described by Rijpkema *et al.* (1995). For every 20 tick samples, 1 negative control (dH<sub>2</sub>O), 1 negative homogenate and 1 positive control (*Borrelia lusitaniae* DNA; this genospecies is currently not known in northern Europe) were processed. Later, the genospecies determination of positive samples showed that none were *B. lusitaniae* so any contamination/false positive results were unlikely.

#### Identification of *B. burgdorferi* s.l. genospecies

The genospecies of *Borrelia* from 82 (all those that successfully amplified) positive nymphs was determined by multi-loci sequence typing of up to 8 genes which are diagnostic for genospecies (Margos *et al.* 2008). Eight housekeeping loci were analysed; *clpA*, *clpX*, *pepX*, *pyrG*, *recG*, *rplB* and *uvrA*. PCRs were performed under the same conditions (cycling conditions, primer concentrations and reagents) as the *B. burgdorferi* s.l. PCR (i.e. 5S-23S IGS). MLST products were cleaned and sequenced on an ABI automated DNA sequencer. Forward and reverse sequences were compared, aligned and trimmed using *Sequence Editor* (version 1.0.3, Macintosh computers). Consensus sequences were then searched on the MLST website ([www.mlst.net](http://www.mlst.net)) to determine the genospecies.

#### Environmental variables from GIS

Environmental variables including growing degree days (day-by-day sum of the mean number of degrees by which the air temperature is more than 5.5 °C), average annual precipitation, average mean temperature, snow-lying days, ground-frost days and growing season length were obtained for each of the 25 woodland sites from a set of 1 km and 5 km gridded climate data rasters derived from the UK Met Office 1971–2000 Long Term Average climate data (Met Office, 2009; Perry and Hollis, 2005). The values were extracted using the Intersect Point tool from Hawth's Analysis Tools for ArcGIS v3.27 (Beyer, 2004) in ArcMap v9.3 (ESRI, 2008).

#### Statistical analysis

Because ecological and tick count data are complex, have non-normal data distributions and many of the factors are inter-related, state-of-the-art statistical techniques were necessary. All statistical analyses were performed in R version 2.11.1 (R Development

Core Team, 2010). We first explored the data for the response variable (nymph count or *B. burgdorferi* s.l. presence in nymphs) and explanatory variables (environmental factors) in terms of data distribution and selection of explanatory variables for initial inclusion in the models. It is essential to use a model that fits the data as well as possible, so we then constructed several types of models and selected the model family that best fit the data. Please see Online Supplementary Material for details of the variable and model selection procedures.

Both relative abundance index of questing nymphs and infection of *B. burgdorferi* s.l. in nymphs were analysed at the level of the individual blanket drag. For the relative abundance index of questing nymphs, the model family that we selected for use was a negative binomial hurdle model, since it best fitted the data. For infection of *B. burgdorferi* s.l. in nymphs, (which could either be positive or negative to *B. burgdorferi* s.l.), the model selected for use was binomial.

Selected explanatory variables initially entered as fixed effects into the full models for both questing nymph abundance index and *B. burgdorferi* s.l. presence/absence were as follows: at the spatial scale of individual blanket drags we entered time, altitude, temperature, ground vegetation height and dominant ground vegetation category. At the scale of each site we entered the following variables which were measured at the time of sampling: Julian day, forest type (coniferous or deciduous/mixed), red deer abundance index, roe deer abundance index, mean relative humidity. At the scale of each site we also entered the following climatic variables (10-year averages) from a GIS database: annual mean precipitation, annual mean temperature, growing degree days, snow-lying days, and ground-frost days. Using a backwards stepwise procedure we eliminated non-significant ( $P > 0.05$ ) explanatory variables such that the final models included only significant variables.

Finally, we tested whether questing *I. ricinus* nymph abundance correlated with *B. burgdorferi* s.l. prevalence in questing nymphs in order to test whether questing nymph abundance can be used as a proxy for prevalence. At the site level, we regressed *B. burgdorferi* s.l. prevalence (number of nymphs testing positive/total number of nymphs tested) at each site against the mean number of nymphs counted per blanket drag at each site, using a Spearman Rank Correlation.

#### RESULTS

During the spring and summers of 2007 and 2008, 2397 blanket drags were conducted and 13 250 nymphs were counted and collected. A subsample of 2000 nymphs was identified to species level and all were found to be *I. ricinus*.



Fig. 1. Prevalence of *Borrelia burgdorferi* s.l. in questing nymph *Ixodes ricinus* ticks in woodlands in Scotland. Gray scale denotes prevalence, ranging from the lowest (white) at 0.8% to the highest (black) at 13.9%.

Nymph *I. ricinus* ticks were assayed from 1218 individual blanket drags for *B. burgdorferi* s.l. (we did not have the time and resources to assay nymphs from all drags, but we ensured that we assayed a minimum of 50 nymphs from each site). The prevalence of *B. burgdorferi* s.l. at each site was calculated as the number of positive nymphs/number of nymphs assayed per site. The overall prevalence in questing nymphs from all 25 sites was 5.6% ( $\pm 1.0\%$ , range 0.8–13.9%, Fig. 1). Of those positive samples that we successfully identified to genospecies level, 48% were *B. afzelii*, 36% *B. garinii*, 7% *B. burgdorferi* s.s., 8% *B. valaisiana* and 1% carried mixed infections of 2 or more genospecies. The number of samples successfully identified to each genospecies was not large enough for conducting meaningful statistical analyses to test for predictive environmental variables, so this was done only for *B. burgdorferi* s.l. prevalence as a whole, at the site level.

#### *Environmental factors influencing questing nymph abundance*

Questing nymph abundance was analysed using a negative binomial hurdle model that gives a dual output, i.e. 2 patterns are fitted to the data. In this case, the hurdle model fitted relationships for (i) nymph presence/absence (i.e. zero counts versus non-zero counts) and (ii) nymph abundance of at least 1 nymph per blanket drag (i.e. non-zero counts); see Table 2. Nymphs were more likely to be present on blanket drags conducted in areas with higher

abundance indices of roe and red deer, where the ground vegetation comprised mainly ericaceous species/*Vaccinium* or mosses or grasses/herbaceous species (compared to bracken/ferns as the baseline), in deciduous/mixed woods (compared to coniferous as the baseline), lower relative humidity and at lower altitudes. For those blanket drags where nymphs were present, there was a very strong positive relationship between nymph abundance and red deer abundance index and deciduous/mixed woods (compared to coniferous). In addition, questing nymph abundance increased with increasing ground vegetation height, decreasing Julian day (i.e. more questing nymphs earlier in the questing season) and decreasing annual precipitation. The following variables were eliminated from the model due to non-significance during the backwards stepwise procedure: time of day, temperature at the time of sampling, growing degree days, mean annual temperature, snow-lying days, ground-frost days and growing season length and there were no significant interaction terms.

#### *Environmental factors influencing B. burgdorferi s.l. infection*

The environmental variables that were statistically associated with *B. burgdorferi* s.l. infection in questing nymphs (Table 3) had general similarities with those associated with questing nymph abundance. There was a strong positive correlation between *B. burgdorferi* s.l. infection and the index of relative

Table 2. Output from the final negative binomial hurdle model describing the association between questing nymph presence/absence and abundance (counts of more than zero on blankets) and environmental variables (Non-significant variables are not shown as they were eliminated during the backwards stepwise procedure. The estimate for deciduous wood is in comparison to the coniferous wood baseline, and the ground vegetation categories are compared with ferns/bracken baseline. CI denotes the 95% confidence interval.)

	Estimate	Upper CI	Lower CI	z-value	P-value
Questing nymph presence/absence					
Deciduous wood	0.0708	0.0474	0.0901	1.23	<0.001
Altitude	-0.0004	-0.0006	-0.0002	1	<0.001
Ground veg: ericaceous	0.1231	0.0983	0.1394	1.45	<0.001
Ground veg: grass/herbs	0.1154	0.0911	0.1324	1.42	<0.001
Ground veg: moss	0.1215	0.0927	0.1397	1.43	<0.001
Roe deer index	0.1505	0.1179	0.1623	1.46	<0.001
Red deer index	0.1637	0.1531	0.1671	1.52	<0.001
Relative humidity	-0.0034	-0.0046	-0.0022	0.99	<0.001
Questing nymph abundance					
Deciduous wood	13.0250	9.7872	16.7925	2.13	<0.001
Julian day	-0.0678	-0.0835	-0.0521	0.99	<0.001
Ground vegetation height	0.0716	0.0264	0.1169	1	0.001
Red deer index	31.0184	19.3605	47.7668	3.47	<0.001
Annual precipitation	-0.0033	-0.0049	-0.0017	1	<0.001

Table 3. Output from the final binomial model describing the associations between the presence/absence of *Borrelia burgdorferi* s.l. infection in questing nymphs and environmental parameters

(Non-significant variables are not shown as they were eliminated during the backwards stepwise procedure. The estimate for deciduous wood is in comparison to the coniferous wood baseline. CI denotes the 95% confidence interval.)

<i>B. burgdorferi</i> s.l.	Estimate	Upper CI	Lower CI	z-value	P-value
Deciduous wood	0.0341	-0.0007	0.0873	1.952	0.051
Altitude	-0.0002	-0.0003	-0.0001	-2.954	0.003
Julian day	0.0007	0.0003	0.0012	3.112	0.002
Relative humidity	-0.0012	-0.0021	-0.0002	-2.393	0.017
Ground vegetation height	-0.0010	-0.0019	-0.0001	-2.173	0.03
Red deer index	0.0861	0.0054	0.2385	2.211	0.027

abundance of red (but not roe) deer. Nymphs were also more likely to be infected in deciduous/mixed (compared to coniferous) woods. The likelihood of questing nymphs being infected with *B. burgdorferi* s.l. decreased with Julian day (i.e. nymphs were more likely to be infected later in the season), decreasing altitude, increasing ground vegetation height and increasing relative humidity (Table 3).

#### Does nymph abundance predict *B. burgdorferi* s.l. prevalence?

There was not a significant relationship between nymph abundance and *B. burgdorferi* s.l. prevalence (Fig. 2), Spearman Rank correlation  $\rho_s = -0.040$ ,  $n = 25$  sites within the ranges of abundances encountered.

#### DISCUSSION

Our main aim was to identify the environmental factors influencing both questing *I. ricinus*

abundance and the presence of *B. burgdorferi* s.l. in those ticks. We also used the data to test for an association between questing nymph abundance and *B. burgdorferi* s.l. prevalence.

#### Questing nymph abundance

We found a strong positive association between questing nymph abundance index (and nymph presence/absence) and the abundance indices of both red and roe deer, as we predicted. This is unsurprising, since deer (roe and red) are the primary 'reproduction' hosts for *I. ricinus* in Scotland, and many previous studies have found a correlation between tick and deer abundance (e.g. Jensen *et al.* 2000; Ruiz-Fons and Gilbert, 2010).

As predicted, we found a strong effect of woodland type, with semi-natural mixed/deciduous woodlands having more nymphs than coniferous woodland. We also found further effects of ground vegetation type (fewer nymphs in bracken/ferns) and height




Fig. 2. The relationship between questing nymph abundance and *Borrelia burgdorferi* s.l. prevalence at the site level.

(more nymphs in tall ground vegetation). Several previous studies have found more *I. ricinus* in deciduous woodlands (e.g. Estrada-Peña, 2001; Guerra *et al.* 2002; Lindström and Jaenson, 2003; Brown *et al.* 2006), although Walker *et al.* (2001) found more ticks in coniferous than in deciduous woods in southern Scotland. It is likely that more ticks are found in habitats with more hosts and that, for our particular study sites, tick abundance reflected areas with an understory vegetation of preference to a range of vertebrates that host immature ticks. Habitats with dense understory may also be associated with higher tick numbers because of the mild and humid micro-climate produced by the thick ground vegetation and the deep leaf litter in deciduous/mixed woodland habitats (e.g. Lindström and Jaenson, 2003).

More questing ticks were predicted in areas with a warmer climate and higher rainfall (or higher humidity) because of the temperature and humidity thresholds on tick questing behaviour and survival. We did find more questing ticks in woodlands at lower altitudes which, given that the model allowed for variance due to hosts and vegetation, is likely to be partly a climatic effect. A strong effect of climate and altitude on questing *I. ricinus* nymphs was recently found for open moorland areas in Scotland (Gilbert, 2010). However, we found the opposite effect to that predicted for rainfall or humidity: questing nymphs were more likely to be present at times of low relative humidity and questing nymphs were more abundant in areas with lower annual rainfall. In general, previous studies have found that tick questing activity increases with increasing humidity (Randolph *et al.* 2000; Piesman and Gern, 2004), but it might be that conditions can be too wet for ticks (e.g. they tend to quest less in the rain). Perhaps in regions with a cool, wet climate, such as Scotland and western Scandinavia, the drier times and drier areas may still be well within the humid conditions needed

for *I. ricinus* nymphs to quest, while the wetter times and wetter areas may be too wet.

#### *Borrelia burgdorferi* s.l. infection in questing nymphs

We predicted that questing *I. ricinus* nymphs would be more likely to be infected with *B. burgdorferi* s.l. in conditions compatible for the presence of competent transmission hosts (small mammals and birds), such as in semi-natural mixed/deciduous (rather than coniferous plantation) woods, or depending on ground vegetation type. This prediction was upheld: nymphs were more likely to be infected with *B. burgdorferi* s.l. in semi-natural mixed/deciduous (compared to coniferous) woodlands and also in woodlands with higher ground vegetation (which may relate to small mammal and bird abundance, although further surveys would be needed to test this for our particular sites). While bank voles, wood mice and common shrews share broadly similar habitat preferences, such as preferring semi-natural woodland to dense coniferous plantations, they differ in their efficiency of transmitting *B. burgdorferi* s.l. (Kurtenbach *et al.* 1994). Therefore, the proportion of, say, wood mice to bank voles in one year might be expected to influence the infection prevalence of questing *I. ricinus* nymphs with *B. burgdorferi* s.l. in the following year. Our study merely used broad habitat type as a proxy for the likely relative abundances of transmission hosts in general, and detailed further studies would be needed to test the effect of actual host community composition on *B. burgdorferi* s.l. prevalence.

Interestingly, it was found that nymphs were more likely to be infected with *B. burgdorferi* s.l. in woodlands with higher red deer abundance indices. Deer cannot transmit *B. burgdorferi* s.l. and could be considered dilution hosts if more deer result in more immature ticks feeding on deer instead of on competent transmission hosts. However, what may possibly be happening at our sites could be that deer may act primarily as tick reproduction hosts (see Gray, 1998) by feeding mainly adult ticks. Therefore, more deer would result in more immature ticks that feed primarily on the small transmission hosts, thereby increasing the likelihood of transmission occurring between infected nymphs and uninfected larvae feeding simultaneously on an individual transmission host. Indeed, a recent study in Italy found that the number of ticks feeding on rodents initially increased with deer density, reaching a peak at intermediate deer densities and then decreased as deer densities got higher (Cagnacci *et al.* 2012). In accordance, models of tick-borne encephalitis virus and Louping ill virus predict an increase in tick-borne pathogen prevalence with increasing deer up to a point, and then at very high deer densities the dilution effect is predicted to occur (Bolzoni *et al.* 2012; Gilbert *et al.* 2001). It is intriguing that, unlike

red deer abundance indices, we found no such positive effect of roe deer on *B. burgdorferi* s.l. This could be a statistical artefact, e.g. if there was less statistical power due to roe deer being less abundant than red deer. Alternatively, it could be a real biological effect due to unmeasured factors, e.g. differences between roe and red deer in spatial distribution with respect to transmission host distribution within sites; or differences between red and roe deer in their burdens of different life stages of *I. ricinus*. Roe deer can carry large numbers of all stages of *I. ricinus*, particularly nymphs (Kiffner *et al.* 2010; Vor *et al.* 2010) while red deer in Scotland carry mainly adults (L. Gilbert, unpublished data available at [http://www.macaulay.ac.uk/deerlarder/a\\_deer\\_ticks.php](http://www.macaulay.ac.uk/deerlarder/a_deer_ticks.php) last accessed 13th June 2012).

It is interesting that the relationship of *B. burgdorferi* s.l. with Julian day was the opposite to that for questing nymph abundance, i.e. questing nymph abundance decreased over the season whereas the incidence of *B. burgdorferi* s.l. infection in nymphs increased over the season. One potential explanation could be that *B. burgdorferi* s.l. transmission hosts increase in abundance over the season, as small mammals and birds produce young over the summer. In addition, it is likely that the infection prevalence of small mammals increases over the season as they are bitten by ticks throughout the season. There is currently no available information on the timings of diapause and moulting of Scottish *I. ricinus* stages, but if there are earlier and later cohorts of each stage of ticks, e.g. larvae that feed later in the season become nymphs that quest later the following season, then this could be a potential mechanism explaining the above contradictory effects of questing nymphs and *B. burgdorferi* s.l. infection over the season. However, this is pure conjecture and further research is required on the seasonality of cohorts and the timing of diapause and moulting of larvae and nymphs in Scottish woodlands to test this idea. Previous studies have found seasonal patterns of *B. burgdorferi* s.l. infection in *I. ricinus* nymphs that differ from our finding, e.g. peaks of infection in late spring in Sweden (Tälleklint and Jaenson, 1996), further indicating that the biological mechanisms behind seasonal patterns in *B. burgdorferi* s.l. prevalence remain obscure.

Nymphs were also more likely to be infected with *B. burgdorferi* s.l. in surveys conducted at lower altitudes (although the effect was not strong), and this could occur if *B. burgdorferi* s.l. transmission hosts are more likely to be found at lower level sites than at higher, more exposed, sites. Several other studies have found this negative association between *B. burgdorferi* s.l. prevalence and altitude in forests, as in Switzerland (Burri *et al.* 2007; Jouda *et al.* 2004; Cadenas *et al.* 2007).

Previous studies have also found associations between *B. burgdorferi* s.l. infection and hosts,

habitat and climate. Altobelli *et al.* (2008), using data from a GIS, found correlations between *B. burgdorferi* s.l. infected *I. ricinus* and roe deer abundance and mean annual temperature in north-eastern Italy. Eisen *et al.* (2010), also using GIS, found correlations between *B. burgdorferi* s.l. infected *I. scapularis* nymphs and spring/summer temperature, variability in water vapour and forest type in the western USA. Jensen *et al.* (2000) found positive associations between *I. ricinus* density and roe deer density and soil water capacity in woodland habitats, yet these factors were not associated with *B. burgdorferi* s.l. prevalence in those ticks. Maetzel *et al.* (2005) did not find a difference in the *B. burgdorferi* s.l. infection prevalence between wooded and non-woodland sites.

While we did not have sufficient statistical power to analyse environmental variables with respect to each different genospecies of *B. burgdorferi* s.l., the fact that almost half the *I. ricinus* we assayed tested positive for *B. afzelii* is noteworthy. Previous studies of *I. ricinus* in other parts of the British Isles have found *B. garinii* and *B. valaisiana* to be predominant (e.g. Kurtenbach *et al.* 1998), although Vollmer *et al.* (2011) found *B. afzelii* at just over half their English sites. Ling *et al.* (2000) analysed 12 positive *I. ricinus* from the Scottish Highlands and, like our study, identified that almost half of these were *B. afzelii*. The reasons for any difference in genospecies predominance between Scotland and the rest of the British Isles warrants further research.

#### *Relationship between questing nymph abundance and B. burgdorferi* s.l. prevalence

We found no evidence for a positive association between questing nymph abundance and *B. burgdorferi* s.l. prevalence at the site level within the ranges encountered. For a given density of transmission hosts, theoretically, higher tick abundance may mean more ticks feeding on the same individual transmission host while it is infectious, thereby leading to higher pathogen prevalence. However, as densities of transmission hosts vary, and the transmission efficiency of different hosts also varies, this theoretical relationship will not necessarily be realised empirically. Some authors suggest using nymph abundance as an indicator of *B. burgdorferi* s.l. prevalence (Jaenson *et al.* 2009) but this lack of an association at our sites suggests that nymph abundance cannot necessarily be used as a proxy for *B. burgdorferi* s.l. prevalence. However, for a given *B. burgdorferi* s.l. prevalence, if there are more nymphs questing, the risk of human infection will be higher. Therefore, all things being equal, more ticks may sometimes mean higher infection risk to humans. However, in reality, as the infection prevalence varies so widely (0.8–13.9% in Scottish woodlands) we caution against

using questing nymph abundance as a proxy for human infection risk.

### Conclusion

The mechanistic causes of *B. burgdorferi* s.l. infection in ticks relate to the abundance of transmission hosts and the burden of larvae, nymphs and adult ticks on transmission hosts. However, collecting such information, while possible for small numbers of sites and small areas, is very labour intensive and is rarely feasible over multiple sites over large areas. Therefore, studies such as ours and the previous studies cited are useful in predicting potential disease risk. Our study and the previous studies share some consistent findings, such as the positive association of deer with questing tick abundance and *B. burgdorferi* s.l. infection, and general associations with habitat (such as woodland type) that may reflect transmission host abundance, and with climate variables (that may reflect hosts and tick abundance).

In summary, we have carried out an extensive study with large sample sizes of sites, blanket drags and ticks assayed, and have identified which environmental variables are associated with questing *I. ricinus* nymphs and the incidence of *B. burgdorferi* s.l. infection in questing nymphs in Scottish woodlands. Woodland type and deer abundance were the primary determinants of both nymph abundance and *B. burgdorferi* s.l. infection. Ground vegetation and weather/climate variables also played a role. Most of our results supported our initial predictions which were made in accordance with tick reproduction hosts and which habitats are generally preferred by transmission hosts (small mammals and birds). These results could be used in future studies to predict which areas might have high tick abundances and *B. burgdorferi* s.l. prevalence based on environmental factors on a broader scale (e.g. using a country-wide GIS). In addition, it may be possible to predict what may happen to pathogen distributions under future environmental changes such as woodland expansion, deer management and climate change.

### ACKNOWLEDGEMENTS

The authors are grateful for field assistance from E. Packer, A. Wiebe, J. Low, E. Stephen and J. Arthur, for laboratory assistance from K. Reay, and three anonymous reviewers for useful comments on the manuscript.

### FINANCIAL SUPPORT

M.C.J. was funded by a Biotechnology and Biological Sciences Research Council (BBSRC) Doctoral Training Grant with CASE support from the Macaulay Development Trust awarded to A.S.B. and L.G., and L.G. and J.McL. were supported by the Scottish Government's Rural and Environment Science and Analytical Services Division (RESAS).

### REFERENCES

- Altobelli, A., Boemo, B., Mignozzi, K., Bandi, M., Floris, R., Menardi, G. and Cinco, M. (2008). Spatial Lyme borreliosis risk assessment in north-eastern Italy. *International Journal of Medical Microbiology* **298**, (Suppl. 1), 125–128.
- Beyer, H. L. (2004). *Hawth's Analysis Tools for ArcGIS*. Available at [www.spatial ecology.com/htools](http://www.spatial ecology.com/htools). Last accessed November 2009.
- Bolzoni, L., Rosà, R., Cagnacci, F. and Rizzoli, A. (2012). Effect of deer density on tick infestation of rodents and the hazard of tick-borne encephalitis. Part II: population and infection models. *International Journal for Parasitology* **42**, 373–381.
- Brown, R. N., Peot, M. A. and Lane, R. S. (2006). Sylvatic maintenance of *Borrelia burgdorferi* (Spirochaetales) in northern California: Untangling the web of transmission. *Journal of Medical Entomology* **43**, 743–751.
- Burri, C., Cadenas, F. M., Douet, V., Moret, J. and Gern, L. (2007). *Ixodes ricinus* density and infection prevalence of *Borrelia burgdorferi* sensu lato along a north-facing altitudinal gradient in the Rhone valley (Switzerland). *Vector-Borne and Zoonotic Diseases* **7**, 50–58.
- Cadenas, F. M., Rais, O., Jouda, F., Douet, V., Humair, P.-F., Moret, J. and Gern, L. (2007). Phenology of *Ixodes ricinus* and infection with *Borrelia burgdorferi* sensu lato along north- and south-facing altitudinal gradient on Chaumont mountain, Switzerland. *Journal of Medical Entomology* **44**, 683–693.
- Cagnacci, F., Bolzoni, L., Rosà, R., Carpi, G., Haufler, H. C., Valent, M., Tagliapietra, V., Kazimirova, M., Koci, J., Stanko, M., Lukan, M., Henttonen, H. and Rizzoli, A. (2012). Effect of deer density on tick infestation of rodents and the hazard of tick-borne encephalitis. Part I: Empirical assessment. *International Journal for Parasitology* **42**, 365–372.
- Eisen, L., Eisen, R. J. and Lane, R. S. (2002). Seasonal activity patterns of *Ixodes pacificus* nymphs in relation to climatic conditions. *Medical and Veterinary Entomology* **16**, 235–244.
- Eisen, R. J., Eisen, L., Girard, Y. A., Fedorova, N., Mun, J., Slikas, B., Leonhard, S., Kitron, U. and Lane, R. S. (2010). A spatially-explicit model of acarological risk of exposure to *Borrelia burgdorferi*-infected *Ixodes pacificus* nymphs in northwestern California based on woodland type, temperature, and water vapor. *Ticks and Tick-Borne Diseases* **1**, 35–43.
- ESRI (2008). *ArcMap v9.3*. Environmental Systems Research Institute, Redlands, CA, USA.
- Estrada-Peña, A. (2001). Distribution, abundance, and habitat preferences of *Ixodes ricinus* (Acari: Ixodidae) in northern Spain. *Journal of Medical Entomology* **38**, 361–370.
- Falco, R. C., McKenna, D. F., Daniels, T. J., Nadelman, R. B., Nowakowski, J., Fish, D. and Wormser, G. P. (1999). Temporal relation between *Ixodes scapularis* abundance and risk for Lyme disease associated with erythema migrans. *American Journal of Epidemiology* **149**, 771–776.
- Flowerdew, J. R. (1993). *Mice and Voles*. British Natural History Series. Whittet Books Ltd, London, UK.
- Gilbert, L. (2010). Altitudinal patterns of tick and host abundance: a potential role for climate change in regulating tick-borne diseases? *Oecologia* **162**, 217–225.
- Gilbert, L., Norman, R., Laurenson, K. M., Reid, H. W. and Hudson, P. J. (2001). Disease persistence and apparent competition in a three-host community: an empirical and analytical study of large-scale wild populations. *Journal of Animal Ecology* **70**, 1053–1061.
- Gray, J. S. (1998). The ecology of ticks transmitting Lyme borreliosis. *Experimental and Applied Acarology* **22**, 249–258.
- Gray, J. S. and Lohan, G. (1982). The development of a sampling method for the tick *Ixodes ricinus* and its use in a redwater fever area. *Annals of Applied Biology* **101**, 421–427.
- Guerra, M., Walker, E., Jones, C., Paskewitz, S., Roberto Cortinas, M., Ashley Stancil, L. B., Bobo, M. and Kitron, U. (2002). Predicting the risk of Lyme disease: Habitat suitability for *Ixodes scapularis* in the north central United States. *Emerging Infectious Diseases* **8**, 289–297.
- Guy, E. C. and Stanek, G. (1991). Detection of *Borrelia burgdorferi* in patients with Lyme disease by the polymerase chain reaction. *Journal of Clinical Pathology* **44**, 610–611.
- Hanincová, K., Schäfer, S. M., Etti, S., Sewell, H., Taragelová, V., Ziak, D., Labuda, M. and Kurtenbach, K. (2003b). Association of *Borrelia afzelii* with rodents in Europe. *Parasitology* **126**, 11–20.
- Hanincová, K., Taragelová, V., Koci, J., Schäfer, S. M., Hails, R., Ullmann, A. J., Piesman, J., Labuda, M. and Kurtenbach, K. (2003a). Association of *Borrelia garinii* and *B. valaisiana* with songbirds in Slovakia. *Applied Environmental Microbiology* **69**, 2825–2830.
- Health Protection Scotland (2011). [www.documents.hps.scot.nhs.uk/giz/10-year-tables/lyme.pdf](http://www.documents.hps.scot.nhs.uk/giz/10-year-tables/lyme.pdf) (last accessed October 2011).
- Hillyard, P. (1996). *Ticks of North West Europe*. Field Studies Council, Shrewsbury, UK.

- Hubálek, Z. and Halouzka, J. (1998). Prevalence rates of *Borrelia burgdorferi* sensu in host-seeking *Ixodes ricinus* ticks in Europe. *Parasitology Research* **84**, 167–172.
- Humair, P. F., Rais, O. and Gern, L. (1999). Transmission of *Borrelia afzelii* from *Apodemus* mice and *Clethrionomys* voles to *Ixodes ricinus* ticks: Differential transmission pattern and overwintering maintenance. *Parasitology* **118**, 33–42.
- Jaenson, T. G. T., Eisen, L., Comstedt, P., Mejlom, H. A., Lindgren, E., Bergström, S. and Olsen, B. (2009). Risk indicators for the tick *Ixodes ricinus* and *Borrelia burgdorferi* sensu lato in Sweden. *Medical and Veterinary Entomology* **23**, 226–237.
- Jaenson, T. G. and Tälleklint, L. (1992). Incompetence of roe deer as reservoirs of the Lyme borreliosis spirochete. *Journal of Medical Entomology* **29**, 813–817.
- James, M. C. (2010). The ecology, genetic diversity and epidemiology of Lyme Borreliosis in Scotland. Ph.D. thesis. University of Aberdeen, Aberdeen, Scotland.
- James, M. C., Furness, R. W., Bowman, A. S., Forbes, K. J. and Gilbert, L. (2011). The importance of passerine birds as tick hosts and in the transmission of *Borrelia burgdorferi*, the agent of Lyme disease. A case study from Scotland. *Ibis* **153**, 293–302.
- Jensen, P. M., Hansen, H. and Frandsen, F. (2000). Spatial risk assessment for Lyme borreliosis in Denmark. *Scandinavian Journal of Infectious Diseases* **32**, 545–550.
- Jouda, F., Perret, J. and Gern, L. (2004). Density of questing *Ixodes ricinus* nymphs and adults infected by *Borrelia burgdorferi* sensu lato in Switzerland: Spatio-temporal pattern at a regional scale. *Vector-Borne Zoonotic Diseases* **4**, 23–32.
- Kalsbeek, V. and Frandsen, F. (1996). The seasonal activity of *Ixodes ricinus* ticks in Denmark. *Anzeiger für Schädlingskunde* **69**, 160–161.
- Kiffner, C., Lödige, C., Alings, Vor, T. and Rühle, F. (2010). Abundance estimate of *Ixodes* ticks (Acari: Ixodidae) on roe deer (*Capreolus capreolus*). *Experimental and Applied Acarology* **52**, 73–84.
- Killilea, M. E., Swei, A., Lane, R. S., Briggs, C. J. and Ostfeld, R. S. (2008). Spatial dynamics of Lyme disease: A review. *EcoHealth* **5**, 167–195.
- Kirby, A. D., Smith, A. A., Benton, T. G. and Hudson, P. J. (2004). Rising burden of immature sheep ticks (*Ixodes ricinus*) on red grouse (*Lagopus lagopus scoticus*) chicks in the Scottish uplands. *Medical and Veterinary Entomology* **18**, 67–70.
- Kurtenbach, K., Dizij, A., Seitz, H. M., Margos, G., Moter, S. E., Kramers, M. D., Wallich, R., Schaible, U. E. and Simon, M. M. (1994). Differential immune responses to *Borrelia burgdorferi* in European wild rodent species influence spirochete transmission to *Ixodes ricinus* L. (Acari: Ixodidae). *Infection and Immunity* **62**, 5344–5352.
- Kurtenbach, K., Hanincová, K., Tsao, J. I., Margos, G., Fish, D. and Ogden, N. H. (2006). Fundamental processes in the evolutionary ecology of Lyme borreliosis. *Nature Reviews Microbiology* **4**, 660–669.
- Kurtenbach, K., Peacey, M., Rijpkema, S. G., Hoodless, A. N., Nuttall, P. A. and Randolph, S. E. (1998). Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England. *Applied Environmental Microbiology* **64**, 1169–1174.
- Lindström, A. and Jaenson, T. G. T. (2003). Distribution of the common tick, *Ixodes ricinus* (Acari: Ixodidae), in different vegetation types in southern Sweden. *Journal of Medical Entomology* **40**, 375–378.
- Ling, C. L., Joss, A. W. L., Davidson, M. M. and Ho-Yen, D. O. (2000). Identification of different *Borrelia burgdorferi* genomic groups from Scottish ticks. *Molecular Pathology* **53**, 94–98.
- Maetzel, D., Maier, W. A. and Kampen, H. (2005). *Borrelia burgdorferi* infection prevalences in questing *Ixodes ricinus* ticks (Acari: Ixodidae) in urban and suburban Bonn, western Germany. *Parasitology Research* **95**, 5–12.
- Margos, G., Gatewood, A. G., Aanensen, D. M., Hanincová, K., Terekhova, D., Vollmer, S. A., Cornet, M., Piesman, J., Donaghy, M., Bormane, A., Hurn, M. A., Feil, E. J., Fish, D., Casjens, S., Wormser, G. P., Schwartz, I. and Kurtenbach, K. (2008). MLST of housekeeping genes captures geographic population structure and suggests a European origin of *Borrelia burgdorferi*. *Proceedings of the National Academy of Sciences, USA* **105**, 8730–8735.
- Met Office (2009). [www.metoffice.gov.uk/climate/uk/2007/summer/averages.html](http://www.metoffice.gov.uk/climate/uk/2007/summer/averages.html). Last accessed March 2010.
- Ostfeld, R. S. and Keesing, F. (2000). Biodiversity and disease risk: the case of Lyme disease. *Conservation Biology* **14**, 1–7.
- Perry, M. and Hollis, D. (2005). The development of a new set of long-term climate averages for the UK. *International Journal of Climatology* **25**, 1023–1039.
- Piesman, J. and Gern, L. (2004). Lyme borreliosis in Europe and North America. *Parasitology* **129**, S191–S220.
- Prusinski, M. A., Chen, H., Drobnack, J. M., Kogut, S. J., Means, R. G., Howard, J. J., Oliver, J., Lukacik, G., Backenson, P. B. and White, D. J. (2006). Habitat structure associated with *Borrelia burgdorferi* prevalence in small mammals in New York State. *Environmental Entomology* **35**, 308–319.
- R Development Core Team (2010). *R: A Language and Environment for Statistical Computing*. [www.R-project.org](http://www.R-project.org). Vienna, Austria.
- Randolph, S. E. (2004). Tick ecology: processes and patterns behind the epidemiological risk posed by ixodid ticks as vectors. *Parasitology* **129**, S1–S29.
- Randolph, S. E., Green, R. M., Peacey, M. F. and Rogers, D. J. (2000). Seasonal synchrony: The key to tick-borne encephalitis foci identified by satellite data. *Parasitology* **121**, 15–23.
- Rijpkema, S. G. T., Molkenboer, M. J. C. H., Schouls, L. M., Jongejan, F. and Schellekens, J. F. P. (1995). Simultaneous detection and genotyping of three genomic groups of *Borrelia burgdorferi* sensu lato in Dutch *Ixodes ricinus* ticks by characterization of the amplified intergenic spacer region between 5S and 23S rRNA genes. *Journal of Clinical Microbiology* **33**, 3091–3095.
- Robertson, J. N., Gray, J. S. and Stewart, P. (2000). Tick bite and Lyme borreliosis risk at a recreational site in England. *European Journal of Epidemiology* **16**, 647–652.
- Ruiz-Fons, F. and Gilbert, L. (2010). The role of deer as vehicles to move ticks, *Ixodes ricinus*, between contrasting habitats. *International Journal for Parasitology* **40**, 1013–1020.
- Sergeant, G. and Morris, P. (2003). *How to Find and Identify Mammals*. 2nd Edn. The Mammal Society, London, UK.
- Tälleklint, L. and Jaenson, T. G. (1996). Seasonal variations in density of questing *Ixodes ricinus* (Acari: Ixodidae) nymphs and prevalence of infection of *B. burgdorferi* s.l. in south central Sweden. *Journal of Medical Entomology* **33**, 592–597.
- Telford, S. R., III, Mather, T. N., Moore, S. I., Wilson, M. L. and Spielman, A. (1988). Incompetence of deer as reservoirs of the Lyme disease spirochete. *American Journal of Tropical Medicine and Hygiene* **39**, 105–109.
- Vollmer, S. A., Bormane, A., Dinnis, R. E., Seelig, F., Dobson, A. D. M., Aanensen, D. M., James, M. C., Donaghy, M., Randolph, S. E., Feil, E. J., Kurtenbach, K. and Margos, G. (2011). Host migration impacts on the phylogeography of Lyme borreliosis spirochaete species in Europe. *Environmental Microbiology* **13**, 184–192.
- Vor, T., Kiffner, C., Hagedorn, P., Niedrig, M. and Rühle, F. (2010). Tick burden on European roe deer (*Capreolus capreolus*). *Experimental and Applied Acarology* **51**, 405–417.
- Walker, A. R., Alberdi, M. P., Urquhart, K. A. and Rose, H. (2001). Risk factors in habitats of the tick *Ixodes ricinus* influencing human exposure to *Ehrlichia phagocytophila* bacteria. *Medical and Veterinary Entomology* **15**, 40–49.