



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 1999

**Electron microscopic analysis reveals that replication factor C is sequestered by
single-stranded DNA.**

Keller, R C ; Mossi, R ; Maga, G ; Wellinger, R E ; Hübscher, U ; Sogo, J M

DOI: <https://doi.org/10.1093/nar/27.17.3433>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-785>

Journal Article

Published Version

Originally published at:

Keller, R C; Mossi, R; Maga, G; Wellinger, R E; Hübscher, U; Sogo, J M (1999). Electron microscopic analysis reveals that replication factor C is sequestered by single-stranded DNA. *Nucleic Acids Research*, 27(17):3433-3437.

DOI: <https://doi.org/10.1093/nar/27.17.3433>

Electron microscopic analysis reveals that replication factor C is sequestered by single-stranded DNA

Robert C. Keller, Romina Mossi¹, Giovanni Maga², Ralf E. Wellinger, Ulrich Hübscher¹ and José M. Sogo*

Institute for Cell Biology, ETH-Hönggerberg, CH-8049 Zürich, Switzerland, ¹Department of Veterinary Biochemistry, University Zürich-Irchel, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland and ²Institute of Biochemical and Evolutionary Genetics IGBE-CNR, National Research Council, via Abbiategrosso 207, Pavia I-27100, Italy

Received June 8, 1999; Revised and Accepted July 16, 1999

ABSTRACT

Replication factor C (RF-C) is a eukaryotic heteropentameric protein required for DNA replication and repair processes by loading proliferating cell nuclear antigen (PCNA) onto DNA in an ATP-dependent manner. Prior to loading PCNA, RF-C binds to DNA. This binding is thought to be restricted to a specific DNA structure, namely to a primer/template junction. Using the electron microscope we have examined the affinity of human heteropentameric RF-C and the DNA-binding region within the large subunit of RF-C from *Drosophila melanogaster* (dRF-Cp140) to heteroduplex DNA. The electron microscopic data indicate that both human heteropentameric RF-C and the DNA-binding region within dRF-Cp140 are sequestered by single-stranded DNA. No preferential affinity for the 3' or 5' transition points from single- to double-stranded DNA was evident.

INTRODUCTION

In vitro studies of the simian virus 40 replication system have identified several auxiliary replication proteins which are essential for eukaryotic DNA replication (1–3). Replication factor C (RF-C) (4) is a heteropentameric protein (reviewed in 5) which is required to load the homotrimeric proliferating cell nuclear antigen (PCNA) (6) in an ATP-dependent manner onto DNA. Upon association of PCNA with either DNA polymerase (pol) δ (7) or ϵ (8) processive DNA synthesis commences (9).

The binding affinity of RF-C for DNA has been directly and indirectly described in numerous studies (2,3,10–18). These reports, however, contain a certain degree of discrepancy. In an early study it was shown that RF-C could bind to a hairpin DNA oligomer of 96 nt total length with a 3' recessed end and, in addition, that DNA binding by RF-C was not sequence specific, but dependent on the DNA structure (2). Later, Podust *et al.* showed that RF-C was able to load PCNA onto double-stranded and gapped M13 DNA (11). They also noted that linearisation of the DNA led to the dissociation of the RF-C–PCNA complex suggesting an ability of the complex to track along

DNA. In a series of studies involving DNA oligomers, either single-stranded or hybridised to form DNA duplexes, the large subunit RF-Cp140 and its DNA-binding region were shown to preferentially bind double- and not single-stranded DNA (12,13,19). This was contradicted in a study where the binding of RF-C and RF-Cp140 to hairpin DNA was not competed by double-stranded DNA and RF-Cp140 displayed, albeit weak, binding to single-stranded DNA oligomers (16). Allen *et al.* (17) have indicated that the minimal DNA-binding region of the large subunit of RF-C from *Drosophila melanogaster* (dRF-Cp140) as well as human RF-C preferentially bind 5' phosphate groups, recessed or duplexed, over 3' hydroxyl groups, and Mossi *et al.* have confirmed these results (R.Mossi, R.C.Keller and U.Hübscher, manuscript in preparation). Recently, Podust *et al.* (18) implicated a sequestering of RF-C by single-stranded M13 when single-strand binding protein was omitted from the reaction, similar to the observation made by Tinker *et al.* (20).

In this paper we have used the transmission electron microscope to visualise protein–DNA complexes. The data suggest that human heteropentameric RF-C as well as the DNA-binding region of dRF-Cp140 efficiently bind to single-stranded DNA and that neither show any preference for either 3' hydroxyl or 5' phosphate groups at the transition points from single- to double-stranded DNA.

MATERIALS AND METHODS

Reagents

All chemical reagents were commercially obtained from Fluka (Chemie AG, Buchs, Switzerland) unless noted otherwise.

Proteins

RF-C from HeLa cells was purified according to Hübscher *et al.* (21) and its activity determined in a RF-C dependent replication assay. One RF-C unit is defined as the incorporation of 1 nmol dNTPs in 60 min at 37°C. The pmaRF-C162–314 deletion construct of the DNA-binding region of dRF-Cp140 was kindly provided by Susan H. Hardin (University of Houston, TX, USA) and the protein purified according to Allen *et al.* (17).

*To whom correspondence should be addressed. Tel: +41 1 633 3342; Fax: +41 1 633 1069; Email: sogo@cell.biol.ethz.ch

DNA heteroduplexes

Single-stranded and replicative form I (RF I) M13mp18 DNA was obtained from New England BioLabs, Inc. (NEB; Beverly, MA, USA). The M13mp18 Δ 5000 heteroduplex DNA was made as follows: a 2455 bp DNA fragment was amplified from ssM13mp18 in 35 PCR rounds with primers U4591 (5'-TGT TCC GCA AAA TGA TAA TG-3') and R7046 (5'-CAA TGC CTG AGT AAT GTG TA-3') (Microsynth, Balgach, Switzerland) (Fig. 2) at an annealing temperature of 52°C. After separation over a 0.8% agarose gel and extraction with AgarACE (Promega, Catalys AG, Zürich, Switzerland) the amplified fragment was hybridised to ssM13mp18 at a ratio of 1:2 (w/w) in annealing buffer (10 mM Tris-HCl, pH 8, 10 mM EDTA, 100 mM NaCl). The M13mp18 Δ 1000 heteroduplex DNA was made as follows: 1 μ g RF I M13mp18 was double digested with 15 U each of *Ava*II and *Bgl*II (Boehringer Mannheim, Rotkreuz, Switzerland) in buffer A [33 mM Tris-acetate, pH 7.9, 10 mM magnesium acetate, 66 mM potassium acetate, 0.5 mM dithiothreitol (DTT)] for 2 h at 37°C. The 6229 bp fragment was gel purified and hybridised to ssM13mp18 as described above. In order to eliminate the salt the buffer was exchanged for 1 mM triethanolamine-HCl (TEA-HCl), pH 7.4, over a Quickspin Sephadex G-50 column (Boehringer Mannheim).

Electron microscopy

Reactions of 200 ng DNA with 1.5 μ g of the DNA-binding region of dRF-Cp140 were incubated for 1 h at 37°C in a total volume of 5 μ l in buffer 4 (20 mM Tris-acetate, pH 7.9, 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT; NEB). Five units of *Eco*RI (Boehringer Mannheim) were added to the sample after 30 min of incubation to linearise M13mp18 Δ 5000. Samples were fixed and spread as described below. Reactions of 200 ng M13mp18 Δ 5000 or 100 ng M13mp18 Δ 1000 with 0.6 mU human heteropentameric RF-C were incubated for 17 min at 37°C in a total volume of 5.5 μ l in TEAD buffer (40 mM TEA-HCl, pH 7.5, 10 mM MgCl₂, 1 mM DTT) in the presence of 1 mM ATP. Five units of *Eco*RI (Boehringer Mannheim) or *Bsp*HI (NEB) were added after 2 min of incubation to linearise the DNA. Spun-down samples were fixed in 3% (v/v) formaldehyde buffered in 15 mM TEA (pH not adjusted) for 15 min at room temperature. A prior fixation with 0.1% glutaraldehyde did not result in any significant change in the yield of protein-DNA complexes and was therefore deemed superfluous. In order to enhance the unfolding of the single-stranded DNA without disrupting the duplex DNA regions an incubation for 15 min at 37°C was followed by a brief chilling in ice water. Benzyltrimethylammonium chloride [BAC; Bayer, AG, Leverkusen, Germany; 0.2% (w/v)] in 100% formamide was diluted 10-fold in water and 0.1 vol was added to the sample before spreading on a hypophase of bi-distilled water. The surface of carbon-coated grids (Stork Veco B.V., The Netherlands) was treated on water droplets containing 33 μ g/ml ethidium bromide for 10 min at room temperature, dried and briefly brought into contact with the monolayer at the water-air interface formed with the spreading solution containing the protein-DNA complexes. Subsequently the grids were positively stained in 1 mM uranyl acetate, 1 mM HCl in 90% ethanol, rinsed in undiluted ethanol, and air dried. The grids were rotary shadowed at an angle of 1.5° to the vertical




Figure 1. Sequestration of the DNA-binding region of the large subunit of RF-C from *D.melanogaster* by single-stranded M13mp18. An aliquot of 1.5 μ g of the DNA-binding region of dRF-Cp140 was incubated with 100 ng single-stranded and 100 ng replicative form I M13mp18 DNA and spread as described in Materials and Methods. The bar corresponds to 1 kb.

with 900 Hz carbon-platinum as measured on a quartz thin crystal monitor from Balzers. Electron micrographs were taken at magnifications of 13 500 \times and 18 500 \times on a Philips EM420 at an accelerating voltage of 100 kV. For DNA length measurements electron micrographs were enlarged 3-fold and dots on DNA filaments were mapped with computerised tracing software (developed by David Sargent, ETH Zürich).

RESULTS

The DNA-binding region of the large subunit of RF-C from *D.melanogaster* is sequestered by single-stranded DNA

The DNA-binding region of dRF-Cp140 was incubated with equal amounts of single-stranded and RF I M13mp18 DNA. Due to the fused maltose binding protein which increased the molecular weight to ~60 kDa the DNA-binding region of dRF-Cp140 could be clearly visualised under the electron microscope. Circular RF I M13mp18 molecules were mainly free of dots whereas the single-stranded DNA was slightly collapsed and coated with numerous, distinct dots (Fig. 1). Similar results were obtained with human RF-C (data not shown) suggesting that the DNA-binding region of dRF-Cp140 and human heteropentameric RF-C preferentially bind to single-stranded DNA.

The DNA-binding region of the large subunit of RF-C from *D.melanogaster* preferentially binds to a single-stranded gap

We next examined the binding affinity in DNA molecules containing both single- and double-stranded DNA regions. These heteroduplexes also inherently contained 3' and 5' transition points from single- to double-stranded DNA which allowed the simultaneous detection of specific binding to primer/template DNA structures. In earlier experiments RF-C and the DNA-binding region of dRF-Cp140 had been shown to bind to single-stranded DNA regions of down to 30 nt and 420 nt length respectively (data not shown). However, due to the small gap size it was not possible to reliably determine the exact position of the protein-DNA complex. Thus, we designed a DNA heteroduplex with a larger gap of 5000 nt




Figure 2. The DNA-binding region of the large subunit of RF-C from *D.melanogaster* preferentially binds single-stranded DNA. (A) Schematic diagram of DNA heteroduplex M13mp18Δ5000. (B) An aliquot of 200 ng circular M13mp18Δ5000 was incubated with 1.5 μg of the DNA-binding region of dRF-Cp140 and spread as described in Materials and Methods. (C) As in (B) but in addition the DNA was asymmetrically linearised with *EcoRI* (see schematic below), subsequently fixed in 3% formaldehyde and spread as described in Materials and Methods. The bar corresponds to 0.5 kb.

through PCR amplification of a 2.4 kb M13mp18 fragment which was subsequently annealed to single-stranded M13mp18 (Fig. 2A) as described in Materials and Methods. In order to orient the 3' and 5' transition points between single- and double-stranded DNA M13mp18Δ5000 was asymmetrically linearised with *EcoRI* in the polylinker region (Fig. 2C, bottom). Upon incubation of this heteroduplex with the DNA-binding region of dRF-Cp140 the resulting dots appeared in the single-stranded region (Fig. 2B) leaving the double-stranded region devoid of dots. When M13mp18Δ5000 was linearised with *EcoRI* during the incubation with the DNA-binding region of dRF-Cp140 and subsequently fixed, the dots remained in the single-stranded region and did not dissociate from the template (Fig. 2C). Due to the partial collapse of the single-stranded DNA a mapping of the DNA-binding region of dRF-Cp140 in the gap was very difficult. Visual inspection of the protein-DNA complexes, however, did not reveal any preferential binding of the DNA-binding region of dRF-Cp140 to the 3' or 5' transition points.

Human heteropentameric RF-C binds to single-stranded regions of M13mp18Δ5000 and does not show preference for the 3' or 5' transition points from single- to double-stranded DNA

As the DNA-binding region of dRF-Cp140 may not have sufficed for DNA structure recognition we expanded the experiments to examine the binding properties of human heteropentameric RF-C. When incubated with M13mp18Δ5000 preferential binding to the single-stranded gap in the heteroduplex DNA molecule was evident (Fig. 3A-C).




Figure 3. Human heteropentameric RF-C randomly binds in the 5 kb single-stranded gap. An aliquot of 200 ng M13mp18Δ5000 was linearised with *EcoRI* and incubated with human RF-C (0.6 mU) as described in Materials and Methods. One or more dots are located within the single-stranded gap (A), at the 5' transition point (B) or at the 3' transition point (C) between single- and double-stranded DNA. In addition, single-stranded DNA can be seen with numerous, distinct dots (B, top). (D) The histogram has been normalised to show the relative position of the dots on their respective DNA molecules. Human heteropentameric RF-C (white columns) preferentially binds to the single-stranded gap (light grey area). The double-stranded DNA with the 5' transition point at R7046 (dark grey area) and the 3' transition point at U4591 (black area) are situated to the left and right of the gap respectively (see schematic below). The bar corresponds to 1 kb.

Occasionally, more than one dot could be detected in the gap (Fig. 3A). In addition complexes could be found located at either the 3' or the 5' transition points (Fig. 3B and C). Occasionally, single-stranded DNA, which was not eliminated from the samples after heteroduplex formation, was observed with

numerous, distinct dots (Fig. 3B, top), similar to the results obtained with the DNA-binding region of dRF-Cp140 (Fig. 1). When the protein : DNA ratio was increased 3-fold compared to Figure 3A–C the single-stranded region collapsed resulting in a protein aggregate with two or more protruding double-stranded DNA filaments corresponding to the non-collapsed double-stranded DNA (data not shown). In order to detect a preference for either transition point the concentration of human RF-C was reduced. Although the yield of protein–DNA complexes was lower under these conditions, the unfolding of the single-stranded region was sufficient for mapping the position of the resulting dots. The histogram in Figure 3D does not show any preferential binding to either transition point. It is, however, evident that human RF-C predominantly binds to the single-stranded region of the DNA heteroduplex.

Human heteropentameric RF-C binds to single-stranded regions of M13mp18Δ1000 and does not show preference for the 3' or 5' transition points from single- to double-stranded DNA

As it is conceivable that the binding to single-stranded DNA was due to its higher ratio over double-stranded DNA, we repeated the experiment using a second template with a reduced gap size of 1 kb. Human RF-C was predominantly located in the single-stranded gap. Dots were located at either transition point or randomly distributed throughout the single-stranded gap (Fig. 4A and B). In Figure 4C the histogram reveals a slightly higher frequency of human RF-C at the 5' transition point.

DISCUSSION

Biochemical studies on RF-C and its subunits with a variety of DNA templates have been performed over the past decade (2,3,10–18). As the observed binding of RF-C and its large subunit to DNA substrates is varied, and it has been hypothesised that RF-C is sequestered by single-stranded DNA in plasmid templates (18,20), we decided to determine whether RF-C preferentially bound single- or double-stranded DNA in large DNA molecules. Furthermore, we tried to clarify whether a preference for either the 3' hydroxyl or the 5' phosphate group was discernible.

Incubation of the DNA-binding region of dRF-Cp140 with single- and double-stranded M13mp18 at a 1:1 ratio clearly showed an almost exclusive binding to single-stranded DNA (Fig. 1). In initial experiments this binding invariably led to the collapse of the single-stranded DNA resulting in amorphous aggregates (data not shown). Upon adjustment of the protein concentration these dots were resolved into aggregates with single-stranded DNA protrusions. As protein-free single- and double-stranded DNA molecules were completely unfolded, the collapse was most likely due to the bound protein aggregates. It is interesting to note that despite an absence of primer/template structures the DNA-binding region of dRF-Cp140 was able to bind single-stranded DNA under the conditions used (Fig. 1).

Having shown that the DNA-binding region of dRF-Cp140 binds to single-stranded DNA molecules, we wanted to examine its binding affinity to heteroduplex molecules containing both single- and double-stranded DNA regions. Furthermore, we wanted to see which transition point it would preferentially




Figure 4. Human heteropentameric RF-C randomly binds in the 1 kb single-stranded gap. An aliquot of 100 ng M13mp18Δ1000 was linearised with *Bsp*HI and incubated with human RF-C (0.6 mU) as described in Materials and Methods. One or more dots are located within the gap (A) or towards the 5' transition point between single- and double-stranded DNA (B). (C) The histogram shows preferential binding of human RF-C (white columns) to the single-stranded gap (light grey area), with the 5' transition point at the *Bgl*III (6935) (dark grey area) and the 3' transition point at the *Ava*II (5914) (black area) restriction sites to the left and right respectively (see schematic below). The bar corresponds to 1 kb.

bind to, when given a choice between a 3' hydroxyl and a 5' phosphate group. In a M13mp18 heteroduplex with a large single-stranded gap of ~5 kb it was virtually impossible to sufficiently unfold the single-stranded region in order to accurately map the position of the protein on the DNA filament. Most dots, however, were located within the single-stranded gap and, curiously, not over either transition point (Fig. 2A and B). This was not entirely unexpected, as it has been shown that the DNA-binding region of RF-Cp140 binds with less specificity and is not required for RF-C binding and subsequent PCNA loading (16,18). Therefore, its function could be to recognise and bind any single-stranded DNA encountered, further actions being determined by the presence or absence of primer/template junctions, which may be detected in concert with the smaller subunits. Thus, we next decided to examine the affinity of human heteropentameric RF-C for the DNA heteroduplex. In the presence of ATP human RF-C randomly bound along the entire length of the single-stranded gap showing no preference for either transition point (Fig. 3C). To exclude the possibility of

human RF-C binding to single-stranded DNA due to its higher availability over double-stranded regions we designed a further M13mp18 heteroduplex with a smaller single-stranded gap of ~1 kb. Despite a 6-fold increase in double- over single-stranded DNA human RF-C still preferentially bound in the single-stranded gap (Fig. 4C). The slightly higher frequency of molecules bound at the 5' transition point supports the findings of Allen *et al.* (17) and Mossi *et al.* (R.Mossi, R.C.Keller and U.Hübscher, manuscript in preparation) but cannot be considered significant. Whether human RF-C directly binds to and remains immobile in the gap, or binds to a transition point and then tracks along the single-stranded DNA could not be elucidated as human RF-C binding after 0.5, 1, 2 and 10 min revealed no significant shift in position in the single-stranded gap (data not shown). The interesting question arises of what effect PCNA may have on the binding of human RF-C to specific DNA structures. Despite an unavoidable dilution of the sample caused by its dispersion on the water hypophase, the protein-free BAC spreading technique applied for the work presented here is the most adequate to study this problem since it facilitates the spreading of large single-stranded DNA regions with bound protein complexes. However, an additionally added protein would necessitate gel filtration in order to reduce the background. This would result in a substantial primary dilution of the samples which was sufficiently compensated in the mica-replica assay described by Podust *et al.* (11) by a subsequent concentration of the sample at the air-water interface of the water droplets before transfer to the mica. Furthermore, it would be desirable to identify and confirm the presence of protein complexes bound to DNA by immuno-gold tagging. The effects of replication accessory proteins on the binding affinity of RF-C for specific DNA structures will form the basis for future binding studies when the assay has been suitably improved.

In conclusion, we have taken advantage of the electron microscope to examine the influence of primer/template structures on a DNA-binding protein in large single-stranded DNA gaps of 1–5 kb. We have demonstrated that the DNA-binding region of dRF-Cp140 and human heteropentameric RF-C are sequestered by single-stranded DNA in M13mp18 DNA molecules and do not bind to double-stranded DNA. Furthermore, no preferential affinity for 3' hydroxyl or 5' phosphate groups in the absence of PCNA were apparent. RF-C may therefore be strongly attracted by single-stranded DNA and upon PCNA binding specifically seek a template/primer junction to load the clamp in order for processive DNA synthesis by pol δ or ϵ to commence.

ACKNOWLEDGEMENTS

We thank Heide Mayer-Rosa for her technical assistance. This work was supported by the Swiss Cancer League (grant no. KFS 164-9-1995 to R.C.K., U.H. and J.M.S.), by the UBS 'im Auftrag eines Kunden' to R.M., by the TMR (grant no. ERB-MRXCT 970125 to G.M. and U.H.), by the Swiss National Science Foundation (grant no. 31-52246.97 to J.M.S.) and by the Kanton of Zürich.

REFERENCES

1. Fairman, M.P., Prelich, G. and Stillman, B. (1987) *Phil. Trans. R. Soc. Lond. B Biol. Sci.*, **317**, 495–505.
2. Tsurimoto, T. and Stillman, B. (1991) *J. Biol. Chem.*, **266**, 1950–1960.
3. Tsurimoto, T. and Stillman, B. (1991) *J. Biol. Chem.*, **266**, 1961–1968.
4. Tsurimoto, T. and Stillman, B. (1989) *Mol. Cell. Biol.*, **9**, 609–619.
5. Mossi, R. and Hübscher, U. (1998) *Eur. J. Biochem.*, **254**, 209–216.
6. Prelich, G., Kostura, M., Marshak, D.R., Mathews, M.B. and Stillman, B. (1987) *Nature*, **326**, 471–475.
7. Hindges, R. and Hübscher, U. (1997) *J. Biol. Chem.*, **378**, 345–362.
8. Hübscher, U. and Thommes, P. (1992) *Trends Biochem. Sci.*, **17**, 55–58.
9. Podust, V.N., Georgaki, A., Strack, B. and Hübscher, U. (1992) *Nucleic Acids Res.*, **20**, 4159–4165.
10. Podust, L.M., Podust, V.N., Floth, C. and Hübscher, U. (1994) *Nucleic Acids Res.*, **22**, 2970–2975.
11. Podust, L.M., Podust, V.N., Sogo, J.M. and Hübscher, U. (1995) *Mol. Cell. Biol.*, **15**, 3072–3081.
12. Cai, J., Uhlmann, F., Gibbs, E., Flores-Rozas, H., Lee, C.G., Phillips, B., Finkelstein, J., Yao, N., O'Donnell, M. and Hurwitz, J. (1996) *Proc. Natl Acad. Sci. USA*, **93**, 12896–12901.
13. Fotedar, R., Mossi, R., Fitzgerald, P., Rousselle, T., Maga, G., Brickner, H., Messier, H., Kasibhatla, S., Hübscher, U. and Fotedar, A. (1996) *EMBO J.*, **15**, 4423–4433.
14. Yao, N., Turner, J., Kelman, Z., Stukenberg, P.T., Dean, F., Shechter, D., Pan, Z.Q., Hurwitz, J. and O'Donnell, M. (1996) *Genes Cells*, **1**, 101–113.
15. Cai, J., Gibbs, E., Uhlmann, F., Phillips, B., Yao, N., O'Donnell, M. and Hurwitz, J. (1997) *J. Biol. Chem.*, **272**, 18974–18981.
16. Uhlmann, F., Cai, J., Gibbs, E., O'Donnell, M. and Hurwitz, J. (1997) *J. Biol. Chem.*, **272**, 10058–10064.
17. Allen, B.L., Uhlmann, F., Gaur, L.K., Mulder, B.A., Posey, K.L., Jones, L.B. and Hardin, S.H. (1998) *Nucleic Acids Res.*, **26**, 3877–3882.
18. Podust, V.N., Tiwari, N., Stephan, S. and Fanning, E. (1998) *J. Biol. Chem.*, **273**, 31992–31999.
19. Tsurimoto, T. and Stillman, B. (1990) *Proc. Natl Acad. Sci. USA*, **87**, 1023–1027.
20. Tinker, R.L., Kassavetis, G.A. and Geiduschek, E.P. (1994) *EMBO J.*, **13**, 5330–5337.
21. Hübscher, U., Mossi, R., Ferrari, E., Stucki, M. and Jonsson, Z.O. (1999) In Cotterill, S. (ed.), *Eukaryotic DNA Replication: A Practical Approach*. Oxford University Press, Oxford, pp. 119–137.