



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2014

High-resolution analysis of a QTL for resistance to *Stagonospora nodorum* glume blotch in wheat reveals presence of two distinct resistance loci in the target interval

Shatalina, Margarita ; Messmer, Monika ; Feuillet, Catherine ; Mascher, Fabio ; Paux, Etienne ;
Choulet, Frédéric ; Wicker, Thomas ; Keller, Beat

DOI: <https://doi.org/10.1007/s00122-013-2240-4>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-100446>

Journal Article

Published Version

Originally published at:

Shatalina, Margarita; Messmer, Monika; Feuillet, Catherine; Mascher, Fabio; Paux, Etienne; Choulet, Frédéric; Wicker, Thomas; Keller, Beat (2014). High-resolution analysis of a QTL for resistance to *Stagonospora nodorum* glume blotch in wheat reveals presence of two distinct resistance loci in the target interval. TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik, 127(3):573-586. DOI: <https://doi.org/10.1007/s00122-013-2240-4>

High-resolution analysis of a QTL for resistance to *Stagonospora nodorum* glume blotch in wheat reveals presence of two distinct resistance loci in the target interval

Margarita Shatalina · Monika Messmer ·
Catherine Feuillet · Fabio Mascher · Etienne Paux ·
Frédéric Choulet · Thomas Wicker · Beat Keller

Received: 11 July 2013 / Accepted: 19 November 2013 / Published online: 4 December 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract *Stagonospora nodorum* glume blotch (SNG), caused by the necrotrophic fungus *Stagonospora nodorum*, is one of the economically important diseases of bread wheat (*Triticum aestivum* L.). Resistance to SNG is known to be quantitative and previous studies of a recombinant inbred line (RIL) population identified a major quantitative trait locus (QTL) for resistance to SNG on the short arm of chromosome 3B. To localize this QTL (*QSnq.sfr-3BS*) with high resolution, we constructed a genetic map for the QTL target region using information from sequenced flow-sorted chromosomes 3B of the two parental cultivars ‘Arina’ and ‘Forno’, the physical map of chromosome 3B of cultivar ‘Chinese Spring’ and BAC-clone sequences. The mapping population of near-isogenic lines (NIL) was evaluated for SNG resistance in field infection tests. NILs segregated for

disease resistance as well as for plant height; additionally, we observed a high environmental influence on the trait. Our analysis detected a strong negative correlation of SNG resistance and plant height. Further analysis of the target region identified two linked loci associated with SNG resistance. One of them was also associated with plant height, revealing an effect of *QSnq.sfr-3BS* on plant height that was hidden in the RIL population. This result demonstrates an unexpectedly high genetic complexity of resistance controlled by *QSnq.sfr-3BS* and shows the importance of the study of QTL in mendelized form in NILs.

Introduction

Bread wheat (*Triticum aestivum* L.) is globally one of the most important crops in terms of cultivated area and food source. In 2010, FAO reported a production of approximately 654 million tons of wheat worldwide (FAO stat 2010). *Phaeosphaeria nodorum* (E. Müller) Hedjaroude (anamorph *Stagonospora nodorum* (Berk.) Castellani and Germano) is a necrotrophic fungus which infects both wheat glumes, causing *Stagonospora nodorum* glume blotch (SNG), and leaves, causing *Stagonospora nodorum* leaf blotch (SNL). SNG and SNL lead to substantial yield losses in many wheat-growing areas, including Europe, North America and Australia (Duczek et al. 1999; Halama 2002; Oliver et al. 2012). It was shown that the infection progress highly depends on the environmental conditions. For example, the analysis of the large dataset from Rothamsted Broadbalk experiment archive, which included the records from 1844 to 2003, suggested that fluctuations in *S. nodorum* epidemics depend on the amount of rainfall, temperature and SO₂ emissions (Shaw et al. 2008). The importance of wetness (relative humidity and precipitations) and

Communicated by S. Dreisigacker.

Electronic supplementary material The online version of this article (doi:10.1007/s00122-013-2240-4) contains supplementary material, which is available to authorized users.

M. Shatalina · T. Wicker · B. Keller (✉)
Institute of Plant Biology, University of Zurich,
Zollikerstrasse 107, 8008 Zurich, Switzerland
e-mail: bkeller@botinst.uzh.ch

M. Messmer
Research Institute of Organic Agriculture,
5070 Frick, Switzerland

C. Feuillet · E. Paux · F. Choulet
INRA, Genetics, Diversity and Ecophysiology of Cereals,
63100 Clermont Ferrand, France

F. Mascher
Research Station Agroscope Changins-Wädenswil ACW,
1260 Nyon, Switzerland

temperature for the SNG progress were demonstrated in field and growth chamber experiments (Wolf and Francl 2000; Shah and Bergstrom 2002).

Resistance to SNG and SNL in bread wheat has been shown to be genetically distinct (Laubscher et al. 1966; Broennimann 1975; Nelson and Gates 1982; Fried and Meister 1987; Bostwick et al. 1993; Van Ginkel and Rajaram 1999; Wicki et al. 1999). To date, only quantitative resistance to SNG has been reported. It has been identified in several wheat populations independently (Schnurbusch et al. 2003; Aguilar et al. 2005). Schnurbusch et al. (2003) performed a QTL mapping study on a population of 240 recombinant inbred lines (RIL) developed from a cross of cultivars ‘Arina’ and ‘Forno’. This study identified two important QTLs providing resistance to SNG: a major QTL, *QSng.sfr-3BS*, was detected on the short arm of chromosome 3B and explained 31.2 % of the phenotypic variance for the disease resistance within the population. The second QTL *QSng.sfr-4BL*, explaining 19.1 % of phenotypic variance, was found on the long arm of chromosome 4B. Both detected QTLs were not co-localized with any QTLs for plant height. Additionally, an association mapping study on 44 wheat cultivars of different origin showed a significant association between the genetic marker *sun2* from the *QSng.sfr-3BS* region and the resistance to SNG (Tommasini et al. 2007).

It has been shown that plant height is often associated with resistance to splashed dispersed fungal pathogens (Eriksen et al. 2003; Draeger et al. 2007; Srinivasachary et al. 2009). Shorter plants show increased susceptibility and taller plants tend to be more resistant. There are several explanations of this negative correlation. First, in the case of a natural infection, since it progresses from the bottom to the top of the plant, taller plants get an advantage of having larger distance between the leaves (Mesterhazy 1995). Second, it was suggested that microclimate conditions for infection development on tall plants are often less favorable in comparison to the short plants (Scott et al. 1985). Third, the genes influencing plant height and disease resistance might be genetically linked (Gervais et al. 2003; Schnurbusch et al. 2003). Besides plant height, SNG resistance was shown to be associated with morphological traits such as flowering time (Broennimann et al. 1973; Scott et al. 1982; Van Ginkel and Rajaram 1999; Wicki et al. 1999) or the wax layer of glumes (Wicki et al. 1999; Aguilar et al. 2005).

Genetic mapping in bread wheat is challenging as wheat has a large (17 Gb), hexaploid and highly repetitive (80–90 %) genome (Bennett and Smith 1976; Hollister and Gaut 2009; Wanjugi et al. 2009). Furthermore, low level of polymorphism between elite wheat cultivars (Bryan et al. 1999) hinders the development of genetic markers. Recent advances in wheat genomics, such as reducing complexity

by sorting individual chromosome arms using flow cytometry, building chromosome-specific BAC libraries and physical maps, partial genome sequencing, and adopting diploid relatives as models for characterization of the genome, contributed to the easier and faster identification of polymorphisms. In addition, they revealed that, at a high-resolution level, synteny breaks, inversions and interspersed non-syntenic genes are frequent (Choulet et al. 2010), thus highlighting the limitations of using model genomes for map-based cloning in wheat. The isolation of target genes from wheat chromosome 3B benefits from the availability of a physical map (Paux et al. 2008; Rustenholz et al. 2011) that was developed using fingerprints from a bacterial artificial chromosomes (BAC) library of the wheat landrace ‘Chinese Spring’. More than 4,000 markers including simple sequence repeats (SSR) and insertion site based polymorphisms (ISBP) markers derived from BAC-end sequences have been anchored to the 3B physical map and can now be used in various mapping projects. In addition, the anchoring of numerous genes (Rustenholz et al. 2011) on the physical map enables to use the synteny between the regions in wheat chromosome 3B and corresponding regions on rice (*Oryza sativa* L.) chromosome 1 (The International Rice Genome Sequencing Project 2005).

Here, we report on the evaluation of a mapping population of near-isogenic lines for SNG resistance in the field which revealed a segregation for both disease resistance and plant height. Our analysis detected a negative correlation between SNG resistance and plant height. Two genetic regions associated with SNG resistance were found in the previously identified target region. One of them was associated with both, plant height and SNG resistance, suggesting that the same QTL controls both traits.

Materials and methods

Plant material

The mapping population was developed from a cross between the cultivars ‘Arina’ and ‘Forno’, two Swiss winter bread wheat cultivars with contrasting SNG phenotypes: ‘Arina’ shows high levels of resistance to *S. nodorum* glume blotch (SNG) and is a tall variety (120 cm) whereas ‘Forno’ is highly susceptible to SNG and is shorter (105 cm).

F1 progeny of this cross was selfed and back-crossed to the susceptible parent ‘Forno’ twice. The resulting BC2 plants were tested using markers *barc133* and *gwm389*, flanking the target interval of *QSng.sfr-3BS* (Schnurbusch et al. 2003; Tommasini et al. 2007), and markers *barc75* and *sun2* located within the interval. Additionally, BC2 plants were also selected for the presence of the ‘Forno’ allele of the second QTL region on chromosome 4B with the marker

gwm251. The selected BC2 line was back-crossed again to ‘Forno’ and the resulting BC3F2 plants were screened for recombinants in the target interval between the flanking markers. The selected BC3F2 plants were repeatedly selfed and multiplied in the greenhouse to obtain near-isogenic lines (NILs). BC3F5 and BC3F6 plants of these NILs were used in the field tests.

Construction of the genetic map

We mapped in the NIL population microsatellite markers *gwm1034*, *barc75* and *sun2* which were previously mapped in a population of recombinant inbred lines from a cross of ‘Arina’ × ‘Forno’ (Tommasini et al. 2007). Genetic markers (SSR, ISBP) from the 3B physical map were amplified according to the conditions published by Paux et al. (2008). Additionally, new SSR markers were developed based on the sequence information from ctg344 (Choulet et al. 2010) which is located in the QTL interval. PCR products from SSR markers were analyzed on an ABI 3730 sequencer (Applied Biosystems, USA). Microsatellite data was analyzed with the GeneMapper 4.0 software for polymorphisms (Applied Biosystems, USA). ISBP markers were detected using both gel-electrophoresis and high-resolution melting curve analysis (HRM) on a CFX96™ Real Time PCR System (Bio-Rad Laboratories, Hercules, USA). Six BAC clones (TaaCsp3BFhA_0035M17, TaaCsp3BFhA_0112H11, TaaCsp3BFhA_0078F20, TaaCsp3BFhA_0027A18, TaaCsp3BFhA_0163M03, TaaCsp3BFhA_0091M22) were ordered from INRA CNRGV (Toulouse, France) and were sequenced with 454 Titanium technology by GATC (Konstanz, Germany). Newly developed BAC sequence-based InDel and CAPS markers were amplified and analyzed by gel electrophoresis (Table S1). Flow-sorted chromosomes 3B of Arina and Forno were prepared and sequenced as described in Shatalina et al. (2013). New SNP and InDel markers derived from the sequence analysis were amplified using the following cycling conditions: 10 s denaturation at 95°, annealing during 30 s at 61° and 30 s extension at 72°, for 35 cycles. The samples were initially denaturated for 3 min and the cycling was followed by an extension step of 7 min. We used gel electrophoresis to detect the amplified fragments for genotyping. SNP marker *swm01810bd* was detected using HRM on a CFX96™ Real Time PCR System (Bio-Rad Laboratories, Hercules, USA).

Disease phenotyping

Field trials with the NIL population were conducted in 2011 and 2012 at three different locations in Switzerland: Agroscope Reckenholz-Tänikon Research Station, near Zurich (environment Reckenholz); Agroscope

Table 1 Field assessments of SNG resistance and plant height in the NIL mapping population

Year	Environment	AUDPC	Plant height	Ear emergence
2011	Changins	67 NILs	67 NILs	67 NILs
2011	Vouvry	67 NILs	NA	67 NILs
2011	Reckenholz	Excluded	92 NILs	92 NILs
2012	Changings	Excluded	NA	NA
2012	Vouvry	Excluded	NA	NA
2012	Reckenholz	28 NILs	28 NILs	28 NILs

NA, not assessed

Changins-Wädenswil Research Station, close to Nyon (environment Changins); Vouvry in canton Valais, near Lake Geneva (environment Vouvry). In Zurich, the trial was located at 443 m above sea level on loamy soils with an average precipitation of 1,000 mm and an average temperature of 7.9 °C. The Vouvry site is situated at about 381 m above sea level on loamy sandy soils with an average precipitation of 881 mm and an average temperature of 8.7 °C. The Changins site is located at 430 m above sea level on well-drained brown soil (calcaric cambisol) with an average precipitation of 654 mm and a mean temperature of 9.5 °C. Each NIL was tested in randomized 5-row plots in two replications in every environment. In 2011, 92 homozygous NILs were tested in Reckenholz and, because of the insufficient number of seeds from some NILs, only 67 lines were evaluated for resistance to SNG in Changins and Vouvry environments (Table 1). In 2012, due to the limited resources, a subset of 28 NILs was tested in all three locations. Artificial infection was done using a motor sprayer. The infection cocktail contained a mixture of *S. nodorum* isolates from the collection of Agroscope Changins-Wädenswil. Plants were infected twice: a first infection was made at the stage before the emergence of spikes with 5 million spores/ml in 300 l water/ha; and a second infection was made at the flowering stage with 1 million spores/ml in 300 l water/ha. Disease severity was scored according to the Broennimann scale (Broennimann 1968). This method allows scoring the percentage of infected glume area and has a scale of 0, 5, 10, 25, 50, 75 and 100 %. For each environment disease development and severity was scored at least four times during its progress. Final scores were transformed to Area Under the Disease Progress Curve (AUDPC) as described in Jeger and Viljanen-Rollinson (2001). Plant height (cm) was measured in environments Reckenholz and Changins in 2011 and Reckenholz-2012 for both replications of NILs. Time to ear emergence was counted as number of days after the January, 1 of corresponding year.

Statistical analysis of the phenotypic assessments

Broad-sense heritability of the traits was estimated according to Hallauer and Miranda Fo (1981):

$$h^2 = \sigma_g^2 / (\sigma_e^2/re + \sigma_{ge}^2/e + \sigma_g^2)$$

where σ_g^2 genotypic variance, σ_e^2 environmental variance, σ_{ge}^2 variance of genotype/environment interaction, e number of environments, r number of replicas.

The variances were estimated from the mixed linear model:

$$Y = X\beta + Zu + Tv + \varepsilon$$

where Y is a vector of phenotypic observations, β is a vector of fixed genotype effects, u is a vector of random environmental effects, v a vector of random effects from interaction of genotype and environment and ε vector of residual errors. X , Z and T are coefficients of corresponding fixed and random effects. Mixed linear model was analyzed using JMP 9.0 software (SAS institute, Cary, USA).

Single marker analysis (SMA) was conducted in the R environment for statistical computing (R development core team 2013) using functions ‘lme’ and ‘lmer’ following the mixed linear model:

$$Y = X\beta + Zu + Tv + \varepsilon.$$

Y is a vector of phenotypic data (SNG resistance or plant height), β is a vector of fixed marker effects, u is a vector of random environmental effects, v a vector of random effects from interaction of marker and environment and ε vector of residual errors. X , Z and T are coefficients of corresponding fixed and random effects. Following the guidelines for model optimization (Crawley 2007), we eliminated the non-significant marker \times environment interaction from the model for both AUDPC and plant height. The calculation was done using the reduced model:

$$Y = X\beta + Zu + \varepsilon.$$

Plant height was included as a fixed cofactor in the general linear model (‘glm’ function in the R environment) to test its effect on AUDPC:

$$Y = X\beta + Zu + \varepsilon.$$

Y is a vector of phenotypic data (SNG resistance), β is a vector of fixed marker effects, u is a vector of fixed effects from plant height and ε vector of residual errors. X and Z are coefficients of corresponding fixed and random effects.

Calculations of Spearman rank correlation coefficient (r) for the traits between different environments and replicas were done using JMP 9.0 (SAS institute, Cary, USA). Percentage of explained variance for environment and plant height was estimated using function ‘ICC1’ of package

‘multilevel’ in the R environment for statistical analysis. t tests were conducted in R.

Results

High-resolution mapping of the *QSng.sfr-3BS* target region supported by the 3B physical map and whole chromosome sequencing

Schnurbusch et al. (2003) identified *QSng.sfr-3BS* as the major QTL for resistance to SNG in the wheat cultivar ‘Arina’. This QTL is located in the telomeric part of chromosome 3B in the interval flanked by the SSR markers *gwm389* and *barc133*. To further characterize this target interval and the QTL, we developed a high-resolution mapping population consisting of NILs derived from a cross between the susceptible cultivar ‘Forno’ as recurrent parent and the resistant cultivar ‘Arina’ as donor. Among 1,320 NILs (BC3F2), we selected 145 lines showing recombination between the flanking markers of the target interval. Ninety-two of them were homozygous for the flanking markers. The size of the target region for the *QSng.sfr-3B* in our NIL population was 11 cM.

Several approaches were used to construct a high-density genetic map in the target interval. In a first step, we mapped markers *gwm1034*, *barc75* and *sun2* from the previous map of the ‘Arina’ \times ‘Forno’ RIL population (Pailard et al. 2003; Tommasini et al. 2007). We also used the availability of the chromosome 3B physical map for cultivar ‘Chinese Spring’ (Paux et al. 2008) which allowed us to estimate the physical size of the region of interest as well as to map additional markers developed for anchoring the physical map. According to the physical map, the 11 cM target interval spanned a region of approximately 10 Mbp on chromosome 3B (Fig. 1). We mapped a set of SSR and ISBP markers, originally derived by Paux et al. (2008) for the anchoring of the physical map of chromosome 3B. Unfortunately, a vast majority of the tested ISBP markers from the physical map were not polymorphic in the ‘Arina’ \times ‘Forno’ NIL population. From a total of 33 tested markers only *cfp3062* was polymorphic and was successfully added to the map. In addition, two new SSR markers (*swm105*, *swm106*) were designed based on the sequence information from the sequenced physical contig 344 of chromosome 3B (Choulet et al. 2010). These markers were polymorphic and genetically mapped to our target interval.

We sequenced five BAC clones from the physical contigs of the target region on chromosome 3B (TaaCsp3BFhA_0112H11 from ctg3042, TaaCsp3BFhA_0078F20 from ctg1525, TaaCsp3BFhA_0027A18 from ctg516, TaaCsp3BFhA_0163M03 from ctg6 and TaaCsp3BFhA_0091M22 from ctg1793). These BAC clones were selected based on their




Fig. 1 Comparative analysis of the target region for *Qsng.sfr-3BS* on the genetic map of the NIL population and the physical map of wheat chromosome 3B. The genetic map of the fine-mapping NIL population (92 lines) of ‘Arina’ × ‘Forno’ (in dark gray) is shown: numbers represent the number of identified recombinants between given markers, in blue-markers developed based on the sequence analysis of BAC-clones and a partial sequence of ctg344, in green-markers developed from SNPs and InDels identified between sequences up- and downstream of coding regions in sequenced flow-sorted chromosomes 3B of Arina and Forno; numbers in the “green” marker names correspond to the numbers of *B. distachyon* gene homologs. Physical map of wheat cv. ‘Chinese Spring’: light gray bars represent physical contigs with their numbers on the left side, short blue bars represent sequenced BAC-clones from corresponding physical contigs. The red bars A represent the interval significantly associated with both SNG resistance and plant height, B the interval associated with SNG resistance. Wheat chromosome 3B is shown in light gray with the target interval indicated in darker gray shade

position on the physical map to cover the large gaps on the genetic map (blue lines, Fig. 1). The BAC sequences were used in two approaches for developing new markers. First, we designed primers to amplify the regions containing microsatellites to develop SSR markers. Second, non-repetitive regions up- and downstream of the coding sequences were amplified from both parents to develop insertion–deletion (InDel) and cleaved amplified polymorphic sequences (CAPS) markers.

Three new InDel markers (*swm101*, *swm102*, *swm103*) and one CAPS marker (*swm104*) were obtained by this approach.

To further increase marker density, we sequenced flow-sorted chromosomes 3B of both parental cultivars and this sequence information was used to design single nucleotide polymorphism (SNP) and InDel markers as described in Shatalina et al. (2013). We selected only contigs containing genes which corresponded to the *Brachypodium distachyon* homologs from the region *Bd2g00010-Bd2g02000* of chromosome 2, showing synteny with the target region of wheat chromosome 3B. InDel and SNP markers were based on small 2–3 bp polymorphisms in intergenic and coding regions of sequences between the two cultivars as well as on the presence and absence of gene fragments. As a result, five new markers (*swm00640 bp*, *swm00850bd*, *swm01020bd*, *swm01210bd* and *swm01820bd*) were identified and integrated in the genetic map. After integration of all newly developed markers, a total number of 31 genetic markers were mapped to the QTL target interval. The two largest regions of the map not covered by markers corresponded to 26 and 28 recombinants in the NIL population.

NIL population segregated for plant height and SNG resistance

The NIL population was tested for resistance to *Stagonospora nodorum* glume blotch (SNG) under field conditions during the years 2011 and 2012 at three locations in Switzerland: Reckenholz, Changins and Vouvry (Table 1). We estimated the infection quality based on good differentiation of the disease scores of the parental cultivars. In 2011, the infection developed well on the field sites in Vouvry and Changins (average AUDPC scores of ‘Forno’ were 526.5- and 423 compared to ‘Arina’ 153.3 and 93, respectively), whereas in 2012 the Reckenholz environment supported a good infection (Table 2). Three other datasets (Reckenholz 2011, Vouvry 2012 and Changins 2012) were excluded from the analysis due to the insufficient development of the infection. Therefore, we included in the analysis only the phenotypic data of 67 NILs from the environments Changins and Vouvry for the infection tests in 2011 and data of 28 NILs from Reckenholz in 2012 (Table 1; Fig. 2).

To test whether the SNG resistance phenotypes were reproducible under different field conditions, we calculated the Spearman correlation coefficient (r) between the replications of the same genotypes within the individual field site as well as correlation between the different fields (Table S2). The correlation of the AUDPC values between the NIL replications within individual environments varied from 0.47 (Changins-2011) to 0.81 (Reckenholz-2012) and between the environments from 0.56 (correlation between Changins-2011 and Vouvry-2011) to 0.84—correlation between Reckenholz-2012 and Changins-2011.

Fig. 2 Graphical genotypes of the 67 and 28 NILs tested in the field. The genetic map shown on the right is based on a total of 92 NILs. In *blue*-‘Arina’ allele, in *red*-‘Forno’ allele. The *dark colored blue* and *red bars* with the *green asterisks* indicate the 28 NILs tested in field trials in 2012. *Red bars* on the genetic map are indicating the intervals associated with SNG resistance (**b**) and SNG resistance and plant height (**a**)




Table 2 Plant height and AUDPC values of parental lines ‘Arina’ and ‘Forno’ and fine-mapping NIL population

Trait	Environment	No of NILs tested	QTL donor line Arina	Recurrent parent Forno	Average NIL	Min NIL	Max NIL
AUDPC	Vouvry-2011	67	153.3 ± 30.8	526.5 ± 76.5	460.4 ± 78.5	295	701.5
	Changins-2011	67	93 ± 8.7	423 ± 73.6	329.4 ± 76.9	187	575
	Reckenholz-2012	28	67.4 ± 6.4	224.3 ± 68.7	143.2 ± 60.1	59	295.5
Plant height (cm)	Reckenholz-2011	67	115 ± 7.06	103.3 ± 5.6	103.2 ± 5.8	90	118
	Changins-2011	67	109.9 ± 2.4	101.8 ± 4.4	101.6 ± 4.6	90	115
	Reckenholz-2012	28	124.4 ± 1.9	107.6 ± 1.4	109.4 ± 4.9	100	120
Ear emergence (days)	Changins-2011	67	139	139.5 ± 0.6	141.7 ± 0.7	141	144
	Reckenholz-2011	67	138.5 ± 0.5	138.5 ± 0.5	139.1 ± 0.6	138	141
	Vouvry-2011	67	142.6 ± 0.9	142.1 ± 0.8	142.6 ± 0.7	141.3	144
	Reckenholz-2012	28	140.5 ± 0.7	140.6 ± 1.3	143.1 ± 0.6	141.2	145.8


The last three columns show average, minimum and maximum values of the corresponding trait in the NIL population

Moderate to high correlation of the SNG resistance scores indicated their reproducibility across the environments. Due to the continuous distribution of AUDPC values, the NILs could not be grouped into two contrasting disease response classes (resistant or susceptible). This observation suggested that the disease response in the NIL population has to be analyzed as a quantitative trait and not as a single Mendelian gene. We also analyzed time to ear emergence

and its correlation with plant susceptibility (Tables 1, 2). The calculated Spearman rank correlation coefficient showed weak (in one environment) or no significant correlation (several environments) between the traits (Table S2).

In addition to a segregating disease response, we observed a segregation for plant height in our high-resolution mapping population. While several lines were taller than the recurrent parent of the NILs (cultivar ‘Forno’), and more similar

Fig. 3 Average values of plant height and AUDPC score of 67 NILs (a) and 28 NILs (b). Average values were calculated from one environment for 67 NILs and two environments for 28 NILs. *Black dots* represent average value of recurrent parent (cultivar Forno)



to SNG-QTL donor cultivar ‘Arina’, some NILs were even shorter than cultivar ‘Forno’. Overall, the tested lines showed a distribution of plant height from 90 to 120 cm (Table 2). The rank correlation (r) of plant height and resistance to SNG ranged from -0.53 (average AUDPC and average plant height in Changins 2011) to -0.78 (average AUDPC and plant height in Reckenholz 2012) and revealed a negative correlation between these two traits (Table S2). This trend is depicted on the plot of SNG resistance (AUDPC value) against plant height of 67 NILs (Fig. 3a), and becomes more pronounced in the combined dataset of 28 NILs from Reckenholz-2012, Changins-2011 and Vouvy-2011 (Fig. 3b). The rank correlation of plant height between the environments varied from a moderate correlation of 0.36 in 67 NILs to a high correlation of 0.77 in 28 NILs (Table S2). Broad-sense heritability of SNG resistance was 0.71 for 67 NILs tested in 2011 in two environments—Changins and Vouvy. Heritability of plant height was 0.63 for 67 NILs, tested in Changins-2011 and Reckenholz-2011.

Single marker analysis (SMA) defines regions associated with SNG resistance and plant height

To localize *Qsng.sfr-3BS* on the genetic map, we applied single marker analysis (using mixed linear model) to the two datasets (67 NILs and the subset of 28 NILs, Table 3). We included the phenotypic data from Changins-2011 and Vouvy-2011 for the analysis of 67 NILs. The SMA of this dataset detected no significant association between the AUDPC value for SNG and any of genetic markers from the target region. Analysis of the subset of 28 NILs was done using the combined data from Changins-2011, Vouvy-2011 and Reckenholz-2012. It identified an association between AUDPC and the interval *swm00850bd-wmm756* (region A, Fig. 1). According to the physical map of chromosome 3B (Fig. 1), region A has a size of 4.5 Mb (6 physical contigs).

Since the NIL population segregated for plant height, we checked whether plant height is associated with any of the markers in the target region of *Qsng.sfr-3BS*. Similarly to the SMA of AUDPC, we analyzed two datasets of 67 and 28 NILs. For the analysis of 67 NILs, we included the data from Changins-2011 and Reckenholz-2011. Analysis of mixed linear model detected a significant association between the trait and the genetic interval *swm00850bd-wmm756* (region A, Fig. 1), with the most significant F -value for the marker *wmm756* (Table 3). The analysis of 28 NILs included the data from Reckenholz-2012 and 2011, and Changins-2011 and revealed associations of plant height with the genetic interval *gwm389-wmm756*, as well as with the marker *nw1644* which co-segregates with the proximal flanking marker *barc133*. We consider the association of *barc133* to be an artifact, because of the selection of the BC3F2 plants for recombination between the proximal markers *gwm389* and *barc133*, the mapping population consisted only of lines recombining in the target region. Therefore, the parental alleles for *barc133* were exactly opposite to the parental alleles of *gwm389* in all recombinants. Thus, as there is a lower AUDPC score of the ‘Arina’ allele for marker *gwm389*, there has also to be a lower AUDPC value for the ‘Forno’ allele at marker *barc133*.

Since plant height correlated with disease resistance, the next step was to conduct SMA of disease resistance using plant height as a cofactor in the analysis. Two datasets were analyzed: 67 NILs with the phenotypic data from Changins-2011 and 28 NILs with the data from Changins-2011 and Reckenholz-2012, where both AUDPC and plant height were measured (Table 4). Surprisingly, SMA for 67 NILs detected one significant association of SNG resistance with the marker *swm01210bd*, which was not detected before in SMA of AUDPC without plant height as a cofactor. Thus, SMA indicates the presence of potentially two intervals associated with resistance to SNG within the target region for *Qsng.sfr-3BS* (regions A and B, Fig. 1).

Table 3 Single marker analysis of SNG resistance and plant height

	Marker	Plant height				AUDPC			
		67 NILs ^a		28 NILs ^b		67 NILs ^c		28 NILs ^d	
		F value ^e	p value ^f	F value ^e	p value ^f	F value ^e	p value ^f	F value ^e	p value ^f
	gwm389	1.324	0.2508	5.593	0.0192*	0.369	0.544	1.250	0.2653
	swm00640bd	1.324	0.2508	5.593	0.0192*	0.369	0.544	1.250	0.2653
	swm00850bd	5.233	0.023*	6.172	0.014*	0.359	0.5498	6.498	0.0117*
	swm101	5.233	0.023*	6.172	0.014*	0.359	0.5498	6.498	0.0117*
	swm102	5.233	0.023*	6.172	0.014*	0.359	0.5498	6.498	0.0117*
The most significant marker <i>wmm756</i> is indicated in bold	swm01020bd	5.233	0.023*	6.172	0.014*	0.359	0.5498	6.498	0.0117*
	gwm1034	5.233	0.023*	6.172	0.014*	0.359	0.5498	6.498	0.0117*
^a SMA (mixed linear model) of the 67 NILs with two replications from 2 environments (Changins-2011 and Reckenholz-2011)	cfb3530	5.116	0.0245*	6.172	0.014*	0.637	0.4256	6.498	0.0117*
	swm103	5.598	0.0187*	6.172	0.014*	0.177	0.6742	6.498	0.0117*
	barc75	5.598	0.0187*	6.172	0.014*	0.177	0.6742	6.498	0.0117*
	swm104	6.239	0.0131*	6.172	0.014*	0.004	0.9501	6.498	0.0117*
^b SMA (mixed linear model) of the 28 NILs with two replications from 3 environments (Changins-2011, Reckenholz-2011 and Reckenholz-2012)	cfb3417	6.366	0.0122*	6.172	0.014*	0.069	0.7935	6.498	0.0117*
	wmm756	9.126	0.0028**	6.172	0.014*	1.171	0.2803	6.498	0.0117*
	sun2	2.196	0.1396	0.131	0.7178	0.025	0.8741	1.086	0.2989
	cfb5000	2.814	0.0947	0.131	0.7178	0.615	0.4336	1.086	0.2989
	cfb5009	2.814	0.0947	0.131	0.7178	0.615	0.4336	1.086	0.2989
^c SMA (mixed linear model) of the 67 NILs with two replications from 2 environments (Changins-2011 and Vouvry-2011)	cfb5025	2.814	0.0947	0.131	0.7178	0.615	0.4336	1.086	0.2989
	cfb5008	2.047	0.1538	0.131	0.7178	0.123	0.7262	1.086	0.2989
	cfb5021	2.814	0.0947	0.131	0.7178	0.615	0.4336	1.086	0.2989
	cfb5026	2.814	0.0947	0.131	0.7178	0.615	0.4336	1.086	0.2989
^d SMA (mixed linear model) of the 28 NILs with two replications from 3 environments (Changins-2011, Vouvry-2011 and Reckenholz-2012)	cfb5010	2.814	0.0947	0.131	0.7178	0.615	0.4336	1.086	0.2989
	cfp3062	1.984	0.1601	0.131	0.7178	0.600	0.4395	1.086	0.2989
	nw1821	2.814	0.0947	0.131	0.7178	0.615	0.4336	1.086	0.2989
	swm106	3.976	0.0472*	1.229	0.2693	0.353	0.5528	0.966	0.3271
^e F value of the given marker effect	swm01210bd	2.248	0.135	2.825	0.0948	1.177	0.279	0.058	0.8100
	swm01810bd	0.24	0.6246	2.384	0.1246	1.128	0.2892	0.345	0.558
^f Significance thresholds of the given marker effect * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$	nw1644	1.324	0.2508	5.593	0.0192*	0.369	0.544	1.250	0.2653
	barc133	1.324	0.2508	5.593	0.0192*	0.369	0.544	1.250	0.2653

The analysis of 28 NILs detected no significant association between disease resistance and any of the markers in the QTL target region; however, the genetic interval *swm00850bd–nw1821* had t values close to significance (Table 4). For both datasets (28 and 67 NILs) plant height as a fixed cofactor in the model was highly significant. The calculated percentage of the total variance of SNG AUDPC explained by plant height was 34.5 % for 67 NILs. Therefore, plant height and disease resistance to SNG for the region A might have the same genetic basis.

Two genetically distinct regions are contributing to SNG resistance in NILs

Single marker analysis indicated the presence of two loci associated with SNG resistance. We wanted to study if additional analysis could further substantiate this finding. Therefore, we analyzed the distributions of the average

AUDPC and plant height of two environments for 67 NILs in the two allele classes for individual markers (Table 5). Marker *wmm756* showed the largest height difference of 2 cm for ‘Arina’ and ‘Forno’ classes as expected from the SMA described above. Similarly, the allele classes for AUDPC values revealed largest differences for markers *wmm756*, *swm01210bd* and *swm01810bd*. The ‘Arina’ class had the lowest AUDPC values for marker *wmm756* from the region A and marker *swm01210bd* from region B, which corresponded to the resistance loci identified by SMA.

Next, we divided the 67 NILs into four groups based on their genotype for the markers *wmm756* and *swm01210bd* to test the hypothesis of two independent SNG loci (Fig. 4). The distributions of the phenotypic values of the four groups AA-‘Arina’ alleles for the two markers *wmm756* and *swm01210bd*, AF-‘Arina’ for *wmm756* and ‘Forno’ for *swm01210bd*, FA-‘Forno’ *wmm756* and ‘Arina’

Table 4 Single marker analysis of SNG resistance with plant height as a cofactor

Marker	AUDPC		AUDPC	
	67 NIL ^a		28 NIL ^b	
	<i>t</i> -value ^c	<i>p</i> -value ^d	<i>t</i> -value ^c	<i>p</i> -value ^d
gwm389	-0.519	0.605	0.886	0.378
swm00640bd	-0.519	0.605	0.886	0.378
swm00850bd	0.007	0.995	1.960	0.0526
swm101	0.007	0.995	1.960	0.0526
swm01020bd	0.007	0.995	1.960	0.0526
gwm1034	0.007	0.995	1.960	0.0526
cfb3530	0.020	0.984	1.960	0.0526
swm103	0.180	0.858	1.960	0.0526
barc75	0.180	0.858	1.960	0.0526
swm104	0.431	0.667	1.960	0.0526
cfb3417	0.168	0.867	1.960	0.0526
wmm756	-0.119	0.905	1.960	0.0526
sun2	-0.171	0.865	1.978	0.0504
cfb5000	0.273	0.786	1.978	0.0504
cfb5009	0.273	0.786	1.978	0.0504
cfb5025	0.273	0.786	1.978	0.0504
cfb5008	0.328	0.743	1.978	0.0504
cfb5021	0.273	0.786	1.978	0.0504
cfb5026	0.273	0.786	1.978	0.0504
cfb5010	0.273	0.786	1.978	0.0504
cfp3062	0.419	0.676	1.978	0.0504
nw1821	0.273	0.786	1.978	0.0504
swm106	-0.183	0.855	1.160	0.249
swm01210bd	2.334	0.0212*	0.911	0.364
swm01810bd	0.074	0.941	-0.803	0.424
nw1644	0.519	0.605	-0.886	0.378
barc133	0.519	0.605	-0.886	0.378

The significant marker *swm01210bd* is indicated in bold

^a SMA (mixed linear model) of the 67 NILs with two replications from Changins-2011

^b SMA (mixed linear model) of the 28 NILs with two replications from 2 environments (Changins-2011 and Reckenholz-2012)

^c *t*-value of the given marker effect

^d Significance thresholds of the given marker effect * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

swm01210bd and FF-‘Forno’ for both markers are shown in Fig. 4. A *t* test did not detect significant differences between the groups, but when considering the group means there was a clear trend. The average values from the two environments Changins-2011 and Reckenholz-2011 for plant height showed that the AA and AF groups were taller (Fig. 4a). A smaller size of boxes and whiskers on the plot for the AA and AF groups in comparison to the FA and FF groups also indicated that plant height values were distributed within a smaller range in the first two groups with the

Table 5 Plant height and AUDPC for 67 NILs (average of two environments) divided into ‘Arina’ and ‘Forno’ classes for each of the markers (average value of the class is shown)

Marker	AUDPC		Plant height	
	Forno allele	Arina allele	Forno allele	Arina allele
gwm389	391.9	398.0	102.0	102.8
swm103	393.0	396.6	101.7	103.2
wmm756	399.4	388.6	101.5	103.5
cfb5000	389.9	398.0	101.9	103.0
swm01210bd	398.4	387.7	102.9	101.9
swm01810bd	400.8	389.9	102.2	102.5
nw1644	398.0	391.9	102.8	102.0

‘Arina’ allele of marker *wmm756*, whereas in the absence of *wmm756* ‘Arina’ allele plant height might be more strongly influenced by environmental factors.

In agreement with the hypothesis of two resistance loci present in the target region, the average AUDPC values of 67 NILs from the two environments Changins-2011 and Vouvy-2011 were lowest for the AA group (Fig. 4b). The short length of the upper whisker for the AA group suggested that except for the two outliers, this group showed a clear tendency for a stronger resistance as the AUDPC values fall into the same range of low values. In contrast, the FF group showed the highest AUDPC of both median and mean values among all genotypic groups. The sizes of boxes and whiskers for the AF, FA and FF groups were larger than for the AA group, which indicated a wider distribution of the AUDPC values in these groups. The wider distribution of the AUDPC in the groups with no (FF) or only one resistance locus (AF and FA) suggested that the presence of two resistance loci (group AA) indeed resulted in strongest resistance. The AF, FA and FF groups might be more influenced by other genetic components and environmental factors. Thus, we conclude that there were two resistance loci in the target interval and the presence of both loci provided the strongest resistance response. The region A resistance locus was associated with both plant height and SNG resistance. The effect of the resistance locus B linked to *swm01210bd* seemed to be larger than the effect of the resistance locus A linked to *wmm756*.

Discussion

Field infection tests of a NIL population revealed a quantitative distribution for both SNG resistance and plant height. The two traits were reproducible between different tested environments and showed a moderate to high correlation. Further analysis of the target region identified two loci: region A (*swm00850bd* – *wmm756*) and B (*swm01210bd*),




Fig. 4 Box-plots of the average AUDPC and plant height of 67 NILs in four genotypic groups. 67 NILs were divided into four groups based on genotypes of two markers (*wmm756* and *swm01210bd*). AA ‘Arina’ allele for *wmm756* and ‘Arina’ allele for *swm01210bd*, AF ‘Arina’ *wmm756* and ‘Forno’ *swm01210bd*, FA ‘Forno’ *wmm756* and

‘Arina’ *swm01210bd* and FF both markers have ‘Forno’ allele. The black line is a median of each group and black diamond-shaped dot is a group mean. **a** Average plant height score among 2 environments; **b** average AUDPC score. The whiskers depict the minimum and maximum values

associated with SNG resistance. Region A was also associated with plant height.

Fine mapping of the target interval for SNG QTL in the ‘Arina’ × ‘Forno’ NIL population

The physical map of chromosome 3B allowed us to improve the genetic map for the target interval of *QSng.sfr-3BS* by integrating recently developed markers as well as by establishing additional new markers. In contrast to the results in other populations, we detected a very low level of polymorphism in ISBP markers in our NIL population (Paux et al. 2010). In fact, ISBP markers are based on differences in the insertion sites of (retro)transposons and since our NIL population is derived from a cross of two Swiss winter varieties, there might be less diversity in these repeat junctions than in populations derived from wide crosses, like ‘Chinese Spring’ × ‘Renan’, ‘Courtot’ × ‘Chinese Spring’ or ITMI populations, which were used for mapping of ISBPs to anchor the physical map of chromosome 3B. Another problem was the presence of gaps in the genetic map (regions not covered with genetic markers), which could not be predicted or explained by the physical map or synteny with *B. distachyon*. Those gaps might be recombination hot spots (Faris et al. 2000; Akhunov et al. 2003) or large chromosomal insertions specific to the ‘Arina’ × ‘Forno’ population. A larger number of mapped markers will be needed to further test the “hot spot”-hypothesis. Recently developed genomic resources such as physical maps can be helpful for high-resolution mapping purposes, but individual differences between the parental lines and the cultivars used for the construction of the

physical map have to be considered. Sequenced flow-sorted chromosome 3B of the parental lines for the NIL population was a successful approach to increase the marker density. Single nucleotide polymorphism markers derived from sequences of the particular parental cultivars combined with the synteny information are very specific and enabled us to target selected regions. Synteny with *B. distachyon* allowed us to predict the possible position and to choose the potential markers specifically for the critical parts of the genetic map. Both resources, the physical map and the sequence information from ‘Arina’ and ‘Forno’, might be extremely valuable for the identification of candidate genes when the target interval is further narrowed.

SNG resistance is strongly influenced by environmental conditions

Successful development of artificially induced SNG infection was difficult to obtain, as it depends strongly on environmental conditions such as temperature and humidity (Eyal et al. 1987; De Wolf and Francl 2000; Shah and Bergstrom 2002). In agreement with these earlier observations, among six conducted field infection tests (2 years, 3 environments) only three trials had optimal conditions for disease development, while three trials showed insufficient disease development and, therefore, had to be discarded from the analysis. The trials with successful infection also showed a significant impact of environmental factors (58.6 % of the variance) on the disease development resulting in continuous distribution of AUPDC. The range of correlation coefficients of AUDPC values (0.47–0.81) between and within individual environments indicated on

the one hand, presence of the proposed QTL in the selected NILs, and on the other hand, a strong influence of environmental factors on SNG resistance. Reproducibility of the results was demonstrated by the correlation between three environments (0.56–0.84) and heritability of the trait (0.71 for 67 NILs). These results are very similar to the results observed previously in the ‘Arina’ × ‘Forno’ RIL population: correlation of SNG resistance between environments varied from 0.59–0.84 and heritability of the trait was 0.80 (Schnurbusch et al. 2003). These results from phenotyping show that our data are reliable, but too variable for clearly assigning a genotype to a “resistant” or “susceptible” class. Future experiments to obtain resistance data should possibly be done at a larger number of field sites and over several years to get data of sufficient quality for high-resolution mapping and ultimately for cloning of the resistance gene(s).

Is the SNG resistance QTL linked to *wmm756* a plant height QTL?

In addition to the segregation for SNG resistance, we observed segregation for plant height in our NIL population. This segregation was reproducible and heritable, and SNG resistance and plant height showed a moderate to high negative correlation in all tested environments. A correlation between plant height and resistance to SNG was previously shown under field conditions suggesting disease escape for the tall plants (Broennimann et al. 1973; Scott et al. 1982; Van Ginkel and Rajaram 1999; Wicki et al. 1999). Association of plant height and SNG resistance was also identified in the previous QTL study on the RIL population of ‘Arina’ × ‘Forno’ (Schnurbusch et al. 2003) and the RIL population of ‘Forno’ × ‘Oberkulmer’ (Aguilar et al. 2005). However, Schnurbusch et al. (2003) did not detect any QTL for plant height in the target region for *QSng.sfr-3BS*. Several QTLs for plant height were detected on chromosomes 5BL, 5AL, 7BS, 3BL and 2AL on the genetic map of ‘Arina’ × ‘Forno’ RILs. The *QSng.sfr-3BS* interval did not show any significant association with other traits such as plant height or flowering time (Schnurbusch et al. 2003). The fact that we detected a region associated with plant height in the NIL population which was not detected in RILs before might be explained by epistatic effects of other, stronger plant height QTLs in the RIL ‘Arina’ × ‘Forno’ population or two nearby QTLs with opposite allele effects. Looking at the mean (Table 5) and median (Fig. 4) of marker classes, the “Arina” allele of region A (*wmm756*), increases plant height by 2 cm, while the marker classes of “Arina” allele of region B (*swm01210*) slightly decreases plant height by 1 cm. Previously, *Rht5*—a gene modifying plant height was mapped to the same deletion bin on chromosome 3B linked to the

marker *barc102* (Ellis et al. 2005). Therefore, we cannot exclude that the gene underlying the plant height locus in the target region is *Rht5*.

Segregation for plant height and its correlation with SNG resistance in our NIL population was unexpected; we used artificial infection after spike emergence in the field and did not rely on natural infections. Artificial infection by spraying plants from above should have decreased the disease escape due to plant height, because in these conditions tall and short plants received equal amount of spores directly to the spikes in contrast to natural infection which usually is spreading from infected ground debris from the lower leaves to the ears by rain splashes. Nevertheless, tall and short plants could also differ in microclimate conditions of the spikes which are known to be important for disease development (Scott et al. 1985). Measuring of additional phenotypic traits such as spike morphology or distance between spikes and flag leaves in future field trials might help to clarify whether plant height *per se* is influencing SNG resistance or if any other minor features of the tested NILs contribute to the disease escape.

Further evidence for the correlation of plant height and SNG resistance was found by including plant height as a factor in general linear model analysis which significantly influenced the SMA results. Furthermore, the previously detected significance for the genetic markers from the interval *swm00850bd–wmm756* (region A, Fig. 1) became less pronounced, indicating that a large part of the variance was explained by plant height. The correlation of plant height and SNG resistance could have several explanations. SNG resistance in region A could be a gene controlling plant height or it might contain both a resistance gene as well as a plant height gene within a gene cluster, which would explain a co-segregation of the traits. However, the most likely explanation is that the region A controls plant height and results in a change of microclimate at the spike level, with SNG resistance improvement as a pleiotropic effect of the plant height gene.

Two genetically distinct SNG resistance loci are present in the *QSng.sfr-3BS* target interval

The absence of any influence from the genetic background in a NIL population theoretically allows to “Mendelize” the trait of interest and to divide the phenotypic response of the fine-mapping population in two contrasting classes—“resistant” and “susceptible” (Keurentjes et al. 2007; Pea et al. 2009). *QSng.sfr-3B* did not manifest itself as Mendelian trait in our NIL population and was scored quantitatively. The identification of two resistance loci in the *QSng.sfr-3BS* target interval as well as the strong environmental influence possibly explains the observed quantitative response to *S. nodorum* infection. The two loci were

detected by SMA: marker *wmm756* (region A) and marker *swm01210bd* (region B) were both significantly associated with disease resistance. The hypothesis of two segregating QTL in the target region would explain why the explained variation of SNG detected by SMA was not as great (Table 5) as expected based on the QTL analysis of the RILs of Arina x Forno (explaining more than 30 % of the variation for SNG; Schnurbusch et al. 2003), although the NILs in our study were on average much more resistant than the recurrent parent Forno (Table 2; Fig. 3). The analysis of genotypic groups, taking into account the allelic composition of both markers for region A and B, supported this hypothesis. However, no statistically significant differences were found between the four marker classes. This absence of significant differences could be due to the relatively low number of NILs in the four genotypic groups as well as the high environmental dependence of the resistance trait. Since the NIL population was developed based on the hypothesis of one QTL localized in the target interval, the number of recombinants was clearly not sufficient for the unexpected need to identify two genetically distinct loci. To separate the effects of two loci and isolate each of them in the ‘Forno’ genetic background, the number of NILs has to be at least doubled in comparison to the 67 lines analyzed in our study. Additionally, some variation among the NILs for both plant height and AUDPC which probably was caused by the remaining 6.25 % genetic background from ‘Arina’ might also have influenced the results of the analysis.

Tightly linked QTLs controlling the same trait have previously been described for other quantitative traits. In tomato, two QTLs *Brix9-2-5* and *PW9-2-5* located 0.3 cM apart are both modifying sugar content (Fridman et al. 2002). In rice, QTL *Hd3* consisted of two linked QTLs *Hd3a* and *Hd3b* controlling heading date (Monna et al. 2002). A QTL for flowering time *dth1.1* consisted of at least two distinct sub-QTLs *dth1.1a* and *dth1.1b* (Thomson et al. 2006). Another example is a growth-rate QTL in *Arabidopsis thaliana*, consisting of two tightly linked QTLs located within a 210 kb interval (Kroymann and Mitchell-Olds 2005). The SNG resistance analysis presented in this study focusing on *QSng.sfr-3BS* revealed an additional case for two closely linked QTL. Furthermore, the pleiotropic effect of the two involved loci on plant height further complicated the analysis. These findings indicate an unexpectedly high genetic complexity of SNG resistance in cultivar ‘Arina’ and the need for the development of larger populations where the effects of the two linked QTL are separated from each other in a sufficiently large number of individual genotypes. Only in such a mendelized form in even more identical backgrounds a future molecular cloning can be envisaged.

Acknowledgments We thank Bea Senger and Stefan Kellenberger for excellent technical assistance with the field experiments. We thank Philip Strecken for the guidance on SNB field scorings. We also thank Bea Senger for the technical help with the plant material in the greenhouse. We thank Hana Simkova and Jaroslav Dolezel for isolation of chromosome 3B by flow cytometry. We also thank Matthias Helling for the help with the statistical analysis. The research leading to these results received funding from the European Community’s Seventh Framework Programme (FP7/2007-2013) under the grant agreement FP7-212019 (Triticeae Genome) and was supported in the framework of the European Cooperation in Science and Technology FA0604 (Tritigen). This work was also supported by a grant from the Swiss National Science Foundation 31003A_127061. The experiments in this study comply with the current laws of Switzerland.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Aguilar V, Stamp P, Winzeler M, Winzeler H, Schachermayr G, Keller B, Zanetti S, Messmer MM (2005) Inheritance of field resistance to *Stagonospora nodorum* leaf and glume blotch and correlations with other morphological traits in hexaploid wheat (*Triticum aestivum* L.). *Theor Appl Genet* 111:325–336
- Akhunov ED, Goodyear AW, Geng S, Qi L-L, Echalié B, Gill BS, Miftahudin, Gustafson JP, Lazo G, Chao S, Anderson OD, Linkiewicz AM, Dubcovsky J, Rota ML, Sorrells ME, Zhang D, Nguyen HT, Kalavacharla V, Hossain K, Kianian SF, Peng J, Lapitan NLV, Gonzalez-Hernandez JL, Anderson JA, Choi D-W, Close TJ, Dilbirligi M, Gill KS, Walker-Simmons MK, Steber C, McGuire PE, Qualset CO, Dvorak J (2003) The organization and rate of evolution of wheat genomes are correlated with recombination rates along chromosome arms. *Genome Res* 13:753–763
- Bennett MD, Smith JB (1976) Nuclear DNA amounts in angiosperms. *Philos Trans R Soc Lond B Biol Sci* 274:227–274
- Bostwick DE, Ohm HW, Shaner G (1993) Inheritance of *Septoria glume blotch* resistance in wheat. *Crop Sci* 33:439–443
- Broennimann A (1968) Untersuchungen über *Septoria nodorum* Berk. des Weizens. *Mitteil Schweiz Landw* 16:65–76
- Broennimann A (1975) Contributions to genetics of tolerance towards *Septoria nodorum* Berk. in wheat (*Triticum aestivum* L.). *J Plant Breed* 75:138–160
- Broennimann A, Fossati A, Hani F (1973) Spreading of *Septoria nodorum* Berk. and damage to artificially induced mutants of winter wheat cultivar Zenith (*Triticum aestivum* L.). *J Plant Breed* 70:230–245
- Bryan GJ, Stepherson P, Collins A, Kirby J, Smith JB, Gale MD (1999) Low level of DNA sequence variation among adapted genotypes of hexaploid wheat. *Theor Appl Genet* 99:192–198
- Choulet F, Wicker T, Rustenholz C, Paux E, Salse J, Leroy P, Schlub S, Paslier M-CL, Magdelenat G, Gonthier C, Couloux A, Budak H, Breen J, Pumphrey M, Liu S, Kong X, Jia J, Gut M, Brunel D, Anderson JA, Gill BS, Appels R, Keller B, Feuillet C (2010) Megabase level sequencing reveals contrasted organization and evolution patterns of the wheat gene and transposable element spaces. *Plant Cell* 22:1686–1701
- Crawley MJ (2007) *The R book*. Wiley, England
- Draeger R, Gosman N, Steed A, Chandler E, Thomsett M, Srinivasachary, Schondelmaier J, Buerstmayr H, Lemmens M, Schmolke M, Mesterhazy A, Nicholson P (2007) Identification of QTLs for resistance to Fusarium head blight, DON accumulation

- and associated traits in the winter wheat variety Arina. *Theor Appl Genet* 115:617–625
- Duczek LJ, Sutherland KA, Reed SL, Bailey KL, Lanford GP (1999) Survival of leaf spot pathogens on crop residues of wheat and barley in Saskatchewan. *Can J Plant Pathol* 21:165–173
- Ellis MH, Rebetzke GJ, Azanza F, Richards RA, Spielmeyer W (2005) Molecular mapping of gibberellin-responsive dwarfing genes in bread wheat. *Theor Appl Genet* 111:423–430
- Eriksen L, Borum F, Jahoor A (2003) Inheritance and localisation of resistance to *Mycosphaerella graminicola* causing septoria tritici blotch and plant height in the wheat (*Triticum aestivum* L.) genome with DNA markers. *Theor Appl Genet* 107:515–527
- Eyal Z, Scharen AL, Prescott JM and van Ginkel M (1987) The *Septoria* diseases of wheat: concepts and methods of disease management. CIMMYT, Mexico, DF
- FAO stat (2010) Food and Agriculture organization of the United Nations. Available at <http://faostat.fao.org>
- Faris JD, Haen KM, Gill BS (2000) Saturation mapping of a gene-rich recombination hot spot region in wheat. *Genetics* 154:823–835
- Fridman E, Liu YS, Carmel-Goren L, Gur A, Shoresh M, Pleban T, Eshed Y, Zamir D (2002) Two tightly linked QTLs modify tomato sugar content via different physiological pathways. *Mol Genet Genomics* 266:821–826
- Fried PM, Meister E (1987) Inheritance of leaf and head resistance of winter wheat to *Septoria nodorum* in a diallel cross. *Phytopathology* 77:1371–1375
- Gervais L, Dedryver F, Morlais J-Y, Bodusseau V, Negre S, Bilous M, Groos C, Trottet M (2003) Mapping of quantitative trait loci for field resistance to Fusarium head blight in an European winter wheat. *Theor Appl Genet* 106:961–970
- Halama P (2002) Mating relationships between isolates of *Phaeosphaeria nodorum*, (anamorph *Stagonospora nodorum*) from geographical locations. *Eur J Plant Pathol* 108:593–596
- Hallauer AR, Miranda Fo JB (1981) Quantitative genetics in maize breeding. The Iowa University Press, Ames
- Hollister JD, Gaut BS (2009) Epigenetic silencing of transposable elements: a trade-off between reduced transposition and deleterious effects on neighboring gene expression. *Genome Res* 19:1419–1428
- Jeger MJ, Viljanen-Rollinson SLH (2001) The use of the area under the disease progress curve (AUDPC) to assess quantitative disease resistance in crop cultivars. *Theor Appl Genet* 102:32–40
- Keurentjes JJB, Bentsink L, Alonso-Blanco C, Hanhart CJ, Vries HB-D, Effgen S, Vreugdenhil D, Koornneef M (2007) Development of a near-isogenic line population of *Arabidopsis thaliana* and comparison of mapping power with a recombinant inbred line population. *Genetics* 175:891–905
- Kroymann J, Mitchell-Olds T (2005) Epistasis and balanced polymorphism influencing complex trait variation. *Nature* 435:95–98
- Laubscher FX, von Wechmar B, van Schalkwyk D (1966) Heritable resistance of wheat varieties to Glume Blotch (*Septoria nodorum* Berk.). *J Phytopathol* 56:260–264
- Mesterhazy A (1995) Types and components of resistance to Fusarium head blight of wheat. *Plant Breed* 114:377–386
- Monna L, Lin X, Kojima S, Sasaki T, Yano M (2002) Genetic dissection of a genomic region for a quantitative trait locus, Hd3, into two loci, Hd3a and Hd3b, controlling heading date in rice. *Theor Appl Genet* 104:772–778
- Nelson LR, Gates CE (1982) Genetics of host plant resistance of wheat to *Septoria nodorum*. *Crop Sci* 22:771–773
- Oliver RP, Friesen TL, Faris JD, Solomon PS (2012) *Stagonospora nodorum*: from pathology to genomics and host resistance. *Annu Rev Phytopathol* 50:23–43
- Paillard S, Schnurbusch T, Winzeler M, Messmer M, Sourdille P, Abderhalden O, Keller B, Schachermayr G (2003) An integrative genetic linkage map of winter wheat (*Triticum aestivum* L.). *Theor Appl Genet* 107:1235–1242
- Paux E, Sourdille P, Salse J, Saintenac C, Choulet F, Leroy P, Korol A, Michalak M, Kianian S, Spielmeyer W, Lagudah E, Somers D, Kilian A, Alaux M, Vautrin S, Bergès H, Eversole K, Appels R, Safar J, Simkova H, Doležel J, Bernard M, Feuillet C (2008) A physical map of the 1-gigabase bread wheat chromosome 3B. *Science* 322:101–104
- Paux E, Faure S, Choulet F, Roger D, Gauthier V, Martinant J-P, Sourdille P, Balfourier F, Paslier M-CL, Chauveau A, Cakir M, Gandon B, Feuillet C (2010) Insertion site-based polymorphism markers open new perspectives for genome saturation and marker-assisted selection in wheat. *Plant Biotechnol J* 8:196–210
- Pea G, Paulstephenraj P, Canè MA, Sardaro MLS, Landi P, Morgante M, Porceddu E, Pè ME, Frascaroli E (2009) Recombinant near-isogenic lines: a resource for the mendelization of heterotic QTL in maize. *Mol Genet Genomics* 281:447–457
- Rustenholtz C, Choulet F, Laugier C, Šafař J, Šimková H, Doležel J, Magni F, Scalabrin S, Cattonaro F, Vautrin S, Bellec A, Bergès H, Feuillet C, Paux E (2011) A 3,000-loci transcription map of chromosome 3B unravels the structural and functional features of gene islands in hexaploid wheat. *Plant Physiol* 157:1596–1608
- R Development Core Team (2013) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>. ISBN 3-900051-07-0
- Schnurbusch T, Paillard S, Fossati D, Messmer M, Schachermayr G, Winzeler M, Keller B (2003) Detection of QTLs for *Stagonospora glume blotch* resistance in Swiss winter wheat. *Theor Appl Genet* 107:1226–1234
- Scott PR, Benedikz PW, Cox CJ (1982) A genetic study of the relationship between height, time of ear emergence and resistance to *Septoria nodorum* in wheat. *Plant Pathol* 31:45–60
- Scott PR, Benedikz PW, Zones HG, Ford MA (1985) Some effects of canopy structure and microclimate on infection of tall and short wheats by *Septoria nodorum*. *Plant Pathol* 34:587–593
- Shah DA, Bergstrom GC (2002) A rainfall-based model for predicting the regional incidence of wheat seed infection by *Stagonospora nodorum* in New York. *Phytopathology* 92:511–518
- Shatalina M, Wicker T, Buchmann JP, Oberhaensli S, Simková H, Doležel J, Keller B (2013) Genotype-specific SNP map based on whole chromosome 3B sequence information from wheat cultivars Arina and Forno. *Plant Biotechnol J* 11:23–32
- Shaw MW, Beachell SJ, Fitt BDL, Fraaije BA (2008) Long-term relationships between environment and abundance in wheat of *Phaeosphaeria nodorum* and *Mycosphaerella graminicola*. *New Phytol* 177:229–238
- Srinivasachary Gosman N, Steed A, Hollins TW, Bayles R, Jennings P, Nicholson P (2009) Semi-dwarfing Rht-B1 and Rht-D1 loci of wheat differ significantly in their influence on resistance to Fusarium head blight. *Theor Appl Genet* 118:695–702
- The International Rice Genome Sequencing Project (2005) The map-based sequence of the rice genome. *Nature* 436:793–800
- Thomson MJ, Edwards JD, Septiningsih EM, Harrington SE, McCouch SR (2006) Substitution mapping of dth1.1, a flowering-time quantitative trait locus (QTL) associated with transgressive variation in rice, reveals multiple sub-QTL. *Genetics* 172:2501–2514
- Tommasini L, Schnurbusch T, Fossati D, Mascher F, Keller B (2007) Association mapping of *Stagonospora nodorum* blotch resistance in modern European winter wheat varieties. *Theor Appl Genet* 115:697–708
- Van Ginkel M, Rajaram S (1999) Breeding for resistance to the *Septoria/Stagonospora* blights of wheat. In: van Ginkel M, McNab A, Krupinsky JM (eds) *Septoria and Stagonospora*

- diseases of cereals: a compilation of global research. CIMMYT, Mexico
- Wanjugi H, Coleman-Derr D, Huo N, Kianian SF, Luo M-C, Wu J, Anderson O, Gu YQ (2009) Rapid development of PCR-based genome-specific repetitive DNA junction markers in wheat. *Genome* 52:576–587
- Wicki W, Winzeler M, Schmid JE, Stamp P, Messmer M (1999) Inheritance of resistance to leaf and glume blotch caused by *Septoria nodorum* Berk. in winter wheat. *Theor Appl Genet* 99:1265–1272
- Wolf EDD, Francl LJ (2000) Neural network classification of tan spot and *Stagonospora* blotch infection periods in a wheat field environment. *Phytopathology* 90:108–113