Microclimate affects soil chemical and mineralogical properties of cold alpine soils of the Altai Mountains (Russia)

Egli, Markus ; Lessovaia, Sofia N ; Chistyakov, Kirill ; Inozemzev, Svyatoslav ; Polekhovsky, Yurii ; Ganyushkin, Dmitry

Abstract: Purpose: The present work focuses on cold alpine soils of the Altai Mountains (Siberia, Russia). Permafrost is widespread and often occurs at a depth of about 100 cm. The area is characterised by extremely cold winters and cool summers: the aim was consequently to find out whether weathering could be more intense on thermally less unfavoured conditions or whether the abundance of water could be a more important factor. Materials and methods: We investigated 10 soils in a very small area close to a local glacier tongue. Five of the investigated soils were south-facing and the other five north-facing. The soils have the same parent material (mica-rich till), altitude, topography and soil age. The vegetation is alpine grassland that is partially intersected with some juniper and mosses. Soil chemical properties such as organic C, N, soil organic matter quality (using DRIFT), pH value, (oxy)hydroxides, total elemental contents (XRF) and soil micromorphology and mineralogy (using diagnostic treatments and XRD) were determined. The age constraint of the site was given by geomorphic studies together with ¹⁴C dating of a nearby peat bog and the stable organic matter fraction of the soils. Results and discussion: The soils have a Holocene age. The results showed astonishingly clearly—similarly to the European Alps—that the north-facing soils have a higher weathering state. This is expressed by lower pH values, higher oxalate and dithionite extractable Fe, Al, Mn and Si contents, higher C and N concentrations and stocks when compared to the south-facing sites. No statistically significant differences with respect to weathering indexes could be detected. The geochemical evolution of the soils seems to be enhanced at north-facing sites, even though very severe climatic conditions prevail. Furthermore, biodegradation seems to be less pronounced on north-facing compared to south-facing sites as poorly degraded organic matter is accumulated. This gives rise to more organic ligands that promote metal binding and their subsequent eluviation along the soil profile. Conclusions: We consequently must assume that weathering is not limited by low temperatures in the active layer but is rather controlled by soil moisture that seems to be higher during the warmer period in the north-facing soils.

DOI: <https://doi.org/10.1007/s11368-013-0838-4>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-102808>

Journal Article

Accepted Version

Originally published at:

Egli, Markus; Lessovaia, Sofia N; Chistyakov, Kirill; Inozemzev, Svyatoslav; Polekhovsky, Yurii; Ganyushkin, Dmitry (2015). Microclimate affects soil chemical and mineralogical properties of cold alpine soils of the Altai Mountains (Russia). *Journal of Soils and Sediments*, 15(6):1420-1436.

DOI: <https://doi.org/10.1007/s11368-013-0838-4>

1 SOIL FORMATION AND WEATHERING IN TIME AND SPACE

2
3 **Microclimate affects soil chemical and mineralogical properties of cold-alpine soils of**
4 **the Altai Mountains (Russia)**

5
6
7
8
9
10 **Markus Egli • Sofia N. Lessovaia • Kirill Chistyakov • Svyatoslav Inozemzev • Yuri**
11 **Polekhovsky • Dmitry Ganyushkin**

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

9 M. Egli (✉)
10 Department of Geography, University of Zurich, Winterthurerstrasse 190, 8057 Zurich,
11 Switzerland
12 e-mail: markus.egli@geo.uzh.ch

14 S. N. Lessovaia • K. Chistyakov • S. Inozemzev • D. Ganyushkin
15 St. Petersburg State University, Faculty of Geography and Geoecology, V.O. 10 line d 33,
16 199178 St. Petersburg, Russia, lessovaia@yahoo.com

18 Y. Polekhovsky
19 St. Petersburg State University, Geological Faculty, Universitetskaya nab., 7/9, St.
20 Petersburg, 199034, Russia

21
22 (✉) **Corresponding author:**
23 Markus Egli
24 e-mail: markus.egli@geo.uzh.ch

27 **Abstract**

1
2 28 *Purpose* The present work focuses on cold-alpine soils of the Altai Mountains (Siberia,
3
4 29 Russia). Permafrost is widespread and often occurs at a depth of about 100 cm. The area is
5
6 30 characterised by extremely cold winters and cool summers: the aim was consequently to find
7
8 31 out whether weathering could be more intense on thermally less unfavoured conditions or
9
10 32 whether the abundance of water could be a more important factor.

11
12 33 *Materials and methods* We investigated ten soils in a very small area close to a local glacier
13
14 34 tongue. Five of the investigated soils were south-facing and the other five north-facing. The
15
16 35 soils have the same parent material (mica-rich till), altitude, topography and soil age. The
17
18 36 vegetation is alpine grassland that is partially intersected with some juniper and mosses. Soil
19
20 37 chemical properties such as organic C, N, soil organic matter quality (using DRIFT), pH
21
22 38 value, (oxy)hydroxides, total elemental contents (XRF) and soil micromorphology and
23
24 39 mineralogy (using diagnostic treatments and XRD) were determined. The age constraint of
25
26 40 the site was given by geomorphic studies together with ^{14}C dating of a nearby peat bog and
27
28 41 the stable organic matter fraction of the soils.

29
30 42 *Results and discussion* The soils have a Holocene age. The results showed astonishingly
31
32 43 clearly – similarly to the European Alps – that the north-facing soils have a higher weathering
33
34 44 state. This is expressed by lower pH-values, higher oxalate and dithionite extractable Fe, Al,
35
36 45 Mn and Si contents, higher C and N concentrations and stocks when compared to the south-
37
38 46 facing sites. No statistically significant differences with respect to weathering indexes could
39
40 47 be detected.

41
42 48 The geochemical evolution of the soils seems to be enhanced at north-facing sites, even
43
44 49 though very severe climatic conditions prevail. Furthermore, biodegradation seems to be less
45
46 50 pronounced on north-facing compared to south-facing sites as poorly-degraded organic
47
48 51 matter is accumulated. This gives rise to more organic ligands that promote metal-binding
49
50 52 and their subsequent eluviation along the soil profile.

51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Conclusions We consequently must assume that weathering is not limited by low temperatures in the active layer but is rather controlled by soil moisture that seems to be higher during the warmer period in the north-facing soils.

Keywords Clay minerals • Organic carbon • Oxyhydroxides • Permafrost • Water availability • Weathering

1 Introduction

Precipitation and temperature particularly influence soil properties by affecting the type and rates of chemical, biological, and physical processes (Dahlgren et al. 1997; Birkeland 1999). Vegetation growth and decomposition, which depend on temperature and the other environmental factors, influence weathering reactions through the production of acidity and organic ligands that may promote chemical weathering and subsequent elemental leaching. Weathering rates, in general, are not only driven by climate as known from the paradigm of Dokuchaev (1883) and, in an extended form, of Jenny (1941). Chemical weathering intensities depend on: (i) the lithology (e.g. highly reactive minerals such as carbonates and sulphates vs. crystalline rocks; see e.g. Simas et al., 2006), (ii) biology (Gorbushina 2007) and the development of organic matter (Conen et al. 2007), (iii) the topography and - related to it - the rate of supply of fresh regolith through physical weathering and erosion (Dixon et al. 2012; Dixon and von Blanckenburg 2012), (iv) the age of exposure and (v) the character of the hydrological system.

Several studies have shown the influence of slope aspect and the resulting microclimate on soil weathering and development (Klemmedson 1964; Macyk et al. 1978; Carter and Ciolkosz 1991; Rech et al. 2001; Egli et al. 2007). Slope aspect is directly related to the incoming solar energy, particularly relevant for permafrost-affected soils of continental climate (Osawa et al. 2010). There is no doubt that temperature (energy) determines mineral and soil weathering – even in the long-term over thousands of years as shown by Williams et

al. (2010). Nevertheless, there is no unanimous agreement whether weathering in cold regions is driven more by the temperature or the availability of water (White and Blum 1995; White et al. 1999; Egli et al. 2006; 2011; Dixon et al. 2009; Williams et al. 2010; Rasmussen et al. 2011; etc.). As shown by previous investigations in the European Alps (Egli et al. 2008), higher temperatures do not necessarily lead to higher weathering rates in cold alpine regions. Water fluxes through the soils seem to be at least equally important. In general terms, the explanation of weathering in cold regions often identified three basic tenets (Hall et al. 2002): (i) weathering is dominated by mechanical processes, (ii) the predominant mechanical process is freeze-thaw and (iii) due to the low temperatures, chemical weathering is not a significant element of cold region processes. Hall et al. (2002) showed that even at low temperature weathering rates can be high.

The scope of this paper was thus to test if the hypothesis that moisture availability predominantly determines weathering can also be applied to a very cold region having extreme climatic conditions. In order to accomplish this, the soils must have (almost) identical conditions (parent material, age, climate in general, vegetation and topography).

2 Study site

To tackle the above-mentioned scope and to test the hypothesis, the sites had to meet the criteria of having a cold climate (in a permafrost region), of showing within short distances north- and south-exposure (leading to a difference in solar energy) and of not being too old (thus avoiding the risk that too many warmer and colder periods could have disturbed the system). Weathering of primary rocks is faster on relatively young surfaces (Mavris et al. 2010). Consequently, differences between north and south-facing sites should be better assignable to thermal differences. The investigated field sites (2380 m asl) are located in the south-eastern part of the Altai Mountains (border region: Russia/Mongolia/China) at 49°49.159' N, 87°50.912' E and are found in an extended proglacial area of the Sophiysky glacier of the South Chuya ridge (Fig. 1). The tendency of glacier degradation since the end of the 19th century is to be at an average rate of 15 – 20 m per year (Ostanin et al. 2004).

The Sophiysky Glacier has retreated 18 m per year over the last 100 years; the altitude of its tongue was 2485 m asl in 2000 (Agatova et.al. 2002). In the Altai Mountains, meteorological stations are mostly found at altitudes < 1000 m asl. Only two are at > 2000 m asl (located in highland) and four are at an altitude between 1000 to 2000 m asl. The annual variation of temperature is in general strongly pronounced. In the highlands, the absolute minimum temperature is around -50 to -56 °C (Marinina and Samoilova 1987). The nearest meteorological station is Kosh-Agach (1830 m asl) that is situated about 55 km east of the investigation site where a minimum air temperature of -62 °C was recorded in 1969 (Narozhniy and Zemtsov 2011). Average soil surface temperature in January is there -30 °C. As an average over the last few decades, a snow cover was measured there for 131 days per year (Osokina et al. 1993). Average snow cover is with 10 cm quite low (which enhances the production of permafrost). According to the geobotanical zonation (Ogureeva 1980), the studied area belongs to the alpine-tundra altitudinal belt (at a height of 2400 – 3200 m asl) of the Chuya – Argutskey region of the Central-Altai sub-province. Typical for this region are meadows and mountainous tundra. Average mean summer temperature varies between +9 °C and +2.0 °C at the lower part (see also Narozhniy and Zemtsov 2011) of the zonal belt to about +5 °C to 1.1 °C in the upper part. Mean annual precipitation is 550 – 1000 mm (Syromyatina 2010). The studied area is located at the boundary between semi-humid and semi-arid mountainous climates. The soil temperature regime is gelic (IUSS Working Group WRB, 2007).

The studied area is located on a ledge – a small ridge (2 – 3 ha) that crosses the Akkol river. The ridge is covered by moraine sediments. The distance from the glacial tongue is ~ 4km. Mostly gneiss-like stones are found close to the edge of the glacier while, on the bank of the river opposite to the ridge, rocks having an alkaline composition are also found.

3 Material and methods

3.1 Sampling strategy

The area of interest was small and covered about 2 – 3 ha. Soil material was collected from

excavated profile pits in undisturbed locations. All sites were fully covered with vegetation. The vegetation is alpine grassland that is partially intersected with some juniper and mosses. Five soils on the north-facing and 5 soils on the south-facing slope were sampled at a depth of 0 – 15 and 15 – 30 cm. In addition, a soil profile including also the BC horizon was dug. They showed the following horizon differentiation: weakly decomposed litter (0 – 5 cm), a humus horizon A (5 – 19 cm), an ABw (19 – 27 cm) horizon that was characterised by a gradual decrease of colour intensity and a BC horizon (27 – 68 cm). Due to the presence of permafrost (often at a depth of about 0.5 to 1.5 m) and some cryogenic features (frost heave), the soils can all be classified as Cambic Cryosols (Table 1). They exhibited a sandy-silt texture (Table 2). Furthermore, some rock fragments in the soil profile and a hard rock (gneiss) in the bank of the Akkol river (opposite to the ridge) were sampled to determine the possible mineral source for soils. One to two kg soil material per sample was taken for the analyses (Hitz et al. 2002). The bulk density (fine earth and soil skeleton) was measured on undisturbed samples. Furthermore, 2 samples from a peat bog were taken and dated in order to have an additional time constraint of the surface and soils. The samples were taken at the transition to the permafrost underground (approx. 1 m depth) and at a depth of 60 cm. This peat bog was located approx. 2 – 3 km further down the valley. A further age indication of the surface was obtained from 2 soil samples of the investigated site by dating the stable organic matter fraction (resistant to an H₂O₂ treatment; see Favilli et al. 2008).

3.2 Soil chemical and physical analyses

Total C and N contents of the soil were measured with a C/H/N analyser (Elementar Vario EL). Soil pH (in 0.01 M CaCl₂) was determined on air-dried fine earth samples using a soil:solution ratio of 1:2.5. As the soils did not contain inorganic carbon, the total C content equals the organic carbon content. LOI (loss-on-ignition) was determined on 2.5g soil material ignited at 950°C during 3h.

After a pre-treatment of the samples with H₂O₂ (3 %), particle size distribution of the soils was measured using a combined method consisting of sieving the coarser particles (2000 - 32 µm) and the measurement of the finer particles (< 32 µm) by means of an X-ray sedimentometer (SediGraph 5100).

Oven-dried (70 °C) samples were sieved to < 2 mm. Fe, Al, Mn and Si concentrations were determined after treatment with NH₄-oxalate (buffered at pH 3, labelled 'o') (McKeague et al. 1971). The extracts were centrifuged for 8 minutes at 4000 rpm and filtered (mesh size 0.45 µm, S&S, filter type 030/20). Element concentrations were measured using atomic absorption spectroscopy (AAnalyst 700, Perkin Elmer). The oxalate treatment extracts both the weakly- and poorly crystalline phases and some of the organic phases, but normally does not dissolve the strong humus-metal complexes (Mizota and van Reeuwijk 1989). Additionally, the dithionite- (Mehra and Jackson 1958) was measured for the elements Fe and Al.

Measurement of the total element content of fine earth and skeleton was done by means of X-ray fluorescence (XRF). Approximately 10 g of soil material were milled to < 50 µm in a tungsten carbide disc swing mill (Retsch® RS1, Germany). 4 g of soil powder was mixed with 0.9 g of Licowax® C Micro-Powder PM (Clariant, Switzerland), pressed into a 32 mm-pellet and analysed using an energy dispersive X-ray fluorescence spectrometer (SPECTRO X-LAB 2000, SPECTRO Analytical Instruments, Germany).

3.3 Weathering indexes

Several indexes have been defined to characterise chemical weathering in soils. The general principle of all these indexes is similar and based on the ratio of the base cations (Ca, Mg, K, Na) to Al and/or Si. We used the 'index A' and 'index B' of Kronberg and Nesbitt (1981). These indexes are defined by the molar ratio of:

$$A = \frac{SiO_2 + CaO + K_2O + Na_2O}{Al_2O_3 + SiO_2 + CaO + K_2O + Na_2O} \quad ; \quad B = \frac{CaO + K_2O + Na_2O}{Al_2O_3 + CaO + K_2O + Na_2O} \quad (1)$$

In addition, also the CIA and CIW-Indexes were tested (Harnois 1988; Nesbitt and Young 1989)

$$CIA = \frac{Al_2O_3}{Al_2O_3 + CaO + Na_2O + K_2O} \quad ; \quad CIW = \frac{Al_2O_3}{Al_2O_3 + CaO + Na_2O} \quad (2)$$

The molar ratio of (K+Ca)/Ti is used as a 'dating' method for rock varnish in semi-arid to arid regions (Harrington and Whitney 1987). As Ti is considered to be an immobile element, the ratio can be used as a weathering index: the lower the ratio, the higher the degree of weathering.

3.4 Stock calculations of C, N and oxyhydroxides

The oxalate-extractable Fe_o, Al_o, Si_o, Mn_o and C and N stocks were calculated according to the following equation:

$$E_{stock} = \sum_{i=1}^n E_i \Delta z_i \rho_i (1 - RM) \quad (3)$$

where E_{stock} denotes the elemental abundance (kg/m²), E the elemental concentration (kg/t), Δz_i the thickness of layer i (m), ρ = soil density (t/m³) and RM the mass proportion of rock fragments.

3.5 Micromorphology and X-ray diffractometry

The rock samples from the soil horizons were studied in thin sections by optical microscopy using Zeiss Axioplan 2 and Polam P-312 microscopes. The micromorphological study of soil horizons was performed using thin sections and an Olympus optical microscope. The fine earth (< 2 mm fraction), the rock fragments and the gneiss were analysed as random samples using X-ray diffraction (MiniFlex ii (Rigaku), CoK α radiation).

After dispersion of the sample by adding some drops of ammonia as peptising agent, the < 1 μ m fraction was separated from the soil material by sedimentation and decantation. Some drops of ammonia as peptising agent were added for dispersion of the soil sample. The

mineralogical composition of the < 1 µm fraction was studied on oriented specimen using X-ray diffractometry (XZG-4A Carl Zeiss Jena) with CuKα radiation and a monochromator in the diffracted beam. Prior to XRD analysis, the following diagnostic treatments were performed: (i) saturation with Mg, (ii) ethylene glycol solvation and (iii) heating at 550°C for 3 h. Digitised X-ray data were corrected for Lorentz and polarisation factors (Moore and Reynolds 1997). To better detect single minerals and mixed-layered phases, diffraction patterns were fitted by the Origin™ PFM program using the Pearson VII algorithm. Background values were calculated by means of a non-linear function (polynomial 2nd order function; Lanson, 1997).

3.6 DRIFT measurements

The samples (< 2 mm fraction) used for DRIFT (Diffuse Reflectance Infrared Fourier Transform Spectroscopy) analyses were homogenised and then dried at 70 °C for 2 hours (prior to analysis). Relative peak intensities were used for DRIFT analysis (Bruker, Tensor 27). Spectra were recorded from 4000 to 250 cm⁻¹. DRIFT measurements were used to check mineralogical aspects and to estimate soil organic matter composition (functional groups).

In order to quantify the relative changes of organic-functional groups in the FT-IR spectra, we divided the values of the relative intensity (area) of each peak by the sum of the relative intensity of all the considered peaks and multiplied it by 100 using the software OPUS 6.5. An integration method was employed to calculate the relative concentration (OPUS 6.5) that used a linear background between the found bases (individual samples) and absorbance values. Major IR absorption bands and functional groups assignments are given in Table 3. Aliphatic compounds were calculated using the IR range 1470-1440 cm⁻¹.

3.7 Radiocarbon dating of organic matter fractions

The turf samples were cleaned using an acid-alkali-acid (AAA) treatment. The soil samples to be dated underwent a H₂O₂ treatment over 1 week (Favilli et al. 2008). The samples were

then heated under vacuum in quartz tubes with CuO (oxygen source) to remove any absorbed CO₂ in the CuO. The tubes were then evacuated, sealed and heated in the oven at 900 °C to obtain CO₂. The CO₂ of the combusted sample was mixed with H₂ (1:2.5) and catalytically reduced over iron powder at 535°C to elemental carbon (graphite). After reduction, the mixture was pressed into a target and carbon ratios were measured by Accelerator Mass Spectrometry (AMS) using the tandem accelerator of the Laboratory of Ion Beam Physics at the Swiss Federal Institute of Technology Zurich (ETHZ). The calendar ages were obtained using the OxCal 4.2 calibration program (Bronk Ramsey 2001; 2009) based on the IntCal 09 calibration curve (Reimer et al. 2009). Calibrated ages are given in the 2 σ range (minimum and maximum value for each).

3.8 Statistics

The individual datasets were checked for normal distribution using a Shapiro-Wilk test (SigmaPlot 11.0 (Systat Software Inc.); Jann 2005). As the data did not always show a normal distribution, differences in mean values were tested using the U-test (Mann–Whitney). For datasets having a normal distribution, the t-test was applied. This procedure was checked using a two-tailed test for significance.

4 Results

4.1 Petrographic characteristics and soil micromorphology

Included rock fragments in the soil showed a polymineral and acid composition. The rock fragments were a modified biotite-plagioclase gneiss having a sub-parallel arrangement of minerals (Fig. 2a). The semi-quantitative composition was as follows: quartz 40 – 50 %; 15 – 25 % plagioclase; 8 – 10 % biotite; 5 – 7 % chlorite; 2 – 3 % epidote; 1 – 2 % ore mineral and leucoxene and single grains of amphiboles and tourmaline. The surface of these rocks was crossed by numerous veins having a quartz, quartz-albite and chalcedony composition (Fig. 2b). The veins reflected several generations of their formation; the last of them were chalcedony veins. Biotite was a predominant mineral among the micas. Muscovite or illite

(so-called 'white' mica) were also found but they were much less frequent. Sometimes the replacement of biotite by chlorite was identified (Fig. 2c). The studied rock initially seems to be a sedimentary rock that was later metamorphosised.

The thin sections of the soil material from the upper part of the A horizon (3 – 8 cm) showed samples having a loose microstructure, coarse to fine silty plasmic fabrics and a fragmentary organic-ferric plasma that is distributed like flocculus and clots (Fig. 3). The 'coarse' fraction was dominated by silt grains with a predominance of quartz and feldspars. In addition, biotite and hornblende were also present. Biotite particles did not show typical alteration features. However, some ferruginous flocculent films have been identified on their surface. Raw and weakly to moderately decomposed organic matter was detected. The soil material was generally slightly transformed by pedogenetic processes and weakly aggregated. The lower part of the A horizon (10 – 15 cm) is weakly aggregated, mainly by a silt-plasma fabric having features of zoogenic transformation. The coarse particles were mostly composed of slightly-rounded quartz. Furthermore, biotite and amphiboles were found. Pedogenetic features were evident. The ABw horizon (sample from 26 – 30 cm depth) was weakly aggregated, showing a silt-plasma fabric. In general, the coarse particles showed a composition similar to that in the overlying horizon.

Similar to the other soil horizons, the BC material was weakly aggregated, having mainly a silt- plasmic fabric that was slightly transformed by pedogenetic processes. The coarse fraction was slightly rounded and composed predominantly of quartz. Compared to the other horizons, an increasing proportion of biotite was detected.

4.2 General physical and chemical soil properties

The investigated soils are acidic to moderately acidic. The pH ranged from 4.4 – 5.4 in the top layer to 6.1 in the subsoil. No carbonates were present in the soils. The grain size in all soils and horizons was similar and consisted of a sandy loam, silt loam or loam (Table 2). Typically, a maximum grain size near to 120 µm was measured (data not shown). A second but much minor maximum was detectable around a grain size of 15 – 25 µm. Bulk density

was quite uniform and varied in between 0.9 to 1.1 g/cm³. At the north-facing sites, the C concentrations varied between 19 and 50 gC/kg and at the south-facing sites between 12 and 30 gC/kg. Similarly, the N contents were in general higher at north-facing sites (1.1 – 3.1 gN/kg) than at the south-facing sites (1 – 2.3 gN/kg). The differences between north- and south-facing sites with respect to the C and N concentrations and also the corresponding stocks (0 – 15cm, 15 – 30 cm and 0 – 30 cm; except N stocks)) were statistically significant ($p < 0.05$; Figs 4 and 5). At north-facing sites, there was a distinct accumulation of soil organic matter. The C/N ratio of the soil organic matter varied in the range of 12.9 – 17.9 at north-facing and 11.1 – 13.7 at south-facing sites (Table 2, Fig. 4). Also here, distinct differences ($p < 0.05$) between north- and south-facing sites could be measured – north-facing sites having higher C/N values.

The north-facing sites furthermore showed significantly lower pH-values ($p < 0.05$; Fig. 4).

4.3 Soil organic matter characteristics

The DRIFT spectra of the bulk SOM delivered information about its chemical characteristics. Independent of the exposure, the bulk organic matter consisted mostly of aliphatic compounds (data not shown), secondary alcohols, phenolic functional groups and aromatic rings and amides and to a lesser extent of polysaccharides and other compounds. Also regarding the chemical composition of the soil organic matter, partially distinct differences between north- and south-facing sites could be detected. North-facing sites showed a distinctly ($p < 0.05$) higher relative (and absolute) proportion of carboxyl-groups (-COOH) and less secondary alcohols, but a significantly higher amount of polysaccharides (Fig. 6). Humification processes were related to the preferential oxidation of plant polysaccharides and phenolic hydroxyl groups and were further characterised by an enrichment of secondary alcohols and N-containing compounds in the SOM. The variations between the C and N contents, the C/N ratio and also the functional groups in the north- and south-facing sites would seem to indicate that there were quite distinct differences in the decomposition processes.

329

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

4.4 Pedogenetic oxyhydroxides

The oxalate- and dithionite-extractable contents are given in Table 4. The concentrations of all compounds were relatively low. The weathering stage of the soils is therefore not very advanced. In most cases the Al_o content was equal to the Al_d content. Sometimes the Al_o concentration exceeded Al_d which could be an indication of the presence of ITM (imogolite type materials). The possible presence of ITM was calculated using the molar ratio Al_o/Si_o and counter-checked using DRIFT. The molar ratio should be close to 1 for allophanes and allophane-like minerals ($2SiO_2 \cdot Al_2O_3 \cdot 2.5H_2O$; Lorenzoni et al. 1995) and close to 2 for imogolite-like minerals ($SiO_2 \cdot Al_2O_3 \cdot 2.5H_2O$). The likelihood of ITM is very low if the molar ratio is < 0.75 or > 2.4 . However, neither the DRIFT spectra nor the molar ratios (that were in the range of 3.7 – 20.5) gave an indication of the presence of ITM.

Similar to the pH and SOM, also here quite distinct differences, in most cases significant (except Fe_d and N), between north- and south-facing sites could be measured (Fig. 7). The concentrations of Fe_o , Fe_d , Al_o , Al_d , Mn_o and also the corresponding stocks were significantly (except Fe_d) higher at the north-facing sites (Table 4, Figs 5 and 7) and indicated a more intense weathering and a higher release of metals (Zanelli et al. 2007). The Fe_o/Fe_d ratio was also higher at north-facing sites showing the formation of iron oxides with a lower degree of crystallinity; this reflects the effect of the higher content of organics that impede the crystallization of iron oxides (Cornell and Schwertmann 2003).

4.5 Mineralogy of the fine earth and clay fraction

Using the DRIFT spectra, an approximate and general mineralogical characterisation of all soil samples (fraction < 2 mm) was made possible. Various regions in the IR spectrum contained OH vibrational information. In the OH-bending and M-O region ($1200 - 300\text{ cm}^{-1}$), the bands associated with hydroxyl groups could be discriminated from each other, and band assignment is straightforward. Furthermore, this region is not affected by the presence of

residual water molecules (Vantelon et al. 2001). Additional indications, in some cases straightforward, can be obtained from the OH-stretching region (4000 – 3000 cm⁻¹). In all samples, primary minerals such as quartz and mica and to a lesser extent muscovite could be detected. Quartz was identified based on 1164 cm⁻¹ and 1080 cm⁻¹ Si-O asymmetrical stretching vibrations, the typical doublet at 800 cm⁻¹ and 780 cm⁻¹ (Si-O symmetrical stretching vibrations), the 700 cm⁻¹ Si-O symmetrical bending vibrations and the 514 cm⁻¹ and 465 cm⁻¹ Si-O asymmetrical bending vibration (Hlavay et al., 1978). The vibration at 3624 cm⁻¹ indicated the presence of dioctahedral silicate (Madejova and Komadel, 2005), probably dioctahedral mica as suggested by the micro-morphology analyses (Fig. 8). In many samples, some kaolinite was discernible (based on weak absorptions at 3694 cm⁻¹ using DRIFT). As a weathering product, some gibbsite seemed also to be present. In most samples, some chlorite was also detected (although the chloritic phases may comprise weathering products such as HIV). Often a peak near 670 cm⁻¹ was detectable: this points either to the 2:1 layer in talc (Schroeder 2002; Klopprogge 2004) or epidote (Langer and Raith 1974). The chloritic components and epidote (or the probable presence of talc) point to a small admixture of basic material. The material that derives from the adjacent valley slopes was interspersed (glacial abrasion) during glacial advance with the gneissic glacial material that derived from the upper part of the valley.

The XRD patterns (Fig. 8) of the fine earth (the < 2 mm fraction) revealed the following minerals (for all soil horizons): (i) chlorite (peaks with *d* values of 1.42, 0.71, 0.47 (very minor peak, not indicated in the XRD pattern), 0.35 nm, etc), (ii) mineral(s) of the mica group (1.00; 0.50). The peak at 0.389 nm denotes muscovite (Brindley and Brown, 1980); (iii) feldspars (0.64, 0.404, 0.377 nm etc.) and (iv) quartz (0.425, 0.344, 0.229, 0.224 nm, etc.). The peak with a *d* value of 0.85 nm could be attributed to amphibole.

When comparing the XRD peak intensity, a relative decrease of chlorite and mica towards the surface could be detected. Mica seems to show the lowest concentration in the ABw horizon. Chlorite, mica, feldspar, quartz and traces of amphibole were also found in the gneiss (rock sample of the river bank) and the rock fragments in the soil. Some variability

could be measured regarding the mica and chlorite proportion (Fig. 8).

In all samples, chlorite was iron rich because of the ratio of odd to even peaks: the $d(002)$ 0.71 nm and $d(004)$ 0.354 nm peaks are considerably higher than the $d(001)$ 1.42 nm and $d(003)$ 0.472 nm peaks) (Moore and Reynolds, 1997). The contribution of kaolinite to the 0.71 nm peak was low, as evidenced by its weak absorption at 3694 cm^{-1} in the DRIFT spectra.

In the fine-size fraction ($< 1\mu\text{m}$) chlorite and biotite were identified by XRD, as were also the products of biotite transformation represented by mixed-layered biotite-vermiculite having a minor proportion of vermiculitic layers. Chlorite was identified based on the d_{001} and d_{003} reflections at 1.40 nm and 0.471 – 0.473 nm, respectively. The d_{001} reflection at 1.40 nm, in fact, remained stable after ethylene glycol solvation and showed a slight contraction to 1.38 nm after heating at $550\text{ }^{\circ}\text{C}$. Mica group minerals were detected for their d_{001} and d_{002} basal reflections at 1.00 nm, which remained stable after the ethylene glycol and heating treatments. The low intensity of the peak at 0.50 nm, in the fine earth random samples, indicated a trioctahedral nature of this mineral. Together with the results of the thin sections, mica could be classified as biotite. The weathering product of biotite was a mixed-layered biotite-vermiculite having a minor proportion of vermiculite. The mixed-layered clay minerals were recognised by the asymmetries towards lower angles of the 1.00 nm peak of the XRD patterns of the Mg-saturated samples. No shift of these peaks occurred after ethylene glycol solvation (consequently, smectitic weathering products were not present). In the clay-size fraction $\leq 1\text{ }\mu\text{m}$, biotite and chlorite were found in all horizons. Compared to the other horizons, the proportion of mixed-layered biotite-vermiculite seemed to be highest in the ABw horizon (Fig. 8).

4.6 Total elemental contents and weathering indexes

The total elemental contents reflect the gneissic/granitic character of the soil material with SiO_2 contents between 65 and 69 % (Table 5). The concentration of base cations (Ca, Mg, K and Na) is relatively high.

The molar ratio of (K+Ca)/Ti is used as a dating method for rock varnish in semi-arid to arid regions (Harrington and Whitney 1987). As Ti is considered to be an immobile element, the ratio can be used as a weathering index: the lower it is, the higher the degree of weathering. The comparison between north- and south-facing sites is given in Fig. 9. Although the differences seem to be small and not significant, a slight trend towards a lower ratio at north-facing sites can be measured. A similar observation is made for the A- and B-index. Particularly for the A-index, a (non-significant) trend towards lower values at north-facing sites can be detected (Fig. 9). It seems that a stronger enrichment of Al over Ca, K and Na occurs at the north-facing sites.

4.7 Age of the sites

There is no detailed geomorphic mapping of the area of interest. The sites are in between the maximum extent of the glacier during the Little Ice Age and the Lateglacial position of the moraines (Younger and Oldest Dryas). Consequently, the surface of interest has been exposed to weathering in the time span > 200 yr and < 15 kyr. Some more detailed age indications are obtained by the dating of a nearby peat bog that gives a minimum age of 1304 – 1415 yr cal BP (Table 6). The stable soil organic matter has a maximum age of 5317 – 5468 cal BP. With this, the age of the soils can be further narrowed down to about 1400 – 5400 yr. The rather low weathering stage of the soils is in agreement with these age indications.

5 Discussion

The investigated soils have a relatively young age and are consequently rather weakly developed. Nonetheless, active pedogenesis is taking place despite the harsh and very cold climatic conditions (Lessoavaia and Polekhovsky 2009). The micro-morphology predominantly showed: (i) an accumulation of weakly and moderately decomposed remains of organic material, (ii) the formation of humus and (iii) an initial development of iron clots, films and ferruginous plasma *in situ*. The orientation of biotite particles, especially at the bottom of the

profile, indicates that the parent substrate mostly shows a fluvial, subglacial origin and sluggish sedimentation. The proportion of biotite was slightly decreasing towards the surface of the soil which can be due to weathering processes or some inhomogeneities of the parent material. The mineralogy of fine-size fractions ($< 1 \mu\text{m}$) showed the presence of chlorite, biotite and irregularly mixed-layered biotite-vermiculite (with a low proportion of vermiculite layers) which is in good agreement with the thin section data. Chlorite and biotite were both identified also in the rock fragments from the soil horizons. This indicated that glacier material is the main source of fine-size fraction of the soil. In addition, material from the river bank and adjacent slopes was interspersed with the glacially abraded material (as shown by the chlorite and mica proportion of the river bank and the rock fragments in the soil). The fine-size fraction finally was the most sensitive to weathering and pedogenesis. Biotite transformation into mixed-layered biotite-vermiculite having a minor proportion of vermiculite reflects first steps of weathering under acid conditions and cold environmental conditions (Vogt et al. 2010). In the studied soil profile, the most weathered horizon (from a mineralogical point of view) seems to be the AB_w horizon. It might be that some aeolian inputs gave rise to a 'rejuvenation' of the uppermost soil horizon. Many experimental and field observations of permafrost conditions demonstrate active weathering, mineral transformation processes and changes in the structure of (clay) minerals (e.g. Konishchev and Rogov 1993; Lessovaia et al. 2012). Often, chlorite seems to be the first silicate mineral to disappear (Vogt et al. 2010). In a literature review, Vogt et al. (2010) showed that hornblende and biotite are the next silicates to be disintegrated and feldspar grains may be covered by yellowish brown fine silty aggregates that are secondary clay minerals (Yershov 1998). At our sites, chlorite and mica reacted most. This can be explained by the selective destruction and transformation of such phyllosilicates in acidic soils of the northern taiga zone (Tonkonogov et al. 1987). Also smectites may be formed by mineral transformation reactions (Bender Koch et al. 1995; Falsone et al. 2012). Vogt et al. (2010) confirmed that cryogenic processes can be at least as efficient as those related to warm and temperate climates, with clay authigenesis being able to take place in relatively short periods of time.

Consequently, the absence of smectitic clays in the studied soils confirms their weak
evolutional stage and indicates the relatively short time of pedogenesis.

The differences between north- and south-facing soils were astonishingly clear. Although the
soils do not have a high age and are not very well developed, significant differences of pH, C
and N concentrations and stocks, C/N ratio, the chemical structure of SOM and the Al_o , Fe_o ,
 Al_d , Fe_d , Mn_o concentrations and stocks could be identified between the north- and south-
exposure sites.

As regards the oxalate- and dithionite-extractable fractions of Al, in general a slight
concentration increase could be noted in the lower layer when compared to the surface layer.
For Fe, both increase and a decrease of the concentrations was measured. This increase in
Al was more evident in the north-facing soils. This probably may point to the beginning of an
eluvation or to inhomogeneities in the soil material (input of some aeolian material? or
disturbances due to cryoturbation?).

Also due to the relatively young age, the C-stocks in the soils are not very high (Fig. 5;
average: 6 – 8 kg C/m²). The C abundance however only refers to a depth of 30 cm and
probably covers only about 60 – 70 % of the total mass. Arctic and subarctic lowland
permafrost soils usually have higher organic carbon stocks. Hugelius and Kuhry (2009), for
example, reported organic carbon stocks of 38 kg/m² (> 100 cm soil depth) in taiga and
tundra permafrost soils of north-eastern European Russia. In northern circumpolar
permafrost soils, Tarnocai et al. (2009) even found carbon abundances of 32.2 – 69.6 kg/m²
(0 – 100 cm soil depth). Wetland conditions (water retention due to underlying permafrost or
reduced evapotranspiration; peatland formation) and cryoturbation, particularly in permafrost
soils, are major soil-forming processes in arctic and alpine regions and may contribute to a
higher long-term storage of C in soils (Ugolini et al. 2006; Kaiser et al. 2007). Using the
alpine soil chronofunctions given in Dahms et al. (2012) for Fe_o , Al_o , Mn_o and Si_o , the
investigated soils here would fit very well for soils having age in the range of 1 to 10 kyr,
which confirms their Holocene age.

Even though the climate can be classified as very harsh, the fact that chemical weathering

seems to be more intense in the north-facing than in the south-facing soils fits with findings in the European Alps where in well-developed (Egli et al. 2010) as well as young soils (Egli et al. 2006, 2011) distinct differences between north- and south-facing slopes were found. The evolution of soils, chemical weathering and formation and transformation of (clay) minerals always seemed to proceed faster on north-facing slopes. In several parts of the European Alps, eluviation and illuviation processes of Al and Fe – and consequently podzolisation processes – within a soil profile are more intense on north-facing slopes (Egli et al. 2010). Especially in the subalpine range, the stemflow as well as the accumulation of coniferous tree litter leads to an intensified acidification of the soil (Certini et al. 1998; Egli et al. 2010) and to a higher production of organic ligands in the soil solution. A higher amount of acid-soluble organic compounds and consequently –COOH functional groups was present on north-facing sites of the Italian Alps (Egli et al. 2010). Organic acids and ligands (Stumm and Wieland 1990; Lundström et al. 2000) particularly accelerate weathering processes on north-facing sites – not only in the European Alps but obviously also in the rather dry and very cold conditions of the Altai Mountains.

North-facing sites are characterised by lower temperatures, a lower evapotranspiration, a higher humidity and higher acidity. The higher humidity and the less favourable thermal conditions lead to an accumulation of labile, weakly-degraded organic matter (e.g. as seen by the higher relative proportion of polysaccharides in the investigated north-facing soils; Fig. 6) and consequently to a higher production of soluble organic ligands and organic acids (higher proportion of carboxylic groups on north-facing sites; Fig. 6) that enhance weathering. This is in agreement with Ascher et al. (2012) who found that the $\delta^{13}\text{C}$ signatures of the PLFA (phospholipid fatty acid analysis) markers in alpine soils suggested a lower decomposition rate at the cooler north-facing sites, resulting in a lower respiratory loss and an accumulation of weakly-decomposed organic material. DGGE (denaturing gradient gel electrophoresis) data supported the PLFA results.

6 Conclusions

Coming back to the initially-made hypothesis, we can confirm that even under very harsh and cold conditions the availability of water is crucial and that temperature, as a factor per se, seems to play a less important role. This statement partially fits with the hypothesis of Brantley et al. (2011; hypothesis no. 9) that the export of weathering products (and consequently weathering in general) predominantly depends on precipitation and evapotranspiration. This, however, may partially conflict with another hypothesis concerning the biota-facilitated chemical weathering (i.e. that the solar-to-chemical conversion of energy by plants controls the location and extent of biological weathering) given in Brantley et al. (2011).

Water availability predominantly determines weathering also in very cold regions of the Altai Mountains and agrees with results from the European Alps (Egli et al. 2006). Due to the less favourable thermal conditions microbial activity is most likely reduced at north-facing sites (a hypothesis that still has to be tested and confirmed). This leads to an accumulation of labile, weakly-degraded organic matter and consequently to a higher production of soluble organic ligands and organic acids. Together with the higher soil moisture, these circumstances enhance weathering mechanisms.

Acknowledgements The XRD measurements of the soil and rock samples were carried out at the X-ray Diffraction Centre of the St. Petersburg State University. This research was supported by a grant of the Scientific & Technological Cooperation Programme Switzerland-Russia.

References

- Agatova AR, Hule VV, Mistrykov AA (2002) Dynamics of Sophiysky glacier (south-eastern Altay): the late glacial maximum-XX century. *Geomorphology (Geomorphologia)* 2:92-105
- Ascher J, Sartori G, Graefe U, Thornton B, Ceccherini MT, Pietramellara G, Egli M (2012) Are humus forms, mesofauna and microflora in subalpine forest soils sensitive to thermal conditions? *Biol Fertil Soils* 48:709-725

- Bender Koch C, Mørup S, Madsen MB, Vistisen L (1995) Iron-containing weathering products of basalt in a cold, dry climate. *Chem Geol* 122:109–112
- Birkeland PW (1999) *Soils and geomorphology*. Oxford University Press, New York
- Brantley SL, Megonigal JP, Scatena FN, Balogh-Brunstad Z, Barnes RT, Bruns MA, van Cappellen P, Dontsova K, Hartnett HE, Hartshorn AS, Heimsath A, Herndorn E, Jin L, Keller CK, Leake JR, McDowell WH, Meinzer FC, Mozdzer TJ, Petsch S, Pett-Ridge J, Pregitzer KS, Raymond PA, Riebe CS, Shumaker K, Sutton-Grier A, Walter R, Yoo K (2011) Twelve testable hypotheses on the geobiology of weathering. *Geobiology* 9:140–165
- Brindley GW, Brown G (1980) *Crystal Structures of Clay Minerals and their X-ray Identification* (Eds) Monograph 5, Mineralogical Society, London
- Bronk Ramsey C (2001) Development of the radiocarbon calibration program OxCal. *Radiocarbon* 43:355–363
- Bronk Ramsey C (2009) Bayesian analysis of radiocarbon dates. *Radiocarbon* 51:337–360
- Carter BJ, Ciolkosz EJ (1991) Slope gradient and aspect effects on soils developed from sandstone in Pennsylvania. *Geoderma* 49:199–213
- Certini G, Ugolini FC, Corti G, Agnelli A (1998) Early stages of podzolization under Corsican pine (*Pinus nigra* Arn. ssp. *larico*). *Geoderma* 83:103–125
- Conen F, Yakutin MV, Zumbunn T, Leifeld J (2007) Organic carbon and microbial biomass in two soil development chronosequences following glacial retreat. *Eur J Soil Sci* 58:758–762
- Cornell RM, Schwertmann U (2003) *The iron oxides structure, properties, reactions, occurrences and uses*, 2nd edition, Wiley-VCH GmbH & Co., Weinheim
- Dahlgren RA, Boettinger JL, Huntington GL, Amundson RG (1997) Soil development along an elevational transect in the western Sierra Nevada, California. *Geoderma* 78:207–236
- Dahms D, Favilli F, Krebs R, Egli M (2012) Soil weathering and accumulation rates of oxalate-extractable phases from alpine chronosequences of up to 1 Ma in age. *Geomorphology* 151–152:99–113

- Dixon JL, von Blanckenburg F (2012) Soils as pacemakers and limiters of global silicate weathering. *CR Geosci* 344:596-609
- Dixon JL, Heimsath AM, Anmundson R (2009) The critical role of climate and saprolite weathering in landscape evolution. *Earth Surf Proc Land* 34:1507-1521
- Dixon JL, Hartshorn AS, Heimsath AM, DiBiase RA, Whipple KX (2012) Chemical weathering response to tectonic forcing: A soils perspective from the San Gabriel Mountains, California. *Earth Planet Sc Lett* 323-324:40-49
- Dokuchaev VV (1883) Russian Chernozem. In: Kaner N. (ed) Selected works of V. V. Dokuchaev (1967), Jerusalem, Int. Program Sci. Transl., pp 1–419
- Egli M, Wernli M, Kneisel C, Haeberli W. (2006) Melting glaciers and soil development in the proglacial area Morteratsch (Swiss Alps): I Soil type chronosequence. *Arc Antarct Alp Res* 38:499-509
- Egli M, Mirabella A, Sartori G, Giaccari D, Zanelli R, Plötze M (2007) Effect of slope aspect on transformation of clay minerals in Alpine soils. *Clay Miner* 42:375–401
- Egli M, Mirabella A, Sartori G (2008) The role of climate and vegetation in weathering and clay mineral formation in late Quaternary soils of the Swiss and Italian Alps. *Geomorphology* 102:307-324
- Egli M, Sartori G, Mirabella A (2010) The effects of exposure and climate on the weathering of late Pleistocene and Holocene Alpine soils. *Geomorphology* 114:466-482
- Egli M, Wernli M, Burga C, Kneisel C, Mavris C, Valboa G, Mirabella A, Plötze M, Haeberli W (2011) Fast but spatially scattered smectite-formation in the proglacial area Morteratsch: an evaluation using GIS. *Geoderma* 164:11-21
- Falsone G, Celi L, Caimi A, Simonov G, Bonifacio E (2012) The effect of clear cutting on podzolisation and soil carbon dynamics in boreal forests (Middle Taiga zone, Russia). *Geoderma* 177-178:27-38
- Favilli F, Egli M, Cherubini P, Sartori G, Haeberli W, Delbos E (2008) Comparison of different methods of obtaining a resilient organic matter fraction in Alpine soils. *Geoderma* 145:355-369

- Gorbushina AA (2007) Life on the rocks. Minireview. *Environ Microbiol* 9:1613–1631
- Hall K, Thorn CE, Matsuoka N, Prick A (2002) Weathering in cold regions: some thoughts and perspectives. *Prog Phys Geog* 26:577-603
- Harnois L (1988) The CIW index: a chemical index of weathering. *Sediment Geol* 55:319-322
- Harrington CD, Whitney JW (1987) Scanning electron microscope method of rock-varnish dating. *Geology* 15:967-970
- Hitz C, Egli M, Fitze P (2002) Determination of the sampling volume for representative analysis of alpine soils. *Z Pflanz Bodenkunde* 165:326-331
- Hlavay J, Jonas K, Elek S, Inczedy J (1978) Characterization of the particle size and the crystallinity of certain minerals by IR spectrophotometry and other instrumental methods; II, Investigations on quartz and feldspar. *Clays Clay Miner* 26:139–143
- Hugelius G, Kuhry P (2009) Landscape partitioning and environmental gradient analyses of soil organic carbon in a permafrost environment. *Global Biogeochem Cy* 23:GB3006
- IUSS Working Group WRB (2007) World Reference Base for Soil Resources (2nd ed.). World Soil Resources Reports No. 103, First update, FAO, Rome
- Jann B (2005) Einführung in die Statistik. 2. bearbeitete Auflage, Oldenburg Wissenschaftsverlag, München
- Jenny H (1941) Factors of soil formation. Mc Graw-Hill Book Company, New York
- Kaiser C, Meyer H, Biasi C, Rusalimova O, Barsukov P, Richter A (2007) Conservation of soil organic matter through cryoturbation in arctic soils in Siberia. *J Geophys Res-Bioge* 112:G02017
- Klemmedson JO (1964) Topofunction of soils and vegetation in a landscape. *Am Soc Agron* 5:176–189
- Kloprogge JT, Hickey L, Frost RL (2004) FT-Raman and FT-IR spectroscopic study of the local structure of synthetic Mg/Zn/Al-hydrotalcites. *J Raman Spectrosc* 35:967-974
- Konishchev VN, Rogov VV (1993) Investigations of cryogenic weathering in Europe and Northern Asia. *Permafrost Periglac* 4:49–64

- 635 Kronberg GI, Nesbitt HW (1981) Quantification of weathering of soil chemistry and soil
636 fertility. *J Soil Sci* 32:453-459
- 637 Langer K, Raith M (1974) Infrared Spectra of Al-Fe(III)-Epidotes and Zoisites, $\text{Ca}_2(\text{Al}_{1-p}\text{Fe}^{3+}_p)\text{Al}_2\text{O}(\text{OH})[\text{Si}_2\text{O}_7][\text{SiO}_4]$. *Am Mineral* 59:1249-1258
- 638
639 Lanson B (1997) Decomposition of experimental X-ray diffraction patterns (profile fitting): a
640 convenient way to study clay minerals. *Clays Clay Miner* 45: 132-146.
- 641 Lessovaia SN, Polekhovsky YuS (2009) Mineralogical composition of shallow soils on basic
642 and ultrabasic rocks of east Fennoscandia and of the Ural mountains, Russia. *Clays
643 Clay Miner* 57:476-485
- 644 Lessovaia S., Dultz S, Polekhovsky Yu, Krupskaya V, Vigasina M, Melchakova L (2012)
645 Rock control of pedogenic clay mineral formation in a shallow soil from serpentinous
646 dunite in the Polar Urals, Russia. *Appl Clay Sci* 64:4–11
- 647 Lorenzoni P, Mirabella A, Bidini C, Lulli L (1995) Soil genesis on trachytic and leucitic lavas
648 of Cimini volcanic complex (Latium, Italy). *Geoderma* 68:79-99
- 649 Lundström US, van Breemen N, Bain D (2000) The podzolization process. A review.
650 *Geoderma* 94:91–107
- 651 Macyk TM, Pawluk S, Lindsay D (1978) Relief and microclimate as related to soil properties.
652 *Can J Soil Sci* 58:421–438
- 653 Madejova J, Komadel P (2005) Information available from infrared spectra of the fine fraction
654 of bentonites. In: Klopogge JT (ed) The application of vibration spectroscopy of clay
655 minerals and layered double hydroxides, CMS Workshop Lectures, vol. 13. The Clay
656 Mineral Society, Aurora, CO, pp 65-98
- 657 Marinina AM, Samoilova GS (1987) Physical geography of the Mountainous (Phizicheskaya
658 geographiya Gornogo Altaya) Barnaul BGPE (in Russian)
- 659 Mavris C, Egli M, Plötze M, Blum J.D, Mirabella A, Giaccari D, Haeberli W (2010) Initial
660 stages of weathering and soil formation in the Morteratsch proglacial area (Upper
661 Engadine, Switzerland). *Geoderma* 155:359-371
- 662 McKeague JA, Brydon JE, Miles NM (1971) Differentiation of forms of extractable iron and

- aluminium in soils. Soil Sci Soc Am Pro 35:33-38
- Mehra OP, Jackson ML (1958) Iron oxide removal from soils and clays by a dithionite-citrate system buffered with sodium bicarbonate. Clay Clay Miner 7:317-327
- Mizota C, van Reeuwijk LP (1989) Clay mineralogy and chemistry of soils formed in volcanic material in diverse climat regions. International Soil Reference and Information Centre, Soil Monograph, vol. 2, Wageningen
- Moore, D.M., Reynolds, R.C., 1997. X-ray diffraction and the identification and analysis of clay minerals. 2nd edition, Oxford University Press, New York, NY
- Narozhniy Y, Zemtsov V (2011) Current State of the Altai Glaciers (Russia) and Trends Over the Period of Instrumental Observations 1952–2008. Ambio 40:575–588
- Nesbitt HW, Young GM (1989) Formation and diagenesis of weathering profiles. J Geol 97:129-147
- Ogureeva GN (1980) Botanical geography of Altai (Botanicheskaya geographia Altaya). Nauka, Moscow (in Russian)
- Osawa A, Zyryanova OA, Matsuura Y, Kajimoto T, Wein RW (eds) (2010) Permafrost Ecosystems: Siberian Larch Forests, Ecological studies, Vol. 209, Springer Science+Business Media B.V.
- Osokina V.G (managing ed.), Babaeva L.I., Sazonova T.N., Fomichkina T.M., Shevchenko M.A., Prokudina T.M., Volodina K. A., Ivanova E.S., Shestel T.V. (1993). Scientific climate handbook of the USSR. Series 3, Multi-annual data, part 1 – 6, issue 20, Regions of Tomsk, Novosibirsk, Altai and province of Kemerovo. Directorate of Hydrometeorology, Hydrometeorological Centre of the West-Siberian territory, Hydrometeoisdat, Sankt-Petersburg (in Russian).
- Ostanin OV, Mikhailov NN, Arkhipov SM (2004) Change of glaciers of south and central Altai from the end of XIX century and tendencies of its development in XXI (Izmenenie lednikov uuzhnogo e zentral'nogo Altaya s konza XIX v. e tendenzii ekh razvitiya v XXI v.) Geography and nature management of Siberia (Geographiya e prirodopol'zovabie Sibiri) 7:172-182, Barnaul (in Russian)

- 691 Piccolo A, Mirabella A (1985) Effetto di differenti estraenti inorganici e organici su alcune
692 proprietà chimiche e chimico-fisiche di un acido umico. Annali dell'Istituto Sperimentale
693 per lo Studio e Difesa Suolo 16:159–168
- 694 Rasmussen C, Brantley SL, Richter D, Blum A, Dixon J, White A (2011) Strong climate
695 control on plagioclase weathering in granitic terrain. Earth Planetary Sc Lett 301:521-530
- 696 Rech JA, Reeves RW, Hendricks D (2001) The influence of slope aspect on soil weathering
697 processes in the Springerville volcanic field. Arizona. Catena 43:49–62
- 698 Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Buck
699 CE, Burr GS, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Hajdas I, Heaton TJ,
700 Hogg AG, Hughen KA, Kaiser KF, Kromer B, McCormac FG, Manning SW, Reimer RW,
701 Richards DA, Southon JR, Talamo S, Turney CSM, van der Plicht J, Weyhenmeyer CE
702 (2009) INTCAL09 and MARINE09 radiocarbon age calibration curves, 0-50,000 years cal
703 BP. Radiocarbon 51:1111-1150
- 704 Schroeder PA (2002) Infrared Spectroscopy in clay science. In: Rule A, Guggenheim S (eds)
705 CMS Workshop Lectures, Vol. 11, Teaching Clay Science, The Clay Mineral Society,
706 Aurora, CO, pp. 181-206
- 707 Senesi N, D'Orazio V, Ricca G (2003) Humic acids in the first generation of EUROSOLS.
708 Geoderma 116:325–344
- 709 Simas FNB, Schaefer CEGR, Melo VF, Guerra MBB, Saunders M, Gilkes RJ (2006) Clay
710 sized minerals in permafrost-affected soils (Cryosols) from King George Island, Antarctica,
711 Clays Clay Miner 54:721-736
- 712 Stevenson FJ (1994) Humus chemistry: genesis, composition, reactions. Wiley, New York,
713 NY
- 714 Stumm W, Wieland E (1990) Dissolution of oxide and silicate minerals: rates depend on
715 surface speciation. In: Stumm W (Ed.), Aquatic Chemical Kinetics: Reaction Rates of
716 Processes in Natural Waters. Wiley-Interscience, New York, pp. 367–398
- 717 Syromyatina MV (2010) Recent changes of climate and elements of high-altitude belts of
718 Altai landscapes (Sovremennyye izmeneniya klimata b elementov vysotnoi poyasnosti

- landshaftov Altaya). PhD thesis, St-Petersburg (in Russian)
- 720 Tan KH (2003) Humic matter in soil and the environment: Principles and controversies.
Marcel Dekker, Inc., New York – Basel
- 722 Tarnocai C, Canadell JG, Schuur EAG, Kuhry P, Mazhitova G, Zimov S (2009) Soil organic
carbon pools in the northern circumpolar permafrost region. *Global Biogeochem Cy*
23:GB2023
- 725 Tonkonogov VD, Gradusov BP, Rubilina NE, Targulian VO, Chizhikova NP (1987) On the
differentiation of the mineralogical and chemical compositions of Soddy_Podzolic and
Podzolic Soils (K differenziatsii mineralogicheskogo e khimicheskogo sostava derno-
podzolistykh e podzolistykh pochv). *Pochvovedenie* 3:68–81 (in Russian)
- 729 Ugolini FC, Corti G, Certini G (2006) Pedogenesis in the sorted patterned ground of Devon
plateau, Devon Island, Nunavut, Canada. *Geoderma* 136:87-106
- 731 Vantelon D, Pelletier M, Michot LJ, Barres O, Thomas F (2001) Fe, Mg and Al distribution in
the octahedral sheet of montmorillonites. An infrared study in the OH-bending region. *Clay
Miner* 36:369-379
- 734 Vogt T, Clauer N, Larqué P (2010) Impact of climate and related weathering processes on
the authigenesis of clay minerals: Examples from circum-Baikal region, Siberia. *Catena*
80:53-64
- 737 White AF, Blum AE (1995), Effects of climate on chemical weathering in watersheds.
Geochim Cosmochim Acta 59:1729-1747
- 739 White AF, Blum AE, Bullen TD, Vivit DV, Schulz M, Fitzpatrick J (1999) The effect of
temperature on experimental and natural chemical weathering rates of granitoid rocks.
Geochim Cosmochim Acta 63:3277–3291
- 742 Whitney D, Evans BW (2010) Abbreviations for names of rock-forming minerals. *Am Mineral*
95:185–187
- 744 Williams JZ, Bandsra JZ, Pollard D, Brantley SL (2010) The temperature dependence of
feldspar dissolution determined using a coupled weathering-climate model for Holocene-
aged loess soils. *Geoderma* 156:11-19

747 Yershov ED (1998) General Geocryology. Cambridge University Press, Cambridge

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

748 Zanelli R, Egli M, Mirabella A, Giaccai A, Abdelmoula M (2007) Vegetation effects on
749 pedogenetic forms of Fe, Al and Si and on clay minerals in soils in southern Switzerland
750 and northern Italy. Geoderma 141, 119-129

Table 1 Characteristics of the study sites. The sites are within an area of about 2-3 ha

Site	Exposure °N	Slope °	Altitude m asl	Parent material	Vegetation	Soil group (IUSS Working Group WRB, 2007)
North-facing sites						
3	340	10	2378	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol
4	0	8	2378	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol
5	8	14	2376	Glacial deposits; silt loam or loam	Juniper, Alpine grassland	Cambic Cryosol
6	10	6	2376	Glacial deposits; silt loam or loam	Juniper, Alpine grassland	Cambic Cryosol
7	303	7	2382	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol
South-facing sites						
8	195	13	2383	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol
9	155	9	2379	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol
10	154	12	2376	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol
11	154	12	2376	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol
12	130	10	2376	Glacial deposits; silt loam or loam	Alpine grassland	Cambic Cryosol

Table 2 General physical and chemical characteristics of the soil material

Site	Depth cm	Skeleton content weight-%	Bulk density g/cm ³	Sand %	Silt %	Clay %	Texture ^a	pH (CaCl ₂)	C g/kg	N g/kg	C/N
North-facing											
Alt 3	0-15	0.3	0.90	46.7	46.2	7.1	L	5.2	41.40	2.86	14.5
Alt 3	15-30	0.0	0.95	54.6	39.0	6.4	SL	5.1	21.30	1.63	13.1
Alt 4	0-15	0.2	0.92				SiL	5.0	30.69	2.25	13.6
Alt 4	15-30	0.2	1.05				SiL	5.1	22.83	1.78	12.9
Alt 5	0-15	0.0	0.83	35.5	51.4	13.1	SiL	4.4	47.09	2.70	17.4
Alt 5	15-30	0.0	0.79	26.6	63.4	10.0	SiL	4.5	49.92	3.08	16.2
Alt 6	0-15	0.0	0.73				SiL	4.4	43.90	2.55	17.2
Alt 6	15-30	0.0	1.10				SiL	4.5	19.11	1.07	17.9
Alt 7	0-15	0.1	1.17				SiL	4.7	22.41	1.74	12.9
Alt 7	15-30	0.0	0.97				SiL	5.4	22.22	1.64	13.5
South-facing											
Alt 8	0-15	0.1	0.98	38.4	51.2	10.4	SiL	5.6	22.49	1.81	12.4
Alt 8	15-30	0.1	0.93	50.2	43.5	6.3	L	6.1	20.96	1.68	12.5
Alt 9	0-15	0.0	1.08				SiL	5.0	16.54	1.39	11.9
Alt 9	15-30	0.0	0.95				SiL	5.2	15.15	1.36	11.1
Alt 10	0-15	0.8	0.91	40.9	51.8	7.7	SiL	5.4	20.63	1.61	12.8
Alt 10	15-30	0.1	0.95	29.6	62.3	8.1	SiL	5.7	12.24	1.04	11.8
Alt 11	0-15	0.0	1.00				SiL	5.3	29.56	2.16	13.7
Alt 11	15-30	0.0	1.20				SiL	5.2	14.69	1.14	12.9
Alt 12	0-15	0.1	0.89				SiL	4.5	29.99	2.27	13.2
Alt 12	15-30	0.0	0.99				SiL	5.4	27.65	2.10	13.2
Profile/Horizon											
A	5-19	0.2	0.92				SiL	5.2	31.19	2.60	12.0
ABw	19-27	0.2	1.05				SiL	5.1	36.41	2.92	12.5
BC	27-68	0.1	1.10				SiL	4.9	6.62	0.43	15.4

^a SiL = Silt loam, SL = sandy loam, L = loam

Table 3 Major IR absorption bands and assignments (Piccolo and Mirabella, 1985; Stevenson, 1994; Senesi et al., 2003; Tan, 2003)

Band	Wave number cm ⁻¹	Assignment
1	2980 - 2880	Aliphatic C-H stretching (aliphatic methyl and methylene groups)
2	1725 - 1710	C=O stretching of COOH, aldehydes and ketones
3	1660 - 1630	C=O stretching of amide groups, quinone C=O and/or C=O of H- bonded conjugated ketones
4	1620 - 1600	Aromatic C=C, strongly H-bonded C=O of conjugated ketones
5	1535 - 1500	Aromatic rings, amide II vibration
6	1495 - 1470	N-H stretching of proteic amides
7	1470 - 1440	Aliphatic C-H bending
8	1413 - 1333	OH deformation and C-O stretching of phenolic groups
9	1280 - 1200	C-O stretching and OH deformation of COOH, C-O stretching of aryl ethers and phenols
10	1190 - 1127	C-OH stretching of aliphatic alcohols
11	1116 - 1050	Secondary alcohols
12	1080 - 1030	C-O stretching of polysaccharides

Table 4 Oxalate (o)- and dithionite(d)-extractable fractions of Al, Fe, Mn and Si in the fine earth

Site	Depth cm	Al(o) g/kg	Al(d) g/kg	Fe(o) g/kg	Fe(d) g/kg	Mn(o) g/kg	Si(o) g/kg
North-facing							
Alt 3	0-15	0.62	0.55	0.90	2.71	0.042	0.081
Alt 3	15-30	0.86	0.86	0.77	2.51	0.090	0.078
Alt 4	0-15	0.74	0.72	1.07	3.01	0.131	0.090
Alt 4	15-30	1.27	1.31	1.30	3.31	0.072	0.124
Alt 5	0-15	0.78	0.94	1.29	3.20	0.050	0.077
Alt 5	15-30	2.05	2.30	1.95	3.98	0.018	0.130
Alt 6	0-15	0.82	1.00	2.40	3.94	0.029	0.110
Alt 6	15-30	0.85	0.74	1.34	2.87	0.045	0.129
Alt 7	0-15	0.61	0.51	1.17	2.63	0.057	0.087
Alt 7	15-30	0.70	0.66	0.67	1.92	0.040	0.078
South-facing							
Alt 8	0-15	0.53	0.41	0.68	2.21	0.047	0.094
Alt 8	15-30	0.62	0.50	0.72	2.32	0.060	0.110
Alt 9	0-15	0.58	0.36	0.64	2.15	0.057	0.068
Alt 9	15-30	0.64	0.63	0.66	2.10	0.062	0.060
Alt 10	0-15	0.60	0.63	0.81	2.79	0.015	0.169
Alt 10	15-30	0.87	0.89	0.99	2.53	0.043	0.244
Alt 11	0-15	0.65	0.65	1.22	2.82	0.015	0.143
Alt 11	15-30	0.84	0.77	1.08	2.75	0.059	0.077
Alt 12	0-15	0.63	0.67	0.96	2.68	0.036	0.032
Alt 12	15-30	0.73	0.71	0.93	2.41	0.050	0.158
Profile/Horizon							
A	5-19	1.01	1.05	1.36	3.23	0.065	0.104
ABw	19-27	2.04	1.85	1.78	3.76	0.077	0.174
BC	27-68	1.20	1.29	1.23	2.86	0.024	0.134

Table 5

Table 5 Total elemental contents (major components) in the fine earth of the LOI-free fraction (inorganic part; normalised to 100%)

Site	Depth cm	SiO ₂ %	Al ₂ O ₃ %	Fe ₂ O ₃ %	CaO %	MgO %	MnO %	Na ₂ O %	K ₂ O %	TiO ₂ %	P ₂ O ₅ %
North-facing											
Alt 3	0-15	66.7	14.0	6.2	2.7	3.9	0.109	2.8	2.3	0.97	0.36
Alt 3	15-30	67.8	14.1	6.0	2.8	3.5	0.118	2.4	1.9	0.97	0.47
Alt 4	0-15	66.4	14.1	6.5	2.6	4.0	0.129	2.5	2.3	1.07	0.36
Alt 4	15-30	68.3	14.0	6.0	2.5	3.2	0.107	2.6	1.8	1.03	0.42
Alt 5	0-15	64.5	14.9	6.8	2.5	4.5	0.114	2.6	2.7	1.06	0.42
Alt 5	15-30	65.0	15.5	6.5	2.4	4.1	0.099	2.6	2.4	1.03	0.45
Alt 6	0-15	64.4	15.6	6.1	2.4	4.9	0.103	2.5	2.7	0.96	0.31
Alt 6	15-30	64.9	15.5	6.1	2.3	4.8	0.108	2.5	2.6	0.96	0.34
Alt 7	0-15	66.6	14.1	6.3	2.6	3.9	0.115	2.7	2.3	1.03	0.38
Alt 7	15-30	68.3	13.7	5.7	2.8	3.4	0.099	2.8	1.9	0.94	0.39
South-facing											
Alt 8	0-15	66.4	14.2	6.1	2.7	3.9	0.110	2.6	2.6	0.96	0.39
Alt 8	15-30	67.1	13.9	5.8	2.9	3.7	0.110	2.6	2.4	1.00	0.46
Alt 9	0-15	67.4	14.0	5.9	2.4	3.8	0.105	2.7	2.3	0.96	0.38
Alt 9	15-30	68.5	13.5	5.8	2.6	3.3	0.107	2.7	2.0	0.99	0.49
Alt 10	0-15	66.7	14.5	5.9	2.4	4.3	0.103	2.4	2.4	0.95	0.37
Alt 10	15-30	66.2	14.9	6.1	2.4	4.1	0.106	2.5	2.3	0.99	0.38
Alt 11	0-15	65.8	14.8	6.1	2.5	4.6	0.102	2.4	2.5	0.98	0.33
Alt 11	15-30	67.8	14.2	5.8	2.5	3.8	0.106	2.4	2.0	0.99	0.45
Alt 12	0-15	67.2	14.1	6.2	2.6	3.8	0.107	2.5	2.1	1.04	0.36
Alt 12	15-30	67.8	13.9	5.8	2.7	3.7	0.107	2.5	2.1	0.97	0.40
Profile/Horizon											
A	5-19	67.0	13.9	6.3	2.5	3.8	0.116	2.6	2.3	1.07	0.37
ABw	19-27	65.8	15.2	6.3	2.5	4.0	0.111	2.5	2.2	1.02	0.46
BC	27-68	65.9	15.0	6.4	2.3	3.9	0.109	2.6	2.3	1.05	0.32

Table 6

Table 6 Radiocarbon dating of the stable organic carbon fraction (H₂O₂ treated sample) of an investigated soil and a nearby peat bog

Sample/Site	Depth (cm)	UZH No	ETH No	Uncalibrated ages (yr uncal. BP)	$\delta^{13}\text{C}$ (‰)	Calibrated ages (yr cal BP) 2 σ range
Peat (Alt2)	60	6105	47904	585 $\pm$ 30	-23.10 $\pm$ 1.1	535 - 652
	100	6106	47905	1485 $\pm$ 30	-27.90 $\pm$ 1.1	1304 - 1415
Soil, stable OM						
Alt4	0-15	6103	48764	3295 $\pm$ 25	-26.11 $\pm$ 1.1	3453 - 3579
Alt4	15-30	6104	48765	4665 $\pm$ 25	-21.72 $\pm$ 1.1	5317 - 5468

Figure captions

Fig. 1 Location of the study sites (soils and peat bog) in the south-eastern part of the Altai mountains (border region: Russia/Mongolia/China). Impression of the environment and sampling sites together with a typical soil profile

Fig. 2 Thin sections of rock samples from the soil profile: (a) gneiss structure with sub-parallel arrangement of minerals; (b) quartz vein crossing the rock; (c) development of chlorite on biotite. Determination on thin sections using (||) plain polars and (X) crossed Nicols polarised lights. Indexes are according to Whitney and Evans (2010), Qz – quartz, Chl – chlorite (group), Bt – biotite (group).

Fig. 3 Thin sections of soil samples using plain polars: (a) weakly rounded quartz grain with cleavages and frost cracks from the A horizon; (b) slightly decomposed organic matter and raw humus (A horizon); (c) fragments of zoogenic transformation (A horizon); and (d) weakly disintegrated biotite grain without features of chemical weathering (A horizon), (e) horizontally oriented biotite particles of the BC horizon and (f) sandy – silty plasmic fabric of the BC horizon using crossed Nicols

Fig. 4 Comparison (box plot; with median, 25th- and 75th-percentile values, min. and max. values) of pH, org. C and N content and C/N ratio between north- and south-facing sites. The differences are significant ($p < 0.05$) for pH(CaCl_2), org. and the C/N ratio

Fig. 5 Abundance (stocks) of oxyhydroxides and C and N (given for 0-15 and 15-30 and as single values of 0-30 cm soil depth) as a function of exposure. The differences are significant ($p < 0.05$) for all parameters except Fe_d and N

Fig. 6 Relative proportion of functional groups as a function of exposure. 9 = C-O stretching and OH deformation of COOH, C-O stretching of aryl ethers and phenols; 10 = C-OH stretching of aliphatic alcohols; 11 = Secondary alcohols; 12 = C-O stretching of polysaccharides. The differences are significant ($p < 0.05$) for all parameters except for 10

Fig. 7 Comparison (box plot; with median, 25th- and 75th-percentile values, min. and max. values) of the dithionite- and oxalate-extractable fractions and Fe_o/Fe_d ratio between north- and south-facing sites. The differences are significant ($p < 0.05$) for all parameters except Al_o and Fe_o/Fe_d

Fig. 8 Mineralogy of the soil profiles: X-ray patterns of the $< 1 \mu m$ fraction of the A, ABw and BC horizons. The XRD-curves are corrected for Lorentz and polarisation factors and d -spacings are given in nm (A –C). Mg = Mg-saturated sample, EG = Ethylene glycol solvation and corresponding heating treatment ($550^\circ C$). (D) shows the X-ray data up to $> 60^\circ 2\theta$ from powder samples of the fine earth fraction ($< 2mm$) of the soil profile (A, ABw and BC horizon), rock fragments in the soil and the gneiss sample (Akkol river), (E) and (F) show the FTIR spectra (absorbance) of the $< 2 mm$ fractions for the range $4000 - 2600 cm^{-1}$ and $1200 - 400 cm^{-1}$ with indication of peak positions

Fig. 9 Weathering indexes (A and B index; according to Kronberg and Nesbitt 1981) and $(K+Ca)/Ti$ ratio as a function of exposure

Figure 1
[Click here to download high resolution image](#)

Figure 2
[Click here to download high resolution image](#)

Figure 3
[Click here to download high resolution image](#)

Figure 4
[Click here to download high resolution image](#)

Figure 5
[Click here to download high resolution image](#)

Figure 6
[Click here to download high resolution image](#)

Figure 7
[Click here to download high resolution image](#)

Figure 8
[Click here to download high resolution image](#)

Figure 9
[Click here to download high resolution image](#)