



---

Year: 2015

---

**Measurement of the ratio of the production cross sections times branching  
fractions of  $B_c^\pm \rightarrow J\psi\pi^\pm$  and  $B_c^\pm \rightarrow J\psi K^\pm$  and  
 $\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm\pi^\pm\pi^\mp) / \mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm)$  in pp collisions at  $\sqrt{s} = 7$  TeV**

CMS Collaboration ; Canelli, M F ; Chiochia, V ; Kilminster, B ; Robmann, P ; et al

Abstract: The ratio of the production cross sections times branching fractions  $(\sigma(B_c^\pm)\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm)) / (\sigma(B^\pm)\mathcal{B}(B^\pm \rightarrow J/\psi\pi^\pm))$  is studied in proton-proton collisions at a center-of-mass energy of 7 TeV with the CMS detector at the LHC. The kinematic region investigated requires  $B_c^\pm$  and  $B^\pm$  mesons with transverse momentum  $p_T > 15$  GeV and rapidity  $|y| < 1.6$ . The data sample corresponds to an integrated luminosity of 5.1 inverse femtobarns. The ratio is determined to be  $[0.48 \pm 0.05$  (stat)  $\pm 0.03$  (syst)  $\pm 0.05$  ( $\tau_{B_c}$ )]

DOI: [https://doi.org/10.1007/JHEP01\(2015\)063](https://doi.org/10.1007/JHEP01(2015)063)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-108051>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 3.0 Unported (CC BY 3.0) License.

Originally published at:

CMS Collaboration; Canelli, M F; Chiochia, V; Kilminster, B; Robmann, P; et al (2015). Measurement of the ratio of the production cross sections times branching fractions of  $B_c^\pm \rightarrow J\psi\pi^\pm$  and  $B_c^\pm \rightarrow J\psi K^\pm$  and  $\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm\pi^\pm\pi^\mp) / \mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm)$  in pp collisions at  $\sqrt{s} = 7$  TeV. *Journal of High Energy Physics*, 2015(63):online.

DOI: [https://doi.org/10.1007/JHEP01\(2015\)063](https://doi.org/10.1007/JHEP01(2015)063)

# Measurement of the ratio of the production cross sections times branching fractions of $B_c^\pm \rightarrow J/\psi\pi^\pm$ and $B^\pm \rightarrow J/\psi K^\pm$ and $\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm\pi^\pm\pi^\mp)/\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm)$ in pp collisions at $\sqrt{s} = 7$ TeV



## The CMS collaboration

*E-mail:* [cms-publication-committee-chair@cern.ch](mailto:cms-publication-committee-chair@cern.ch)

**ABSTRACT:** The ratio of the production cross sections times branching fractions  $(\sigma(B_c^\pm)\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm))/(\sigma(B^\pm)\mathcal{B}(B^\pm \rightarrow J/\psi K^\pm))$  is studied in proton-proton collisions at a center-of-mass energy of 7 TeV with the CMS detector at the LHC. The kinematic region investigated requires  $B_c^\pm$  and  $B^\pm$  mesons with transverse momentum  $p_T > 15$  GeV and rapidity  $|y| < 1.6$ . The data sample corresponds to an integrated luminosity of  $5.1 \text{ fb}^{-1}$ . The ratio is determined to be  $[0.48 \pm 0.05 (\text{stat}) \pm 0.03 (\text{syst}) \pm 0.05 (\tau_{B_c})]\%$ . The  $B_c^\pm \rightarrow J/\psi\pi^\pm\pi^\pm\pi^\mp$  decay is also observed in the same data sample. Using a model-independent method developed to measure the efficiency given the presence of resonant behaviour in the three-pion system, the ratio of the branching fractions  $\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm\pi^\pm\pi^\mp)/\mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm)$  is measured to be  $2.55 \pm 0.80 (\text{stat}) \pm 0.33 (\text{syst})_{-0.01}^{+0.04} (\tau_{B_c})$ , consistent with the previous LHCb result.

**KEYWORDS:** Hadron-Hadron Scattering, Branching fraction, B physics

ARXIV EPRINT: [1410.5729](https://arxiv.org/abs/1410.5729)

---

**Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>CMS detector</b>	<b>2</b>
<b>3</b>	<b>Event selection</b>	<b>2</b>
<b>4</b>	<b><math>B_c^+ \rightarrow J/\psi\pi^+</math> and <math>B^+ \rightarrow J/\psi K^+</math> decays</b>	<b>4</b>
<b>5</b>	<b><math>R_{c/u}</math> measurement</b>	<b>4</b>
<b>6</b>	<b><math>B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-</math> decay</b>	<b>6</b>
<b>7</b>	<b><math>R_{B_c}</math> measurement</b>	<b>6</b>
<b>8</b>	<b>Summary</b>	<b>9</b>
	<b>The CMS collaboration</b>	<b>13</b>

---

**1 Introduction**

The pseudoscalar  $B_c^+$  ( $B_c^-$ ) meson, the ground state of the  $\bar{b}c$  ( $b\bar{c}$ ) system, is the lightest particle containing two heavy quarks of different flavors and thus represents a unique laboratory in which to study heavy-quark dynamics. The investigation of the  $B_c^+$  meson properties (charge conjugation is implied throughout this paper) is of special interest compared to the flavor symmetric heavy-quarkonium ( $b\bar{b}$ ,  $c\bar{c}$ ) states, and provides a new testing ground for predictions in the context of effective models inspired by quantum chromodynamics [1]. The decay processes of the  $B_c^+$  meson can be generically divided into three classes: those involving the decay of the b quark, the decay of the c quark, and the annihilation of the b and c quarks [1–3]. The  $b \rightarrow c$  transition, accounting for about 20% of the decay rate [1], offers an easily accessible experimental signature, having a high probability of producing a  $J/\psi$  meson. Consequently, the first  $B_c^+$  observation was made in the semileptonic channel  $B_c^+ \rightarrow J/\psi\ell^+\nu$  ( $\ell = e, \mu$ ) by the CDF Collaboration [4].

Information on the production and decays of the  $B_c^+$  meson and its possible excited states is quite limited. Production characteristics have been studied at the Tevatron for  $p_T > 4$  GeV and  $|y| < 1$  and by LHCb in the kinematic region  $p_T > 4$  GeV,  $2.5 < \eta < 4.5$ .

The advent of the CERN LHC has opened a new era for  $B_c^+$  investigations; a rich program of measurements involving new decay modes is being carried out by the LHCb Collaboration [5–11]. The ATLAS experiment has recently observed a new state whose mass is consistent with the predicted mass for the second S-wave state of the  $B_c^+$  meson [12]. The CMS experiment, owing to its excellent muon identification system and

tracking detectors, is particularly well suited to the study of final states containing  $J/\psi$  mesons, where  $J/\psi \rightarrow \mu^+\mu^-$ .

In this Letter, the  $B_c^+$  production cross section relative to that of  $B^+$  at 7 TeV center of mass energy is presented in a kinematic region complementary to that accessible to LHCb. This is done by comparing the  $B_c^+ \rightarrow J/\psi\pi^+$  mode to  $B^+ \rightarrow J/\psi K^+$ , which has a similar vertex topology. The ratio of their production cross sections times branching fractions  $R_{c/u} \equiv (\sigma(B_c^+)\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+))/(\sigma(B^+)\mathcal{B}(B^+ \rightarrow J/\psi K^+))$  is measured for  $B_c^+$  transverse momentum  $p_T > 15$  GeV and in the central rapidity region,  $|y| < 1.6$ . This measurement contributes to a more complete understanding of  $B_c^+$  production in pp collisions.

We also observe the decay  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$ , previously measured by LHCb [10]. The three-pion system exhibits resonant structure that is not well-determined, but can affect the reconstruction efficiency. A model-independent efficiency correction method is developed and implemented in this analysis, which can be considered for future high-statistics analyses of multibody decays. The ratio of the branching fractions  $R_{B_c} \equiv \mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-)/\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)$ , which is independent of the poorly known  $B_c^+$  production cross section, is also presented; our measurement provides the first confirmation of the LHCb Collaboration's result [10].

## 2 CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter and 3.8 T field. Within the superconducting solenoid volume are silicon pixel and strip trackers, a crystal electromagnetic calorimeter, and a brass/scintillator hadron calorimeter. Muons are measured in gas-ionization detectors embedded in the steel return yoke. The subdetectors relevant for this analysis are the silicon tracker and the muon systems. The inner tracker measures charged particles within the pseudorapidity range  $|\eta| < 2.5$ . It consists of layers totaling 66 million  $100 \times 150 \mu\text{m}^2$  silicon pixels and 9.6 million silicon strips with pitches ranging from 80 to  $183 \mu\text{m}$ . Muons are measured in the pseudorapidity range  $|\eta| < 2.4$  with detection planes constructed using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. The first level of the CMS trigger system, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select the most interesting events in a fixed time interval of less than  $4 \mu\text{s}$ . The high level trigger processor farm further decreases the event rate from around 100 kHz to approximately 400 Hz before data storage. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in ref. [13].

## 3 Event selection

This analysis is based on pp data collected with the CMS detector at a center-of-mass energy of 7 TeV in 2011. Events selected with displaced-vertex dimuon triggers are considered, corresponding to an integrated luminosity of  $5.1 \text{ fb}^{-1}$ . The analysis is driven by the  $J/\psi$  meson reconstruction. The dimuon triggers apply topological and kinematic selections

on dimuon candidates:  $\cos \alpha > 0.9$ , where  $\alpha$  is the pointing angle, in the transverse plane, between the dimuon momentum and the direction from the mean pp collision position (beam spot) to the dimuon vertex;  $L_{xy}/\sigma_{xy} > 3$ , where  $L_{xy}$  is the transverse distance between the beam spot and the dimuon vertex, and  $\sigma_{xy}$  is the corresponding uncertainty. In addition, the two muons must have opposite charges, dimuon  $p_T > 6.9$  GeV, satisfy an invariant mass requirement  $2.9 < m(\mu^+\mu^-) < 3.3$  GeV, and have a mutual distance of closest approach in the transverse plane of less than 0.5 cm. Trigger selection requirements on the  $\chi^2$  probability of the dimuon kinematic fit ( $P_{\text{VTX}}$ ) and muon transverse momentum  $p_T(\mu)$  were made more restrictive as the luminosity increased and ranged from  $P_{\text{VTX}} > 0.5\%$  to 15% and up to  $p_T(\mu) > 4$  GeV, respectively. A requirement on the muon pseudorapidity,  $|\eta(\mu)| < 2.2$ , is also applied.

Monte Carlo (MC) simulations are employed to design the offline selection, assess the reconstruction efficiency, and study systematic effects. The  $B_c^+$  signal events are simulated using a dedicated generator (BCVEGPY) [14, 15] interfaced with the PYTHIA simulation program (version 6.424, Z2 tune [16, 17]), which hadronizes the whole event. Unstable particle decays are simulated with EVTGEN [18] and the detector response with GEANT4 [19].

The offline selection starts from  $J/\psi$  candidates reconstructed from pairs of oppositely charged muons. Muons are identified through the standard CMS muon reconstruction procedure [20] and are required to have a track matched with at least one muon segment, a track fit  $\chi^2$  per degree of freedom less than 1.8, at least 11 hits in the tracker with at least 2 from the pixel detector, and a transverse (longitudinal) impact parameter less than 3 cm (30 cm). Offline requirements on the dimuon pair are tightened, with respect to those of the trigger, requiring a dimuon  $p_T > 7.1$  GeV and  $L_{xy}/\sigma_{xy} \geq 5$ .

The  $B_c^+ \rightarrow J/\psi\pi^+$  ( $B^+ \rightarrow J/\psi K^+$ ) candidates are formed by combining a  $J/\psi$  candidate with one track, assuming that it is a pion (kaon). The track must not be identified as a muon. The  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  candidates are analogously formed by combining a  $J/\psi$  candidate with three tracks, assuming that they are pions and requiring that the total charge is +1. The pion (kaon) candidates are required to have a track fit  $\chi^2$  less than three times the number of degrees of freedom;  $\geq 6$  tracker hits;  $\geq 2$  pixel hits;  $|\eta| < 2.4$ ; and  $p_T > 0.9$  GeV. The three-dimensional impact parameter between each pion (kaon) and the  $J/\psi$  vertex is required to be less than 6 times its uncertainty to reduce combinatorial background and the effect of the number of simultaneous pp interactions per bunch crossing (pileup). The decay vertex is reconstructed using a kinematic vertex fit [21], which constrains the invariant mass of the two muons to the nominal  $J/\psi$  mass. After the vertex fit, the track parameters are re-estimated at the fitted vertex to improve their resolution. To reduce backgrounds, only the highest-transverse-momentum  $B_c^+$  or  $B^+$  candidate is retained per event. This method has been studied using the MC samples and found to select the right candidate in 99.3% (99.2%) of the  $B_c^+ \rightarrow J/\psi\pi^+$  ( $B^+ \rightarrow J/\psi K^+$ ) events and 91% of the  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  events. In the  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  data sample, this requirement reduces the background by more than a factor of four.

Additional topological requirements are made to improve the signal-to-background ratio, as discussed below.

## 4 $B_c^+ \rightarrow J/\psi\pi^+$ and $B^+ \rightarrow J/\psi K^+$ decays

The selection criteria for the  $B_c^+ \rightarrow J/\psi\pi^+$  decay have been optimized in the kinematic region  $p_T > 15 \text{ GeV}$  and  $|y| < 1.6$  by maximizing  $S/\sqrt{(S+B)}$  as a figure of merit, where  $S$  is the signal yield obtained from a Gaussian fit to the MC reconstructed events and  $B$  is the amount of background extrapolated from the  $J/\psi\pi^+$  invariant mass sidebands in the data. The two sideband regions are defined as being between  $5\sigma_{m(B_c)}$  and  $8\sigma_{m(B_c)}$  of the world-average  $B_c$  mass [22], where  $\sigma_{m(B_c)}$  is the resolution of the signal as determined in simulation.


The procedure results in the following requirements:  $B_c^+$  vertex probability  $>6\%$ ,  $\cos\alpha' > 0.9$ , where  $\alpha'$  is the angle between the candidate  $B_c^+$  momentum vector and the displacement between the beam spot and the decay vertex evaluated in the plane transverse to the beam;  $p_T(\pi) > 2.7 \text{ GeV}$ , and  $\Delta R(J/\psi, \pi) < 1$ , where  $\Delta R$  is the distance in the  $(\eta, \phi)$  plane between the  $J/\psi$  and pion momentum vector. The  $B_c^+ \rightarrow J/\psi\pi^+$  invariant mass distribution is shown in figure 1 (left). The  $B^+ \rightarrow J/\psi K^+$  signal is obtained with the same selections and is shown in figure 1 (right). The  $B_c^+ \rightarrow J/\psi\pi^+$  and the  $B^+ \rightarrow J/\psi K^+$  invariant mass distributions are fit with an unbinned maximum likelihood estimator. The  $B_c^+$  signal is fit with a Gaussian distribution and the background with a second-order Chebyshev polynomial. The  $B_c^+ \rightarrow J/\psi\pi^+$  signal has a yield of  $176 \pm 19$ , a mass of  $6.267 \pm 0.003 \text{ GeV}$ , and a resolution of  $0.025 \pm 0.003 \text{ GeV}$  (statistical uncertainties only). Contamination from other  $B_c^+$  decay modes in the  $B_c^+ \rightarrow J/\psi\pi^+$  channel has been investigated. A possible reflection of the Cabibbo-suppressed  $B_c^+ \rightarrow J/\psi K^+$  mode in the  $J/\psi\pi^+$  mass spectrum has been modeled from a simulated sample of  $B_c^+ \rightarrow J/\psi K^+$  events and its contribution constrained using the value of the relative branching fraction to  $J/\psi\pi^+$  [9]. Furthermore, the effect due to a possible undetected  $\pi^0$  from  $B_c^+ \rightarrow J/\psi\pi^+\pi^0$  decay has been modeled from a dedicated MC sample. The partially reconstructed  $J/\psi\pi^+$  mass spectrum obtained from the simulated events has been fit with an ARGUS function [23] convolved with a Gaussian function describing the detector resolution. The resulting parametrization, added to a linear function, has been used to describe the background on the left of the signal peak in the fit of the  $J/\psi\pi^+$  mass spectrum in data. No significant variation of the  $B_c^+ \rightarrow J/\psi\pi^+$  signal yield is found.

The  $B^+$  invariant mass distribution is fit with a sum of two Gaussian distributions with a common mean for the signal and a second-order Chebyshev polynomial for the background. Additional contributions from partially reconstructed  $B^0$  and  $B^+$  decays are parametrized with functions determined from inclusive  $B^+ \rightarrow J/\psi X$  and  $B^0 \rightarrow J/\psi X$  MC samples.

## 5 $R_{c/u}$ measurement

The ratio  $R_{c/u}$  of the production cross sections times branching fractions is obtained from the relation

$$R_{c/u} = \frac{\sigma(B_c^+) \mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)}{\sigma(B^+) \mathcal{B}(B^+ \rightarrow J/\psi K^+)} = \frac{Y_{B_c^+ \rightarrow J/\psi\pi^+}}{Y_{B^+ \rightarrow J/\psi K^+}}, \quad (5.1)$$



**Figure 1.** The  $J/\psi\pi^+$  (left) and  $J/\psi K^+$  (right) invariant mass distributions. The result of the fit is superimposed. The lines represent the signal-plus-background fit (solid) and the background-only component (dashed).

where  $Y_{B_c^+ \rightarrow J/\psi\pi^+}$  and  $Y_{B^+ \rightarrow J/\psi K^+}$  are the signal yields extracted from the efficiency-corrected invariant mass distributions for the  $B_c^+ \rightarrow J/\psi\pi^+$  and  $B^+ \rightarrow J/\psi K^+$  channels, respectively, in the kinematic region  $p_T > 15 \text{ GeV}$  and  $|y| < 1.6$ . The efficiencies for the two channels are evaluated from MC simulations and include geometrical acceptance, reconstruction, selection, and trigger effects.

The simulation of the two-body  $B_c^+ \rightarrow J/\psi\pi^+$  and  $B^+ \rightarrow J/\psi K^+$  decays takes into account the spins of the particles. The efficiencies are evaluated as a function of the  $B^+$  or  $B_c^+$  candidate's  $p_T$  and computed in  $p_T$  bins, whose sizes are determined by the available size of the  $B_c^+ \rightarrow J/\psi\pi^+$  and  $B^+ \rightarrow J/\psi K^+$  MC samples. Data are corrected event-by-event according to the candidate's  $p_T$ .

Possible systematic uncertainties introduced by different trigger and pileup conditions and analysis selections have been investigated by dividing the data and evaluating the statistical consistency [22] of the independent samples; the resulting systematic uncertainties are found to be insignificant. Uncertainties from the different signal and background fit functions and fit ranges have been evaluated through a “fit variant” approach [24] and account for a 5.3% uncertainty. The finite size of the MC samples introduces a systematic uncertainty of 2.1% and the choice of the  $p_T$  binning in the efficiency calculation an additional 3.1%. The total systematic uncertainty in the ratio is 6.5%.

Recently, the LHCb Collaboration published a new, more precise  $B_c^+$  lifetime measurement [25], which is significantly higher than the previous world average [22]. The  $B_c^+ \rightarrow J/\psi\pi^+$  reconstruction efficiency has a dependence on the  $B_c^+$  lifetime. To determine the systematic uncertainty associated with the uncertainty in the  $B_c^+$  lifetime, the efficiency is evaluated while changing the  $B_c^+$  lifetime in the simulation to cover the range from the world average minus its one standard deviation uncertainty, to the new LHCb measurement. The resulting variation in the  $R_{c/u}$  ratio is quoted separately as a lifetime systematic uncertainty ( $\sigma(\tau_{B_c})$ ) and is  $\pm 10.4\%$ . The different contributions to the systematic uncertainty are listed in table 1.

Systematic source	%
Fit variant	5.3
MC sample size	2.1
Efficiency binning	3.1
Total uncertainty	6.5
$B_c$ lifetime	10.4

**Table 1.** Systematic uncertainties in the measurement of  $R_{c/u}$ .

The measurement of the ratio, including all the uncertainties, is

$$R_{c/u} = [0.48 \pm 0.05 \text{ (stat)} \pm 0.03 \text{ (syst)} \pm 0.05 \text{ } (\tau_{B_c})] \% . \quad (5.2)$$

## 6 $B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-$ decay


The same figure of merit  $S/\sqrt{(S+B)}$  is maximized in the selection of the  $B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-$  signal in the same kinematic phase space as defined for the  $B_c^+ \rightarrow J/\psi \pi^+$  decay, i.e.,  $p_T(B_c^+) > 15 \text{ GeV}$  and  $|y(B_c^+)| < 1.6$ . The optimized selection requirements are:  $\chi^2$  probability of the five-track kinematic fit  $> 20\%$ ;  $\cos \alpha' > 0.99$ ;  $p_T(\pi_1) > 2.5 \text{ GeV}$ ;  $p_T(\pi_2) > 1.7 \text{ GeV}$ ;  $p_T(\pi_3) > 0.9 \text{ GeV}$ , where the three pions are referred to as  $\pi_1$ ,  $\pi_2$ , and  $\pi_3$  from highest to lowest  $p_T$ ; and  $\Delta R(J/\psi, \pi_S) < 0.5$ , where  $\pi_S$  is the sum of the momentum vectors of the three pions. The resulting  $B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-$  invariant mass distribution is shown in figure 2. A fit is performed with an unbinned maximum likelihood estimator. The signal is parametrized as a Gaussian distribution and the background as a second-order Chebyshev polynomial. The signal yield is  $92 \pm 27$  events and the fitted mass and resolution values are  $6.266 \pm 0.006 \text{ GeV}$  and  $0.021 \pm 0.006 \text{ GeV}$ , respectively, where the uncertainties are statistical only. Possible contamination from other  $B_c^+$  decay modes in the  $B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-$  channel has been investigated. No  $B_c^+ \rightarrow J/\psi K^+ K^- \pi^+$  decays are observed in the data with the applied selection cuts. The effect from a possible undetected  $\pi^0$  in the decay  $B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^- \pi^0$  has been modeled with a dedicated MC sample. The partially reconstructed  $J/\psi \pi^+ \pi^+ \pi^-$  mass spectrum obtained from the simulated events has been fit with an ARGUS function convolved with a Gaussian function describing the detector resolution. The resulting parametrization, added to a linear polynomial function, has been used to describe the background on the left of the signal peak in the fit of the  $J/\psi \pi^+ \pi^+ \pi^-$  mass spectrum in data. No significant variation in the  $B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-$  signal yield is found.

## 7 $R_{B_c}$ measurement

The ratio  $R_{B_c}$  is defined as

$$R_{B_c} = \frac{\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-)}{\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)} = \frac{Y_{B_c^+ \rightarrow J/\psi \pi^+ \pi^+ \pi^-}}{Y_{B_c^+ \rightarrow J/\psi \pi^+}}, \quad (7.1)$$






**Figure 2.** The  $J/\psi\pi^+\pi^+\pi^-$  invariant mass distribution. The result of the fit is superimposed; the lines represent the signal-plus-background fit (solid) and the background-only component (dashed).

where  $Y_{B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-}$  and  $Y_{B_c^+ \rightarrow J/\psi\pi^+}$  are the signal yields extracted from the efficiency-corrected invariant mass distributions for the  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  and  $B_c^+ \rightarrow J/\psi\pi^+$  channels, respectively, in the kinematic region  $p_T > 15$  GeV and  $|y| < 1.6$ . Efficiency corrections of the  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  and  $B_c^+ \rightarrow J/\psi\pi^+$  data include geometrical acceptance, reconstruction, selection, and trigger effects. The efficiencies for the two channels are evaluated from MC simulations.

The efficiency for the  $B_c^+ \rightarrow J/\psi\pi^+$  channel is evaluated as a function of the candidate’s  $p_T$ , as explained in section 5.

The  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  decay can involve intermediate resonant states; indeed, the  $\pi^+\pi^+\pi^-$  and  $\pi^+\pi^-$  invariant mass projections from data show evidence for the presence of  $a_1(1260)$  and  $\rho(770)$  in the decay (figure 3). No hint of either  $\psi(2S)(\rightarrow J/\psi\pi^+\pi^-)$  or  $X(3872)(\rightarrow J/\psi\pi^+\pi^-)$  is detected in the  $\mu^+\mu^-\pi^+\pi^-$  mass projections. The quantitative determination of the resonant contributions and their interferences in the decay requires a sophisticated amplitude analysis which is not feasible with the available amount of data. However, the reconstruction efficiency for this five-body decay could be affected by the decay dynamics; thus, a model-independent efficiency treatment is needed.

A five-body decay of a spinless particle can be fully described in its center-of-mass frame by eight independent mass combinations of the type  $m_{ij}^2$  ( $i \neq j$ ), where  $m_{ij}^2$  is the squared invariant mass of the pair of particles  $i$  and  $j$  in the final state (Dalitz plot representation). In the present case, the additional  $J/\psi$  mass constraint reduces the number of independent  $m_{ij}^2$  to seven. The following seven mass combinations have been chosen:  $x = m^2(\mu^+\pi^+)_{\text{low}}$ ,  $y = m^2(\pi^+\pi^-)_{\text{high}}$ ,  $z = m^2(\mu^+\pi^-)$ ,  $w = m^2(\pi^+\pi^+)$ ,  $r = m^2(\mu^-\pi^+)_{\text{low}}$ ,  $t = m^2(\mu^-\pi^+)_{\text{high}}$ , and  $v = m^2(\mu^-\pi^-)$ ; the “low” and “high” subscripts refer to the lower and higher invariant mass combination where a  $\pi^+$  is involved. A  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  nonresonant MC has been produced to access all the phase-space configurations. The efficiency is parametrized as a



**Figure 3.** Background-subtracted invariant mass projections for  $\pi^+\pi^+\pi^-$  (left),  $(\pi^+\pi^-)_{\text{high}}$  (center), and  $(\pi^+\pi^-)_{\text{low}}$  (right) from the  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  candidate events. Since two same-sign pions are present in the final state, we indicate with  $(\pi^+\pi^-)_{\text{low}}$  the  $\pi^+\pi^-$  pair with the lower invariant mass and with  $(\pi^+\pi^-)_{\text{high}}$  the higher invariant mass combination.

linear function of these seven mass combinations:

$$\epsilon = |p_0 + p_1 \cdot x + p_2 \cdot y + p_3 \cdot z + p_4 \cdot w + p_5 \cdot r + p_6 \cdot t + p_7 \cdot v|, \quad (7.2)$$

where  $p_i$  are free parameters to be determined via an unbinned maximum likelihood fit to the generated events in the seven-dimensional space using a binomial probability density function. The absolute value is required to prevent the function from assuming negative values. The resulting efficiency function is used to weight the data event-by-event. The data efficiency-corrected invariant mass for the  $B_c^+ \rightarrow J/\psi\pi^+$  and  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  channels is fit with a function consisting of a Gaussian distribution for the signal and a second-order Chebyshev polynomial for the background. An unbinned maximum likelihood estimator is used to extract the  $Y_{B_c^+ \rightarrow J/\psi\pi^+}$  and  $Y_{B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-}$  yields. The resulting measurement of the branching fraction ratio is  $R_{B_c} = 2.55 \pm 0.80$ , where the uncertainty is statistical only.

The same sources of systematic uncertainties considered in section 5 have been evaluated for this measurement as well; the biggest uncertainty comes from the fit variant and accounts for a 9.4 % effect. An additional contribution is considered coming from the choice of the seven-dimensional efficiency parametrization of eq. (7.2). To estimate this contribution, data are alternatively weighted according to the efficiency distribution directly obtained from the MC samples, binned in the seven two-body submasses. The difference between the ratio measured using the binned efficiency distribution and the function in eq. (7.2) is taken as a systematic uncertainty (1.0%). In the evaluation of  $R_{B_c}$ , two different multiplicity final states are compared. Assuming a tracking efficiency uncertainty for each pion track of 3.9% [26], a global 7.8% uncertainty is included in the final systematic evaluation. The total systematic uncertainty in the ratio, obtained by adding all the contributions in quadrature, is 13.1%.

The systematic uncertainty from the  $B_c^+$  lifetime uncertainty is quoted separately in the ratio and is  $(\sigma(\tau_{B_c})) = {}^{+1.6}_{-0.4}\%$ . The sources of systematic uncertainty are summarized in table 2.

Systematic source	%
Fit variant	9.4
MC sample size	4.1
Efficiency fit function	1.0
Efficiency binning	1.9
Tracking efficiency	7.8
Total uncertainty	13.1
Lifetime	+1.6 -0.4

**Table 2.** Systematic uncertainties in the measurement of  $R_{B_c}$ .

The resulting ratio, including all uncertainties, is

$$R_{B_c} = 2.55 \pm 0.80 \text{ (stat)} \pm 0.33 \text{ (syst)}^{+0.04}_{-0.01} (\tau_{B_c}). \quad (7.3)$$

## 8 Summary

A measurement of the ratio of the cross sections times branching fractions for  $B_c^+ \rightarrow J/\psi\pi^+$  and  $B^+ \rightarrow J/\psi K^+$  has been presented based on pp collision data at a center-of-mass energy of 7 TeV collected by the CMS experiment and corresponding to an integrated luminosity of  $5.1 \text{ fb}^{-1}$ . The analysis, performed for  $B_c^+$  and  $B^+$  mesons with  $p_T > 15 \text{ GeV}$  and in the central rapidity region  $|y| < 1.6$ , gives a measured ratio of

$$R_{c/u} = [0.48 \pm 0.05 \text{ (stat)} \pm 0.03 \text{ (syst)} \pm 0.05 (\tau_{B_c})]\%. \quad (8.1)$$

A similar measurement from LHCb in the kinematic region  $p_T > 4 \text{ GeV}$ ,  $2.5 < \eta < 4.5$  gives  $[0.68 \pm 0.10 \text{ (stat)} \pm 0.03 \text{ (syst)} \pm 0.05 (\tau_{B_c})]\%$  [27]. The two measurements, performed in different kinematic regions, are expected to differ because of the softer  $p_T$  distribution of the  $B_c^+$  with respect to that of the  $B^+$ , implying a lower value of the ratio at higher  $p_T$ . The measurements are consistent with this expectation. Measurements of the production cross section times branching fraction for  $B_c^+ \rightarrow J/\psi\ell^+\nu$  relative to that for  $B^+ \rightarrow J/\psi K^+$  are also available from the CDF experiment [4] in the kinematic region  $p_T > 4 \text{ GeV}$  and  $|y| < 1$ . With the present  $B_c^+$  ( $p_T, |y|$ ) coverage, these experimental results can give guidance to improve the theoretical calculations still affected by large uncertainties and constrain the various  $B_c^+$  production models.

The ratio of the  $B_c^+ \rightarrow J/\psi\pi^+\pi^+\pi^-$  and  $B_c^+ \rightarrow J/\psi\pi^+$  branching fractions has been measured to be

$$R_{B_c} = 2.55 \pm 0.80 \text{ (stat)} \pm 0.33 \text{ (syst)}^{+0.04}_{-0.01} (\tau_{B_c}), \quad (8.2)$$

which is in good agreement with the result from the LHCb experiment,  $2.41 \pm 0.30 \text{ (stat)} \pm 0.33 \text{ (syst)}$  [10], and represents its first confirmation. This measurement can be compared with the theoretical predictions, which assume factorization into  $B_c^+ \rightarrow J/\psi W^{+*}$  and  $W^{+*} \rightarrow n\pi^+$  ( $n = 1, 2, 3, 4$ ). In particular, ref. [28] predicts 1.5 for the ratio, whereas

ref. [29] predicts three different values, 1.9, 2.0, and 2.3, depending on the chosen set of  $B_c^+$  meson form factors. More precise measurements are needed to determine if one of the predictions is favored by the data. The model-independent method implemented for the efficiency evaluation of the five-body final state can be considered in future high-statistics analyses to reduce systematic uncertainties associated with the unknown multibody decay dynamics.

## Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC and the CMS detector provided by the following funding agencies: BMFWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES and CSF (Croatia); RPF (Cyprus); MoER, ERC IUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); OTKA and NIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); NRF and WCU (Republic of Korea); LAS (Lithuania); MOE and UM (Malaysia); CINVESTAV, CONACYT, SEP, and UASLP-FAI (Mexico); MBIE (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Dubna); MON, RosAtom, RAS and RFBR (Russia); MESTD (Serbia); SEIDI and CPAN (Spain); Swiss Funding Agencies (Switzerland); MST (Taipei); ThEPCenter, IPST, STAR and NSTDA (Thailand); TUBITAK and TAEK (Turkey); NASU and SFFR (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie programme and the European Research Council and EPLANET (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Council of Science and Industrial Research, India; the HOMING PLUS programme of Foundation for Polish Science, cofinanced from European Union, Regional Development Fund; the Compagnia di San Paolo (Torino); the Consorzio per la Fisica (Trieste); MIUR project 20108T4XTM (Italy); the Thalís and Aristeia programmes cofinanced by EU-ESF and the Greek NSRF; and the National Priorities Research Program by Qatar National Research Fund.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] QUARKONIUM WORKING GROUP collaboration, N. Brambilla et al., *Heavy quarkonium physics*, [hep-ph/0412158](#) [[INSPIRE](#)].
- [2] I.P. Gouz et al., *Prospects for the  $B_c$  studies at LHCb*, *Phys. Atom. Nucl.* **67** (2004) 1559 [[hep-ph/0211432](#)] [[INSPIRE](#)].
- [3] H.-M. Choi and C.-R. Ji, *Non-leptonic two-body decays of the  $B_c$  meson in light-front quark model and QCD factorization approach*, *Phys. Rev. D* **80** (2009) 114003 [[arXiv:0909.5028](#)] [[INSPIRE](#)].
- [4] CDF collaboration, F. Abe et al., *Observation of the  $B_c$  meson in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, *Phys. Rev. Lett.* **81** (1998) 2432 [[hep-ex/9805034](#)] [[INSPIRE](#)].
- [5] LHCb collaboration, *Observation of the decay  $B_c \rightarrow J/\psi K^+ K^- \pi^+$* , *JHEP* **11** (2013) 094 [[arXiv:1309.0587](#)] [[INSPIRE](#)].
- [6] LHCb collaboration, *Observation of the decay  $B_c^+ \rightarrow B_s^0 \pi^+$* , *Phys. Rev. Lett.* **111** (2013) 181801 [[arXiv:1308.4544](#)] [[INSPIRE](#)].
- [7] LHCb collaboration, *Observation of  $B_c^+ \rightarrow J/\psi D_s^+$  and  $B_c^+ \rightarrow J/\psi D_s^{*+}$  decays*, *Phys. Rev. D* **87** (2013) 112012 [[arXiv:1304.4530](#)] [[INSPIRE](#)].
- [8] LHCb collaboration, *Observation of the decay  $B_c^+ \rightarrow \psi(2S)\pi^+$* , *Phys. Rev. D* **87** (2013) 071103 [[arXiv:1303.1737](#)] [[INSPIRE](#)].
- [9] LHCb collaboration, *First observation of the decay  $B_c^+ \rightarrow J/\psi K^+$* , *JHEP* **09** (2013) 075 [[arXiv:1306.6723](#)] [[INSPIRE](#)].
- [10] LHCb collaboration, *First observation of the decay  $B_c^+ \rightarrow J/\psi \pi^+ \pi^- \pi^+$* , *Phys. Rev. Lett.* **108** (2012) 251802 [[arXiv:1204.0079](#)] [[INSPIRE](#)].
- [11] LHCb collaboration, *Evidence for the decay  $B_c^+ \rightarrow J/\psi 3\pi^+ 2\pi^-$* , *JHEP* **05** (2014) 148 [[arXiv:1404.0287](#)] [[INSPIRE](#)].
- [12] ATLAS collaboration, *Observation of an excited  $B_c^\pm$  meson state with the ATLAS detector*, *Phys. Rev. Lett.* **113** (2014) 212004 [[arXiv:1407.1032](#)] [[INSPIRE](#)].
- [13] CMS collaboration, *The CMS experiment at the CERN LHC, 2008* *JINST* **3** S08004 [[INSPIRE](#)].
- [14] C.-H. Chang, C. Driouiichi, P. Eerola and X.G. Wu, *BCVEGPY: an event generator for hadronic production of the  $B_c$  meson*, *Comput. Phys. Commun.* **159** (2004) 192 [[hep-ph/0309120](#)] [[INSPIRE](#)].
- [15] C.-H. Chang, J.-X. Wang and X.-G. Wu, *BCVEGPY2.0: an upgraded version of the generator BCVEGPY with the addition of hadroproduction of the P-wave  $B_c$  states*, *Comput. Phys. Commun.* **174** (2006) 241 [[hep-ph/0504017](#)] [[INSPIRE](#)].
- [16] T. Sjöstrand, S. Mrenna and P.Z. Skands, *PYTHIA 6.4 physics and manual*, *JHEP* **05** (2006) 026 [[hep-ph/0603175](#)] [[INSPIRE](#)].
- [17] R. Field, *Early LHC underlying event data — Findings and surprises*, [arXiv:1010.3558](#) [[INSPIRE](#)].
- [18] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].

- [19] GEANT4 collaboration, V.N. Ivanchenko, *Geant4 toolkit for simulation of HEP experiments*, *Nucl. Instrum. Meth. A* **502** (2003) 666 [INSPIRE].
- [20] CMS collaboration, *Performance of CMS muon reconstruction in pp collision events at  $\sqrt{s} = 7$  TeV*, 2012 *JINST* **7** P10002 [arXiv:1206.4071] [INSPIRE].
- [21] G.E. Forden and D.H. Saxon, *Improving vertex position determination using a kinematic fit*, *Nucl. Instrum. Meth. A* **248** (1986) 439 [INSPIRE].
- [22] PARTICLE DATA GROUP collaboration, J. Beringer et al., *Review of particle physics*, *Phys. Rev. D* **86** (2012) 010001 [INSPIRE].
- [23] ARGUS collaboration, H. Albrecht et al., *Search for hadronic  $b \rightarrow u$  decays*, *Phys. Lett. B* **241** (1990) 278 [INSPIRE].
- [24] FOCUS collaboration, J.M. Link et al., *Study of the Cabibbo suppressed decay modes  $D^0 \rightarrow \pi^- \pi^+$  and  $D^0 \rightarrow K^- K^+$* , *Phys. Lett. B* **555** (2003) 167 [hep-ex/0212058] [INSPIRE].
- [25] LHCb collaboration, *Measurement of the  $B_c^+$  meson lifetime using  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu X$  decays*, *Eur. Phys. J. C* **74** (2014) 2839 [arXiv:1401.6932] [INSPIRE].
- [26] CMS collaboration, *Measurement of tracking efficiency*, CMS-PAS-TRK-10-002 (2010).
- [27] LHCb collaboration, *Measurements of  $B_c^+$  production and mass with the  $B_c^+ \rightarrow J/\psi \pi^+$  decay*, *Phys. Rev. Lett.* **109** (2012) 232001 [arXiv:1209.5634] [INSPIRE].
- [28] A. Rakitin and S. Koshkarev, *Hadronic  $B_c$  decays as a test of  $B_c$  cross section*, *Phys. Rev. D* **81** (2010) 014005 [arXiv:0911.3287] [INSPIRE].
- [29] A.K. Likhoded and A.V. Luchinsky, *Light hadron production in  $B_c \rightarrow J/\psi + X$  decays*, *Phys. Rev. D* **81** (2010) 014015 [arXiv:0910.3089] [INSPIRE].

**The CMS collaboration****Yerevan Physics Institute, Yerevan, Armenia**

V. Khachatryan, A.M. Sirunyan, A. Tumasyan

**Institut für Hochenergiephysik der OeAW, Wien, Austria**

W. Adam, T. Bergauer, M. Dragicevic, J. Erö, C. Fabjan<sup>1</sup>, M. Friedl, R. Frühwirth<sup>1</sup>, V.M. Ghete, C. Hartl, N. Hörmann, J. Hrubec, M. Jeitler<sup>1</sup>, W. Kiesenhofer, V. Knünz, M. Krammer<sup>1</sup>, I. Krätschmer, D. Liko, I. Mikulec, D. Rabady<sup>2</sup>, B. Rahbaran, H. Rohringer, R. Schöfbeck, J. Strauss, A. Taurok, W. Treberer-Treberspurg, W. Waltenberger, C.-E. Wulz<sup>1</sup>

**National Centre for Particle and High Energy Physics, Minsk, Belarus**

V. Mossolov, N. Shumeiko, J. Suarez Gonzalez

**Universiteit Antwerpen, Antwerpen, Belgium**

S. Alderweireldt, M. Bansal, S. Bansal, T. Cornelis, E.A. De Wolf, X. Janssen, A. Knutsson, S. Luyckx, S. Ochesanu, R. Rougny, M. Van De Klundert, H. Van Haevermaet, P. Van Mechelen, N. Van Remortel, A. Van Spilbeeck

**Vrije Universiteit Brussel, Brussel, Belgium**

F. Blekman, S. Blyweert, J. D'Hondt, N. Daci, N. Heracleous, J. Keaveney, S. Lowette, M. Maes, A. Olbrechts, Q. Python, D. Strom, S. Tavernier, W. Van Doninck, P. Van Mulders, G.P. Van Onsem, I. Villella

**Université Libre de Bruxelles, Bruxelles, Belgium**

C. Caillol, B. Clerbaux, G. De Lentdecker, D. Dobur, L. Favart, A.P.R. Gay, A. Grebenyuk, A. Léonard, A. Mohammadi, L. Pernie<sup>2</sup>, T. Reis, T. Seva, L. Thomas, C. Vander Velde, P. Vanlaer, J. Wang, F. Zenoni

**Ghent University, Ghent, Belgium**

V. Adler, K. Beernaert, L. Benucci, A. Cimmino, S. Costantini, S. Crucy, S. Dildick, A. Fagot, G. Garcia, J. Mccartin, A.A. Ocampo Rios, D. Ryckbosch, S. Salva Diblen, M. Sigamani, N. Strobbe, F. Thyssen, M. Tytgat, E. Yazgan, N. Zaganidis

**Université Catholique de Louvain, Louvain-la-Neuve, Belgium**

S. Basegmez, C. Beluffi<sup>3</sup>, G. Bruno, R. Castello, A. Caudron, L. Ceard, G.G. Da Silveira, C. Delaere, T. du Pree, D. Favart, L. Forthomme, A. Giammanco<sup>4</sup>, J. Hollar, A. Jafari, P. Jez, M. Komm, V. Lemaitre, C. Nuttens, D. Pagano, L. Perrini, A. Pin, K. Piotrkowski, A. Popov<sup>5</sup>, L. Quertenmont, M. Selvaggi, M. Vidal Marono, J.M. Vizan Garcia

**Université de Mons, Mons, Belgium**

N. Belyi, T. Caebergs, E. Daubie, G.H. Hammad

**Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil**

W.L. Aldá Júnior, G.A. Alves, L. Brito, M. Correa Martins Junior, T. Dos Reis Martins, C. Mora Herrera, M.E. Pol

**Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil**

W. Carvalho, J. Chinellato<sup>6</sup>, A. Custódio, E.M. Da Costa, D. De Jesus Damiao, C. De Oliveira Martins, S. Fonseca De Souza, H. Malbouisson, D. Matos Figueiredo, L. Mundim, H. Nogima, W.L. Prado Da Silva, J. Santaolalla, A. Santoro, A. Sznajder, E.J. Tonelli Manganote<sup>6</sup>, A. Vilela Pereira

**Universidade Estadual Paulista <sup>a</sup>, Universidade Federal do ABC <sup>b</sup>, São Paulo, Brazil**

C.A. Bernardes<sup>b</sup>, S. Dogra<sup>a</sup>, T.R. Fernandez Perez Tomei<sup>a</sup>, E.M. Gregores<sup>b</sup>, P.G. Mercadante<sup>b</sup>, S.F. Novaes<sup>a</sup>, Sandra S. Padula<sup>a</sup>

**Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria**

A. Aleksandrov, V. Genchev<sup>2</sup>, P. Iaydjiev, A. Marinov, S. Piperov, M. Rodozov, S. Stoykova, G. Sultanov, M. Vutova

**University of Sofia, Sofia, Bulgaria**

A. Dimitrov, I. Glushkov, R. Hadjiiska, V. Kozhuharov, L. Litov, B. Pavlov, P. Petkov

**Institute of High Energy Physics, Beijing, China**

J.G. Bian, G.M. Chen, H.S. Chen, M. Chen, R. Du, C.H. Jiang, R. Plestina<sup>7</sup>, F. Romeo, J. Tao, Z. Wang

**State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China**

C. Asawatangtrakuldee, Y. Ban, Q. Li, S. Liu, Y. Mao, S.J. Qian, D. Wang, W. Zou

**Universidad de Los Andes, Bogota, Colombia**

C. Avila, L.F. Chaparro Sierra, C. Florez, J.P. Gomez, B. Gomez Moreno, J.C. Sanabria

**University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**

N. Godinovic, D. Lelas, D. Polic, I. Puljak

**University of Split, Faculty of Science, Split, Croatia**

Z. Antunovic, M. Kovac

**Institute Rudjer Boskovic, Zagreb, Croatia**

V. Brigljevic, K. Kadija, J. Luetic, D. Mekterovic, L. Sudic

**University of Cyprus, Nicosia, Cyprus**

A. Attikis, G. Mavromanolakis, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis

**Charles University, Prague, Czech Republic**

M. Bodlak, M. Finger, M. Finger Jr.<sup>8</sup>

**Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**

Y. Assran<sup>9</sup>, A. Ellithi Kamel<sup>10</sup>, M.A. Mahmoud<sup>11</sup>, A. Radi<sup>12,13</sup>



**National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**

M. Kadastik, M. Murumaa, M. Raidal, A. Tiko

**Department of Physics, University of Helsinki, Helsinki, Finland**

P. Eerola, G. Fedi, M. Voutilainen

**Helsinki Institute of Physics, Helsinki, Finland**

J. Härkönen, V. Karimäki, R. Kinnunen, M.J. Kortelainen, T. Lampén, K. Lassila-Perini, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, T. Peltola, E. Tuominen, J. Tuominiemi, E. Tuovinen, L. Wendland

**Lappeenranta University of Technology, Lappeenranta, Finland**

J. Talvitie, T. Tuuva

**DSM/IRFU, CEA/Saclay, Gif-sur-Yvette, France**

M. Besancon, F. Couderc, M. Dejardin, D. Denegri, B. Fabbro, J.L. Faure, C. Favaro, F. Ferri, S. Ganjour, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, E. Locci, J. Malcles, J. Rander, A. Rosowsky, M. Titov

**Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France**

S. Baffioni, F. Beaudette, P. Busson, C. Charlot, T. Dahms, M. Dalchenko, L. Dobrzynski, N. Filipovic, A. Florent, R. Granier de Cassagnac, L. Mastrolorenzo, P. Miné, C. Mironov, I.N. Naranjo, M. Nguyen, C. Ochando, P. Paganini, S. Regnard, R. Salerno, J.B. Sauvan, Y. Sirois, C. Veelken, Y. Yilmaz, A. Zabi

**Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France**

J.-L. Agram<sup>14</sup>, J. Andrea, A. Aubin, D. Bloch, J.-M. Brom, E.C. Chabert, C. Collard, E. Conte<sup>14</sup>, J.-C. Fontaine<sup>14</sup>, D. Gelé, U. Goerlach, C. Goetzmann, A.-C. Le Bihan, P. Van Hove

**Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France**

S. Gadrat

**Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France**

S. Beauceron, N. Beaupere, G. Boudoul<sup>2</sup>, E. Bouvier, S. Brochet, C.A. Carrillo Montoya, J. Chasserat, R. Chierici, D. Contardo<sup>2</sup>, P. Depasse, H. El Mamouni, J. Fan, J. Fay, S. Gascon, M. Gouzevitch, B. Ille, T. Kurca, M. Lethuillier, L. Mirabito, S. Perries, J.D. Ruiz Alvarez, D. Sabes, L. Sgandurra, V. Sordini, M. Vander Donckt, P. Verdier, S. Viret, H. Xiao

**Institute of High Energy Physics and Informatization, Tbilisi State University, Tbilisi, Georgia**

Z. Tsamalaidze<sup>8</sup>

**RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**

C. Autermann, S. Beranek, M. Bontenackels, M. Edelhoff, L. Feld, O. Hindrichs, K. Klein, A. Ostapchuk, A. Perieanu, F. Raupach, J. Sammet, S. Schael, H. Weber, B. Wittmer, V. Zhukov<sup>5</sup>

**RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**

M. Ata, M. Brodski, E. Dietz-Laursonn, D. Duchardt, M. Erdmann, R. Fischer, A. Güth, T. Hebbeker, C. Heidemann, K. Hoepfner, D. Klingebiel, S. Knutzen, P. Kreuzer, M. Merschmeyer, A. Meyer, P. Millet, M. Olschewski, K. Padeken, P. Papacz, H. Reithler, S.A. Schmitz, L. Sonnenschein, D. Teyssier, S. Thüer, M. Weber

**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**

V. Cherepanov, Y. Erdogan, G. Flügge, H. Geenen, M. Geisler, W. Haj Ahmad, A. Heister, F. Hoehle, B. Kargoll, T. Kress, Y. Kuessel, A. Künsken, J. Lingemann<sup>2</sup>, A. Nowack, I.M. Nugent, L. Perchalla, O. Pooth, A. Stahl

**Deutsches Elektronen-Synchrotron, Hamburg, Germany**

I. Asin, N. Bartosik, J. Behr, W. Behrenhoff, U. Behrens, A.J. Bell, M. Bergholz<sup>15</sup>, A. Bethani, K. Borras, A. Burgmeier, A. Cakir, L. Calligaris, A. Campbell, S. Choudhury, F. Costanza, C. Diez Pardos, S. Dooling, T. Dorland, G. Eckerlin, D. Eckstein, T. Eichhorn, G. Flucke, J. Garay Garcia, A. Geiser, P. Gunnellini, J. Hauk, M. Hempel<sup>15</sup>, D. Horton, H. Jung, A. Kalogeropoulos, M. Kasemann, P. Katsas, J. Kieseler, C. Kleinwort, D. Krücker, W. Lange, J. Leonard, K. Lipka, A. Lobanov, W. Lohmann<sup>15</sup>, B. Lutz, R. Mankel, I. Marfin<sup>15</sup>, I.-A. Melzer-Pellmann, A.B. Meyer, G. Mittag, J. Mnich, A. Mussgiller, S. Naumann-Emme, A. Nayak, O. Novgorodova, E. Ntomari, H. Perrey, D. Pitzl, R. Placakyte, A. Raspereza, P.M. Ribeiro Cipriano, B. Roland, E. Ron, M.Ö. Sahin, J. Salfeld-Nebgen, P. Saxena, R. Schmidt<sup>15</sup>, T. Schoerner-Sadenius, M. Schröder, C. Seitz, S. Spannagel, A.D.R. Vargas Trevino, R. Walsh, C. Wissing

**University of Hamburg, Hamburg, Germany**

M. Aldaya Martin, V. Blobel, M. Centis Vignali, A.R. Draeger, J. Erfle, E. Garutti, K. Goebel, M. Görner, J. Haller, M. Hoffmann, R.S. Höing, H. Kirschenmann, R. Klanner, R. Kogler, J. Lange, T. Lapsien, T. Lenz, I. Marchesini, J. Ott, T. Peiffer, N. Pietsch, J. Poehlsen, T. Poehlsen, D. Rathjens, C. Sander, H. Schettler, P. Schleper, E. Schlieckau, A. Schmidt, M. Seidel, V. Sola, H. Stadie, G. Steinbrück, D. Troendle, E. Usai, L. Vanelderren, A. Vanhoefer

**Institut für Experimentelle Kernphysik, Karlsruhe, Germany**

C. Barth, C. Baus, J. Berger, C. Böser, E. Butz, T. Chwalek, W. De Boer, A. Descroix, A. Dierlamm, M. Feindt, F. Frensch, M. Giffels, F. Hartmann<sup>2</sup>, T. Hauth<sup>2</sup>, U. Husemann, I. Katkov<sup>5</sup>, A. Kornmayer<sup>2</sup>, E. Kuznetsova, P. Lobelle Pardo, M.U. Mozer, Th. Müller, A. Nürnberg, G. Quast, K. Rabbertz, F. Ratnikov, S. Röcker, H.J. Simonis, F.M. Stober, R. Ulrich, J. Wagner-Kuhr, S. Wayand, T. Weiler, R. Wolf

**Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece**

G. Anagnostou, G. Daskalakis, T. Gerasis, V.A. Giakoumopoulou, A. Kyriakis, D. Loukas, A. Markou, C. Markou, A. Psallidas, I. Topsis-Giotis

**University of Athens, Athens, Greece**

A. Agapitos, S. Kesisoglou, A. Panagiotou, N. Saoulidou, E. Stiliaris

**University of Ioánnina, Ioánnina, Greece**

X. Aslanoglou, I. Evangelou, G. Flouris, C. Foudas, P. Kokkas, N. Manthos, I. Papadopoulos, E. Paradas

**Wigner Research Centre for Physics, Budapest, Hungary**

G. Bencze, C. Hajdu, P. Hidas, D. Horvath<sup>16</sup>, F. Sikler, V. Veszpremi, G. Vesztergombi<sup>17</sup>, A.J. Zsigmond

**Institute of Nuclear Research ATOMKI, Debrecen, Hungary**

N. Beni, S. Czellar, J. Karancki<sup>18</sup>, J. Molnar, J. Palinkas, Z. Szillasi

**University of Debrecen, Debrecen, Hungary**

P. Raics, Z.L. Trocsanyi, B. Ujvari

**National Institute of Science Education and Research, Bhubaneswar, India**

S.K. Swain

**Panjab University, Chandigarh, India**

S.B. Beri, V. Bhatnagar, R. Gupta, U.Bhawandeep, A.K. Kalsi, M. Kaur, R. Kumar, M. Mittal, N. Nishu, J.B. Singh

**University of Delhi, Delhi, India**

Ashok Kumar, Arun Kumar, S. Ahuja, A. Bhardwaj, B.C. Choudhary, A. Kumar, S. Malhotra, M. Naimuddin, K. Ranjan, V. Sharma

**Saha Institute of Nuclear Physics, Kolkata, India**

S. Banerjee, S. Bhattacharya, K. Chatterjee, S. Dutta, B. Gomber, Sa. Jain, Sh. Jain, R. Khurana, A. Modak, S. Mukherjee, D. Roy, S. Sarkar, M. Sharan

**Bhabha Atomic Research Centre, Mumbai, India**

A. Abdulsalam, D. Dutta, S. Kailas, V. Kumar, A.K. Mohanty<sup>2</sup>, L.M. Pant, P. Shukla, A. Topkar

**Tata Institute of Fundamental Research, Mumbai, India**

T. Aziz, S. Banerjee, S. Bhowmik<sup>19</sup>, R.M. Chatterjee, R.K. Dewanjee, S. Dugad, S. Ganguly, S. Ghosh, M. Guchait, A. Gurtu<sup>20</sup>, G. Kole, S. Kumar, M. Maity<sup>19</sup>, G. Majumder, K. Mazumdar, G.B. Mohanty, B. Parida, K. Sudhakar, N. Wickramage<sup>21</sup>

**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**

H. Bakhshiansohi, H. Behnamian, S.M. Etesami<sup>22</sup>, A. Fahim<sup>23</sup>, R. Goldouzian, M. Khakzad, M. Mohammadi Najafabadi, M. Naseri, S. Paktinat Mehdiabadi, F. Rezaei Hosseinabadi, B. Safarzadeh<sup>24</sup>, M. Zeinali

University College Dublin, Dublin, Ireland

M. Felcini, M. Grunewald

**INFN Sezione di Bari <sup>a</sup>, Università di Bari <sup>b</sup>, Politecnico di Bari <sup>c</sup>, Bari, Italy**

M. Abbrescia<sup>a,b</sup>, L. Barbone<sup>a,b</sup>, C. Calabria<sup>a,b</sup>, S.S. Chhibra<sup>a,b</sup>, A. Colaleo<sup>a</sup>, D. Creanza<sup>a,c</sup>, N. De Filippis<sup>a,c</sup>, M. De Palma<sup>a,b</sup>, L. Fiore<sup>a</sup>, G. Iaselli<sup>a,c</sup>, G. Maggi<sup>a,c</sup>, M. Maggi<sup>a</sup>, S. My<sup>a,c</sup>, S. Nuzzo<sup>a,b</sup>, A. Pompili<sup>a,b</sup>, G. Pugliese<sup>a,c</sup>, R. Radogna<sup>a,b,2</sup>, G. Selvaggi<sup>a,b</sup>, A. Sharma, L. Silvestris<sup>a,2</sup>, R. Venditti<sup>a,b</sup>, G. Zito<sup>a</sup>

**INFN Sezione di Bologna <sup>a</sup>, Università di Bologna <sup>b</sup>, Bologna, Italy**

G. Abbiendi<sup>a</sup>, A.C. Benvenuti<sup>a</sup>, D. Bonacorsi<sup>a,b</sup>, S. Braibant-Giacomelli<sup>a,b</sup>, L. Brigliadori<sup>a,b</sup>, R. Campanini<sup>a,b</sup>, P. Capiluppi<sup>a,b</sup>, A. Castro<sup>a,b</sup>, F.R. Cavallo<sup>a</sup>, G. Codispoti<sup>a,b</sup>, M. Cuffiani<sup>a,b</sup>, G.M. Dallavalle<sup>a</sup>, F. Fabbri<sup>a</sup>, A. Fanfani<sup>a,b</sup>, D. Fasanella<sup>a,b</sup>, P. Giacomelli<sup>a</sup>, C. Grandi<sup>a</sup>, L. Guiducci<sup>a,b</sup>, S. Marcellini<sup>a</sup>, G. Masetti<sup>a</sup>, A. Montanari<sup>a</sup>, F.L. Navarria<sup>a,b</sup>, A. Perrotta<sup>a</sup>, F. Primavera<sup>a,b</sup>, A.M. Rossi<sup>a,b</sup>, T. Rovelli<sup>a,b</sup>, G.P. Siroli<sup>a,b</sup>, N. Tosi<sup>a,b</sup>, R. Travaglini<sup>a,b</sup>

**INFN Sezione di Catania <sup>a</sup>, Università di Catania <sup>b</sup>, CSFNSM <sup>c</sup>, Catania, Italy**

S. Albergo<sup>a,b</sup>, G. Cappello<sup>a</sup>, M. Chiorboli<sup>a,b</sup>, S. Costa<sup>a,b</sup>, F. Giordano<sup>a,2</sup>, R. Potenza<sup>a,b</sup>, A. Tricomi<sup>a,b</sup>, C. Tuve<sup>a,b</sup>

**INFN Sezione di Firenze <sup>a</sup>, Università di Firenze <sup>b</sup>, Firenze, Italy**

G. Barbagli<sup>a</sup>, V. Ciulli<sup>a,b</sup>, C. Civinini<sup>a</sup>, R. D'Alessandro<sup>a,b</sup>, E. Focardi<sup>a,b</sup>, E. Gallo<sup>a</sup>, S. Gozzi<sup>a,b</sup>, V. Gori<sup>a,b,2</sup>, P. Lenzi<sup>a,b</sup>, M. Meschini<sup>a</sup>, S. Paoletti<sup>a</sup>, G. Sguazzoni<sup>a</sup>, A. Tropiano<sup>a,b</sup>

**INFN Laboratori Nazionali di Frascati, Frascati, Italy**

L. Benussi, S. Bianco, F. Fabbri, D. Piccolo

**INFN Sezione di Genova <sup>a</sup>, Università di Genova <sup>b</sup>, Genova, Italy**

R. Ferretti<sup>a,b</sup>, F. Ferro<sup>a</sup>, M. Lo Vetere<sup>a,b</sup>, E. Robutti<sup>a</sup>, S. Tosi<sup>a,b</sup>

**INFN Sezione di Milano-Bicocca <sup>a</sup>, Università di Milano-Bicocca <sup>b</sup>, Milano, Italy**

M.E. Dinardo<sup>a,b</sup>, P. Dini<sup>a</sup>, S. Fiorendi<sup>a,b,2</sup>, S. Gennai<sup>a,2</sup>, R. Gerosa<sup>a,b,2</sup>, A. Ghezzi<sup>a,b</sup>, P. Govoni<sup>a,b</sup>, M.T. Lucchini<sup>a,b,2</sup>, S. Malvezzi<sup>a</sup>, R.A. Manzoni<sup>a,b</sup>, A. Martelli<sup>a,b</sup>, B. Marzocchi<sup>a,b</sup>, D. Menasce<sup>a</sup>, L. Moroni<sup>a</sup>, M. Paganoni<sup>a,b</sup>, D. Pedrini<sup>a</sup>, N. Redaelli<sup>a</sup>, T. Tabarelli de Fatis<sup>a,b</sup>

**INFN Sezione di Napoli <sup>a</sup>, Università di Napoli 'Federico II' <sup>b</sup>, Università della Basilicata (Potenza) <sup>c</sup>, Università G. Marconi (Roma) <sup>d</sup>, Napoli, Italy**

S. Buontempo<sup>a</sup>, N. Cavallo<sup>a,c</sup>, S. Di Guida<sup>a,d,2</sup>, F. Fabozzi<sup>a,c</sup>, A.O.M. Iorio<sup>a,b</sup>, L. Lista<sup>a</sup>, S. Meola<sup>a,d,2</sup>, M. Merola<sup>a</sup>, P. Paolucci<sup>a,2</sup>

**INFN Sezione di Padova <sup>a</sup>, Università di Padova <sup>b</sup>, Università di Trento (Trento) <sup>c</sup>, Padova, Italy**

P. Azzi<sup>a</sup>, N. Bacchetta<sup>a</sup>, D. Bisello<sup>a,b</sup>, A. Branca<sup>a,b</sup>, R. Carlin<sup>a,b</sup>, P. Checchia<sup>a</sup>, M. Dall'Osso<sup>a,b</sup>, T. Dorigo<sup>a</sup>, U. Dosselli<sup>a</sup>, M. Galanti<sup>a,b</sup>, U. Gasparini<sup>a,b</sup>, P. Giubilato<sup>a,b</sup>,

A. Gozzelino<sup>a</sup>, K. Kanishchev<sup>a,c</sup>, S. Lacaprara<sup>a</sup>, M. Margoni<sup>a,b</sup>, A.T. Meneguzzo<sup>a,b</sup>,  
F. Montecassiano<sup>a</sup>, M. Passaseo<sup>a</sup>, J. Pazzini<sup>a,b</sup>, N. Pozzobon<sup>a,b</sup>, P. Ronchese<sup>a,b</sup>,  
F. Simonetto<sup>a,b</sup>, E. Torassa<sup>a</sup>, M. Tosi<sup>a,b</sup>, P. Zotto<sup>a,b</sup>, A. Zucchetta<sup>a,b</sup>

**INFN Sezione di Pavia <sup>a</sup>, Università di Pavia <sup>b</sup>, Pavia, Italy**

M. Gabusi<sup>a,b</sup>, S.P. Ratti<sup>a,b</sup>, V. Re<sup>a</sup>, C. Riccardi<sup>a,b</sup>, P. Salvini<sup>a</sup>, P. Vitulo<sup>a,b</sup>

**INFN Sezione di Perugia <sup>a</sup>, Università di Perugia <sup>b</sup>, Perugia, Italy**

M. Biasini<sup>a,b</sup>, G.M. Bilei<sup>a</sup>, D. Ciangottini<sup>a,b</sup>, L. Fanò<sup>a,b</sup>, P. Lariccia<sup>a,b</sup>, G. Mantovani<sup>a,b</sup>,  
M. Menichelli<sup>a</sup>, A. Saha<sup>a</sup>, A. Santocchia<sup>a,b</sup>, A. Spiezia<sup>a,b,2</sup>

**INFN Sezione di Pisa <sup>a</sup>, Università di Pisa <sup>b</sup>, Scuola Normale Superiore di Pisa <sup>c</sup>, Pisa, Italy**

K. Androsov<sup>a,25</sup>, P. Azzurri<sup>a</sup>, G. Bagliesi<sup>a</sup>, J. Bernardini<sup>a</sup>, T. Boccali<sup>a</sup>, G. Broccolo<sup>a,c</sup>,  
R. Castaldi<sup>a</sup>, M.A. Ciocci<sup>a,25</sup>, R. Dell'Orso<sup>a</sup>, S. Donato<sup>a,c</sup>, F. Fiori<sup>a,c</sup>, L. Foà<sup>a,c</sup>,  
A. Giassi<sup>a</sup>, M.T. Grippo<sup>a,25</sup>, F. Ligabue<sup>a,c</sup>, T. Lomtadze<sup>a</sup>, L. Martini<sup>a,b</sup>, A. Messineo<sup>a,b</sup>,  
C.S. Moon<sup>a,26</sup>, F. Palla<sup>a,2</sup>, A. Rizzi<sup>a,b</sup>, A. Savoy-Navarro<sup>a,27</sup>, A.T. Serban<sup>a</sup>, P. Spagnolo<sup>a</sup>,  
P. Squillacioti<sup>a,25</sup>, R. Tenchini<sup>a</sup>, G. Tonelli<sup>a,b</sup>, A. Venturi<sup>a</sup>, P.G. Verdini<sup>a</sup>, C. Vernieri<sup>a,c,2</sup>

**INFN Sezione di Roma <sup>a</sup>, Università di Roma <sup>b</sup>, Roma, Italy**

L. Barone<sup>a,b</sup>, F. Cavallari<sup>a</sup>, G. D'imperio<sup>a,b</sup>, D. Del Re<sup>a,b</sup>, M. Diemoz<sup>a</sup>, M. Grassi<sup>a,b</sup>,  
C. Jorda<sup>a</sup>, E. Longo<sup>a,b</sup>, F. Margaroli<sup>a,b</sup>, P. Meridiani<sup>a</sup>, F. Micheli<sup>a,b,2</sup>, S. Nourbakhsh<sup>a,b</sup>,  
G. Organtini<sup>a,b</sup>, R. Paramatti<sup>a</sup>, S. Rahatlou<sup>a,b</sup>, C. Rovelli<sup>a</sup>, F. Santanastasio<sup>a,b</sup>,  
L. Soffi<sup>a,b,2</sup>, P. Traczyk<sup>a,b</sup>

**INFN Sezione di Torino <sup>a</sup>, Università di Torino <sup>b</sup>, Università del Piemonte Orientale (Novara) <sup>c</sup>, Torino, Italy**

N. Amapane<sup>a,b</sup>, R. Arcidiacono<sup>a,c</sup>, S. Argiro<sup>a,b</sup>, M. Arneodo<sup>a,c</sup>, R. Bellan<sup>a,b</sup>, C. Biino<sup>a</sup>,  
N. Cartiglia<sup>a</sup>, S. Casasso<sup>a,b,2</sup>, M. Costa<sup>a,b</sup>, A. Degano<sup>a,b</sup>, N. Demaria<sup>a</sup>, L. Finco<sup>a,b</sup>,  
C. Mariotti<sup>a</sup>, S. Maselli<sup>a</sup>, E. Migliore<sup>a,b</sup>, V. Monaco<sup>a,b</sup>, M. Musich<sup>a</sup>, M.M. Obertino<sup>a,c,2</sup>,  
G. Ortona<sup>a,b</sup>, L. Pacher<sup>a,b</sup>, N. Pastrone<sup>a</sup>, M. Pelliccioni<sup>a</sup>, G.L. Pinna Angioni<sup>a,b</sup>,  
A. Potenza<sup>a,b</sup>, A. Romero<sup>a,b</sup>, M. Ruspa<sup>a,c</sup>, R. Sacchi<sup>a,b</sup>, A. Solano<sup>a,b</sup>, A. Staiano<sup>a</sup>,  
U. Tamponi<sup>a</sup>

**INFN Sezione di Trieste <sup>a</sup>, Università di Trieste <sup>b</sup>, Trieste, Italy**

S. Belforte<sup>a</sup>, V. Candelise<sup>a,b</sup>, M. Casarsa<sup>a</sup>, F. Cossutti<sup>a</sup>, G. Della Ricca<sup>a,b</sup>, B. Gobbo<sup>a</sup>,  
C. La Licata<sup>a,b</sup>, M. Marone<sup>a,b</sup>, A. Schizzi<sup>a,b</sup>, T. Umer<sup>a,b</sup>, A. Zanetti<sup>a</sup>

**Kangwon National University, Chunchon, Korea**

S. Chang, A. Kropivnitskaya, S.K. Nam

**Kyungpook National University, Daegu, Korea**

D.H. Kim, G.N. Kim, M.S. Kim, D.J. Kong, S. Lee, Y.D. Oh, H. Park, A. Sakharov,  
D.C. Son

**Chonbuk National University, Jeonju, Korea**

T.J. Kim

**Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea**

J.Y. Kim, S. Song

**Korea University, Seoul, Korea**

S. Choi, D. Gyun, B. Hong, M. Jo, H. Kim, Y. Kim, B. Lee, K.S. Lee, S.K. Park, Y. Roh

**University of Seoul, Seoul, Korea**

M. Choi, J.H. Kim, I.C. Park, G. Ryu, M.S. Ryu

**Sungkyunkwan University, Suwon, Korea**

Y. Choi, Y.K. Choi, J. Goh, D. Kim, E. Kwon, J. Lee, H. Seo, I. Yu

**Vilnius University, Vilnius, Lithuania**

A. Juodagalvis

**National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**

J.R. Komaragiri, M.A.B. Md Ali

**Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**

H. Castilla-Valdez, E. De La Cruz-Burelo, I. Heredia-de La Cruz<sup>28</sup>, A. Hernandez-Almada, R. Lopez-Fernandez, A. Sanchez-Hernandez

**Universidad Iberoamericana, Mexico City, Mexico**

S. Carrillo Moreno, F. Vazquez Valencia

**Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**

I. Pedraza, H.A. Salazar Ibarquen

**Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico**

E. Casimiro Linares, A. Morelos Pineda

**University of Auckland, Auckland, New Zealand**

D. Krofcheck

**University of Canterbury, Christchurch, New Zealand**

P.H. Butler, S. Reucroft

**National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**

A. Ahmad, M. Ahmad, Q. Hassan, H.R. Hoorani, S. Khalid, W.A. Khan, T. Khurshid, M.A. Shah, M. Shoaib

**National Centre for Nuclear Research, Swierk, Poland**

H. Bialkowska, M. Bluj, B. Boimska, T. Frueboes, M. Górski, M. Kazana, K. Nawrocki, K. Romanowska-Rybinska, M. Szleper, P. Zalewski

**Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**

G. Brona, K. Bunkowski, M. Cwiok, W. Dominik, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski, M. Misiura, M. Olszewski, W. Wolszczak

**Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**

P. Bargassa, C. Beirão Da Cruz E Silva, P. Faccioli, P.G. Ferreira Parracho, M. Gallinaro, L. Lloret Iglesias, F. Nguyen, J. Rodrigues Antunes, J. Seixas, J. Varela, P. Vischia

**Joint Institute for Nuclear Research, Dubna, Russia**

S. Afanasiev, P. Bunin, M. Gavrilenko, I. Golutvin, I. Gorbunov, A. Kamenev, V. Karjavin, V. Konoplyanikov, A. Lanev, A. Malakhov, V. Matveev<sup>29</sup>, P. Moisenz, V. Palichik, V. Perelygin, S. Shmatov, N. Skatchkov, V. Smirnov, A. Zarubin

**Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia**

V. Golovtsov, Y. Ivanov, V. Kim<sup>30</sup>, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, V. Sulimov, L. Uvarov, S. Vavilov, A. Vorobyev, An. Vorobyev

**Institute for Nuclear Research, Moscow, Russia**

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, M. Kirsanov, N. Krasnikov, A. Pashenkov, D. Tlisov, A. Toropin

**Institute for Theoretical and Experimental Physics, Moscow, Russia**

V. Epshteyn, V. Gavrilov, N. Lychkovskaya, V. Popov, G. Safronov, S. Semenov, A. Spiridonov, V. Stolin, E. Vlasov, A. Zhokin

**P.N. Lebedev Physical Institute, Moscow, Russia**

V. Andreev, M. Azarkin, I. Dremin, M. Kirakosyan, A. Leonidov, G. Mesyats, S.V. Rusakov, A. Vinogradov

**Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia**

A. Belyaev, E. Boos, M. Dubinin<sup>31</sup>, L. Dudko, A. Ershov, A. Gribushin, V. Klyukhin, O. Kodolova, I. Lokhtin, S. Obraztsov, S. Petrushanko, V. Savrin, A. Snigirev

**State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, Russia**

I. Azhgirey, I. Bayshev, S. Bitioukov, V. Kachanov, A. Kalinin, D. Konstantinov, V. Krychkin, V. Petrov, R. Ryutin, A. Sobol, L. Tourtchanovitch, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

**University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia**

P. Adzic<sup>32</sup>, M. Ekmedzic, J. Milosevic, V. Rekovic

**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**

J. Alcaraz Maestre, C. Battilana, E. Calvo, M. Cerrada, M. Chamizo Llatas, N. Colino, B. De La Cruz, A. Delgado Peris, D. Domínguez Vázquez, A. Escalante Del Valle, C. Fernandez Bedoya, J.P. Fernández Ramos, J. Flix, M.C. Fouz, P. Garcia-Abia, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, E. Navarro De Martino, A. Pérez-Calero Yzquierdo, J. Puerta Pelayo, A. Quintario Olmeda, I. Redondo, L. Romero, M.S. Soares

**Universidad Autónoma de Madrid, Madrid, Spain**

C. Albajar, J.F. de Trocóniz, M. Missiroli, D. Moran

**Universidad de Oviedo, Oviedo, Spain**

H. Brun, J. Cuevas, J. Fernandez Menendez, S. Folgueras, I. Gonzalez Caballero

**Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**

J.A. Brochero Cifuentes, I.J. Cabrillo, A. Calderon, J. Duarte Campderros, M. Fernandez, G. Gomez, A. Graziano, A. Lopez Virto, J. Marco, R. Marco, C. Martinez Rivero, F. Matorras, F.J. Munoz Sanchez, J. Piedra Gomez, T. Rodrigo, A.Y. Rodríguez-Marrero, A. Ruiz-Jimeno, L. Scodellaro, I. Vila, R. Vilar Cortabitarte

**CERN, European Organization for Nuclear Research, Geneva, Switzerland**D. Abbaneo, E. Auffray, G. Auzinger, M. Bachtis, P. Baillon, A.H. Ball, D. Barney, A. Benaglia, J. Bendavid, L. Benhabib, J.F. Benitez, C. Bernet<sup>7</sup>, P. Bloch, A. Bocci, A. Bonato, O. Bondu, C. Botta, H. Breuker, T. Camporesi, G. Cerminara, S. Colafranceschi<sup>33</sup>, M. D'Alfonso, D. d'Enterria, A. Dabrowski, A. David, F. De Guio, A. De Roeck, S. De Visscher, E. Di Marco, M. Dobson, M. Dordevic, N. Dupont-Sagorin, A. Elliott-Peisert, J. Eugster, G. Franzoni, W. Funk, D. Gigi, K. Gill, D. Giordano, M. Girone, F. Glege, R. Guida, S. Gundacker, M. Guthoff, J. Hammer, M. Hansen, P. Harris, J. Hegeman, V. Innocente, P. Janot, K. Kousouris, K. Krajczar, P. Lecoq, C. Lourenço, N. Magini, L. Malgeri, M. Mannelli, J. Marrouche, L. Masetti, F. Meijers, S. Mersi, E. Meschi, F. Moortgat, S. Morovic, M. Mulders, P. Musella, L. Orsini, L. Pape, E. Perez, L. Perrozzi, A. Petrilli, G. Petrucciani, A. Pfeiffer, M. Pierini, M. Pimiä, D. Piparo, M. Plagge, A. Racz, G. Rolandi<sup>34</sup>, M. Rovere, H. Sakulin, C. Schäfer, C. Schwick, A. Sharma, P. Siegrist, P. Silva, M. Simon, P. Sphicas<sup>35</sup>, D. Spiga, J. Steggemann, B. Stieger, M. Stoye, Y. Takahashi, D. Treille, A. Tsirou, G.I. Veres<sup>17</sup>, N. Wardle, H.K. Wöhri, H. Wollny, W.D. Zeuner**Paul Scherrer Institut, Villigen, Switzerland**

W. Bertl, K. Deiters, W. Erdmann, R. Horisberger, Q. Ingram, H.C. Kaestli, D. Kotlinski, U. Langenegger, D. Renker, T. Rohe

**Institute for Particle Physics, ETH Zurich, Zurich, Switzerland**F. Bachmair, L. Bäni, L. Bianchini, M.A. Buchmann, B. Casal, N. Chanon, G. Dissertori, M. Dittmar, M. Donegà, M. Dünser, P. Eller, C. Grab, D. Hits, J. Hoss, W. Luster, B. Mangano, A.C. Marini, P. Martinez Ruiz del Arbol, M. Masciovecchio, D. Meister, N. Mohr, C. Nägeli<sup>36</sup>, F. Nessi-Tedaldi, F. Pandolfi, F. Pauss, M. Peruzzi, M. Quittnat, L. Rebane, M. Rossini, A. Starodumov<sup>37</sup>, M. Takahashi, K. Theofilatos, R. Wallny, H.A. Weber**Universität Zürich, Zurich, Switzerland**C. Amsler<sup>38</sup>, M.F. Canelli, V. Chiochia, A. De Cosa, A. Hinzmann, T. Hreus, B. Kilminster, C. Lange, B. Millan Mejias, J. Ngadiuba, P. Robmann, F.J. Ronga, S. Taroni, M. Verzetti, Y. Yang



**National Central University, Chung-Li, Taiwan**

M. Cardaci, K.H. Chen, C. Ferro, C.M. Kuo, W. Lin, Y.J. Lu, R. Volpe, S.S. Yu

**National Taiwan University (NTU), Taipei, Taiwan**

P. Chang, Y.H. Chang, Y.W. Chang, Y. Chao, K.F. Chen, P.H. Chen, C. Dietz, U. Grundler, W.-S. Hou, K.Y. Kao, Y.J. Lei, Y.F. Liu, R.-S. Lu, D. Majumder, E. Petrakou, Y.M. Tzeng, R. Wilken

**Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand**

B. Asavapibhop, G. Singh, N. Srimanobhas, N. Suwonjandee

**Cukurova University, Adana, Turkey**

A. Adiguzel, M.N. Bakirci<sup>39</sup>, S. Cerci<sup>40</sup>, C. Dozen, I. Dumanoglu, E. Eskut, S. Girgis, G. Gokbulut, E. Gurpinar, I. Hos, E.E. Kangal, A. Kayis Topaksu, G. Onengut<sup>41</sup>, K. Ozdemir, S. Ozturk<sup>39</sup>, A. Polatoz, D. Sunar Cerci<sup>40</sup>, B. Tali<sup>40</sup>, H. Topakli<sup>39</sup>, M. Vergili

**Middle East Technical University, Physics Department, Ankara, Turkey**

I.V. Akin, B. Bilin, S. Bilmis, H. Gamsizkan<sup>42</sup>, G. Karapinar<sup>43</sup>, K. Ocalan<sup>44</sup>, S. Sekmen, U.E. Surat, M. Yalvac, M. Zeyrek

**Bogazici University, Istanbul, Turkey**

E. Gülmez, B. Isildak<sup>45</sup>, M. Kaya<sup>46</sup>, O. Kaya<sup>47</sup>

**Istanbul Technical University, Istanbul, Turkey**

K. Cankocak, F.I. Vardarli

**National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine**

L. Levchuk, P. Sorokin

**University of Bristol, Bristol, United Kingdom**

J.J. Brooke, E. Clement, D. Cussans, H. Flacher, J. Goldstein, M. Grimes, G.P. Heath, H.F. Heath, J. Jacob, L. Kreczko, C. Lucas, Z. Meng, D.M. Newbold<sup>48</sup>, S. Paramesvaran, A. Poll, S. Senkin, V.J. Smith, T. Williams

**Rutherford Appleton Laboratory, Didcot, United Kingdom**

K.W. Bell, A. Belyaev<sup>49</sup>, C. Brew, R.M. Brown, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, E. Olaiya, D. Petyt, C.H. Shepherd-Themistocleous, A. Thea, I.R. Tomalin, W.J. Womersley, S.D. Worm

**Imperial College, London, United Kingdom**

M. Baber, R. Bainbridge, O. Buchmuller, D. Burton, D. Colling, N. Cripps, M. Cutajar, P. Dauncey, G. Davies, M. Della Negra, P. Dunne, W. Ferguson, J. Fulcher, D. Futyan, A. Gilbert, G. Hall, G. Iles, M. Jarvis, G. Karapostoli, M. Kenzie, R. Lane, R. Lucas<sup>48</sup>, L. Lyons, A.-M. Magnan, S. Malik, B. Mathias, J. Nash, A. Nikitenko<sup>37</sup>, J. Pela, M. Pesaresi, K. Petridis, D.M. Raymond, S. Rogerson, A. Rose, C. Seez, P. Sharp<sup>†</sup>, A. Tapper, M. Vazquez Acosta, T. Virdee, S.C. Zenz

**Brunel University, Uxbridge, United Kingdom**

J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, D. Leggat, D. Leslie, W. Martin, I.D. Reid, P. Symonds, L. Teodorescu, M. Turner

**Baylor University, Waco, U.S.A.**

J. Dittmann, K. Hatakeyama, A. Kasmi, H. Liu, T. Scarborough

**The University of Alabama, Tuscaloosa, U.S.A.**

O. Charaf, S.I. Cooper, C. Henderson, P. Rumerio

**Boston University, Boston, U.S.A.**

A. Avetisyan, T. Bose, C. Fantasia, P. Lawson, C. Richardson, J. Rohlf, J. St. John, L. Sulak

**Brown University, Providence, U.S.A.**

J. Alimena, E. Berry, S. Bhattacharya, G. Christopher, D. Cutts, Z. Demiragli, N. Dhingra, A. Ferapontov, A. Garabedian, U. Heintz, G. Kukartsev, E. Laird, G. Landsberg, M. Luk, M. Narain, M. Segala, T. Sinthuprasith, T. Speer, J. Swanson

**University of California, Davis, Davis, U.S.A.**

R. Breedon, G. Breto, M. Calderon De La Barca Sanchez, S. Chauhan, M. Chertok, J. Conway, R. Conway, P.T. Cox, R. Erbacher, M. Gardner, W. Ko, R. Lander, T. Miceli, M. Mulhearn, D. Pellett, J. Pilot, F. Ricci-Tam, M. Searle, S. Shalhout, J. Smith, M. Squires, D. Stolp, M. Tripathi, S. Wilbur, R. Yohay

**University of California, Los Angeles, U.S.A.**

R. Cousins, P. Everaerts, C. Farrell, J. Hauser, M. Ignatenko, G. Rakness, E. Takasugi, V. Valuev, M. Weber

**University of California, Riverside, Riverside, U.S.A.**

K. Burt, R. Clare, J. Ellison, J.W. Gary, G. Hanson, J. Heilman, M. Ivova Rikova, P. Jandir, E. Kennedy, F. Lacroix, O.R. Long, A. Luthra, M. Malberti, H. Nguyen, M. Olmedo Negrete, A. Shrinivas, S. Sumowidagdo, S. Wimpenny

**University of California, San Diego, La Jolla, U.S.A.**

W. Andrews, J.G. Branson, G.B. Cerati, S. Cittolin, R.T. D'Agnolo, D. Evans, A. Holzner, R. Kelley, D. Klein, M. Lebourgeois, J. Letts, I. Macneill, D. Olivito, S. Padhi, C. Palmer, M. Pieri, M. Sani, V. Sharma, S. Simon, E. Sudano, M. Tadel, Y. Tu, A. Vartak, C. Welke, F. Würthwein, A. Yagil

**University of California, Santa Barbara, Santa Barbara, U.S.A.**

D. Barge, J. Bradmiller-Feld, C. Campagnari, T. Danielson, A. Dishaw, V. Dutta, K. Flowers, M. Franco Sevilla, P. Geffert, C. George, F. Golf, L. Gouskos, J. Incandela, C. Justus, N. Mccoll, J. Richman, D. Stuart, W. To, C. West, J. Yoo

**California Institute of Technology, Pasadena, U.S.A.**

A. Apresyan, A. Bornheim, J. Bunn, Y. Chen, J. Duarte, A. Mott, H.B. Newman, C. Pena, C. Rogan, M. Spiropulu, V. Timciuc, J.R. Vlimant, R. Wilkinson, S. Xie, R.Y. Zhu

**Carnegie Mellon University, Pittsburgh, U.S.A.**

V. Azzolini, A. Calamba, B. Carlson, T. Ferguson, Y. Iiyama, M. Paulini, J. Russ, H. Vogel, I. Vorobiev

**University of Colorado at Boulder, Boulder, U.S.A.**

J.P. Cumalat, W.T. Ford, A. Gaz, E. Luiggi Lopez, U. Nauenberg, J.G. Smith, K. Stenson, K.A. Ulmer, S.R. Wagner

**Cornell University, Ithaca, U.S.A.**

J. Alexander, A. Chatterjee, J. Chu, S. Dittmer, N. Eggert, N. Mirman, G. Nicolas Kaufman, J.R. Patterson, A. Ryd, E. Salvati, L. Skinnari, W. Sun, W.D. Teo, J. Thom, J. Thompson, J. Tucker, Y. Weng, L. Winstrom, P. Wittich

**Fairfield University, Fairfield, U.S.A.**

D. Winn

**Fermi National Accelerator Laboratory, Batavia, U.S.A.**

S. Abdullin, M. Albrow, J. Anderson, G. Apollinari, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, G. Bolla, K. Burkett, J.N. Butler, H.W.K. Cheung, F. Chlebana, S. Cihangir, V.D. Elvira, I. Fisk, J. Freeman, Y. Gao, E. Gottschalk, L. Gray, D. Green, S. Grünendahl, O. Gutsche, J. Hanlon, D. Hare, R.M. Harris, J. Hirschauer, B. Hooberman, S. Jindariani, M. Johnson, U. Joshi, K. Kaadze, B. Klima, B. Kreis, S. Kwan, J. Linacre, D. Lincoln, R. Lipton, T. Liu, J. Lykken, K. Maeshima, J.M. Marraffino, V.I. Martinez Outschoorn, S. Maruyama, D. Mason, P. McBride, P. Merkel, K. Mishra, S. Mrenna, Y. Musienko<sup>29</sup>, S. Nahn, C. Newman-Holmes, V. O'Dell, O. Prokofyev, E. Sexton-Kennedy, S. Sharma, A. Soha, W.J. Spalding, L. Spiegel, L. Taylor, S. Tkaczyk, N.V. Tran, L. Uplegger, E.W. Vaandering, R. Vidal, A. Whitbeck, J. Whitmore, F. Yang

**University of Florida, Gainesville, U.S.A.**

D. Acosta, P. Avery, P. Bortignon, D. Bourilkov, M. Carver, T. Cheng, D. Curry, S. Das, M. De Gruttola, G.P. Di Giovanni, R.D. Field, M. Fisher, I.K. Furic, J. Hugon, J. Konigsberg, A. Korytov, T. Kypreos, J.F. Low, K. Matchev, P. Milenovic<sup>50</sup>, G. Mitselmakher, L. Muniz, A. Rinkevicius, L. Shchutska, M. Snowball, D. Sperka, J. Yelton, M. Zakaria

**Florida International University, Miami, U.S.A.**

S. Hewamanage, S. Linn, P. Markowitz, G. Martinez, J.L. Rodriguez

**Florida State University, Tallahassee, U.S.A.**

T. Adams, A. Askew, J. Bochenek, B. Diamond, J. Haas, S. Hagopian, V. Hagopian, K.F. Johnson, H. Prosper, V. Veeraraghavan, M. Weinberg

**Florida Institute of Technology, Melbourne, U.S.A.**

M.M. Baarmand, M. Hohmann, H. Kalakhety, F. Yumiceva

**University of Illinois at Chicago (UIC), Chicago, U.S.A.**

M.R. Adams, L. Apanasevich, V.E. Bazterra, D. Berry, R.R. Betts, I. Bucinskaite, R. Cavanaugh, O. Evdokimov, L. Gauthier, C.E. Gerber, D.J. Hofman, S. Khalatyan, P. Kurt, D.H. Moon, C. O'Brien, C. Silkworth, P. Turner, N. Varelas

**The University of Iowa, Iowa City, U.S.A.**

E.A. Albayrak<sup>51</sup>, B. Bilki<sup>52</sup>, W. Clarida, K. Dilsiz, F. Duru, M. Haytmyradov, J.-P. Merlo, H. Mermerkaya<sup>53</sup>, A. Mestvirishvili, A. Moeller, J. Nachtman, H. Ogul, Y. Onel, F. Ozok<sup>51</sup>, A. Penzo, R. Rahmat, S. Sen, P. Tan, E. Tiras, J. Wetzel, T. Yetkin<sup>54</sup>, K. Yi

**Johns Hopkins University, Baltimore, U.S.A.**

B.A. Barnett, B. Blumenfeld, S. Bolognesi, D. Fehling, A.V. Gritsan, P. Maksimovic, C. Martin, M. Swartz

**The University of Kansas, Lawrence, U.S.A.**

P. Baringer, A. Bean, G. Benelli, C. Bruner, R.P. Kenny III, M. Malek, M. Murray, D. Noonan, S. Sanders, J. Sekaric, R. Stringer, Q. Wang, J.S. Wood

**Kansas State University, Manhattan, U.S.A.**

I. Chakaberia, A. Ivanov, S. Khalil, M. Makouski, Y. Maravin, L.K. Saini, S. Shrestha, N. Skhirtladze, I. Svintradze

**Lawrence Livermore National Laboratory, Livermore, U.S.A.**

J. Gronberg, D. Lange, F. Rebassoo, D. Wright

**University of Maryland, College Park, U.S.A.**

A. Baden, A. Belloni, B. Calvert, S.C. Eno, J.A. Gomez, N.J. Hadley, R.G. Kellogg, T. Kolberg, Y. Lu, M. Marionneau, A.C. Mignerey, K. Pedro, A. Skuja, M.B. Tonjes, S.C. Tonwar

**Massachusetts Institute of Technology, Cambridge, U.S.A.**

A. Apyan, R. Barbieri, G. Bauer, W. Busza, I.A. Cali, M. Chan, L. Di Matteo, G. Gomez Ceballos, M. Goncharov, D. Gulhan, M. Klute, Y.S. Lai, Y.-J. Lee, A. Levin, P.D. Luckey, T. Ma, C. Paus, D. Ralph, C. Roland, G. Roland, G.S.F. Stephans, F. Stöckli, K. Sumorok, D. Velicanu, J. Veverka, B. Wyslouch, M. Yang, M. Zanetti, V. Zhukova

**University of Minnesota, Minneapolis, U.S.A.**

B. Dahmes, A. Gude, S.C. Kao, K. Klapoetke, Y. Kubota, J. Mans, N. Pastika, R. Rusack, A. Singovsky, N. Tambe, J. Turkewitz

**University of Mississippi, Oxford, U.S.A.**

J.G. Acosta, S. Oliveros

**University of Nebraska-Lincoln, Lincoln, U.S.A.**

E. Avdeeva, K. Bloom, S. Bose, D.R. Claes, A. Dominguez, R. Gonzalez Suarez, J. Keller, D. Knowlton, I. Kravchenko, J. Lazo-Flores, S. Malik, F. Meier, G.R. Snow, M. Zvada

**State University of New York at Buffalo, Buffalo, U.S.A.**

J. Dolen, A. Godshalk, I. Iashvili, A. Kharchilava, A. Kumar, S. Rappoccio

**Northeastern University, Boston, U.S.A.**

G. Alverson, E. Barberis, D. Baumgartel, M. Chasco, J. Haley, A. Massironi, D.M. Morse, D. Nash, T. Orimoto, D. Trocino, R.-J. Wang, D. Wood, J. Zhang

**Northwestern University, Evanston, U.S.A.**

K.A. Hahn, A. Kubik, N. Mucia, N. Odell, B. Pollack, A. Pozdnyakov, M. Schmitt, S. Stoynev, K. Sung, M. Velasco, S. Won

**University of Notre Dame, Notre Dame, U.S.A.**

A. Brinkerhoff, K.M. Chan, A. Drozdetskiy, M. Hildreth, C. Jessop, D.J. Karmgard, N. Kellams, K. Lannon, W. Luo, S. Lynch, N. Marinelli, T. Pearson, M. Planer, R. Ruchti, N. Valls, M. Wayne, M. Wolf, A. Woodard

**The Ohio State University, Columbus, U.S.A.**

L. Antonelli, J. Brinson, B. Bylsma, L.S. Durkin, S. Flowers, A. Hart, C. Hill, R. Hughes, K. Kotov, T.Y. Ling, D. Puigh, M. Rodenburg, G. Smith, B.L. Winer, H. Wolfe, H.W. Wulsin

**Princeton University, Princeton, U.S.A.**

O. Driga, P. Elmer, P. Hebda, A. Hunt, S.A. Koay, P. Lujan, D. Marlow, T. Medvedeva, M. Mooney, J. Olsen, P. Piroué, X. Quan, H. Saka, D. Stickland<sup>2</sup>, C. Tully, J.S. Werner, A. Zuranski

**University of Puerto Rico, Mayaguez, U.S.A.**

E. Brownson, H. Mendez, J.E. Ramirez Vargas

**Purdue University, West Lafayette, U.S.A.**

V.E. Barnes, D. Benedetti, D. Bortoletto, M. De Mattia, L. Gutay, Z. Hu, M.K. Jha, M. Jones, K. Jung, M. Kress, N. Leonardo, D. Lopes Pegna, V. Maroussov, D.H. Miller, N. Neumeister, B.C. Radburn-Smith, X. Shi, I. Shipsey, D. Silvers, A. Svyatkovskiy, F. Wang, W. Xie, L. Xu, H.D. Yoo, J. Zablocki, Y. Zheng

**Purdue University Calumet, Hammond, U.S.A.**

N. Parashar, J. Stupak

**Rice University, Houston, U.S.A.**

A. Adair, B. Akgun, K.M. Ecklund, F.J.M. Geurts, W. Li, B. Michlin, B.P. Padley, R. Redjimi, J. Roberts, J. Zabel

**University of Rochester, Rochester, U.S.A.**

B. Betchart, A. Bodek, R. Covarelli, P. de Barbaro, R. Demina, Y. Eshaq, T. Ferbel, A. Garcia-Bellido, P. Goldenzweig, J. Han, A. Harel, A. Khukhunaishvili, G. Petrillo, D. Vishnevskiy

**The Rockefeller University, New York, U.S.A.**

R. Ciesielski, L. Demortier, K. Goulios, G. Lungu, C. Mesropian

**Rutgers, The State University of New Jersey, Piscataway, U.S.A.**

S. Arora, A. Barker, J.P. Chou, C. Contreras-Campana, E. Contreras-Campana, D. Duggan, D. Ferencek, Y. Gershtein, R. Gray, E. Halkiadakis, D. Hidas, S. Kaplan, A. Lath, S. Panwalkar, M. Park, R. Patel, S. Salur, S. Schnetzer, S. Somalwar, R. Stone, S. Thomas, P. Thomassen, M. Walker

**University of Tennessee, Knoxville, U.S.A.**

K. Rose, S. Spanier, A. York

**Texas A&M University, College Station, U.S.A.**

O. Bouhali<sup>55</sup>, A. Castaneda Hernandez, R. Eusebi, W. Flanagan, J. Gilmore, T. Kamon<sup>56</sup>, V. Khotilovich, V. Krutelyov, R. Montalvo, I. Osipenkov, Y. Pakhotin, A. Perloff, J. Roe, A. Rose, A. Safonov, T. Sakuma, I. Suarez, A. Tatarinov

**Texas Tech University, Lubbock, U.S.A.**

N. Akchurin, C. Cowden, J. Damgov, C. Dragoiu, P.R. Duderov, J. Faulkner, K. Kovitangoon, S. Kunori, S.W. Lee, T. Libeiro, I. Volobouev

**Vanderbilt University, Nashville, U.S.A.**

E. Appelt, A.G. Delannoy, S. Greene, A. Gurrola, W. Johns, C. Maguire, Y. Mao, A. Melo, M. Sharma, P. Sheldon, B. Snook, S. Tuo, J. Velkovska

**University of Virginia, Charlottesville, U.S.A.**

M.W. Arenton, S. Boutle, B. Cox, B. Francis, J. Goodell, R. Hirosky, A. Ledovskoy, H. Li, C. Lin, C. Neu, J. Wood

**Wayne State University, Detroit, U.S.A.**

C. Clarke, R. Harr, P.E. Karchin, C. Kottachchi Kankanamge Don, P. Lamichhane, J. Sturdy

**University of Wisconsin, Madison, U.S.A.**

D.A. Belknap, D. Carlsmith, M. Cepeda, S. Dasu, L. Dodd, S. Duric, E. Friis, R. Hall-Wilton, M. Herndon, A. Hervé, P. Klabbers, A. Lanaro, C. Lazaridis, A. Levine, R. Lovelless, A. Mohapatra, I. Ojalvo, T. Perry, G.A. Pierro, G. Polese, I. Ross, T. Sarangi, A. Savin, W.H. Smith, D. Taylor, P. Verwilligen, C. Vuosalo, N. Woods

†: Deceased

1: Also at Vienna University of Technology, Vienna, Austria

2: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

3: Also at Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France

4: Also at National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

5: Also at Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia

6: Also at Universidade Estadual de Campinas, Campinas, Brazil

7: Also at Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France

8: Also at Joint Institute for Nuclear Research, Dubna, Russia

9: Also at Suez University, Suez, Egypt

10: Also at Cairo University, Cairo, Egypt

11: Also at Fayoum University, El-Fayoum, Egypt

12: Also at Ain Shams University, Cairo, Egypt

13: Now at Sultan Qaboos University, Muscat, Oman

- 14: Also at Université de Haute Alsace, Mulhouse, France
- 15: Also at Brandenburg University of Technology, Cottbus, Germany
- 16: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- 17: Also at Eötvös Loránd University, Budapest, Hungary
- 18: Also at University of Debrecen, Debrecen, Hungary
- 19: Also at University of Visva-Bharati, Santiniketan, India
- 20: Now at King Abdulaziz University, Jeddah, Saudi Arabia
- 21: Also at University of Ruhuna, Matara, Sri Lanka
- 22: Also at Isfahan University of Technology, Isfahan, Iran
- 23: Also at University of Tehran, Department of Engineering Science, Tehran, Iran
- 24: Also at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran
- 25: Also at Università degli Studi di Siena, Siena, Italy
- 26: Also at Centre National de la Recherche Scientifique (CNRS) - IN2P3, Paris, France
- 27: Also at Purdue University, West Lafayette, U.S.A.
- 28: Also at Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Mexico
- 29: Also at Institute for Nuclear Research, Moscow, Russia
- 30: Also at St. Petersburg State Polytechnical University, St. Petersburg, Russia
- 31: Also at California Institute of Technology, Pasadena, U.S.A.
- 32: Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia
- 33: Also at Facoltà Ingegneria, Università di Roma, Roma, Italy
- 34: Also at Scuola Normale e Sezione dell'INFN, Pisa, Italy
- 35: Also at University of Athens, Athens, Greece
- 36: Also at Paul Scherrer Institut, Villigen, Switzerland
- 37: Also at Institute for Theoretical and Experimental Physics, Moscow, Russia
- 38: Also at Albert Einstein Center for Fundamental Physics, Bern, Switzerland
- 39: Also at Gaziosmanpasa University, Tokat, Turkey
- 40: Also at Adiyaman University, Adiyaman, Turkey
- 41: Also at Cag University, Mersin, Turkey
- 42: Also at Anadolu University, Eskisehir, Turkey
- 43: Also at Izmir Institute of Technology, Izmir, Turkey
- 44: Also at Necmettin Erbakan University, Konya, Turkey
- 45: Also at Ozyegin University, Istanbul, Turkey
- 46: Also at Marmara University, Istanbul, Turkey
- 47: Also at Kafkas University, Kars, Turkey
- 48: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom
- 49: Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- 50: Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia
- 51: Also at Mimar Sinan University, Istanbul, Istanbul, Turkey
- 52: Also at Argonne National Laboratory, Argonne, U.S.A.
- 53: Also at Erzincan University, Erzincan, Turkey
- 54: Also at Yildiz Technical University, Istanbul, Turkey
- 55: Also at Texas A&M University at Qatar, Doha, Qatar
- 56: Also at Kyungpook National University, Daegu, Korea