



Year: 2016

Simultaneous effects on interneuron and dopaminergic system development in a transgenic rat model for sporadic schizophrenia

Hamburg, Hannah

Abstract: Despite continuous effort, unequivocal neuroanatomical features for sporadic or familial schizophrenia could not yet be defined. A standardized characterization might give rise to issues because of the discrete nature of changes, the heterogeneity of biological causes, and a high intersubject variability. Disrupted-in-schizophrenia 1 (DISC1) is a risk gene for chronic mental illnesses and its overexpression is a vulnerability factor for a subset of sporadic disease types. In this study we report on neuroanatomical abnormalities in transgenic rats overexpressing human full-length, non-mutant DISC1, modeling a subset of sporadic chronic mental illness. A decreased density of dopaminergic neurons in the substantia nigra and reduced dopaminergic fibres in the striatum, but not the amygdala, were detected in tgDISC1 rats. Parvalbumin-interneuron distribution was shifted to the deeper cortical layers and calbindin-interneurons showed a subtle decrease in their total number. In vivo MRI and voxel-wise tensor-based morphometry revealed a slight volumetric decrease of the corpus callosum which could be confirmed microscopically. In this study, we demonstrate that tgDISC1 rats display concurrent neuroanatomical abnormalities in two important neuronal systems. Our analysis gives an insight into neuroanatomical changes of the tgDISC1 rat model that may contribute to a clearer definition of neuroanatomical abnormalities associated with subsets of human sporadic schizophrenia. Eine einheitliche neuroanatomische Charakterisierung sporadischer sowie erblicher Schizophrenieformen konnte bislang trotz stetiger Bemühungen nicht definiert werden. Sie erweist sich als schwierig aufgrund der Feinheit der Veränderungen, der Heterogenität biologischer Ursachen und der hohen interindividuellen Variabilität. Disrupted-in-Schizophrenia 1 (DISC1) ist ein Risikogen für chronisch-mentale Erkrankungen, seine Überexpression prädestiniert für eine Subgruppe sporadischer Erkrankungsformen. In dieser Studie beschreiben wir neuroanatomische Veränderungen eines transgenen Rattenmodells mit Überexpression von nicht-mutiertem, humanem DISC1 in voller Länge. In tgDISC1 Ratten wurden eine verringerte Dichte dopaminerger Neuronen in der Substantia nigra und eine reduzierte dopaminerge Faserdichte im Striatum, jedoch nicht der Amygdala gefunden. Die Verteilung der Parvalbumin-Interneuronen war in die tiefen Kortexschichten verschoben und Calbindin-Interneuronen zeigten eine dezent verringerte Gesamtanzahl. Im in vivo MRT und in Voxel-basierter Morphometrie fiel eine geringe Volumenminderung des Corpus callosum auf, was auch histologisch bestätigt werden konnte. Diese Studie zeigt, dass tgDISC1 Ratten gleichzeitig in zwei wichtigen neuronalen Schaltkreisen Abweichungen aufweisen. Sie gibt einen Einblick in neuroanatomische Veränderungen dieses Modells und kann zu einer klareren Definition histologischer Abweichungen in Subgruppen sporadischer Schizophrenie des Menschen beitragen.

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-131122>

Dissertation

Published Version

Originally published at:

Hamburg, Hannah. Simultaneous effects on interneuron and dopaminergic system development in a transgenic rat model for sporadic schizophrenia. 2016, University of Zurich, Vetsuisse Faculty.



Simultaneous effects on interneuron and dopaminergic system development in a transgenic rat model for sporadic schizophrenia

Hamburg, Hannah

Abstract: Despite continuous effort, unequivocal neuroanatomical features for sporadic or familial schizophrenia could not yet be defined. A standardized characterization might give rise to issues because of the discrete nature of changes, the heterogeneity of biological causes, and a high intersubject variability. Disrupted-in-schizophrenia 1 (DISC1) is a risk gene for chronic mental illnesses and its overexpression is a vulnerability factor for a subset of sporadic disease types. In this study we report on neuroanatomical abnormalities in transgenic rats overexpressing human full-length, non-mutant DISC1, modeling a subset of sporadic chronic mental illness. A decreased density of dopaminergic neurons in the substantia nigra and reduced dopaminergic fibres in the striatum, but not the amygdala, were detected in tgDISC1 rats. Parvalbumin-interneuron distribution was shifted to the deeper cortical layers and calbindin-interneurons showed a subtle decrease in their total number. In vivo MRI and voxel-wise tensor-based morphometry revealed a slight volumetric decrease of the corpus callosum which could be confirmed microscopically. In this study, we demonstrate that tgDISC1 rats display concurrent neuroanatomical abnormalities in two important neuronal systems. Our analysis gives an insight into neuroanatomical changes of the tgDISC1 rat model that may contribute to a clearer definition of neuroanatomical abnormalities associated with subsets of human sporadic schizophrenia. Eine einheitliche neuroanatomische Charakterisierung sporadischer sowie erblicher Schizophrenieformen konnte bislang trotz stetiger Bemühungen nicht definiert werden. Sie erweist sich als schwierig aufgrund der Feinheit der Veränderungen, der Heterogenität biologischer Ursachen und der hohen interindividuellen Variabilität. Disrupted-in-Schizophrenia 1 (DISC1) ist ein Risikogen für chronisch-mentale Erkrankungen, seine Überexpression prädestiniert für eine Subgruppe sporadischer Erkrankungsformen. In dieser Studie beschreiben wir neuroanatomische Veränderungen eines transgenen Rattenmodells mit Überexpression von nicht-mutiertem, humanem DISC1 in voller Länge. In tgDISC1 Ratten wurden eine verringerte Dichte dopaminerger Neuronen in der Substantia nigra und eine reduzierte dopaminerge Faserdichte im Striatum, jedoch nicht der Amygdala gefunden. Die Verteilung der Parvalbumin-Interneuronen war in die tiefen Kortexschichten verschoben und Calbindin-Interneuronen zeigten eine dezent verringerte Gesamtanzahl. Im in vivo MRT und in Voxel-basierter Morphometrie fiel eine geringe Volumenminderung des Corpus callosum auf, was auch histologisch bestätigt werden konnte. Diese Studie zeigt, dass tgDISC1 Ratten gleichzeitig in zwei wichtigen neuronalen Schaltkreisen Abweichungen aufweisen. Sie gibt einen Einblick in neuroanatomische Veränderungen dieses Modells und kann zu einer klareren Definition histologischer Abweichungen in Subgruppen sporadischer Schizophrenie des Menschen beitragen.

Originally published at:

Hamburg, Hannah. Simultaneous effects on interneuron and dopaminergic system development in a transgenic rat model for sporadic schizophrenia. 2016, University of Zurich, Vetsuisse Faculty.

Institut für Veterinärpathologie
der Vetsuisse-Fakultät Universität Zürich

Direktorin: Prof. Dr. Anja Kipar

Arbeit unter wissenschaftlicher Betreuung von
Prof. Dr. Anja Kipar

**Simultaneous effects on interneuron and dopaminergic system development in
a transgenic rat model for sporadic schizophrenia**

Inaugural-Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Hannah Hamburg

Tierärztin
aus Düsseldorf, Deutschland

genehmigt auf Antrag von

Prof. Dr. Anja Kipar, Referentin
Prof. Dr. Anna Oevermann, Korreferentin

2016

Inhaltsverzeichnis

Abstract	2
Zusammenfassung.....	3
Original Article.....	4
Abstract	5
Introduction.....	6
Results	12
Discussion.....	17
Materials and Methods	21
References.....	27
Acknowledgements.....	33
Author contributions statement	33
Additional information	33
Figures	34
Curriculum Vitae.....	44

Vetsuisse Faculty, University of Zurich 2016

Hannah Hamburg

**Institute of Veterinary Pathology
ivpz@vetpath.uzh.ch**

Simultaneous effects on interneuron and dopaminergic system development in a transgenic rat model for sporadic schizophrenia

Abstract

Despite continuous effort, unequivocal neuroanatomical features for sporadic or familial schizophrenia could not yet be defined. A standardized characterization might give rise to issues because of the discrete nature of changes, the heterogeneity of biological causes, and a high intersubject variability.

Disrupted-in-schizophrenia 1 (DISC1) is a risk gene for chronic mental illnesses and its overexpression is a vulnerability factor for a subset of sporadic disease types. In this study we report on neuroanatomical abnormalities in transgenic rats overexpressing human full-length, non-mutant DISC1, modelling a subset of sporadic chronic mental illness.

A decreased density of dopaminergic neurons in the substantia nigra and reduced dopaminergic fibres in the striatum, but not the amygdala, were detected in tgDISC1 rats. Parvalbumin-interneuron distribution was shifted to the deeper cortical layers and calbindin-interneurons showed a subtle decrease in their total number. In vivo MRI and voxel-wise tensor based morphometry revealed a slight volumetric decrease of the corpus callosum which could be confirmed microscopically.

In this study, we demonstrate that tgDISC1 rats display concurrent neuroanatomical abnormalities in two important neuronal systems. Our analysis gives an insight into neuroanatomical changes of the tgDISC1 rat model that may contribute to a clearer definition of neuroanatomical abnormalities associated with subsets of human sporadic schizophrenia.

Key words: Disrupted-in-Schizophrenia 1 (DISC1); dopaminergic system; interneurons; neuroanatomy; schizophrenia

Hannah Hamburg

Institut für Veterinärpathologie
ivpz@vetpath.uzh.ch

Zusammenfassung

Eine einheitliche neuroanatomische Charakterisierung sporadischer sowie erblicher Schizophrenieformen konnte bislang trotz stetiger Bemühungen nicht definiert werden. Sie erweist sich als schwierig aufgrund der Feinheit der Veränderungen, der Heterogenität biologischer Ursachen und der hohen interindividuellen Variabilität. Disrupted-in-Schizophrenia 1 (DISC1) ist ein Risikogen für chronisch-mentale Erkrankungen, seine Überexpression prädestiniert für eine Subgruppe sporadischer Erkrankungsformen. In dieser Studie beschreiben wir neuroanatomische Veränderungen eines transgenen Rattenmodells mit Überexpression von nicht-mutiertem, humanem DISC1 in voller Länge.

In tgDISC1 Ratten wurden eine verringerte Dichte dopaminergener Neuronen in der Substantia nigra und eine reduzierte dopaminerge Faserdichte im Striatum, jedoch nicht der Amygdala gefunden. Die Verteilung der Parvalbumin-Interneuronen war in die tiefen Kortexschichten verschoben und Calbindin-Interneuronen zeigten eine dezent verringerte Gesamtanzahl. Im in vivo MRT und in Voxel-basierter Morphometrie fiel eine geringe Volumenminderung des Corpus callosum auf, was auch histologisch bestätigt werden konnte.

Diese Studie zeigt, dass tgDISC1 Ratten gleichzeitig in zwei wichtigen neuronalen Schaltkreisen Abweichungen aufweisen. Sie gibt einen Einblick in neuroanatomische Veränderungen dieses Modells und kann zu einer klareren Definition histologischer Abweichungen in Subgruppen sporadischer Schizophrenie des Menschen beitragen.

Stichwörter: Disrupted-in-schizophrenia 1 (DISC1); dopaminerges System; Interneuronen; Neuroanatomie; Schizophrenie

Original Article

Simultaneous effects on interneuron and dopaminergic system development in a transgenic rat model for sporadic schizophrenia

Hannah Hamburg^{1,2,3}, Svenja V. Trossbach¹, Verian Bader¹, Caroline Chwiesko⁴, Anja Kipar³, Magdalena Sauvage⁴, William R. Crum⁵, Anthony C. Vernon⁶, Hans Jürgen Bidmon², Carsten Korth^{1*}

¹ Institute of Neuropathology, Heinrich Heine University Düsseldorf, Germany

² Cécile and Oskar Vogt Institute of Brain Research, Heinrich Heine University Düsseldorf, Germany

³ Institute of Veterinary Pathology, Vetsuisse Faculty, University of Zurich, Switzerland

⁴ Functional Architecture of Memory Unit, Mercator Research Group, Faculty of Medicine, Ruhr University Bochum, Bochum, Germany

⁵ King's College London, Institute of Psychiatry, Psychology and Neuroscience, Department of Neuroimaging, London, United Kingdom

⁶ King's College London, Institute of Psychiatry, Psychology and Neuroscience, Department of Basic and Clinical Neuroscience, London, United Kingdom

*corresponding author
Prof. Carsten Korth, MD PhD
Department Neuropathology
Heinrich Heine University Düsseldorf
Moorenstrasse 5
40225 Düsseldorf
Germany
Tel 0211-811-6153
Fax 0211-811-7804
Email ckorth@hhu.de

Abstract

To date, no clear neuroanatomical features have been demonstrated for sporadic or familial schizophrenia. Here, we investigated the neuroanatomical changes in a transgenic rat model of sporadic chronic mental illness (CMI), which modestly overexpresses human full-length, non-mutant DISC1, a vulnerability factor for CMIs. This model shows aberrant dopamine homeostasis consistent with some schizophrenia phenotypes.

Neuroanatomical analysis revealed a reduced density of dopaminergic neurons in the substantia nigra and reduced dopaminergic fibres in the striatum. Parvalbumin-positive interneuron occurrence in the somatosensory cortex was shifted from layers II/III to V/VI, and the number of calbindin-positive interneurons was slightly decreased. Reduced corpus callosum thickness confirmed trend-level observations from *in vivo* MRI and voxel-wise tensor based morphometry.

These neuroanatomical changes help explain functional phenotypes of this animal model, some of which resemble changes observed in human schizophrenia *post mortem* brain tissues. Our findings also demonstrate how a single molecular factor, DISC1 overexpression or misassembly, can account for a variety of seemingly unrelated morphological phenotypes and thus provides a possible explanation for findings observed in schizophrenia patients. Our anatomical investigation of a defined model for sporadic mental illness enables a clearer definition of neuroanatomical changes associated with subsets of human sporadic schizophrenia.

Introduction

The introduction of the diagnosis "schizophrenia" by Bleuler^{1,2,3} followed the previous concept of "dementia praecox" conceived by Kraepelin with the aim of defining a subset of chronic mental illnesses that led to lifelong cognitive impairments^{4,5}, as opposed to affective disorders. This distinction has been maintained for over a century through several editions of the Diagnostic and Statistical Manual of Mental Disorders (DSM,⁶) and the International Statistical Classification of Diseases (ICD,⁷), despite the inability to biologically define any chronic mental illness. Instead, the diagnosis is still based exclusively on a clinical interview.

Even though the clinical success of dopamine antagonists in treating the acute psychotic phase of schizophrenia in a majority of patients has led to the biological "dopamine hypothesis" of schizophrenia (reviewed in ref. 8), neuropathological investigations have failed to clearly define phenotype - clinical disease correlations^{9,10,11}. This is in contrast to, for example, the classification of neurodegenerative diseases, where, apart from distinctive neuronal cell death, each category can be defined by the accumulation or disturbed proteostasis of specific proteins. Strikingly, the proteins that accumulate in the majority of sporadic cases of neurodegenerative diseases are the same as those which are mutated in the familial cases that usually make less than 5% of all cases¹². Notably, however, any neuropathological changes in schizophrenia are likely to be more subtle than those of neurodegenerative diseases and do not comprise significant cell death¹³.

To date, no unequivocal neuropathological features correlating to clinical schizophrenia have been identified, likely due to 1) the subtlety of changes, 2) the heterogeneity of biological causes, and 3) the high inter-subject variability of brain microanatomy in human individuals. Efforts to characterise morphological changes have been made since the clinical description of the condition utilizing both *in vivo* as well as *post mortem* approaches. We will briefly review some reported neuroanatomical changes in schizophrenia.

Early on, multiple pneumoencephalographic studies led to the observation of dilated ventricles in schizophrenia patients, when compared to healthy controls^{14,15}, a finding which was later confirmed by computerized tomography (CT) scans¹⁶ (reviewed in ref. 11). The introduction of magnetic resonance imaging (MRI) enabled differentiation between changes in white and grey matter, respectively. Prospective large-scale meta-analyses of MRI data from more than 2000 cases of schizophrenia and healthy controls each via the ENIGMA consortium confirmed an approximate ~19% increase in ventricular volume of schizophrenia patients with medium effect size (Cohen's $d = 0.37$)¹⁷.

Furthermore, a decrease in cortical volume and disproportionate volume loss of temporal structures have been described (reviewed in ref. 11). Regarding white matter alterations, the corpus callosum is one of the structures which most commonly shows abnormalities^{18,19} with several studies reporting a reduction of its volume^{20,21,22}.

Early MRI analyses assessing striatal size because of its major dopaminergic input and relevance in cognitive, sensory and motor processing yielded mixed results, with the majority of studies reporting an increase in volume (reviewed in ref. 23). Two

recent studies have also demonstrated enlarged striatal sizes in drug-naïve patients suffering from either schizophrenia or schizotypal disorders ^{24,25}. Nevertheless, recent analysis of MRI data from the schizophrenia working group of the ENIGMA consortium reported no differences in either caudate or putamen volume in schizophrenia patients relative to healthy controls, although increased putamen volumes were associated with duration of illness and age ²⁶. This corroborates the findings reported in a large-scale meta-analysis of MRI data (18,000 subjects) by Haijma and colleagues ²⁷ who observed lower caudate volumes only in medication-naïve individuals with schizophrenia compared with controls. Importantly, cumulative exposure to antipsychotic drugs has been linked to either increase or normalization of caudate and putamen volumes, as well as cortical thinning in clinical MRI studies of schizophrenia patients ²⁷ and pre-clinical MRI studies of antipsychotic drug effects on brain volume ^{28,29}, making it difficult to always distinguish definitively between drug effects and disease effects in medicated schizophrenia patients. The inconsistency of neuroanatomical findings in schizophrenia has been attributed to the heterogeneity of underlying biological disease causes.

Alterations in the number and positioning of inhibitory GABAergic neurons have repeatedly been reported to be a phenotype of schizophrenia (reviewed in refs. 30, 31, 32). The most consistent change is a reduced number of interneurons, especially parvalbumin (PV) expressing cells which provide the main inhibitory control for pyramidal output neurons (reviewed in ref. 9).

The conundrum is that even though it is generally recognized that schizophrenia comprises heterogeneous biological causes converging on a final common, stereotypical behavioural pathway, all cases of schizophrenia are still treated as one entity in clinical as well as research practice. This clearly provides the risk of mixing

up potentially different anatomical/neuropathological phenotypes in systematic investigations under the common umbrella of clinically defined "schizophrenia". In addition, the considerable background of microanatomical inter-subject variability between humans renders investigations of subtle microanatomical changes difficult^{33,34}. It is also not clear how the different reported changes could eventually hang together. A solution to this dilemma may be, rather than searching for a common denominator to all clinically defined schizophrenia cases, to instead define subsets of cases with schizophrenia by common biological and neuropathological signatures.

Within this context it has been revealing to investigate mutant genes, originally identified in families affected with schizophrenia, in genetically modified mouse models. Notably, the mutant *Disrupted-in-schizophrenia 1 (DISC1)* gene has been shown to lead to neurodevelopmental changes consistent with the proposed neurodevelopmental origin of schizophrenia^{35,36,37}. A mutant *DISC1* gene has been identified in two independent families to date^{38,39,40} and animal models of mutant *DISC1* have consistently revealed, amongst other phenotypes, neuroanatomical changes including altered interneuron distribution, disturbed dopamine signalling and dendritic alterations (reviewed in ref. 41). Furthermore, *in utero* *DISC1* knockdown studies have revealed deficits in the migration of pyramidal neurons, a reduced number of PV-interneurons and abnormal maturation of dopaminergic neurons^{35,42,43} establishing a neurodevelopmental function for *DISC1*.

DISC1 is known to be involved in the neurodevelopment of cortical progenitor cells via GSK-3b/b-catenin where it acts as a switch from proliferation to migration through a phosphorylation site^{44,45}. However, it has recently been demonstrated that it is also involved in the tangential migration of cortical interneurons in a regulatory manner⁴³. Knockdown of *DISC1* leads to a reduction of PV-positive neurons in the cortex by

interaction with the actin cytoskeleton during nucleokinesis, leading to impaired tangential migration of interneurons originating from the ganglionic eminence ⁴⁶. Reduced numbers or abnormal distribution of PV-positive neurons are a frequent finding in mutant DISC1 mouse models ^{42,47,48}.

One characteristic feature of DISC1 mouse models is the enlargement of the lateral ventricles, with some also displaying a concurrent thinning of the cortex ⁴⁷⁻⁵¹. Further morphological changes include diverse dendritic alterations, such as reduced or enhanced spine density, reduced dendrite length and neurite outgrowth as well as mis-oriented dendrites ^{42,47,48,52-54}.

To date, partial agenesis of the corpus callosum has been described in one mutant DISC1 mouse model ⁴⁷, consistent with the frequent reporting of white matter changes in neuroimaging studies of patients with schizophrenia ¹⁹.

Even though analysis of the effects of mutant genes on neurodevelopment and adult behavioural phenotypes may be revealing, potentially linking the mutant gene to a particular disease-related phenotype, it is not evident how the non-mutant gene could do so in the majority of sporadic cases of schizophrenia, i.e. patients without an obvious genetic defect. One possible way in how non-mutant gene products could nevertheless play a specific role in disease mechanisms would be through posttranslational modifications. In analogy to the fact that proteins subject to proteostatic disturbance in familial cases of neurodegenerative disease are often the same as those in sporadic cases ¹², we previously hypothesized that the DISC1 protein could be insoluble in sporadic cases of chronic mental illness. Indeed, we demonstrated that to be the case in about 10% of *post mortem* brains of patients with chronic psychiatric disease ⁵⁵. Modelling the presence of insoluble DISC1 aggregates

in a transgenic rat model by modestly overexpressing the full length, non-mutant DISC1 protein, we observed ubiquitous perinuclear DISC1 aggregates that were accentuated in dopamine-rich regions such as the dorsal striatum, as well as biochemical and behavioural phenotypes consistent with aberrant dopamine homeostasis ⁵⁶. Specifically, this rat model showed decreased total levels of dopamine in the dorsal striatum, the amygdala and the hippocampus, an increased proportion of high-affinity dopamine D2 receptors and translocated dopamine transporter leading to altered dopamine flow dynamics. On the systemic level, an amphetamine supersensitivity was also observed ⁵⁶.

Based on this behavioural and neurochemical analysis of the tgDISC1 rat, we hypothesized that morphological irregularities in the central nervous system of tgDISC1 rats would be present. We therefore carried out a detailed *post mortem* microscale examination of brains from tgDISC1 rats, as well as littermate controls (LM). Based on prior observations in this tgDISC1 rat as well as mutant DISC1 models, this *post mortem* analysis initially focused on the dopaminergic system and PV-interneurons as well as the corpus callosum. In parallel, macroscale MRI combined with voxel-wise tensor-based morphometry ²⁹ were conducted to provide brain-wide mapping of the anatomy of tgDISC1 rats in comparison to LM. Additional *post mortem* analyses were then carried out guided by the additional results arising from the MRI analysis. We found reduced numbers of dopaminergic neurons in the compact part of the substantia nigra (SNc) and sparser dopaminergic fibre density in the striatum. The distribution of PV-positive neurons was shifted to the deeper cortex layers and the corpus callosum showed a volumetric decrease in tgDISC1 rats, when compared to LM.

Results

Modest overexpression of full-length human DISC1 reduces TH-positive fibre density in the striatum and enlarges the striosomal compartment in tgDISC1 rats.

Since we had previously observed aberrant dopamine homeostasis in the tgDISC1 rat, accentuated in the striatum⁵⁶, the histomorphology of the striatum of the tgDISC1 rat was analyzed by comparing striatal size and tyrosine hydroxylase (TH)-positive fibre density of tgDISC1 rats vs. non-transgenic LM (Figure 1A-C). Grey value for a precise estimation of TH-positive fibres as well as area measurements were performed on TH-immunostained brain sections. Analysis was carried out on the striatum as a whole and also on both striatal compartments: the matrix and the striosomes.

TgDISC1 rats showed a decrease in grey value intensity following TH immunostaining when compared to LM in measurements of the entire striatum ($p = 0.005$, Cohen's $d = 1.66$) as well as in the matrix exclusively without the striosomes ($p = 0.004$, Cohen's $d = 1.73$), indicating sparser TH-positive fibres at this location (Figure 1D-F). There was no significant difference in the total area of the striatum, but there was a trend towards an increase in transgenic animals (Figure 1G). Looking at the individual compartments, the size of the striatal matrix did not differ between the two groups, whereas a striosomal expansion was present in tgDISC1 rats ($p = 0.038$, Cohen's $d = 1.15$, Figure 1H, I). Therefore, our morphological analysis indicates that tgDISC1 rats display lower TH-positive fibre

density, resulting from both fewer absolute numbers of TH-positive fibres and enlargement of the striosomal compartment.

Decreased numbers of TH-positive neurons in the SNC but not in the VTA of tgDISC1 rats

Dopaminergic fibres of the nigrostriatal, mesolimbic and mesocortical pathway arise from neurons in the SNC and the ventral tegmental area (VTA), innervating the dorsal striatum as well as the nucleus accumbens and cortex, respectively ⁵⁷. To assess potential alterations in dopaminergic neuron density in these regions, we performed a neuronal cell count on TH-immunostained brain sections from both tgDISC1 rats and LM. Transgenic animals showed a decrease in TH-positive neurons in the SNC ($p = 0.037$, Cohen's $d = 1.15$) but not in the VTA ($p = 0.557$, Figure 2).

Dopamine fibre density in the amygdala is not affected by DISC1 overexpression.

The amygdala plays a major role in processing affective and especially threatening stimuli as well as emotional memory consolidation. It receives its dopaminergic input from the VTA. To determine whether changes in dopamine fibre density, such as those observed in the striatum, were also present in the amygdala, we examined the size and TH-positive fibre density within the amygdalar complex (Figure 3A-C). The amygdala was divided into lateral, basal, and central nuclei.

Grey value measurement showed no differences between tgDISC1 rats and LM (Figure 3D-F). However, tgDISC1 rats displayed a slight but non-significant trend towards an increased size of the basal nucleus and a trend towards a slight shrinkage of the central nucleus (Supplementary Figure 1)

Altered interneuron distribution in the cortex, but no differences in cortical layer thickness in the tgDISC1 rat

To determine whether DISC1 overexpression causes alterations in quantity or distribution of immunolabelled interneurons, we examined cortical layer thickness as well as PV- and calbindin (CB)-positive interneuron numbers and distribution in tgDISC1 rats compared to LM (Figure 4A-C). Analysis was performed in the primary somatosensory cortex (S1), specifically in both the forelimb region (S1FL) and the barrel field (S1BF). Numbers of interneurons in both groups were compared in each cortical layer as well as the total cell count across all such layers. For the assessment of cortical thickness, adjacent Nissl stained sections were used. No differences in the cortical thickness were observed between the two groups (Figure 4D, E).

In tgDISC1 rats, numbers of PV-positive interneurons were decreased in layers II+III ($p = 0.005$, Cohen's $d = 1.72$) and increased in layers V ($p = 0.011$, Cohen's $d = 1.56$) and VI ($p = 0.019$, Cohen's $d = 1.69$), thus revealing a modified distribution pattern (Figure 4F-I). This shift of PV-positive interneurons to deeper cortical layers was present in both S1BF as well as in the S1FL region, being more pronounced in the S1BF. However, no changes in the total number of PV-positive

interneurons were observed ($p = 0.963$ in the S1FL and $p = 0.732$ in the S1BF, Figure 4).

CB-positive interneurons had similar distribution patterns in both tgDISC1 rats and LM, however, their total amount was slightly decreased in the S1BF but not in the S1FL of tgDISC1 rats (Supplementary Figure 2)

Macroscale neuroanatomy of tgDISC1 rats in comparison to LM

We utilized MR images from live, anesthized tgDISC1 rats ($n=21$) and LM ($n=24$) for automated, voxel-wise, tensor-based morphometry (TBM) analysis to examine neuroanatomical differences on a brain-wide scale. This analysis revealed no significant clusters of volume increase or decrease when comparing tgDISC1 rats and LM after FDR correction ($q = 0.1$), irrespective of controlling for variation in brain volume (Figure 5A, B). We therefore conducted a second-level analysis at an exploratory threshold of $p < 0.05$ (uncorrected for multiple comparisons). This revealed left-lateralized trend-level decreases in the volume of the corpus callosum and external capsule (Figure 5A). These remain, although to a reduced extent, when analysis was performed taking into account the variation in whole brain volume (Figure 5B). The latter analysis also revealed an additional left-lateralized cluster of voxels of increased volume in the hippocampus, particularly, the CA3 sub-field (Figure 5B). Consistent with the MRI investigations outlined above there were no significant changes in the volume of the cortex when comparing tgDISC1 rats and LM.

Increased hippocampal volume and decreased corpus callosum volume in tgDISC1 rats.

Despite the high power, the TBM analysis findings did not survive statistical correction for multiple comparisons. This may in part be due to the anisotropic 2D nature and low resolution of the MR images acquired, which may affect the precision of our image registration and negatively affect our ability to detect subtle anatomical differences between groups using TBM^{58 59}. Nevertheless, TBM demonstrated clear trends that might be regarded as “hypothesis generators” to guide further focused postmortem investigations, not based on any *a priori* hypotheses²⁹. Therefore, to evaluate the changes in the corpus callosum observed by NMR imaging at a finer scale, we calculated volume estimations using Cavalieri’s principle^{60 61} to confirm (or disprove) the trend suggested by the imaging findings. Our calculations confirmed a volumetric reduction of the corpus callosum ($p = 0.013$, Cohen’s $d = 1.3$). We could also confirm a slight increase in hippocampal volume of tgDISC1 rats ($p = 0.042$, Cohen’s $d = 1.3$). Due to the fact that TBM results were available after histological preparation of the brains, section sampling of the hippocampus did only partially allow an unequivocal comparison between animals. Therefore, we rather consider this as a supporting trend.

Discussion

DISC1 is a major vulnerability factor for a wide range of chronic mental illnesses,^{13,62,63,64} including schizophrenia, but without a clear segregation to a specific clinical diagnosis. We generated the tgDISC1 rat as a model for sporadic chronic mental illnesses, in contrast to existing mouse models which variously investigate the Scottish mutation, artificial mutations or partial knockouts of DISC1 (reviewed in ref. 41). The tgDISC1 rat displays protein pathology of which is of relevance for sporadic cases of mental illnesses and has a distinct phenotype of aberrant dopamine homeostasis⁵⁶. In this study, we report morphological aberrations in the tgDISC1 rat, both at light microscopical level and macroscale (MRI).

Our analyses revealed a decrease in dopaminergic neurons in the SNC, reduced fibre density in the striatum, and an increase in striosomal surface area. This is consistent with our previous finding of reduced whole dopamine content in the dorsal striatum⁵⁶. Dopaminergic neurons from the SNC send their neurites to the striatum, forming the nigrostriatal pathway⁵⁷. Both this decrease⁵⁷ and an increased striosomal surface area seem to be relevant factors leading to a reduced dopaminergic fibre density in the striatum. Imbalances between the striosomal compartment and the matrix have been reported to play a role in various neurological disorders, in which they are believed to be a causal factor of mood dysfunctions⁶⁵. Interference of DISC1 with the development of the dopaminergic system has been demonstrated in various DISC1 mouse models using neurochemistry, biochemistry and behavioural

tests (reviewed in ref. 41), however, to the best of our knowledge, in none of these models has a detailed neuroanatomical analysis been performed. Thus we conclude that DISC1 is involved in the development of a mature dopaminergic system and its overexpression and misassembly leads to structural abnormalities in the nigrostriatal pathway in the tgDISC1 rat.

Aberrations of interneurons have been reported in several mutant DISC1 mouse models^{42,47,48}, with a reduced number of PV-positive interneurons being the most frequently reported abnormality (reviewed in ref. 41). Indeed, tgDISC1 rats also show differences when compared to LM. The distribution of PV-interneurons is shifted to the deeper layers in tgDISC1 rats in both the S1BF and the S1FL. Since the thickness of the cortical layers does not change, the abnormal localization pattern is likely the consequence of a migration deficit, as opposed to a layer displacement. The possibility of a mutual relationship existing between altered cortical PV-positive neuronal distribution patterns and alterations in the dopaminergic system cannot be excluded.

Our observations resemble findings of *post mortem* studies on brains of schizophrenic patients which consistently reported alterations in GABAergic interneurons. Typically, PV-interneurons are reduced in the middle layers of the cortex, resulting in disturbed microcircuitry^{66,67}. The reduced PV-interneuron density in the middle layers demonstrated in tgDISC1 rats may potentially have similar effects, resulting in a discrepancy of the most important inhibitory input to pyramidal effector neurons in layers II/III. In contrast, CB-interneurons do not display such prominent changes, with their total number being slightly decreased in the BF. Looking at each individual cortex layer, a very subtle reduction can nevertheless be noted resulting in a discretely declined total cell count. Of note, dopaminergic

receptors seem to be selective for PV-interneurons, which could partially explain the more striking abnormalities in that interneuron subgroup⁶⁸.

TgDISC1 rats have previously been reported to have enlarged lateral ventricles, based on a blinded manual analysis⁵⁶. These findings are consistent with observations in schizophrenia: ventricular enlargement is a common morphological feature of schizophrenia, while a reduction of the corpus callosum has also been reported in several studies (reviewed in refs. 18,19). Furthermore, operator-independent, brain-wide TBM analysis of tgDISC1 rats suggests a trend-level reduction in the volume of the corpus callosum and the external capsule, which we could confirm in our subsequent volumetric measurement on a microscopic scale. Indeed, the enlargement of lateral ventricles and the volume of the corpus callosum are mutually dependent variables since an inverse correlation between the volumes of these structures has been recently demonstrated in an MRI study of schizophrenia patients⁶⁹.

In summary, the tgDISC1 rat displays morphological changes in several areas of the brain, including in dopaminergic cells and projections, cortical interneuron positioning and white matter/ventricle volume. To various extents, similar findings also have been reported in neuroanatomical investigations in *post mortem* brains of patients with schizophrenia. The present results also allow to establish a causal relationship between DISC1 overexpression / misassembly and a variety of seemingly unrelated morphological phenotypes, i.e. decreased dopaminergic cells and fibres, PV-positive interneurons, enlarged ventricles and decreased white matter in the corpus callosum. It is thus conceivable that a single molecular cause can simultaneously be linked to these morphological findings that, in part, have also been reported in patients with schizophrenia. We conclude that the tgDISC1 rat is therefore a face valid animal

model for sporadic schizophrenia, based not only on the described behavioural and physiological phenotype of aberrant dopamine homeostasis but also in terms of its neuroanatomical phenotype.

Materials and Methods

Animals

All experiments were conducted in conformity with the Animal Protection Law and were approved by local authorities (LANUV NRW, Recklinghausen, Germany). Male Sprague-Dawley rats were housed under 12 h light/dark conditions with *ad libitum* access to food and water.

Magnetic resonance imaging

Structural MRI analysis was performed as previously described⁵⁶ on a 7.0 tesla small animal Scanner (Bruker BioSpin, Billerica, MA, USA) with a horizontal bore magnet. We utilized voxel-wise tensor based morphometry (TBM) on coronal scans from tgDISC1 rats ($n=21$) and LM ($n=24$) to provide an unbiased, brain-wide overview of neuroanatomical differences between tgDISC1 rats and LM. TBM analysis was performed as previously described^{29,70,71}. Briefly, all brains in the study were first rigidly aligned using an automated intensity-based group-wise registration approach^{72,73}. A high-dimensional non-rigid registration algorithm was applied to warp each globally aligned scan to the population reference^{72,74,75}. Maps of localized volume difference at each voxel relative to the reference brain were computed from the log of the Jacobian determinant of this non-rigid transformation for each scan. Voxel-wise statistical tests over a brain mask were performed to establish regions of significant volumetric difference between groups. These analyses were performed for both absolute (6 degrees of freedom [dof] registration) and relative (9 dof registration) changes in volume, the latter taking into account population variation in anatomy due to differences in total brain volume⁵⁸. Voxel-wise analyses were corrected for

multiple comparisons using the false discovery rate (FDR) correction⁷⁶ at a threshold of $q=0.1$ (10%)⁵⁹. Additional exploratory analyses were performed at $p<0.05$, uncorrected for multiple comparisons.

Tissue preparation

Fourteen tgDISC1 rats and fourteen LM aged four to five months were deeply anesthetized with sodium-pentobarbital (70 mg/kg) and transcardially perfused with saline and heparin (400 ml, 10000 IU/l) followed by Zamboni fixative (400 ml, 4% paraformaldehyde and 10% picric acid in PBS, pH 7.4).

Brains were removed and post-fixed in Zamboni fixative for 48 h and incubated in sucrose solution (30% in PBS, pH 7.4) over a period of 48 h for cryoprotection. Subsequently, brains were frozen in isopentane at -40°C and stored at -80°C before further processing.

Frozen coronal sections of 50 μm thickness were cut using a microtome cryostat system (Leica SM2000R, Leica Biosystems, Wetzlar, Germany).

For Nissl staining, sections were mounted on tissue slides, air dried, and immersed in 70 % ethanol for 12 h. Treatment with 0.1% cresyl violet solution (Sigma-Aldrich, St. Louis, MO, USA) for 20 min was followed by differentiation in 70% ethanol for 10 min. Then, sections were dehydrated in ascending series of ethanol (70% for 10 min, 96% for 20 min and 100% for 30 min), treated with xylene (100%, twice for 10 min each) and covered with coverslips using DPX Mounting medium (Sigma-Aldrich, St. Louis, MO, USA).

Free floating immunohistochemical staining for each marker was carried out in multiwell plates in parallel on tissue sections from all animals.

Sections were initially incubated in blocking solution (10% normal goat serum, Vector laboratories, Burlingame, CA, USA, S-1000 and 0.3% saponin in PBS, pH 7.4) for 2 h at room temperature to reduce non-specific background staining. This was followed by incubation with the primary antibodies diluted in 2% normal goat serum and 0.1% saponin in PBS for 48h at 4°C. Anti-tyrosine-hydroxylase-antibody (1:100, mouse-monoclonal, Millipore, Billerica, MA, USA, MAB318), anti-parvalbumin-antibody (1:10.000, mouse-monoclonal, Sigma-Aldrich, St. Louis, MO, USA, P-3171) and anti-calbindin-antibody (1:10.000, mouse-monoclonal, Sigma-Aldrich, C-9848) were used as primary antibodies. Rinsing in PBS (pH 7.4) was carried out three times for 30 min. Biotinylated secondary anti-mouse antibodies (1:500, Dianova, Hamburg, Germany, 115-065-166) diluted in PBS (pH 7.4) were used as secondary antibodies and sections incubated for 24 h at 4°C, then rinsed in PBS (pH 7.4) three times for 15 min.

Sections were incubated with ABC reagent for 30 min at room temperature (ABC elite kit, 1:300, Vector laboratories, Burlingame, CA, USA, PK-6100) and washed out for 15 min using PBS (pH 7.4), followed by incubation with trisamine buffer twice for 15 min (pH 8.1, 0.05 M, Trizma[®] base, Sigma-Aldrich, T1503).

Sections were incubated with diaminobenzidine (DAB) (0.05% DAB in trisamine buffer, pH 7.6) for 10 min and hydrogen peroxide (1% H₂O₂, resulting in a final concentration of 0.01% H₂O₂ in the probes) was added for precisely 10 minutes to all sections. Afterwards the sections were rinsed three times in trisamine buffer (pH 7.6). Finally they were mounted on adhesive glass slides (HistoBond^R, P. Marienfeld, Lauda-Königshofen, Germany) and covered with coverslips using DPX Mounting medium (Sigma-Aldrich, St. Louis, MO, USA).

Tissue analysis

Anatomical regions were defined according to *The Rat Brain in Stereotaxic Coordinates* by Paxinos and Watson⁷⁷. Specimens were digitalized at a resolution of 0.01 $\mu\text{m}/\text{pixel}$ using a Zeiss Axio Z1 imager and Zen software (Zeiss, Jena, Germany). Tissue samples were blinded and randomized.

For the analysis of the dopamine pathway, equidistant brain slices of eight animals were used per group. Seven striatal sections between bregma 2.5 and -1.0 mm, six sections of the amygdala between bregma -2.0 and -3.2 mm, ten sections of the SNC between bregma -4.8 and -5.8 mm, and eight sections of the VTA between bregma -5.2 and -6.0 mm were analysed per animal.

Digitalized specimens were converted to 8-bit grey value images using IrfanView (Irfan Skiljan, Vienna, Austria).

The striatal region was digitally defined and the mean grey value and the area were measured using a script of Matlab software (MathWorks, Natick, MA, USA). In order to measure the mean grey value of the striatal matrix exclusively, a threshold of 120 was defined, resulting in the subtraction of the striosomes.

The amygdala was divided into basal, lateral and central nuclei and the mean grey value and area were measured using the above mentioned Matlab script (MathWorks, Natick, MA, USA).

Normalized reciprocals of the grey values are displayed in the charts in order to clarify the correlation between measured grey value and fibre density.

Neuronal cell count in the SNC and the VTA was performed using the multi-point tool in ImageJ (National Institute of Health, Bethesda, MD, USA).

For the analysis of CB- and PV-positive cortical interneurons, six animals were used per group. Analysis was performed in the forelimb region (S1FL) and the barrel field (S1BF) of the primary somatosensory cortex (S1) using eight equidistant sections per neuronal marker and anatomical region between bregma 1.2 and -0.2 and bregma -0.8 and -3.7, respectively.

A region of interest of 500 μm width and variable length comprising all cortical layers was set perpendicular to the white matter at random positions within the above-mentioned anatomical locations. Neurons were counted per cortex layer using the multi-point-tool in ImageJ (National Institute of Health, Methesda, MD, USA).

The thickness of the cortical layers was measured on consecutive Nissl stained sections perpendicularly to the white matter using ImageJ (National Institute of Health, Methesda, MD, USA).

Volumes of the CC and the anterior hippocampus were estimated using the Cavalieri's principle on PV-immunostained slides of six animals per group^{59,60}. Stereological analyses was performed on four sections per animal from bregma 2.0, 0.5, -1.0, and -2.5, in the CC and two sections per animal from bregma 2.7 and 3.6 in the hippocampus. A stereological counting grid was placed over the specimens and volumes were calculated from Cavalieri's principle:

$$V_{hip} = t \times a(p) \times \sum P_{hip}$$

$$V_{CC} = t \times a(p) \times \sum P_{CC}$$

in which V_{hip} and V_{CC} are the volumes of the hippocampus and the CC, t is the slice thickness, $a(p)$ the area associated with each point in the grid and $\sum P_{hip}$ and $\sum P_{CC}$ are the total number of points hitting the hippocampus and the CC, respectively.

Statistical analysis

All data were tested for normal distribution using D'Agostino-Pearson omnibus normality test and Kolmogorov-Smirnov normality test and two-tailed unpaired t-test was applied. Statistical analysis was performed using GraphPad Prism 6 software (GraphPad, La Jolla, CA, USA).

References

1. Bleuler, E. *Dementia Praecox oder Gruppe der Schizophrenien*, (Deuticke, Leipzig, Germany, 1911).
2. Bleuler, E. Die Prognose der Dementia Praecox. *Allgemeine Zeitschrift für Psychiatrie und psychisch-gerichtliche Medizin*. (1908).
3. Heckers, S. Bleuler and the neurobiology of schizophrenia. *Schizophr Bull* **37**, 1131-1135 (2011).
4. Kraepelin, E. *Klinische Psychiatrie. Siebente Auflage. I. Band*, (Barth Verlag, Leipzig, Germany, 1904).
5. Kraepelin, E. *Dementia praecox and Paraphrenia*, (E. & S. Livingstone, Edinburgh, United Kingdom, 1919).
6. Association, A.P. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*, (American Psychiatric Publishing, 2013).
7. Organization, W.H. *International Statistical Classification of Diseases and Related Health Problems*, (World Health Organization, 2004).
8. Howes, O.D. & Kapur, S. The dopamine hypothesis of schizophrenia: version III--the final common pathway. *Schizophr Bull* **35**, 549-562 (2009).
9. Bakhshi, K. & Chance, S.A. The neuropathology of schizophrenia: A selective review of past studies and emerging themes in brain structure and cytoarchitecture. *Neuroscience* **303**, 82-102 (2015).
10. Weinberger, D.R., Wagner, R.L. & Wyatt, R.J. Neuropathological studies of schizophrenia: a selective review. *Schizophr Bull* **9**, 193-212 (1983).
11. Harrison, P.J. The neuropathology of schizophrenia. A critical review of the data and their interpretation. *Brain* **122 (Pt 4)**, 593-624 (1999).
12. Prusiner, S.B. Shattuck lecture--neurodegenerative diseases and prions. *N Engl J Med* **344**, 1516-1526 (2001).
13. Korth, C. Aggregated proteins in schizophrenia and other chronic mental diseases: DISC1opathies. *Prion* **6**, 134-141 (2012).
14. Huber, G. [The pneumoencephalogram at the onset of schizophrenic disease]. *Arch Psychiatr Nervenkr Z Gesamte Neurol Psychiatr* **193**, 406-426 (1955).
15. Haug, J.O. Pneumoencephalographic evidence of brain atrophy in acute and chronic schizophrenic patients. *Acta Psychiatr Scand* **66**, 374-383 (1982).

16. Johnstone, E.C., Crow, T.J., Frith, C.D., Husband, J. & Kreel, L. Cerebral ventricular size and cognitive impairment in chronic schizophrenia. *Lancet* **2**, 924-926 (1976).
17. van Erp, T.G., *et al.* Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol Psychiatry* **21**, 547-553 (2016).
18. Innocenti, G.M., Ansermet, F. & Parnas, J. Schizophrenia, neurodevelopment and corpus callosum. *Mol Psychiatry* **8**, 261-274 (2003).
19. Wheeler, A.L. & Voineskos, A.N. A review of structural neuroimaging in schizophrenia: from connectivity to connectomics. *Front Hum Neurosci* **8**, 653 (2014).
20. Woodruff, P.W., McManus, I.C. & David, A.S. Meta-analysis of corpus callosum size in schizophrenia. *J Neurol Neurosurg Psychiatry* **58**, 457-461 (1995).
21. Hulshoff Pol, H.E., *et al.* Focal white matter density changes in schizophrenia: reduced inter-hemispheric connectivity. *Neuroimage* **21**, 27-35 (2004).
22. Shergill, S.S., *et al.* A diffusion tensor imaging study of fasciculi in schizophrenia. *Am J Psychiatry* **164**, 467-473 (2007).
23. Shenton, M.E., Dickey, C.C., Frumin, M. & McCarley, R.W. A review of MRI findings in schizophrenia. *Schizophr Res* **49**, 1-52 (2001).
24. Emsley, R., *et al.* Dorsal striatal volumes in never-treated patients with first-episode schizophrenia before and during acute treatment. *Schizophrenia Research* **169**, 89-94 (2015).
25. Chemerinski, E., *et al.* Larger putamen size in antipsychotic-naive individuals with schizotypal personality disorder. *Schizophr Res* **143**, 158-164 (2013).
26. van Erp, T.G., *et al.* A multi-scanner study of subcortical brain volume abnormalities in schizophrenia. *Psychiatry Res* **222**, 10-16 (2014).
27. Haijma, S.V., *et al.* Brain volumes in schizophrenia: a meta-analysis in over 18 000 subjects. *Schizophr Bull* **39**, 1129-1138 (2013).
28. Vernon, A.C., *et al.* Contrasting effects of haloperidol and lithium on rodent brain structure: a magnetic resonance imaging study with postmortem confirmation. *Biol Psychiatry* **71**, 855-863 (2012).
29. Vernon, A.C., *et al.* Reduced cortical volume and elevated astrocyte density in rats chronically treated with antipsychotic drugs-linking magnetic resonance imaging findings to cellular pathology. *Biol Psychiatry* **75**, 982-990 (2014).
30. Lewis, D.A. Inhibitory neurons in human cortical circuits: substrate for cognitive dysfunction in schizophrenia. *Curr Opin Neurobiol* **26**, 22-26 (2014).

31. Lewis, D.A., Curley, A.A., Glausier, J.R. & Volk, D.W. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci* **35**, 57-67 (2012).
32. Volk, D.W. & Lewis, D.A. Early developmental disturbances of cortical inhibitory neurons: contribution to cognitive deficits in schizophrenia. *Schizophr Bull* **40**, 952-957 (2014).
33. Zilles, K. & Amunts, K. Individual variability is not noise. *Trends Cogn Sci* **17**, 153-155 (2013).
34. Zilles, K. & Amunts, K. Centenary of Brodmann's map--conception and fate. *Nat Rev Neurosci* **11**, 139-145 (2010).
35. Kamiya, A., *et al.* A schizophrenia-associated mutation of DISC1 perturbs cerebral cortex development. *Nat Cell Biol* **7**, 1167-1178 (2005).
36. Kamiya, A., *et al.* DISC1-NDEL1/NUDEL protein interaction, an essential component for neurite outgrowth, is modulated by genetic variations of DISC1. *Hum Mol Genet* **15**, 3313-3323 (2006).
37. Lewis, D.A. & Levitt, P. Schizophrenia as a disorder of neurodevelopment. *Annu Rev Neurosci* **25**, 409-432 (2002).
38. St Clair, D., *et al.* Association within a family of a balanced autosomal translocation with major mental illness. *Lancet* **336**, 13-16 (1990).
39. Blackwood, D.H., *et al.* Schizophrenia and affective disorders--cosegregation with a translocation at chromosome 1q42 that directly disrupts brain-expressed genes: clinical and P300 findings in a family. *Am J Hum Genet* **69**, 428-433 (2001).
40. Sachs, N.A., *et al.* A frameshift mutation in Disrupted in Schizophrenia 1 in an American family with schizophrenia and schizoaffective disorder. *Mol Psychiatry* **10**, 758-764 (2005).
41. Lipina, T.V. & Roder, J.C. Disrupted-In-Schizophrenia-1 (DISC1) interactome and mental disorders: impact of mouse models. *Neurosci Biobehav Rev* **45**, 271-294 (2014).
42. Niwa, M., *et al.* Knockdown of DISC1 by in utero gene transfer disturbs postnatal dopaminergic maturation in the frontal cortex and leads to adult behavioral deficits. *Neuron* **65**, 480-489 (2010).
43. Steinecke, A., Gampe, C., Valkova, C., Kaether, C. & Bolz, J. Disrupted-in-Schizophrenia 1 (DISC1) is necessary for the correct migration of cortical interneurons. *J Neurosci* **32**, 738-745 (2012).
44. Ishizuka, K., *et al.* DISC1-dependent switch from progenitor proliferation to migration in the developing cortex. *Nature* **473**, 92-96 (2011).

45. Singh, K.K., *et al.* Common DISC1 polymorphisms disrupt Wnt/GSK3beta signaling and brain development. *Neuron* **72**, 545-558 (2011).
46. Steinecke, A., Gampe, C., Nitzsche, F. & Bolz, J. DISC1 knockdown impairs the tangential migration of cortical interneurons by affecting the actin cytoskeleton. *Front Cell Neurosci* **8**, 190 (2014).
47. Shen, S., *et al.* Schizophrenia-related neural and behavioral phenotypes in transgenic mice expressing truncated Disc1. *J Neurosci* **28**, 10893-10904 (2008).
48. Ayhan, Y., *et al.* Differential effects of prenatal and postnatal expressions of mutant human DISC1 on neurobehavioral phenotypes in transgenic mice: evidence for neurodevelopmental origin of major psychiatric disorders. *Mol Psychiatry* **16**, 293-306 (2011).
49. Hikida, T., *et al.* Dominant-negative DISC1 transgenic mice display schizophrenia-associated phenotypes detected by measures translatable to humans. *Proc Natl Acad Sci U S A* **104**, 14501-14506 (2007).
50. Pletnikov, M.V., *et al.* Inducible expression of mutant human DISC1 in mice is associated with brain and behavioral abnormalities reminiscent of schizophrenia. *Mol Psychiatry* **13**, 173-186, 115 (2008).
51. Kvajo, M., *et al.* A mutation in mouse Disc1 that models a schizophrenia risk allele leads to specific alterations in neuronal architecture and cognition. *Proc Natl Acad Sci U S A* **105**, 7076-7081 (2008).
52. Clapcote, S.J., *et al.* Behavioral phenotypes of Disc1 missense mutations in mice. *Neuron* **54**, 387-402 (2007).
53. Lee, F.H., *et al.* Disc1 point mutations in mice affect development of the cerebral cortex. *J Neurosci* **31**, 3197-3206 (2011).
54. Juan, L.W., *et al.* Phenotypic characterization of C57BL/6J mice carrying the Disc1 gene from the 129S6/SvEv strain. *Brain Struct Funct* **219**, 1417-1431 (2014).
55. Leliveld, S.R., *et al.* Insolubility of disrupted-in-schizophrenia 1 disrupts oligomer-dependent interactions with nuclear distribution element 1 and is associated with sporadic mental disease. *J Neurosci* **28**, 3839-3845 (2008).
56. Trossbach, S.V., *et al.* Misassembly of full-length Disrupted-in-Schizophrenia 1 protein is linked to altered dopamine homeostasis and behavioral deficits. *Mol Psychiatry* (2016).
57. Calabresi, P., Picconi, B., Tozzi, A., Ghiglieri, V. & Di Filippo, M. Direct and indirect pathways of basal ganglia: a critical reappraisal. *Nat Neurosci* **17**, 1022-1030 (2014).
58. Lerch, J.P., *et al.* Wanted dead or alive? The tradeoff between in-vivo versus ex-vivo MR brain imaging in the mouse. *Front Neuroinform* **6**, 6 (2012).

59. van Eede, M.C., Scholz, J., Chakravarty, M.M., Henkelman, R.M. & Lerch, J.P. Mapping registration sensitivity in MR mouse brain images. *Neuroimage* **82**, 226-236 (2013).
60. Gundersen, H.J., *et al.* Some new, simple and efficient stereological methods and their use in pathological research and diagnosis. *APMIS* **96**, 379-394 (1988).
61. Michel, R.P. & Cruz-Orive, L.M. Application of the Cavalieri principle and vertical sections method to lung: estimation of volume and pleural surface area. *J Microsc* **150**, 117-136 (1988).
62. Brandon, N.J., *et al.* Understanding the role of DISC1 in psychiatric disease and during normal development. *J Neurosci* **29**, 12768-12775 (2009).
63. Brandon, N.J. & Sawa, A. Linking neurodevelopmental and synaptic theories of mental illness through DISC1. *Nat Rev Neurosci* **12**, 707-722 (2011).
64. Korth, C. DISCopathies: brain disorders related to DISC1 dysfunction. *Rev Neurosci* **20**, 321-330 (2009).
65. Crittenden, J.R. & Graybiel, A.M. Basal Ganglia disorders associated with imbalances in the striatal striosome and matrix compartments. *Front Neuroanat* **5**, 59 (2011).
66. Benes, F.M. The role of stress and dopamine-GABA interactions in the vulnerability for schizophrenia. *J Psychiatr Res* **31**, 257-275 (1997).
67. Lewis, D.A., Pierri, J.N., Volk, D.W., Melchitzky, D.S. & Woo, T.U. Altered GABA neurotransmission and prefrontal cortical dysfunction in schizophrenia. *Biol Psychiatry* **46**, 616-626 (1999).
68. Le Moine, C. & Gaspar, P. Subpopulations of cortical GABAergic interneurons differ by their expression of D1 and D2 dopamine receptor subtypes. *Brain Res Mol Brain Res* **58**, 231-236 (1998).
69. Del Re, E.C., *et al.* Enlarged lateral ventricles inversely correlate with reduced corpus callosum central volume in first episode schizophrenia: association with functional measures. *Brain Imaging Behav* (2015).
70. Vernon, A.C., Crum, W.R., Johansson, S.M. & Modo, M. Evolution of extranigral damage predicts behavioural deficits in a rat proteasome inhibitor model of Parkinson's disease. *PLoS One* **6**, e17269 (2011).
71. Smith, E.J., *et al.* Implantation site and lesion topology determine efficacy of a human neural stem cell line in a rat model of chronic stroke. *Stem Cells* **30**, 785-796 (2012).
72. Crum, W.R., *et al.* A comparison of automated anatomical-behavioural mapping methods in a rodent model of stroke. *J Neurosci Methods* **218**, 170-183 (2013).

73. Crum, W.R., Modo, M., Vernon, A.C., Barker, G.J. & Williams, S.C. Registration of challenging pre-clinical brain images. *J Neurosci Methods* **216**, 62-77 (2013).
74. Crum, W.R., Tanner, C. & Hawkes, D.J. Anisotropic multi-scale fluid registration: evaluation in magnetic resonance breast imaging. *Phys Med Biol* **50**, 5153-5174 (2005).
75. Fox, N.C., *et al.* Imaging of onset and progression of Alzheimer's disease with voxel-compression mapping of serial magnetic resonance images. *Lancet* **358**, 201-205 (2001).
76. Genovese, C.R., Lazar, N.A. & Nichols, T. Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *Neuroimage* **15**, 870-878 (2002).
77. Paxinos, G. & Watson, C. *The Rat Brain in Stereotaxic Coordinates: Hard Cover Edition*, (Elsevier Science, 2006).
78. Tang, Y., Nyengaard, J.R., Pakkenberg, B. & Gundersen, H.J. Age-induced white matter changes in the human brain: a stereological investigation. *Neurobiol Aging* **18**, 609-615 (1997).

Acknowledgements

We thank L. Igdalova for technical assistance, C. Opfermann for imaging support, R. Hübbers for computational help and N. Bradshaw for language editing. Funding was obtained from NARSAD /BBR Independent Investigator Award (#20350), EU-FP7 (MC-ITN “IN-SENS” #607616), and the Research Commission of the Medical Faculty of the Heinrich Heine University Düsseldorf to C. K.

Author contributions statement

C.K., H.J.B. and A.K. supervised the project. Animals breeding, screening and matching was conducted by S. T. *Post mortem* experiments and microscopic analyses were carried out by H.H. MRI measurement was performed by C.C. and M.S. and TBM was carried out by A.V. and W.C. Results were discussed by H.H., S.V., V. B. C.K., H.J.B., A.V. and A.K. H.H. prepared the manuscript, S.V., V. B. C.K., A.V. and A.K. contributed to the script. All authors reviewed the manuscript.

Additional information

The authors declare no competing financial interest.

Correspondence to Carsten Korth.

Figures

Figure legends

Figure 1

Dopaminergic fibre density and striosomal surface in the striatum. (A) Schematic figure highlighting the relevant anatomical structures of (B) a representative section. (C) A close-up is displayed illustrating the striatum with a dashed line. Gray value and area measurements were obtained within this labelled ROI. The position with respect to bregma is indicated in the schematic (lateral) brain illustration. (D) Exemplary TH-immunostained striatal sections of each group with the upper specimen originating from a tgDISC1 rat and the lower specimen from a littermate control. Colours were inverted to better illustrate the fibre density, and close-ups are displayed on the right. Note the subtly sparser matrix and increase of striosomal surface in the tgDISC1 rat. (E) Grey value of the entire striatum. TgDISC1 rats display lower grey intensity in the entire striatum ($p = 0.005$). (F) Grey value of the striatal matrix without striosomes. TgDISC1 rats have a lower grey intensity in the striatal matrix ($p = 0.004$). (G) Area of the entire striatum including matrix and striosomal compartment. No difference in the surface area is detected between the two groups. (H) Area of the striatal matrix. No difference in the surface area of the striatal matrix is detected between the two groups. (I) Area of striosomes. TgDISC1 rats have an increased striosomal area ($p = 0.038$). Surface area is displayed in $\mu\text{m}^2 \pm \text{s.e.m.}$ and normalized reciprocals of the grey values are displayed in percent $\pm \text{s.e.m.}$

Abbreviations: anterior part of the anterior commissure (aca), cortex (C), corpus callosum (cc), external capsule (ec), striatum (St), olfactory tubercle (Tu)

Figure 2

Tyrosine Hydroxylase-positive neurons in the substantia nigra pars compacta and the ventral tegmental area. (A) Schematic figure highlighting the relevant anatomical structures of a (B) TH-immunostained exemplary section. (C) A close-up of these nuclei is displayed. The position with respect to bregma is indicated in the lateral schematic illustration. (D) TH-positive neurons in the SNC. TgDISC1 rats have a decreased number of TH-positive neurons in the SNC ($p = 0.037$) compared to littermate controls. (E) TH-positive neurons in the VTA. No difference in quantity of TH-positive neurons is observed in the VTA ($p = 0.557$). Neuronal cell counts are displayed in number of cells \pm s.e.m.

Abbreviations: cortex (C), cerebral peduncle (cp), hippocampus (H), medial terminal nucleus of the accessory optical tract (MT), substantia nigra, pars compacta (SNC), substantia nigra, pars reticulata (SNR), ventral tegmental area (VTA)

Figure 3

Dopaminergic fibre density in the amygdala. (A) Schematic figure highlighting the relevant anatomical structures of (B) a TH-immunostained exemplary section. (C) A close-up is displayed illustrating the amygdalar nuclei with a dashed line. Grey value and area measurements were carried out in the lateral, the basal and the central nucleus. The position with respect to bregma is indicated in the lateral schematic

illustration. (D-F) Grey value of the lateral, basal and central nucleus, respectively. No differences in grey value of the nuclei of the amygdala are detected between the two groups. The normalized reciprocals of the grey value are displayed in percent \pm s.e.m.

Abbreviations: amygdalostriatal transition area (ASt), basal amygdalar nucleus (BA), cortex (C), corpus callosum (cc), central amygdalar nucleus (CeA), external capsule (ec), dorsal endopiriform nucleus (DEn), Hippocampus (H), lateral amygdalar nucleus (LA), piriform cortex (Pir), striatum (S)

Figure 4

Distribution of Parvalbumin (PV)-positive interneurons in the SSC and cortex layer thickness. (A) Schematic figure highlighting the relevant anatomical structure of (B) a PV-immunostained exemplary section. (C) A close-up of a ROI of the barrel field is illustrated. ROIs of 500 μ m width and variable length covering all cortical layers were placed perpendicularly to the white matter and interneurons were counted per cortex layer. The position with respect to bregma is indicated in the lateral schematic illustration. (D) Thickness of individual cortex layers. No difference in the thickness of each individual cortex layer is observed between the two groups. (E) Cortex thickness including all layers. No difference between the two groups is detected in the thickness of the entire cortex. (F) PV-positive neurons per cortex layer in the BF. TgDISC1 rats show a decrease of PV-positive neurons in layer II+III ($p = 0.005$) and an increase in layers V ($p = 0.011$) and VI ($p = 0.019$) compared to littermate controls. (G) Total cell count of PV-positive neurons in the BF. No difference in total number of PV-positive neurons is detected. (H) PV-positive neurons per cortex layer

in the FL region. In tgDISC1 rats, PV-positive neurons are reduced in layer II+III ($p = 0.014$), but increased in layers V ($p = 0.022$) and VI ($p = 0.015$). (I) Total cell count of PV-positive neurons in the FL region. No difference in total number of PV-positive neurons is detected. Layer thickness is displayed in $\mu\text{m} \pm \text{s.e.m.}$, neuronal cell counts are displayed in number of cells $\pm \text{s.e.m.}$

Abbreviations: amygdala (Am), corpus callosum (cc), external capsule (ec), hippocampus (H), insular cortex (I), lateral ventricle (LV), motor cortex (M), piriform cortex (Pir), retrosplenial cortex (RS), barrel field of the primary somatosensory cortex (S1BF), dysgranular zone of the primary somatosensory cortex (S1DZ), trunk region of the primary somatosensory cortex (S1Tr), upper lip region of the primary somatosensory cortex (S1ULp), secondary somatosensory cortex (S2)

Figure 5

Neuroanatomical differences between LM and DISC1tg rats assessed through TBM.

Maps of voxel-wise local volumetric changes expressed as the delta log scaled jacobian determinant ($\Delta\log J$) in male DISC1tg rats ($n=21$) compared to LM (NC; $n=24$) at 6 months of age. Images are presented as either (A) global volume change (6 dof), or (B) absolute changes corrected for global changes (9 dof). Colour scales are for volume difference (cyan/blue to red/yellow) and the raw t statistical value at each voxel (dark to light green). Data shown are raw values of $\Delta\log J$ and their associated t -statistic at each voxel and following statistical comparisons at both

$p < 0.05$ uncorrected and following corrections for multiple comparisons
(FDR = $q < 0.1$).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

