



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Regulated expression of amyloidogenic immunoglobulin light chains in mice

Nuvolone, Mario ; Sorce, Silvia ; Pelczar, Pawel ; Rushing, Elisabeth ; Lavatelli, Francesca ; Rognoni, Paola ;
Valentini, Veronica ; Palladini, Giovanni ; Merlini, Giampaolo ; Aguzzi, Adriano

DOI: <https://doi.org/10.1080/13506129.2017.1289914>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-137096>


Journal Article

Accepted Version

Originally published at:

Nuvolone, Mario; Sorce, Silvia; Pelczar, Pawel; Rushing, Elisabeth; Lavatelli, Francesca; Rognoni, Paola; Valentini, Veronica; Palladini, Giovanni; Merlini, Giampaolo; Aguzzi, Adriano (2017). Regulated expression of amyloidogenic immunoglobulin light chains in mice. *Amyloid*, 24(Suppl 1):52-53.

DOI: <https://doi.org/10.1080/13506129.2017.1289914>



Regulated expression of amyloidogenic immunoglobulin light chains in mice.

Journal:	<i>Amyloid</i>
Manuscript ID	Draft
Manuscript Type:	Letter to the Editor
Date Submitted by the Author:	n/a
Complete List of Authors:	Nuvolone, Mario; Institute of Neuropathology - University Hospital Zurich; Amyloidosis Research and Treatment Center, Foundation Scientific Institute San Matteo, Department of Molecular Medicine, University of Pavia Sorce, Silvia; Institute of Neuropathology - University Hospital Zurich Pelczar, Pawel; Institute of Laboratory Animal Science, University of Zurich Rushing, Elisabeth; Institute of Neuropathology - University Hospital Zurich Lavatelli, Francesca; Amyloidosis Research and Treatment Center, Foundation Scientific Institute San Matteo, Department of Molecular Medicine, University of Pavia Rognoni, Paola; Amyloidosis Research and Treatment Center, Foundation Scientific Institute San Matteo, Department of Molecular Medicine, University of Pavia Valentini, Veronica; Amyloidosis Research and Treatment Center, Foundation Scientific Institute San Matteo, Department of Molecular Medicine, University of Pavia Palladini, Giovanni; Amyloidosis Research and Treatment Center, Foundation Scientific Institute San Matteo, Department of Molecular Medicine, University of Pavia Merlini, Giampaolo; Amyloidosis Research and Treatment Center, Foundation Scientific Institute San Matteo, Department of Molecular Medicine, University of Pavia Aguzzi, Adriano; Institute of Neuropathology - University Hospital Zurich
Keywords:	AL amyloidosis, Mouse models, Conditional transgenic mice, Immunoglobulin light chains

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For Peer Review Only

Regulated expression of amyloidogenic immunoglobulin light chains in mice.

Mario Nuvolone^{1,2}, Silvia Sorce¹, Pawel Pelczar³, Elisabeth Rushing¹, Francesca Lavatelli², Paola Rognoni², Veronica Valentini², Giovanni Palladini², Giampaolo Merlini² and Adriano Aguzzi¹.

¹*Institute of Neuropathology, University Hospital of Zurich, Zurich, Switzerland.*
²*Amyloidosis Research and Treatment Center, Foundation Scientific Institute San Matteo, Department of Molecular Medicine, University of Pavia, Pavia, Italy.* ³*Institute of Laboratory Animal Science, University of Zurich, Zurich, Switzerland.*

Address for correspondence: Adriano Aguzzi, Institute of Neuropathology, University Hospital of Zurich, Zurich, Switzerland. Email: adriano.aguzzi@usz.ch

Background. In immunoglobulin light chain (AL) amyloidosis, clinical and experimental observations converge towards a direct toxic role of the amyloidogenic light chain (aLC) precursor [1, 2, 3]. However, the molecular underpinnings of AL amyloidosis remain obscure, partly due to the paucity of preclinical models for this disease. Generation of conventional transgenic mice with constitutive overexpression of human aLCs is technically hampered by the potential toxicity of these proteins, possibly resulting in embryoletality and selection against high expressor lines [4]. To overcome these limitations, we have employed the Cre/loxP system to generate a conditional transgenic mouse allowing regulated expression of an aLC.

Materials and methods. The cDNA encoding a human aLC (λ), termed MAB, was subcloned in the pCAG-CAT-Oligo vector [5]. The resulting expression vector consisted of (5'→3'): a ubiquitous promoter (CAG), a reporter gene-stop cassette (CAT) flanked by two equally oriented loxP sites and the aLC (MAB). Transgenic mice obtained by pronuclear injection in C57BL/6J fertilized oocytes (termed CAG-CAT-MAB) were crossed with either mice expressing a constitutively active cre recombinase under the albumin promoter (Alb-cre) [6], or with mice expressing a tamoxifen-inducible cre recombinase under the CAG-promoter (CAGGS-creTM) [7]. Expression of human aLCs was analyzed by RT-PCR, immunohistochemistry, Western blotting and ELISA.

Results. Pronuclear injection of the CAG-CAT-MAB vector resulted in the generation of three transgenic lines. CAG-CAT-MAB^{Tg/+} mice showed ubiquitous expression of the CAT reporter gene, but no expression of aLC, as expected. Crossing with Alb-cre transgenic mice resulted in hepatocyte-restricted expression of aLC, whereas crossing with CAGGS-creTM transgenic mice lead to ubiquitous expression of aLC after tamoxifen administration (Figure 1). Human aLC could be detected in serum, and mice with ubiquitous expression show higher levels than mice with hepatocyte-restricted expression. Mice with hepatocyte-restricted expression of aLC have been monitored for up to two years, with no evidence of amyloid formation.




Figure 1. Regulated expression of amyloidogenic immunoglobulin light chains in mice. Immunohistochemical analysis of human λ expression (in red). Scale bar: 100 μ m.

Discussion and conclusions. We have generated a novel transgenic mouse in which the expression of aLC can be activated in a spatially and temporally regulated manner. In mice with hepatocyte-restricted expression of aLC, all target organs (except the liver) are exposed to aLC exclusively coming from the circulation, resembling the clinical situation. Ubiquitous expression of aLC results in higher levels of circulating aLC. Additional cre transgenic lines can be used to further manipulate the temporal and spatial pattern of aLC expression. This newly developed conditional transgenic mouse could prove instrumental to deepen our current mechanistic understanding of AL amyloidosis.

Declaration of interest. The authors have no conflict of interest to report.

References. 1. Nuvolone M, Merlini G. Systemic amyloidosis: novel therapies and role of biomarkers. *Nephrol Dial Transplant*. 2016.
 2. Liao R, Jain M, Teller P, Connors LH, Ngoy S, Skinner M, et al. Infusion of light chains from patients with cardiac amyloidosis causes diastolic dysfunction in isolated mouse hearts. *Circulation*. 2001;104:1594-7.
 3. Palladini G, Lavatelli F, Russo P, Perlini S, Perfetti V, Bosoni T, et al. Circulating amyloidogenic free light chains and serum N-terminal natriuretic peptide type B decrease simultaneously in association with improvement of survival in AL. *Blood*. 2006;107:3854-8.
 4. Ward JE, Ren R, Toraldo G, Soohoo P, Guan J, O'Hara C, et al. Doxycycline reduces fibril formation in a transgenic mouse model of AL amyloidosis. *Blood*. 2011;118:6610-7.
 5. Araki K, Araki M, Miyazaki J, Vassalli P. Site-specific recombination of a transgene in fertilized eggs by transient expression of Cre recombinase. *Proc Natl Acad Sci U S A*. 1995;92:160-4.
 6. Postic C, Shiota M, Niswender KD, Jetton TL, Chen Y, Moates JM, et al. Dual roles for glucokinase in glucose homeostasis as determined by liver and pancreatic beta cell-specific gene knock-outs using Cre recombinase. *J Biol Chem*. 1999;274:305-15.
 7. Hayashi S, McMahon AP. Efficient recombination in diverse tissues by a tamoxifen-inducible form of Cre: a tool for temporally regulated gene activation/inactivation in the mouse. *Dev Biol*. 2002;244:305-18.