



---

Year: 2017

---

## Prevalence of *Candida albicans* and *Streptococcus aureus* on maxillary obturators, maxillary defects and in saliva: A cross-sectional study

Celakil, Tamer ; Baca, Emrah ; Topcuoglu, Emine Nursen ; Röhlig, Bilge Gökçen ; Evlioglu, Gülümser ; Özcan, Mutlu

**Abstract:** Objective: This study investigated on the incidence of *Candida albicans* and *Staphylococcus aureus* adhesion onto maxillary obturator prostheses with soft lining materials placed by prosthodontists. Material and Methods: Between years 2013 and 2014, patients with maxillary defects due to maxillary tumor surgery and acquired obturator prostheses ( $n = 21$ , study group) were clinically evaluated and microbiological outcomes were compared with complete maxillary denture wearers ( $n = 21$ , control group). Data were collected on demographic features, histological type of cancer, presence of chemotherapy or radiotherapy, identification of soft lining materials used on obturator prostheses, cleansing methods practiced for obturator and maxillary complete prostheses. The presence of *Candida albicans* was determined on soft lining materials or polymethylmethacrylate surfaces and in the oral cavity, and *Staphylococcus aureus* in nasal mucosa and saliva. Data were analyzed using Kruskal Wallis, Fisher-Freeman-Halton tests ( $p < 0.05$ ). Results: The highest amount of *Candida albicans* was found on surfaces of prostheses, in the saliva and on maxillary defects on the hard palate and on definitive soft lining material with 90%, 90%, and 70%, respectively. The highest amount of *Staphylococcus aureus* was observed in the saliva and nasal cavity in the definitive soft lining material group (50%). Conclusion: Short renewal period of lining materials may cause less *Candida albicans* colonisation and denture stomatitis in the oral cavity. Patients who have undergone maxillectomy were more prone to *Staphylococcus aureus* colonisation.

DOI: <https://doi.org/10.14295/bds.2017.v20i3.1366>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-146603>

Journal Article

Accepted Version

Originally published at:

Celakil, Tamer; Baca, Emrah; Topcuoglu, Emine Nursen; Röhlig, Bilge Gökçen; Evlioglu, Gülümser; Özcan, Mutlu (2017). Prevalence of *Candida albicans* and *Streptococcus aureus* on maxillary obturators, maxillary defects and in saliva: A cross-sectional study. *Brazilian Dental Science*, 20(3):44-51.

DOI: <https://doi.org/10.14295/bds.2017.v20i3.1366>

**Prevalence of Candida albicans and Streptococcus aureus on maxillary  
obturators,  
maxillary defects and in saliva: A cross-sectional study**

**Tamer CELAKIL<sup>1</sup>, Emrah BACA<sup>1</sup>, Emine Nursen TOPCUOGLU<sup>2</sup>, Bilge Gokcen  
ROHLIG<sup>1</sup>,  
Gulumser EVLIOGLU<sup>1</sup>, Mutlu ÖZCAN<sup>3</sup>**

1-Istanbul University, Faculty of Dentistry, Department of Prosthodontics, Istanbul,  
Turkey

2-Istanbul University, Faculty of Dentistry, Department of Oral Microbiology, Istanbul,  
Turkey

3-University of Zurich, Division of Dental Materials, Center for Dental and Oral  
Medicine, Clinic for Fixed and Removable Prosthodontics and Dental Materials  
Science, Zurich, Switzerland

**Short title:** *Microorganism colonization on denture materials*

**\*Corresponding author:**

Dr. Tamer Celakil  
Istanbul University  
Faculty of Dentistry  
Department of Prosthodontics  
34093 Istanbul, Turkey  
E-mail: tamer.celakil@istanbul.edu.tr

## **ABSTRACT**

**Objective:** This study investigated on the incidence of *Candida albicans* and *Staphylococcus aureus* adhesion onto maxillary obturator prostheses with soft lining materials placed by prosthodontists. **Materials and Methods:** Between years 2013 and 2014, patients with maxillary defects due to maxillary tumor surgery and acquired obturator prostheses (n=21, study group) were clinically evaluated and microbiological outcomes were compared with complete maxillary denture wearers (n=21, control group). Data were collected on demographic features, histological type of cancer, presence of chemotherapy or radiotherapy, identification of soft lining materials used on obturator prostheses, cleansing methods practiced for obturator and maxillary complete prostheses. The presence of *Candida albicans* was determined on soft lining materials or polymethylmethacrylate surfaces and in the oral cavity, and *Staphylococcus aureus* in nasal mucosa and saliva. Data were analyzed using Kruskal Wallis, Fisher-Freeman-Halton tests ( $p < 0.05$ ). **Results:** The highest amount of *Candida albicans* was found on surfaces of prostheses, in the saliva and on maxillary defects on the hard palate and on definitive soft lining material with 90%, 90%, and 70%, respectively. The highest amount of *Staphylococcus aureus* was observed in the saliva and nasal cavity in the definitive soft lining material group (50%). **Conclusion:** Short renewal period of lining materials may cause less *Candida albicans* colonisation and denture stomatitis in the oral cavity. Patients who have undergone maxillectomy were more prone to *Staphylococcus aureus* colonisation.

## **KEYWORDS**

Candida albicans; Denture liners; Maxillary prosthesis; Palatal obturators; Staphylococcus aureus

## **INTRODUCTION**

Polymeric materials, mainly composed of polymethylmethacrylate (PMMA) or polyethylmethacrylate (PEMA), have been used in dentures for prosthodontic treatment of complete or partial edentulism [1]. However, PMMA based dentures may lead to discomfort in the denture bearing areas [2]. Acrylic or silicon-based materials have been used as soft lining materials for almost sixty years to facilitate comfort for denture wearing patients [2-5]. Such materials could be classified as temporary or definitive in accordance with their chemical composition (acrylic or silicon based) and may be divided into room temperature (cold polymerized) or at high temperature (heat polymerized) polymerizing categories depending on their processing methods [2,6] where biocompatibility is of importance.

Biocompatibility was defined as "ability of a biomaterial to perform its desired function with respect to a medical therapy, without eliciting any undesirable local or systemic effects in the recipient or beneficiary of that therapy but generating the most appropriate beneficial cellular or tissue response to that specific situation, and optimising the clinically relevant performance of that therapy" [7]. Denture base materials are also expected to be biocompatible and durable in the oral cavity environment [2] but such materials are subject of various unfavorable conditions in the oral cavity that may lead to alterations on their chemical, physical and mechanical properties [1,8]. Furthermore, surface properties of soft lining materials may contribute to colonization of microorganisms because loss of substances such

as plasticizers may lead to surface roughness as a result of loss of elasticity [2,9-11]. Surface roughness in particular usually causes irregular micropores and microorganisms adhering to these surfaces are sheltered from shear forces and cleaning procedures [11]. Adhered microorganisms to denture materials may then initiate colonization and development of pathogenesis [12-14]. Many species of microorganisms are colonized in the oral cavity and they usually act as commensal organisms in healthy individuals [15]. However, predisposing conditions, such as the presence of poor fit and/or poor oral hygiene accompanied by *Candida albicans*, may trigger for the development of opportunistic infections [16,17].

Denture stomatitis (DS) is an infection comprised of palatal mucosa under complete or partial removable dentures [18,19] and *Candida* species are generally related with high prevalence of DS that affects up to 65-70% of denture wearers [19-21]. Fungal infection risk due to DS becomes more serious and may yield to septicemia for immunosuppressed patients who have undergone maxillectomy due to tumor and have received radiotherapy or chemotherapy [18,22]. In 2008, in a surveillance study, Health Protection Agency (HPA) reported that *Candida albicans* was isolated in 70 % of the patients who had solid tumours [23].

*Staphylococcus aureus*, methicillin resistant (MRSA) and methicillin sensitive (MSSA) *Staphylococcus aureus*, is a pathogen generally found on the nasal membranes [24]. Naso-oral transmission of *Staphylococcus aureus* may occur through a congenital or acquired defect in the maxilla [24] where dentures may also act as carriage of this pathogen [25]. *Staphylococcus aureus* can survive from weeks to months on dry environmental surfaces such as denture acrylic resin and soft lining materials. Thus, some pharyngeal and respiratory tract infections may occur after this colonization [26-28]. In addition, MRSA may further cause skin

infections, pneumonia, meningitis, osteomyelitis, endocarditis, toxic shock syndrome, bacteremia and sepsis [29-32].

Soft lining materials may affect intraoral tolerance of the patients who have undergone irradiation, tumor surgery and thereafter, maxillary obturator prosthesis due to head and neck tumors [14,33] since when such materials are exposed to postnasal secretion, they act as a reservoir for microorganisms and cause complications in this immunosuppressed patients [14].

The objectives of this study therefore were to investigate *Candida albicans* adhesion onto maxillary obturator prostheses with soft lining materials placed by prosthodontists at university settings and to determine the presence of *Staphylococcus aureus* in these patients who have acquired maxillary defects due to maxillary tumor surgery.

## **MATERIAL AND METHODS**

### **Experimental design**

A cross-sectional study was planned to evaluate patients who had several cancer diagnosis, undergone maxillectomy and treated with maxillary obturator prostheses between years 2013 and 2014. Patients were recalled for microbiological analysis. Randomly selected, systemically healthy, complete denture wearers in the maxilla acted as the control group. All patients in the study and control group received their prostheses at the Department of Maxillofacial Prosthodontics, Istanbul University Turkey. Either prosthodontics specialists or doctorate program students placed dentures. Clinical Research Ethics Committee approved the study and informed content was obtained from all participants.

A total of 46 patients were identified from patient record system of whom 33 were rehabilitated with maxillary obturator prostheses. Seven patients could not be reached due to various reasons such as mortality or incorrect contact information. One patient did not want to participate yielding to a sample of 25 patients. After a thorough clinical assessment, 4 patients appeared not to fulfill the minimum inclusion criteria and final study group was formed with 21 participants (Figure 1). The control group also consisted of 21 patients who wore maxillary complete dentures and were treated between years 2013 and 2014.

All study group participants met the following criteria to be eligible for the study: 1) obturator prosthesis applied between 2013 and 2014, 2) presence of soft lining material on obturator prosthesis surface, 3) ability to clean prosthesis everyday, 4) presence of regularly renewed soft lining material at follow-up, 5) no smoking habit, 6) no antibiotics therapy within the last 8 weeks, 7) ability to read and understand the informed consent. The control group participants met the following criteria to be eligible for the study: 1) PMMA surface without any soft lining material on complete prosthesis surface, 2) ability to clean prosthesis everyday, 3) no smoking habit, 4) no antibiotics therapy within last 8 weeks, 5) ability to read and understand the informed consent.

Data obtained from participants included demographic information (age and gender), histological type of cancer, chemotherapy or radiotherapy history, identification of soft lining materials on the obturator, cleansing method for obturator prostheses or maxillary complete prostheses and the evaluation of the presence of *Candida albicans* on soft lining materials or PMMA surfaces and in the oral cavity (hard palate or maxillary defect and saliva) and *Staphylococcus aureus* in nasal mucosa and saliva. Soft lining materials used on obturator prostheses were

classified as temporary (Visco-gel, Dentsply Detrey GmbH, Konstanz, Germany) or definitive (Molloplast-B, Detax GmbH & Co KG, Ettlingen, Germany) in accordance with their chemical composition. Definitive lining materials and temporary lining materials were renewed respectively once every six months or every two months during routine follow-up. Cleansing method and frequency that participants preferred was recorded as rinsing with water, brushing or using immersion agents (sodium hypochlorite or cleansing tablet) and participants were informed that they could specify one or more of these cleansing method.

Microbiological examinations were performed at the Istanbul University, Department of Oral Microbiology. The presence of *Candida albicans* in the maxillary defects, different surfaces of the obturator prostheses as well as in the saliva, and the presence of *Staphylococcus aureus* in the nasal cavity and saliva were investigated. Samples from the nasal cavity, maxillary defects and different surfaces of the obturator prostheses were taken by sterile cotton swabs, transferred into a vessel with 1 mL saline solution and mixed 20 second for homogenization by means of a Vortex mixer. Aliquots of 0.1 mL samples were serially diluted 10-fold, plated onto Mannitol Salt Agar (Acumedia Manufacturers Inc, Baltimore, Maryland) for staphylococci and onto Sabouroud Dextrose Agar (Oxoid Ltd, Basingstone, UK) for yeasts and incubated aerobically at 37°C for 48 hours. *Candida albicans* isolates were identified by the germ tube test, while *Staphylococcus aureus* isolates were identified by DNase test performed using DNase test agar (Difco Lab, Detroit MI 48232-7058, USA). Methicillin resistance was detected with disc diffusion method on Mueller Hinton agar (Merck KgaA, 64271, Darmstadt, Germany) plates with 4% NaCl using 1µg oxacillin discs (Oxoid Ltd) according to Clinical and Laboratory Standards Institute (CLSI). Paraffin stimulated saliva samples were taken for 5

minutes. Aliquots of 0.1 ml samples were diluted 10-fold serially and plated onto Mannitol Salt Agar (Acumedia Manufacturers Inc) for staphylococci and onto Sabouroud Dextrose Agar (Oxoid Ltd) for yeasts and incubated aerobically at 37°C for 48 hours.

The parameters for presence of microorganisms were observed for temporary (Visco-gel) and definitive soft lining materials (Molloplast-B) and PMMA surfaces (control group). Even though the presence of microorganisms in the saliva, method and frequency of cleansing were recorded, the aim of the study was to focus on the presence of *Candida albicans* and *Staphylococcus aureus* on soft lining materials and PMMA surfaces only.

### **Statistical analysis**

The data were analysed using Number Cruncher Statistical System (2007 Statistical Software NCSS, LLC Kaysville, Utah, USA). Kruskal Wallis test was used for the comparisons of descriptive statistical methods (mean, standard deviation, median, frequency and ratio) as well as intergroup comparison of age parameter that was not normally distributed. Fisher-Freeman-Halton test was used for the comparison of qualitative data. The results were evaluated at 95% confidence interval at a significance level of 0.05.

## **RESULTS**

The mean age of the participants in this study was 58.6±9.6 years. More than half of the males (71.4%) wore obturators or maxillary complete prostheses whereas this accounted for only 28.6% of the women (Table I).

Prostheses types consisted of 11 obturator prostheses with Visco-gel (26.2%), 10 obturator prostheses with Molloplast-B (23.8%), and 21 maxillary complete

prostheses without any soft lining materials (50%). No statistically significant difference was determined between the distributions of mean age and gender of the patients according to the groups ( $p>0.05$ ) (Table II).

All study group participants who had several cancer diagnoses treated with radiotherapy and/or chemotherapy have undergone maxillectomy. The highest incidence cancer was squamous cell carcinoma (66.6%), while the least, invasive ductal carcinoma (4.7%).

The presence of *Candida albicans* on prostheses differed significantly according to type of surfaces (Fisher-Freeman-Halton  $p<0.05$ ) (Table III). No statistically significant difference was found between the incidence of *Candida albicans* in the saliva, hard palate or maxillary defects in all groups ( $p>0.05$ ) but the incidence of *Staphylococcus aureus* was significantly different in the saliva according to the groups (Fisher-Freeman-Halton,  $p>0.05$ ) (Table IV). There was also a statistically significant difference between the frequencies of *Staphylococcus aureus* in the nasal cavity according to the groups ( $p<0.05$ ) (Table V). Cleansing method and frequency did not significantly affect the results (Table VI).

## **DISCUSSION**

Studies present prevalence of denture stomatitis among denture wearers ranging from 15 to over 70%. Poor denture hygiene, *Candida* infection primarily due to *C. albicans*, and continuous wear of dentures are reported to be the main etiological factors for denture stomatitis [20]. In the present study, all obturator prostheses and maxillary complete prostheses were applied between 2013 and 2014. Therefore the age of dentures was similar. In addition, patients who have obturator prostheses or

maxillary complete dentures with poor denture hygiene were excluded from the study. Thus, all participants were able to clean their prostheses daily.

Previous studies emphasized water sorption as one of the main factors that cause dimensional changes in acrylic polymers as a consequence of plastic deformation and surface roughness [34,35]. An increase in the surface roughness may lead to increase in adhesion of microorganisms on polymer surfaces [36]. In addition, higher hydrophilicity of elastomers result in not only water sorption but also consequent loss of strength [37]. Surface roughness of room-temperature polymerized lining materials were reported to be more than heat-polymerized ones. Accordingly, most published in vitro studies reported higher Candida albicans adhesion on the room-temperature polymerized lining materials than heat-polymerised ones [10]. Moreover, soft lining materials were found to be more prone to microorganism adhesion than PMMA used for denture base materials [20]. In this retrospective clinical study, conversely, percentage positivity for the presence of Candida albicans was found similar in temporary lining material (Visco-gel) and PMMA groups. On the other hand, definitive lining materials (Molloplast-B) was more prone to microorganism adhesion than PMMA and temporary lining materials that is most probably related to clinical life of soft lining materials. It can be hypothesised that short renewal period that is 2 months, resulted in less Candida albicans colonisation.

Adherence of Candida albicans to PMMA or soft lining material surfaces may produce inflammation of adjacent mucosa surfaces or microorganism colonisation on the hard tissues that result in denture stomatitis [11]. Hygiene regimens may reduce Candida albicans adhesion and represent a significant improvement in the prevention of denture stomatitis [36]. In contrast to the previous reported studies

[2,38], the findings of this current study demonstrated that the presence of Candida albicans had no significant association with the cleansing method and frequency.

One limitation of this study was the limited number of patients rehabilitated with maxillary obturator prosthesis. Nevertheless, the findings of this study could help general practitioners and prosthodontists in selecting the most favourable type of soft lining materials used in conjunction with maxillary obturator prostheses.

## **CONCLUSIONS**

From this study, the following conclusions could be drawn:

- The highest amount of *Candida albicans* was found on the prosthesis surfaces with definitive soft lining material followed by saliva, hard palate/maxillary defects.
- The highest amount of *Staphylococcus aureus* was found in the saliva and nasal cavity in the definitive soft lining material group.
- Short renewal period of lining materials may cause less *Candida albicans* colonisation and denture stomatitis in the oral cavity.
- Patients who have undergone maxillectomy were more prone to *Staphylococcus aureus* colonization.

## **DISCLOSURE**

The authors claim to have no financial interest, either directly or indirectly, in the products or information presented in the article.

## REFERENCES

1. Bettencourt AF, Neves CB, de Almeida MS, Pinheiro LM, Oliveira SA, Lopes LP, Castro MF. Biodegradation of acrylic based resins: A review. *Dent Mater.* 2010;26:e171-80.
2. Brožek R, Koczorowski R, Rogalewicz R, Voelkel A, Czarnecka B, Nicholson JW. Effect of denture cleansers on chemical and mechanical behavior of selected soft lining materials. *Dent Mater.* 2011;27:281-90.
3. Jagger DC, Harrison A. Complete dentures--the soft option. An update for general dental practice. *Br Dent J.* 1997;182:313-7.
4. Hekimoğlu C, Anil N. The effect of accelerated ageing on the mechanical properties of soft denture lining materials. *J Oral Rehabil.* 1999;26:745-8.
5. Rueggeberg FA. From vulcanite to vinyl, a history of resins in restorative dentistry. *J Prosthet Dent.* 2002;87:364-79.
6. Murata H, Hamada T, Djulaeha E, Nikawa H. Rheology of tissue conditioners. *J Prosthet Dent.* 1998;79:188-99.
7. Williams DF. On the mechanisms of biocompatibility. *Biomaterials* 2008;29:2941-53.
8. Salim N, Satterthwaite JD, Rautemaa R, Silikas N. Impregnation with antimicrobials challenge bonding properties and water sorption behaviour of an acrylic liner. *J Dent.* 2012;40:693-9.
9. Murata H, Haberham RC, Hamada T, Taguchi N. Setting and stress relaxation behavior of resilient denture liners. *J Prosthet Dent.* 1998;80:714-22.
10. Kurtulmus H, Kumbuloglu O, Özcan M, Ozdemir G, Vural C. Candida

albicans adherence on silicone elastomers: effect of polymerisation duration and exposure to simulated saliva and nasal secretion. *Dent Mater.* 2010;26:76-82.

11. Al-Dwairi ZN, Al-Quran FA, Al-Omari OY. The effect of antifungal agents on surface properties of poly(methyl methacrylate) and its relation to adherence of *Candida albicans*. *J Prosthodont Res.* 2012;56:272-80.

12. Segal E. Pathogenesis of human mycoses: role of adhesion to host surfaces. *Microbiol Sci.* 1987;4:344-7.

13. Verran J, Maryan CJ. Retention of *Candida albicans* on acrylic resin and silicone of different surface topography. *J Prosthet Dent.* 1997;77:535-9.

14. Vural C, Ozdemir G, Kurtulmus H, Kumbuloglu O, Özcan M. Comparative effects of two different artificial body fluids on *Candida albicans* adhesion to soft lining materials. *Dent Mater J.* 2010;29:206-12.

15. Cannon RD, Chaffin WL. Oral colonization by *Candida albicans*. *Crit Rev Oral Biol Med.* 1999;10:359-83.

16. Pereira-Cenci T, Deng DM, Kraneveld EA, Manders EM, Del Bel Cury AA, Ten Cate JM, **et al.** The effect of *Streptococcus mutans* and *Candida glabrata* on *Candida albicans* biofilms formed on different surfaces. *Arch Oral Biol.* 2008;53:755-64.

17. Wilkieson C, Samaranayake LP, MacFarlane TW, Lamey PJ, MacKenzie D. Oral candidosis in the elderly in long term hospital care. *J Oral Pathol Med.* 1991;20:13-6.

18. Jeganathan S, Lin CC. Denture stomatitis--a review of the aetiology, diagnosis and management. *Aust Dent J.* 1992;37:107-14.

19. Mima EG, Pavarina AC, Silva MM, Ribeiro DG, Vergani CE, Kurachi C, Bagnato VC. Denture stomatitis treated with photodynamic therapy: five cases. *Oral*

Surg Oral Med Oral Pathol Oral Radiol Endod. 2011;112:602-8.

20. Gendreau L, Loewy ZG. Epidemiology and etiology of denture stomatitis. *J Prosthodont* 2011;20:251-60.

21. Valentini F, Luz MS, Boscato N, Pereira-Cenci T. Biofilm formation on denture liners in a randomised controlled in situ trial. *J Dent*. 2013;41:420-7.

22. Epstein JB. Diagnosis and treatment of oropharyngeal candidiasis. *Oral Maxillofac Surg Clin North Am* 2003;15:91-102.

23. Pereira Gonzales F, Maisch T. Photodynamic inactivation for controlling *Candida albicans* infections. *Fungal Biol* 2012;116:1-10.

24. Tuna EB, Topçuoğlu N, İlhan B, Gençay K, Kulekçi G. *Staphylococcus aureus* transmission through oronasal fistula in children with cleft lip and palate. *Cleft Palate Craniofac J* 2008;45:477-80.

25. Rossi T, Laine J, Eerola E, Kotilainen P, Peltonen R. Denture carriage of methicillin-resistant *Staphylococcus aureus*. *Lancet* 1995;345:1577.

26. Johanson WG Jr, Pierce AK, Sanford JP, Thomas GD. Nosocomial respiratory infections with gram negative bacilli. The significance of colonization of the respiratory tract. *Ann Intern Med*. 1972;77:701-6.

27. Matsuura T, Abe Y, Sato Y, Okamoto K, Ueshige M, Akagawa Y. Prolonged antimicrobial effect of tissue conditioners containing silver-zeolite. *J Dent*. 1997;25:373-7.

28. Cimolai N. MRSA and the environment: implications for comprehensive control measures. *Eur J Clin Microbiol Infect Dis*. 2008;27:481-93.

29. Peacock JE Jr, Marsik FJ, Wenzel RP. Methicillin resistant *Staphylococcus aureus*: introduction and spread within a hospital. *Ann Intern Med*. 1980;93:526-32.

30. Haley RW, Hightower AW, Khabbaz RF, Thornsberry C, Martone WJ, Allen

JR, Hughes JM. The emergence of methicillin-resistant *Staphylococcus aureus* infections in United States hospitals. Possible role of the house staff-patient transfer circuit. *Ann Intern Med.* 1982;97:297-308.

31. Kluytmans J, van Belkum A, Verbrugh H. Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol Rev.* 1997;10:505-20.

32. Cole AM, Tahk S, Oren A, Yoshioka D, Kim YH, Park A, Ganz T. Determinants of *Staphylococcus aureus* nasal carriage. *Clin Diagn Lab Immunol.* 2001;8:1064-9.

33. Gronet PM, Driscoll CF, Hondrum SO. Resiliency of surface-sealed temporary soft denture liners. *J Prosthet Dent.* 1997;77:370-4.

34. Woelfel JB, Paffenbarger GC, Sweeney WT. Some physical properties of organic denture base materials. *J Am Dent Assoc.* 1963;67:499-504.

35. Dixon DL, Breeding LC, Ekstrand KG. Linear dimensional variability of three denture base resins after processing and in water storage. *J Prosthet Dent.* 1992;68:196-200.

36. Lazarin AA, Machado AL, Zamperini CA, Wady AF, Spolidorio DM, Vergani CE. Effect of experimental photopolymerized coatings on the hydrophobicity of a denture base acrylic resin and on *Candida albicans* adhesion. *Arch Oral Biol.* 2013;58:1-9.

37. Liao WC, Pearson GJ, Braden M, Wright PS. The interaction of various liquids with long-term denture soft lining materials. *Dent Mater.* 2012;28:e199-206.

38. Odman PA. The effectiveness of an enzyme containing denture cleanser. *Quintessence Int.* 1992;23:187-90.

**Legends for figures and tables:**

**Figures:**

**Figure 1-** Sequence of selection for the designated cohort.

**Tables:**

**Table I-** Demographic characteristics of the study cohort (t: Visco-gel; d: Molloplast-B; c: Control group).

**Table II-** Evaluation of demographic findings related to groups. <sup>a</sup>Kruskal Wallis test was used for the comparisons of descriptive statistical methods as well as intergroup comparison of age parameter. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data. See Table I for group abbreviations.


**Table III-** Presence of Candida Albicans related to the groups. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data See Table I for group abbreviations.

**Table IV-** Presence of Staphylococcus Aureus related to groups. See Table I for group abbreviations. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data.

**Table V-** Findings related to cleansing methods and frequency of cleansing.

**Table VI-** Evaluation of cleansing methods and frequency findings related to groups. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data.

**Figures:**



**Figure 1-** Sequence of selection for the designated cohort.

**Tables:**

		<b>Min-Max</b>	<b>Mean (SD)</b>
<b>Age (Year)</b>		38-74	58.57(9.64)
		<b>n</b>	<b>%</b>
<b>Gender</b>	<b>Men</b>	30	71.4
	<b>Women</b>	12	28.6
<b>Group</b>	<b>Temporary lining material (t)</b>	11	26.2
	<b>Definitive lining material (d)</b>	10	23.8
	<b>PMMA (control group) (c)</b>	21	50.0

**Table I-** Demographic characteristics of the study cohort (t: Visco-gel; d: Molloplast-B; c: Control group).

		<b>t</b> (n=11)	<b>d</b> (n=10)	<b>c</b> (n=21)	<b>P</b>
<b>Age (Year)</b>	<b>Mean (±SD)</b>	55 (10.13)	59.9 (9.66)	59.81(9.36)	<sup>a</sup> 0.463
	<b>Median</b>	(58)	(56.5)	(63)	
<b>Gender</b>	<b>Men</b>	8 (72.7)	9 (90)	13 (61.9)	<sup>b</sup> 0.268
	<b>Women</b>	3 (27.3)	1 (10)	8 (38.1)	

**Table II-** Evaluation of demographic findings related to groups. <sup>a</sup>Kruskal Wallis test was used for the comparisons of descriptive statistical methods as well as intergroup comparison of age parameter. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data. See Table I for group abbreviations.

	<b>t (%)</b>	<b>d (%)</b>	<b>c (%)</b>	<b>P</b>
<b>Saliva</b>	8 (72.7)	9 (90)	12 (57.1)	0.195
<b>Hard Palate/Maxillary Defect</b>	3 (27.3)	7 (70)	7 (33.3)	0.122
<b>Prosthesis Surface</b>	5 (45.5)	9 (90)	9 (42.9)	0.037

**Table III-** Presence of Candida Albicans related to the groups. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data See Table I for group abbreviations.

	<b>t (%)</b>	<b>d (%)</b>	<b>c (%)</b>	<b>P</b>
<b>Saliva</b>	5 (45,5)	5 (50)	2 (9,5)	<b>0.020</b>
<b>Nasal Cavity</b>	5 (45,5)	5 (50)	2 (9,5)	<b>0.020</b>

**Table IV-** Presence of Staphylococcus Aureus related to groups. See Table I for group abbreviations. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data.

		<b>n (%)</b>
<b>Cleansing methods</b>	Rinsing with water	23 (54.8)
	Brushing	12 (28.6)
	Water+Brushing+Cleaning tablet	3 (7.1)
	Water+Sodium hypochlorite	2 (4.8)
	Water+Cleaning tablet	2 (4.8)
<b>Frequency of cleansing</b>	Once a day	25 (59.5)
	Two or three times a day	17 (40.5)

**Table V-** Findings related to cleansing methods and frequency of cleansing.

		<b>t (%)</b>	<b>d (%)</b>	<b>c (%)</b>	<b>P (%)</b>
<b>Cleansing methods</b>	Rinsing with water	7 (63.6)	8 (80)	8 (38.1)	0.601
	Brushing	4 (36.4)	2 (20)	6 (28.6)	
	Water+Brushing+Cleaning tablet	0	0	3 (14.3)	
	Water+Sodium hypochlorite	0	0	2 (9.5)	
	Water+Cleaning tablet	0	0	2 (9.5)	
<b>Frequency of cleansing</b>	Once a day	7 (63.6)	6 (60)	12 (57.1)	0.938
	Two or three times a day	4 (36.4)	4 (40)	9 (42.9)	

**Table VI-** Evaluation of cleansing methods and frequency findings related to groups. <sup>b</sup>Fisher-Freeman-Halton test was used for the comparison of qualitative data.