**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Brain stimulation studies of social norm compliance: implications for personality disorders?

Ruff, Christian C

Abstract: Several personality disorders involve pathological behaviors that violate social norms, commonly held expectations about what ought to be done in specific situations. These symptoms usually emerge early in development, are persistent and hard to treat, and are often ego-syntonic. Here I present some recent brain stimulation studies suggesting that pathological changes in different aspects of norm-compliant behavior reflect dysfunctions of brain circuits involving distinct prefrontal brain areas. One set of studies shows that transcranial direct current stimulation of the right lateral prefrontal cortex changes the behavioral sensitivity to social incentives for norm-compliant behavior. Crucially, social norm compliance in response to such incentives could even be increased during excitatory stimulation, demonstrating that the affected neural process is a biological prerequisite for appropriate reaction to social signals that trigger norm compliance. In another set of studies, we show that stimulation of a different (more dorsal) part of the right prefrontal cortex enhances honesty in a realistic setting where participants had the opportunity to cheat for real monetary gains. Interestingly, these stimulation-induced increases in both socially cued or purely voluntary norm compliance were not linked to changes in other aspects of decision-making (such as risk or impatience), and they did not reflect changes in beliefs about what is appropriate behavior. These results suggest that disorders of distinct brain circuits may causally underlie egosyntonic changes in norm-compliant behavior. This raises the tantalizing possibility that pathologies of norm-compliant behavior may be ameliorated by interventions targeting the function of these brain circuits.

DOI: <https://doi.org/10.1159/000486898>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-151913>

Journal Article

Accepted Version

Originally published at:

Ruff, Christian C (2018). Brain stimulation studies of social norm compliance: implications for personality disorders? *Psychopathology*, 51(2):105-109.

DOI: <https://doi.org/10.1159/000486898>

Special Issue: Personality Disorders, Functioning and Health

**Brain stimulation studies of social norm compliance:
Implications for personality disorders?**

Christian C. Ruff
Laboratory for Social and Neural Systems Research (SNS-Lab)
Department of Economics
University of Zurich

Address:

Christian C. Ruff
Department of Economics
University of Zurich
Blumlisalpstrasse 10
8006 Zurich
Switzerland
Phone: +41 44 634 5067
Fax: +41 44 634 4907

Abstract:

Several personality disorders involve pathological behaviours that violate social norms, commonly held expectations about what ought to be done in specific situations. These symptoms usually emerge early in development, are persistent and hard to treat, and are often egosyntonic. Here I present some recent brain stimulation studies suggesting that pathological changes in different aspects of norm-compliant behaviour reflect dysfunctions of brain circuits involving distinct prefrontal brain areas. One set of studies shows that transcranial direct current stimulation of right lateral prefrontal cortex changes the behavioural sensitivity to social incentives for norm-compliant behaviour. Crucially, social norm compliance in response to such incentives could even be increased during excitatory stimulation, demonstrating that the affected neural process is a biological prerequisite for appropriate reaction to social signals that trigger norm compliance. In another set of studies, we show that stimulation of a different (more dorsal) part of right prefrontal cortex enhances honesty in a realistic setting where participants had the opportunity to cheat for real monetary gains. Interestingly, these stimulation-induced increases in both socially-cued or purely voluntary norm compliance were not linked to changes in other aspects of decision-making (such as risk or impatience), and they did not reflect changes in beliefs about what is appropriate behaviour. These results suggest that disorders of distinct brain circuits may causally underlie egosyntonic changes in norm-compliant behaviour. This raises the tantalizing possibility that pathologies of norm-compliant behaviour may be ameliorated by interventions targeting the function of these brain circuits.

Main text:

Most aspects of our behaviour are regulated by social norms, widely shared standards specifying how we ought to behave in a given situation (i.e., we should be fair, honest, punctual, etc.). Such norms exist in all known societies and are essential for peaceful and productive interactions^{1,2}, since they minimize the need for explicit coordination between interaction partners. Prevailing social norms are usually known by all members of a given society; nevertheless, the power of norms to guide collective behaviour erodes in the presence of just a few norm-violators, since people follow social norms only if all others comply with it³. Norms are therefore often enforced by social sanction threats that are present both at an official level (e.g., legal codes specifying punishments for norm violations) and at the level of private peer-to-peer interactions (i.e., we know that other people will privately punish norm transgressions). The importance of credible sanction threats for preserving norm compliance is well documented through evolutionary, ethnographic, and laboratory studies¹⁻⁵.

While most people effortlessly comply with prevailing norms, some individuals routinely violate them and even break the law. These individuals are often diagnosed with a personality disorder, specifically if the corresponding behaviour emerges early in life, is stable across time and hard to change by interventions, and is egosyntotic. Depending on the age of emergence, the diagnostic guidelines used, and other specifics of the symptomatology, the corresponding diagnoses may be Antisocial personality disorder, Dissocial personality disorder, Psychopathy, Sociopathy, or Conduct disorder⁶⁻⁸. These disorders are common - with estimated prevalences ranging from 1% - 4%^{9,10} - and they are associated with significant financial and emotional costs to society and the affected individuals.

The origins of personality disorders associated with norm-violations are debated. Both genetic predispositions^{11,12} and environmental factors¹²⁻¹⁴ - such as prenatal influences and early childhood experiences - may play a role. However, it is largely unclear what psychological and/or biological mechanisms may be disrupted to bring about the symptoms of personality disorders. Perhaps even more importantly, very little is known about what psychological and/or biological processes should be targeted with external interventions to remedy pathological norm-violating behaviour (though see¹⁵).

Here I describe the results of brain stimulation studies suggesting that two distinct aspects of pathological norm-violating behaviour may reflect dysfunctions in brain circuits involving specific parts of the prefrontal lobe. Importantly, the experiments also suggest that these aspects of norm-violating behaviour may - at least in principle - be ameliorated by interventions that target the excitability of these brain circuits.

In the first set of studies, young healthy participants faced a series of choices about how to allocate money between themselves (the proposer) and a fully anonymous interaction partner (the recipient). The roles of proposer and recipient were randomly determined and the resulting payoffs were real, resulting in a salient fairness norm of a 50:50 split. In one condition measuring voluntary compliance with this norm, the proposed transfer was implemented without any possible consequences. In line with other findings¹⁶, participants in this condition adhered to the fairness norm only weakly and transferred around 15 % of the total amount¹⁷. However, in another condition, the recipient could react to the transfer by investing some of his/her own payoff to take money away from the proposer. This threat of a possible sanction led to a considerable increase in sanction-induced norm compliance (mean proposed transfers of around 45%). The magnitude of this behavioural reaction to the social sanction threat has been shown to correlate with increased neural activity in the right inferolateral prefrontal cortex¹⁸ (rLPFC), leading us to hypothesize that this brain area may causally underlie the behavioural sensitivity to social incentives for compliance with social norms.

In our experiment, we employed transcranial direct current stimulation (tDCS) to experimentally increase (anodal tDCS) or decrease (cathodal tDCS) the excitability of this rLPFC region (see blue circle in Figure 1C). These activity modulations are implemented by means of weak electrical currents running between two electrodes, which change neural excitability in relatively large swathes of cortex, thereby facilitating or impeding neural activity under the electrodes elicited by behavioural control¹⁹. The stimulation indeed changed sanction-induced norm compliance, with transfer increases that were 30% higher (anodal) or 30% lower (cathodal) than in a placebo stimulation condition (see Figure 1A). Crucially, control experiments showed that these behavioural changes reflected a heightened sensitivity to social incentives for norm compliance, rather than changes in the willingness to share money generally¹⁷. The increased transfers also did not relate to

changes in risk-taking or cognitive abilities necessary to adequately react to the two conditions, demonstrating that the stimulation-induced changes reflect a specific mechanism for social norm compliance and that the tDCS, despite its limited spatial resolution, did not affect other aspects of behavioural control. Interestingly, the tDCS-elicited increase in sanction-induced norm compliance occurred without any change in the participants' beliefs or perceptions of the opponent reactions or the fairness norm itself (see Figure 1B). Thus, the pattern of results suggests that the stimulated prefrontal circuit causally controls the behavioural sensitivity to social incentives for non-compliant behaviour, in a manner that is not directly accessible to conscious report of the participants.

Figure 1: Two prefrontal circuits for norm compliance. **A:** tDCS over the blue-colored region in panel **C** leads to changes in norm-compliance (compared to neurally ineffective sham tDCS) that are motivated by social incentives. When no social sanctions were present (right bar plot), excitatory anodal tDCS (An) decreased and inhibitory cathodal tDCS (Ca) increased voluntary compliance with the fairness norm. However, when social sanctions were present (left bar plot), excitatory anodal tDCS lead to higher transfers (more compliance) and inhibitory cathodal tDCS to lower transfers (less compliance). Please note that the circles are much smaller than electric fields elicited by tDCS, but that the effective resolution of neural excitability changes due to these fields may also depend on the presence of neural activity related to behaviour, which can be much more focal. **B:** The tDCS-induced changes in norm compliance were not accompanied by any changes in beliefs about what constitutes fair behaviour (left line plot) or the possible consequences of norm transgressions (right line plot). For both **A** and **B**, see¹⁷ for further information. **C:** Schematic location of the two prefrontal tDCS target areas employed in¹⁷ (blue circle) and in²¹ (red circle). **D:** Participants with neurally ineffective sham tDCS over the red area in panel **C** lied substantially in the die-rolling game, as indicated by the shift

*towards implausibly high outcomes in the observed distribution of reported outcomes (in gray) compared to the statistically expected distribution for honest reporting (in black). **E**: Honest reporting is increased in participants with excitatory anodal tDCS over the red area in panel **C**. Compared to the sham distribution in panel **D**, the observed distribution of reported outcomes in the anodal group (in red) is systematically shifted back towards the statistically expected distribution (in black). For both **D** and **E**, see²¹ for further information.*

Our second set of experiments highlights that purely voluntary norm compliance, in the explicit absence of any social incentives, may require a brain circuit involving a different prefrontal area. These studies focussed on honesty, a behaviour that is purely motivated by the private determination to comply with this social norm (since lying is not evident to outside observers). We again employed tDCS, but now over a more dorsal area in right prefrontal cortex (see red circle in Figure 1C) that had previously been shown to be activated when participants told the truth despite having the option to cheat for money²⁰. We therefore hypothesized that this region and its interconnected structures may be involved in representing the motivation to remain honest in situations where this motive conflicts with the temptation to lie for financial benefit.

In the experiment, participants completed a series of tasks that they thought measured determinants of luck. One of these tasks offered them the opportunity to cheat in order to gain money. They rolled a die 10 times and entered the corresponding outcomes on a computer screen, which told them explicitly which three of the six possible outcomes was associated with a monetary gain. Participants therefore faced the temptation to misreport unsuccessful die outcomes, and they knew that such cheating could not be detected (since the die rolls were unobserved and all outcomes were equally possible). However, by analysing the distribution of earnings across the different stimulation groups, we could statistically determine how the level of cheating was influenced by brain stimulation.

Participants in the group with placebo brain stimulation lied substantially (on 37% of all possible occasions, see Figure 1D), but this tendency to cheat was reduced by half (to 15%) during excitatory anodal tDCS (see Figure 1E). The tDCS-related increase in honesty was highly specific to situations where the honesty norm conflicted with the motivation for private monetary gain, as the tDCS affected neither purely financial choices (e.g., risky choices or choices with rewards given at different timepoints) nor choices between different moral motives (e.g., lying for someone

else's benefits)²¹. Again, this stimulation effect on compliance with the honesty norm was not accompanied by changes in the perceptions of the social norm or the appropriateness of the behaviour. Thus, in close similarity to the findings concerning sanction-induced norm compliance, our results suggest that the stimulated (different) circuit is a biological prerequisite for compliance with social norms, but now those that are motivated purely internally. Moreover, our results suggest that this circuit also operates in a way that is not consciously accessible to the decision-maker.

That prefrontal cortex dysfunction may generally underlie symptoms of personality disorders has been proposed for a long time²². This general hypothesis concurs with the well-established role of prefrontal cortex in the executive control of behavior²³ and with the timecourse of development of this brain structure that parallels the emergence of relevant symptoms²⁴. Moreover, neuroimaging evidence already points to dysfunctions of both medial and lateral prefrontal circuits in personality disorders^{25,26}. However, our data demonstrate several important and more detailed aspects of this prefrontal involvement. First, they suggest that circuits involving distinct prefrontal areas may underlie selective disruptions of social norm compliance that need not be accompanied by more general behavioural deficits in non-social contexts (e.g., risk-taking, general impulsivity). Second, they suggest that these prefrontal circuits themselves are functionally specialised for norm compliant behaviour that is either motivated by social incentives (right infero-lateral PFC) or purely internally (right dorso-lateral PFC). Third, they show that these circuits operate in a fashion that is dissociated from explicit judgements about norms and appropriate behaviour (see also^{27,28} for potentially convergent findings), which may form a basis for egosyntonic norm-violating behaviours in various personality disorders. More generally, these findings suggest that overt behaviour may reflect behavioural control by brain circuits that are distinct from other neural mechanisms involved in explicit knowledge and verbal reports about appropriate behaviour, a point that may also be relevant for other psychological disorders (e.g., substance abuse and impulse control disorders). Finally, our results show that the behavioural functions of these circuits may, at least in principle, be enhanced by external interventions to enhance norm compliance.

The precise neuroanatomy of all structures involved in these circuits may have to be determined by future studies with combined brain stimulation and fMRI²⁹. In particular, it is unclear which interconnected brain areas may be involved in these

circuits, and what their hemispheric lateralization and spatial extent is. Convergent evidence with studies employing brain imaging³⁰ and lesion methodology³¹ may help to narrow down these candidate regions. Finally, translational studies are needed to establish how clinically useful our findings could be, what the duration of the induced tDCS effects is, and what interventions may best enhance the functions of these circuits. Nevertheless, our findings highlight two prefrontal brain processes that are promising candidate targets to better diagnose, ameliorate, or even prevent personality disorders marked by norm-violating behaviour.

References

1. Elster J: The cement of society: a study of social order, ed 1. Cambridge, Cambridge University Press, 1989.
2. Sober E, Wilson DS: *Unto others: The Evolution and Psychology of Unselfish Behavior*, ed 1. Cambridge, Mass., Harvard University Press, 1999.
3. Fischbacher U, Gächter S, Fehr E: Are people conditionally cooperative? Evidence from a public goods experiment. *Econ Lett*, 2001; 71: 397–404.
4. Boyd R, Gintis H, Bowles S, Richerson PJ: The evolution of altruistic punishment. *PNAS*, 2003; 100: 3531–3535.
5. Fischbacher U, Forsythe R, Horowitz HL, Savin NE, Sefton M: Fairness in simple bargaining experiments. *Games Econ Behav* 1994; 6: 347–369.
6. Hare RD, Hart SD, Harpur, TJ: Psychopathy and the DSM-IV criteria for antisocial personality disorder. *J Abnorm Psychol* 1991; 100, 391-398.
7. Bach B, Sellbom M, Kongerslev M, Simonsen E, Krueger RF, Mulder R. Deriving ICD-11 personality disorder domains from dsm-5 traits: initial attempt to harmonize two diagnostic systems. *Acta Psychiatr Scand* 2017; 136:108-117.
8. Loeber R, Burke JD, Lahey BB: What are adolescent antecedents to antisocial personality disorder? *Criminal Behaviour and Mental Health* 2002; 12, 24-36.
9. Compton WM, Conway KP, Stinson FS, Colliver JD, Grant, BF: Prevalence, Correlates, and Comorbidity of DSM-IV Antisocial Personality Syndromes and Alcohol and Specific Drug Use Disorders in the United States: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry* 2005; 66: 677-685.
10. Coid J, Yang M, Tyrer P, Roberts A, Ullrich S: Prevalence and correlates of personality disorder in Great Britain. *Brit J Psychiatry* 2006; 188: 423-431.
11. Carey G, Goldman D: The genetics of anti-social behaviour; in Stoff DM, Breiling J, Maser JD (eds): *Handbook of antisocial behavior*. Wiley, Chichester, 1997, pp 243–250.
12. Goodman M, New A, Siever L: Trauma, genesis, and the neurobiology of personality disorders. *Ann N Y Acad Sci* 2004; 1032: 104–116.
13. Slutske WS, Heath AC, Dinwiddie SH, et al: Modeling genetic and environmental influences in the etiology of conduct disorder. A study of 2682 adult twin pairs. *J Abnorm Psychol* 1997; 106: 266–279.
14. Stouthamer-Loeber M, Loeber R, Wei E et al: Risk and promotive effects in the explanation of persistent serious delinquency in boys. *J Consult Clin Psychol* 2002; 70: 111–123.

15. Andrews D, Zinger I, Hoge RD et al: Does correctional treatment work? A clinically relevant and psychologically informed meta-analysis. *Criminology* 1990; 28: 369–404.
16. Levitt SD, List JA: What Do Laboratory Experiments Measuring Social Preferences Reveal About the Real World? *J Econ Perspectives* 2007; 21: 153-174.
17. Ruff CC, Ugazio G, Fehr E: Changing social norm compliance with noninvasive brain stimulation. *Science* 2013; 342:482-484.
18. Spitzer M, Fischbacher U, Herrnberger B, Grön G, Fehr E: The neural signature of social norm compliance. *Neuron* 2007; 56: 185–196.
19. Nitsche MA, Paulus W: Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *J Physiol* 2000; 527: 633–639.
20. Greene JD, Paxton JM: Patterns of neural activity associated with honest and dishonest moral decisions. *PNAS* 2009; 106: 12506-12511.
21. Maréchal MA, Cohn A, Ugazio G, Ruff CC: Increasing honesty in humans with noninvasive brain stimulation. *PNAS* 2017; 114: 4360-4364.
22. Koenigs M: The role of prefrontal cortex in psychopathy. *Rev Neurosci* 2012; 23: 253–262.
23. Szczepanski SM, Knight RT: Insights into human behavior from lesions to the prefrontal cortex. *Neuron* 2014; 83: 1002-1018.
24. Diamond A: Normal development of prefrontal cortex from birth to young adulthood: cognitive functions, anatomy, and biochemistry; in Stuss D, Knight R (eds): *Principles of Frontal Lobe Function*. New York: Oxford University Press, 2002, pp. 466–503.
25. Hosking JG, Kastman EK, Dorfman HM, Samanez-Larkin GR, Baskin-Sommers A, Kiehl KA, Newman JP, Buckholz JW. Disrupted Prefrontal Regulation of Striatal Subjective Value Signals in Psychopathy. *Neuron* 2017; 95: 221-231
26. Contreras-Rodríguez O, Pujol J, Batalla I, Harrison BJ, Soriano-Mas C, Deus J, López-Solà M, Macià D, Pera V, Hernández-Ribas R, Pifarré J, Menchón JM, Cardoner N. Functional Connectivity Bias in the Prefrontal Cortex of Psychopaths. *Biol Psychiatry* 2015; 78:647-655.
27. Knoch D, Pascual-Leone A, Meyer K, Treyer V, Fehr E: Diminishing reciprocal fairness by disrupting the right prefrontal cortex. *Science* 2006; 314:829-832.
28. Strang S, Gross J, Schuhmann T, Riedl A, Weber B, Sack AT: Be nice if you have to -the neurobiological roots of strategic fairness. *Soc Cogn Affect Neurosci* 2015; 10: 790-796.
29. Driver J, Blankenburg F, Bestmann S, Vanduffel W, Ruff CC: Concurrent brain-stimulation and neuroimaging for studies of cognition. *Trends Cogn Sci* 2009; 13: 319-327.

30. Steinbeis N, Bernhardt BC, Singer T. Impulse control and underlying functions of the left DLPFC mediate age-related and age-independent individual differences in strategic social behavior. *Neuron* 2012; 73: 1040-1051.
31. Zhu L, Jenkins AC, Set E, Scabini D, Knight RT, Chiu PH, King-Casas B, Hsu M: Damage to dorsolateral prefrontal cortex affects tradeoffs between honesty and self-interest. *Nat Neurosci* 2014; 17: 1319-1321.