



---

Year: 2018

---

## C-reactive protein as a predictor of posttraumatic stress induced by acute myocardial infarction

Bielas, Hannes ; Meister-Langraf, Rebecca E ; Schmid, Jean-Paul ; Barth, Jürgen ; Znoj, Hansjörg ; Schnyder, Ulrich ; Princip, Mary ; von Känel, Roland

**Abstract:** **BACKGROUND:** Acute coronary syndrome (ACS) may cause clinically relevant posttraumatic stress disorder symptoms (PTSS). An inflammatory state might be one mechanism linking PTSS with poor prognosis after ACS. We tested the hypothesis that a change in C-reactive protein (CRP) between hospital admission and 3-month follow-up is an independent predictor of ACS-triggered PTSS. **METHODS:** We assessed 183 patients (median age 59 years; 84% men) with verified myocardial infarction (MI) within 48 h of an acute coronary intervention and three months post-MI for self-rated PTSS. 14 (7.7%) patients fulfilled definition criteria for PTSS caseness. CRP values were categorized according to the predicted risk of cardiovascular disease (CVD) at hospital admission (acute inflammatory response): 0 to <5 mg/L, 5 to <10 mg/L, 10 to <20 mg/L, and ≥ 20 mg/L; and at 3-month follow-up (low-grade inflammation): 0 to <1 mg/L, 1 to <3 mg/L, and ≥ 3 mg/L. Additionally, in a subsample of 84 patients with CRP levels below 20 mg/L at admission, CRP values were log-transformed. **RESULTS:** After adjustment for covariates, less of a reduction or an increase of log CRP values between admission and 3-month follow-up predicted PTSS caseness (OR 6.25, 95% CI 1.25, 31.38), and continuous PTSS (unstandardized B = 0.21, 95% CI 0.07, 4.19; p = 0.043). Less reduction in CRP risk categories predicted both PTSS caseness (OR 4.14, 95% CI 1.89, 9.06) and continuous PTSS (B = 1.80, 95% CI 1.09, 2.51; p < 0.001). **CONCLUSIONS:** Persistently heightened inflammation seems to be predictive for the development of PTSS three months after ACS, so interventions to lower inflammation might be warranted.

DOI: <https://doi.org/10.1016/j.genhosppsy.2018.03.008>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-152709>

Journal Article

Supplemental Material

Originally published at:

Bielas, Hannes; Meister-Langraf, Rebecca E; Schmid, Jean-Paul; Barth, Jürgen; Znoj, Hansjörg; Schnyder, Ulrich; Princip, Mary; von Känel, Roland (2018). C-reactive protein as a predictor of posttraumatic stress induced by acute myocardial infarction. *General Hospital Psychiatry*, 53:125-130.

DOI: <https://doi.org/10.1016/j.genhosppsy.2018.03.008>

**Table 2: Correlations between ASDS, CRP categories and covariates**

Entered variables	ASDS score (coefficients)				
	Sum	Re-experiencing	Avoidance	Arousal	Dissociation
CRP categories	<b>0.20**</b>	-0.07	<b>0.19**</b>	0.13	<b>0.25***</b>
Female gender	-0.06	0.09	0.00	-0.07	<b>-0.18*</b>
Age	-0.05	0.03	-0.06	-0.04	-0.04
Education	0.05	0.02	-0.02	0.12	0.03
Social support	-0.07	-0.07	<b>-0.22**</b>	<b>-0.15*</b>	0.06
Smoking	0.09	0.05	0.01	0.07	0.11
Alcohol consumption	-0.02	-0.03	-0.07	-0.02	0.01
Body mass index	0.01	0.07	-0.02	-0.08	0.10
Physical activity	-0.06	-0.08	-0.01	0.09	<b>-0.15*</b>
STEMI	-0.02	0.07	0.07	0.07	-0.12
WBC count	<b>0.17*</b>	0.10	<b>0.16*</b>	0.08	<b>0.15*</b>
LVEF	0.04	0.04	-0.02	0.07	0.00
Troponin T	-0.12	<b>-0.17*</b>	-0.07	-1.11	0.04
GRACE score	0.05	0.06	0.06	-0.21	0.04
Intervention	-0.08	-0.10	0.01	-0.09	0.12
Acute pain	-0.07	0.03	-0.08	-0.13	0.02

Notes: Significant Spearman-Rho's coefficients are displayed in **bold**: \*, $P < 0.05$ ; \*\*, $P < 0.01$ ; \*\*\*, $P \leq 0.001$ .

Abbreviations: ASDS, acute stress disorder scale; CRP, C-reactive protein; GRACE, Global Registry of Acute Coronary Events; LVEF, Left ventricular ejection fraction; STEMI, ST-elevation myocardial infarction; WBC, white blood cell