



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

**Projection of long-term visual acuity outcomes based on initial treatment response
in neovascular age-related macular degeneration**

Nguyen, Vuong ; Daien, Vincent ; Guymer, Robyn ; Young, Stephanie ; Hunyor, Alex ; Fraser-Bell, Samantha ;
Hunt, Adrian ; Gillies, Mark C ; Barthelmes, Daniel

DOI: <https://doi.org/10.1016/j.ophtha.2018.08.023>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-153377>

Journal Article

Accepted Version

Originally published at:

Nguyen, Vuong; Daien, Vincent; Guymer, Robyn; Young, Stephanie; Hunyor, Alex; Fraser-Bell, Samantha; Hunt, Adrian; Gillies, Mark C; Barthelmes, Daniel (2019). Projection of long-term visual acuity outcomes based on initial treatment response in neovascular age-related macular degeneration. *Ophthalmology*, 126(1):64-74.

DOI: <https://doi.org/10.1016/j.ophtha.2018.08.023>

Accepted Manuscript

Projection of long-term visual acuity outcomes based on initial treatment response in neovascular age-related macular degeneration

Vuong Nguyen, PhD, Vincent Daien, MD PhD, Robyn Guymer, MBBS PhD, Stephanie Young, MBBS, Alex Hunyor, MBBS, Samantha Fraser-Bell, MBBS PhD, Adrian Hunt, MBBS, Mark C. Gillies, MBBS PhD, Daniel Barthelmes, MD PhD

PII: S0161-6420(18)30673-0

DOI: [10.1016/j.ophtha.2018.08.023](https://doi.org/10.1016/j.ophtha.2018.08.023)

Reference: OPHTHA 10444

To appear in: *Ophthalmology*

Received Date: 15 March 2018

Revised Date: 9 August 2018

Accepted Date: 17 August 2018

Please cite this article as: Nguyen V, Daien V, Guymer R, Young S, Hunyor A, Fraser-Bell S, Hunt A, Gillies MC, Barthelmes D, the Fight Retinal Blindness! Study Group, Projection of long-term visual acuity outcomes based on initial treatment response in neovascular age-related macular degeneration, *Ophthalmology* (2018), doi: [10.1016/j.ophtha.2018.08.023](https://doi.org/10.1016/j.ophtha.2018.08.023).

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **Projection of long-term visual acuity outcomes based on initial treatment response in**
2 **neovascular age-related macular degeneration**

3

4 Vuong Nguyen PhD¹, Vincent Daien MD PhD^{1,2,3}, Robyn Guymer MBBS PhD⁴, Stephanie
5 Young MBBS⁵, Alex Hunyor MBBS^{1,6,7}, Samantha Fraser-Bell MBBS PhD^{1,6,7}, Adrian Hunt
6 MBBS⁸, Mark C Gillies MBBS PhD^{1,7} and Daniel Barthelmes MD PhD^{1,9}; the Fight Retinal
7 Blindness! Study Group

8 ¹The Save Sight Institute, Sydney Medical School, The University of Sydney, Sydney, NSW

9 ²Department of Ophthalmology, Gui De Chauliac Hospital, Montpellier, F-34000, France

10 ³Inserm, U1061, Montpellier, F-34093, France

11 ⁴Centre for Eye Research Australia, University of Melbourne, Royal Victorian Eye and Ear
12 Hospital, Australia

13 ⁵Gladesville Retina, NSW, Australia

14 ⁶Retina Associates, Chatswood, NSW, Australia

15 ⁷Sydney Eye Hospital, NSW, Australia

16 ⁸Department of Ophthalmology, Westmead Hospital, NSW, Australia

17 ⁹Department of Ophthalmology, University Hospital Zurich, University of Zurich, Zurich,
18 Switzerland

19 **Financial Interests:** Supported by a grant from the Royal Australian NZ College of
20 Ophthalmologists Eye Foundation (2007-2009), a grant from the National Health and
21 Medical Research Council, Australia (NHMRC 2010-2012) and a grant from the Macular
22 Disease Foundation, Australia. Mark Gillies is a Sydney Medical Foundation Fellow and is
23 supported by an NHMRC practitioner fellowship. Robyn Guymer is supported by an
24 NHMRC research fellowship. Daniel Barthelmes was supported by the Walter and Gertrud
25 Siegenthaler Foundation Zurich, Switzerland and the Swiss National Foundation. Vincent
26 Daien was supported by the research grant of the French Society of Ophthalmology. Funding
27 was also provided by Novartis and Bayer. These supporting organisations had no role in the
28 design or conduct of the research.

29 **Corresponding Author:** Vuong Nguyen, 8 Macquarie Street, The Save Sight Institute,
30 Sydney Medical School, The University of Sydney, Sydney, 2000, NSW, Australia. Email:
31 phuc.nguyen@sydney.edu.au

32 **Abstract**

33 **Purpose:** To explore various methods for assessing the early response to vascular endothelial
34 growth factor (VEGF) inhibitors for neovascular age-related macular degeneration and
35 investigate their association with 3 year visual acuity (VA) outcomes.

36 **Design:** Observational study from a prospectively collected registry.

37 **Participants:** Treatment-naïve eyes in the Fight Retinal Blindness! outcomes registry that
38 commenced anti-VEGF therapy between 1st January 2007 and 1st March 2014 that received
39 3 anti-VEGF injections within the first 3 months.

40 **Methods:** The early response was defined as occurring up until the 4th injection. Various
41 early response metrics, which included both continuous and categorical variables, were
42 explored: 1) achieving good VA (≥ 70 letters [20/40]), 2) absolute change in VA from
43 baseline, 3) time to first grading of the choroidal neovascular lesion as inactive, 4) maximum
44 rate of VA change between successive injections.

45 **Main Outcome Measures:** Proportion of eyes achieving ≥ 70 letters at 3 years.

46 **Results:** This study included 2051 treatment-naïve eyes from 1828 patients. Achieving good
47 vision at 3 years was significantly associated with 1) having good vision by the 4th injection
48 (odds ratio [95% CI]: 9.8 [6.5, 14.7] for $VA \geq 70$ vs. $VA < 70$ letters), 2) small (1-5 letters) or
49 large (> 5 letters) early VA gains (1.8 [1.2, 2.6], $P = 0.002$ and 1.8 [1.3, 2.5], $P < 0.001$ vs.
50 eyes with early VA loss), 3) fewer injections until first grading of lesion inactivity (1.6 [1.2,
51 2.1], $P < 0.001$ for ≤ 3 vs. > 3 injections), 4) gradual change (between -4 and 4 letters) or rapid
52 (> 5 letters) gains between successive injections (1.7 [1.1, 2.6], $P = 0.015$ and 1.6 [1.1, 2.3], P
53 = 0.018 for gradual change and rapid gain vs. rapid loss). Eyes that achieved small or large
54 early gains achieved similar vision at 3 years (65.0 and 64.7 letters respectively), and had
55 better vision than eyes with early VA loss (57.2 letters).

56 **Conclusions:** Attainment of good vision by the 4th injection was strongly associated with 3
57 year visual outcomes, while other early response parameters had a moderate association. The
58 early response during the initial 3 monthly loading doses can be a useful guide for subsequent
59 treatment decisions.

60

61 **Introduction**

62 Large variations in the response to vascular endothelial growth factor (VEGF) inhibitors in
63 patients with neovascular age-related macular degeneration (nAMD), as reported in clinical
64 and observational studies, have been attributed to a number of factors, notably demographic
65 and clinical characteristics at baseline and treatment protocols.¹⁻⁷ Baseline clinical
66 characteristics such as age, lesion size and lesion subtype in particular have been identified in
67 multiple studies as predictive of visual outcomes.^{1, 2, 8, 9} In addition, several studies have
68 assessed the effect of genetic factors on treatment outcomes, but these associations are
69 weaker or non-existent.^{4, 10-12} By contrast, the visual acuity (VA) at presentation is one of the
70 strongest predictors of long-term outcomes, whereby eyes with poor starting VA are more
71 likely to achieve larger gains in vision, but have worse final vision than those that present
72 with good VA.^{4, 5, 8}

73 While VEGF inhibitors have generally been shown to provide good visual outcomes for
74 nAMD, some eyes do not respond well to treatment.⁵ Predictive markers based on an eye's
75 early response to treatment may assist in making subsequent treatment decisions and guiding
76 patient expectations. A post-hoc analysis of the Comparison of AMD Treatments (CATT)
77 cohort identified the 12 week change in VA to provide significantly more predictive power
78 for 2 year outcomes compared with the baseline and 4 week response.¹³ In the present study,
79 we explored various metrics for measuring the early response to treatment with VEGF
80 inhibitors, and assessed their ability to predict 3 year visual outcomes. We also assessed
81 whether these early response markers provided additional predictive power that could not
82 already be inferred from the baseline vision.

83

84 **Methods**

85 This study followed the STROBE checklist items for reporting observational study data.¹⁴

86

87 *Study Design*

88 Observational study using data from a prospectively collected registry.

89

90 ***Setting***

91 Data were obtained from the Fight Retinal Blindness! (FRB!) database, a large international
92 registry that tracks real-world outcomes of treatment of nAMD. The FRB! database is
93 compliant with the International Consortium for Healthcare Outcome Measurement's
94 (ICHOM) minimum standard set of treatment outcomes for macular degeneration.¹⁵ Further
95 details of the FRB! database have been published elsewhere.¹⁶ Ethics approval was obtained
96 from the Human Research Ethics Committees of the Royal Victorian Eye and Ear Hospital,
97 the Royal Australian and New Zealand College of Ophthalmologists, the University of
98 Sydney and the Cantonal Ethics Committee Zurich, Switzerland. This study conformed to the
99 tenets of the Declaration of Helsinki.

100

101 ***Data Sources/Measurements***

102 The FRB! system collects data from each clinical visit, including the number of letters read
103 on a logarithm of the minimum angle of resolution (LogMAR) VA Chart (best of
104 uncorrected, corrected or pinhole), treatment given, choroidal neovascular (CNV) lesion
105 activity, as judged by the treating physician based on funduscopy, optical coherence
106 tomography imaging or fluorescein angiography alone or in combination (an active grading
107 indicated the presence of "intraretinal or subretinal fluid attributable to leak from choroidal
108 neovascularisation lesion or fresh haemorrhage"), and ocular adverse events. Previous
109 treatments received, lesion subtype as determined by the practitioner based on retinal
110 angiography and lesion size (greatest linear dimension, GLD) were recorded during the
111 baseline visit. Treatment decisions, including drug choice and treatment frequency, were at
112 the discretion of the practitioner in consultation with the patient, thereby reflecting real-world
113 practice.

114 We explored several avenues for assessing the early response. Most protocols for treating
115 nAMD generally start with a loading of 3 injections of a VEGF inhibitor at monthly intervals
116 regardless of the treatment regimen.^{17, 18} Thus, the early response was specified to occur at
117 the time the 4th injection was due. The metrics for measuring the early response and the
118 expected relationship with long-term outcomes are described below:

- 119 1. **Achieving good vision**, defined as having ≥ 70 letters (20/40 vision)
- 120 2. **Absolute change in VA from baseline**, defined as the change in VA from baseline,
121 was analysed as a continuous variable and as a categorical variable based on the
122 following groups:
- 123 1. *Early Loss*: ≤ 0 letter improvement i.e. loss of vision or no change in vision
 - 124 2. *Small Early Gain*: 1-5 letter improvement
 - 125 3. *Large Early Gain*: > 5 letter improvement
- 126 3. **Time to CNV Inactivity**, defined by the lesion activity status. Following the
127 definitions from a previous FRB! study,¹⁹ we defined the following groups:
- 128 1. *Short Induction*: Eye required ≤ 3 injections until the first grading of the CNV
129 lesion as inactive
 - 130 2. *Long Induction*: Eye required > 3 injections before the lesion was graded as
131 inactive. This included eyes whose CNV lesion remained active throughout the 3
132 year study period.
- 133 4. **Maximum rate of VA change**, defined as the highest rate of change in VA between
134 two successive injections until the 4th injection was due and converted to a
135 standardised rate of letter change per 4 weeks. This rate of change was analysed as a
136 continuous variable and as a categorical variable based on the following groups:
- 137 1. *Rapid Loss*: Largest VA change between successive injections > 5 letter loss per 4
138 weeks
 - 139 2. *Gradual Change*: Largest VA change between successive injections between -4
140 and 4 letters per 4 weeks
 - 141 3. *Rapid Gain*: Maximum of ≥ 5 letter improvement per 4 weeks.
- 142

143 **Participants**

144 Treatment-naïve eyes with nAMD tracked by the FRB! registry commencing anti-VEGF
145 therapy between 1st January 2007 and 1st March 2014 were considered, thereby allowing all
146 eyes the possibility of completing at least 3 years of follow-up at the time the analysis was
147 conducted. For inclusion, eyes were also required to have received 3 monthly anti-VEGF
148 injections as a loading dose to establish the early response and limit the possibility that poor
149 early response was due to under-treatment. Completers were defined as eyes completing 3

150 years of follow-up while non-completers were eyes that did not complete 3 years of follow-
151 up.

152 *Outcome Measures*

153 The primary outcome was the proportion of eyes achieving VA \geq 70 letters at 3 years.

154 Secondary outcomes included the change in VA at 3 years and non-completion rates.

155

156 *Statistical Analysis*

157 Descriptive data included the mean, standard deviation (SD), median, 25th and 75th

158 percentiles (Q1, Q3), and percentages where appropriate. Baseline demographics were

159 compared using ANOVA, Kruskal-Wallis, t-test, Wilcoxon rank sum and Chi-square tests.

160 Longitudinal generalised additive models were used to plot longitudinal visual outcomes over

161 3 years of treatment and included data from completers and non-completers.^{20, 21}

162 The early response was analysed according to the 4 definitions described above. Logistic

163 regression models were also performed with the VA at 3 years as a categorical variable (<70

164 letters vs. \geq 70 letters) and odds ratios reported. Linear mixed-effects models were used to

165 assess the relationship between the change in VA and final VA at 3 years and early response

166 definitions. Since these early responses were likely to be correlated, separate models were fit

167 for each definition. Injection frequency was analysed using Poisson regression models with

168 an offset for log follow-up duration (days). Cox-proportional hazards models were used to

169 assess non-completion rates and visualised using Kaplan-Meier survival curves.

170 Covariates for linear, Poisson and Cox-proportional hazards models also included

171 adjustments for age, lesion size, lesion type (fixed-effects) and clustering by practice and

172 patient (random-effects).

173 Baseline VA was not included as a covariate to avoid potential multicollinearity with the

174 early responses. Instead, separate models were fitted with baseline VA instead of the early

175 response to determine whether using the early response is better than simply using the

176 baseline VA for predicting outcomes. Models were compared using marginal R² values for

177 mixed-models.²² We also report Akaike's Information Criterion (AIC) for model comparison

178 where smaller values indicate better fit. Sensitivity analyses were conducted in which only

179 one eye per patient was analysed for bilateral patients; either the first presenting eye or the
180 worse presenting eye if both eyes were diagnosed simultaneously.

181 Pairwise comparisons were performed using the Holm-Bonferroni adjustment where
182 appropriate. A p-value of 0.05 was considered statistically significant.

183 All analyses were conducted in R version 3.3.2 using the *lme4* package (V1.1-13) for mixed-
184 effects models and *coxme* package (V2.2-5) for Cox-proportional hazards models.²³⁻²⁵

185

186 **Results**

187 *Study Population*

188 This study included 2051 treatment-naïve eyes from 1828 patients (223 bilateral patients) that
189 initiated treatment between 1st January 2007 and 1st March 2014. There were 762 (37%)
190 eyes that did not complete 3 years of follow-up during the study period. The median (Q1,
191 Q3) days until the 4th injection was 105 (91, 123) days. Baseline demographic characteristics
192 partitioned by the categorical early response definitions set out above are summarised in
193 Table 1.

194 Overall, there were 572 (28%) eyes with good VA (≥ 70 letters, Snellen equivalent 20/40) at
195 baseline; at the 4th injection this number had increased to 882 (43%) eyes. Approximately
196 half of eyes underwent a longer period of monthly injections after the initial 3 loading
197 injections (1067 eyes; 52%), including 222 eyes who either remained active by the end of the
198 3 year follow-up (69 eyes) or at time of non-completion (153 eyes).

199 Eyes in the Large Early Gain group had significantly lower mean [SD] baseline VA (49.6
200 [18.1] letters) compared with the Early Loss (60.1 [19.1] letters, $P < 0.001$) and the Small
201 Early Gain (63.1 [14.4] letters, $P < 0.001$) groups. However, we note that eyes in the Early
202 Loss group had similar baseline VA to the Small Early Gain group ($P = 0.13$). Lesion sizes
203 were significantly smaller in eyes with good VA at 4th injection (median [Q1, Q3]: 1934 μm
204 [1124, 2800] vs. 2500 μm [1500, 3500], $P < 0.001$) and in eyes with shorter time to inactivity
205 (median [Q1, Q3]: 2000 [1298, 3000] μm vs. 2500 [1500, 3500] μm , $P < 0.001$).

206 The overall mean (SD) early change in VA was 6.1 (13.2) letters; the overall mean (SD)
207 maximal change in VA between successive injections was 5.2 (14.6) letters.

208

209 *Achieving Good Vision at 3 Years*

210 The association between categorical early response definitions and achieving good vision
211 (≥ 70 letters, 20/40) at 3 years is summarised in Table 2. Of the 1289 eyes that completed 3
212 years of treatment, 608 (47%) had good VA after 3 years of treatment. Overall, achieving
213 good VA by the 4th injection was the best predictor of good vision at 3 years ($R^2 = 0.30$),
214 outperforming the model using baseline VA ($R^2 = 0.17$).

215 Eyes were significantly more likely to achieve good VA (odds ratio [95%CI]) if they had
216 already achieved good vision by the 4th injection (9.8 [6.5, 14.7], $P < 0.001$ for $VA \geq 70$ vs.
217 $VA < 70$ letters by 4th injection), achieved small early or large early gains (1.8 [1.2, 2.6], $P =$
218 0.002 and 1.8 [1.3, 2.5], $P < 0.001$ for small and large early gains vs. early loss), had a short
219 induction (1.6 [1.2, 2.1], $P < 0.001$ for short vs. long induction), or experienced gradual
220 change or rapid gain (1.7 [1.1, 2.6], $P = 0.015$ and 1.6 [1.0, 2.3], $P = 0.018$ for gradual
221 change and rapid gain vs. rapid loss). However, with the exception of achieving good vision
222 by the 4th injection, the remaining early response definitions failed to outperform the baseline
223 model. Sensitivity analyses including only one eye per patient yielded the same result
224 (supplementary material S1).

225 Approximately three quarters (73.0%) of eyes with good VA at the 4th injection maintained
226 good vision after 3 years of treatment. Encouragingly, an additional 149/1289 (22.6%) eyes
227 that had < 70 letters at the 4th injection achieved > 70 letters at year 3.

228

229 *Visual Acuity Outcomes at 3 Years*

230 The association between early response definitions and change in VA at 3 years is
231 summarised in Table 3. Longitudinal VA outcomes through 3 years for categorical early
232 response variables are shown in Figure 1. Overall, the model using the absolute change in VA
233 at the 4th injection (continuous variable) provided the best fit ($R^2 = 0.37$) for predicting the

234 long-term change in VA, outperforming the model using baseline vision instead of the early
235 response ($R^2 = 0.20$).

236 Eyes in the Early Loss group at the 4th injection had worse vision (mean VA change [95%
237 CI]) at 3 years (-5.9 [-7.5, -4.3] letters) than the Small Early Gain (0.7 [-0.9, 2.3] letters, $P <$
238 0.001) and Large Early Gain groups (12.8 [11.4, 14.1], $P < 0.001$). Applying these same
239 categories for VA change at 3 years (Figure 2), 68% of eyes that experienced Early Loss had
240 VA loss at the end of the third year of treatment; these eyes had a relatively high (mean [SD])
241 baseline VA (64.4 [16.2]). The remaining eyes with Early Loss went on to achieve a small
242 (14%) or large (18%) gain in vision despite this early loss, possibly indicating a delayed
243 response. Similarly, 71% of eyes in the Large Early Gain group maintained their large VA
244 gain at 3 years. Only 20% of eyes in the Small Early Gain group had a 1-5 letter gain at 3
245 years, with the remaining 80% split evenly between VA loss and large gains.

246 Visual acuity at 3 years (mean VA [SD]) was significantly worse for eyes in the Early Loss
247 group (57.4 [20.7]) than the Small Early Gain (65.0 [17.2], $P < 0.001$) and Large Early Gain
248 groups (64.7 [17.6], $P < 0.001$). Eyes in the Large Early Gain group had a significantly
249 greater improvement in vision at 3 years compared with the Small Early Gain group ($P <$
250 0.001) although the VA at 3 years was similar for these 2 groups ($P = 0.826$).

251 When eyes were grouped by early maximal rate of VA change, similar patterns were
252 observed whereby the Rapid Loss, Gradual Change, and Rapid Gain groups performed
253 similarly to the Early Loss, Small Early Gain and Large Early Gain groups respectively
254 (Table 3).

255 Eyes with shorter induction had significantly better VA (mean [SD]) at the end of 3 years
256 (65.3 [17.9] letters vs. 59.6 [20.3] letters, $P < 0.001$) although there was no significant
257 difference in VA change ($P = 0.145$).

258

259 ***Injection Frequency***

260 Overall, eyes completing 3 years of follow-up received a median (Q1, Q3) of 19 (15, 23)
261 injections. More frequent injections were associated with higher VA change at 3 years (model
262 coefficient [95%CI]: 0.31 [0.18, 0.44] letters at 3 years per injection, $P < 0.001$). We did not

263 find an association between VA change at the 4th injection (continuous: $P = 0.750$ and
264 categorical: $P = 0.754$) or maximum change of VA (continuous: $P = 0.088$ and categorical: P
265 $= 0.345$) with the number of injections.

266

267 *Non-completion*

268 Change in VA at time of dropout, non-completion rates and their association with the early
269 response are summarised in Table 4. Overall, 762 (37%) eyes did not complete 3 years of
270 follow-up during the study period. Doctor-reported reasons for non-completion were
271 available for 311 eyes and included patient going to another doctor (100 eyes [32%]), further
272 treatment futile (79 eyes [25%]), patient deceased (57 eyes [19%]), patient declined further
273 treatment (44 eyes [14%]), treatment successful (26 eyes [8%]) and medically contraindicated
274 (5 eyes [2%]).

275 Visual outcomes were generally worse compared with completers, although early response
276 groups followed similar trends. At last visit, higher VA (mean [SD]) was observed in eyes
277 with $VA \geq 70$ letters at the 4th injection (72.1 [13.4] vs. 43.0 [23.0] letters), small or large
278 early VA gains (59.0 [23.4] and 55.6 [22.3] vs. 46.1 [26.2] letters for small and large early
279 gains vs. early loss), short induction (56.0 [24.0] vs. 50.0 [24.7] letters for short vs. long
280 induction) and gradual VA change or rapid VA gains (55.2 [25.3] and 54.3 [23.1] vs. 44.2
281 [25.4] letters for gradual change and rapid gain vs. rapid loss). As with completers, 75% of
282 eyes achieving good vision at the 4th injection retained good vision at time of last
283 observation.

284 Survival curves for non-completion over time by early response group are presented in Figure
285 3. Risk of non-completion (hazards ratio, HR [95% CI]) was significantly reduced when VA
286 was ≥ 70 letters at the 4th injection (0.6 [0.5, 0.7] for $VA \geq 70$ vs. $VA < 70$ at the 4th injection, P
287 < 0.001), VA gains at the 4th injection were greater (0.8 [0.6, 0.9], $P = 0.018$, and 0.9 [0.7,
288 1.0], $P = 0.100$, for Small and Large Early Gain vs. Early Loss; global test, $P = 0.016$).

289

290 **Discussion**

291 This study explored several metrics for describing the early response to anti-VEGF treatment
292 for nAMD and their ability to predict 3-year outcomes. We studied whether these early
293 response definitions might predict long-term visual acuity outcomes better than the baseline
294 visual acuity.

295 Eyes with $VA \geq 70$ letters (Snellen equivalent of 20/40) at the time of the 4th injection were
296 almost 10 times more likely have good vision at 3 years than eyes with $VA < 70$ letters at the
297 4th injection. Furthermore, although baseline vision was also a strong predictor of good
298 visual acuity at 3 years, this relationship was not as strong as the visual acuity at the 4th
299 injection.

300 Eyes that experienced early VA loss or small gain in the present study had somewhat similar
301 baseline VA (60.1 and 63.1 letters respectively) but different outcomes at the 4th injection
302 and at 3 years. Similar observations have been reported previously in DME for early
303 moderate (5-9 letter gain) and suboptimal (< 5 letter gain) VA gain groups.²⁶ Eyes that lost
304 vision by the 4th injection had a mean loss of 1 line of vision at 3 years. For eyes that did not
305 complete 3 years of treatment, there was a loss of almost 2 lines at time of dropout. In
306 contrast, eyes that experienced a small early VA gain finished with the same visual acuity as
307 eyes that achieved large early visual acuity gains (65.0 and 64.7 letters respectively) and were
308 similarly likely to achieve good vision at 3 years. In addition, 18% of eyes that experienced
309 early VA loss went on to gain more than 1 line of vision at 3 years, indicating a delayed
310 response to anti-VEGF treatment. A post-hoc analysis of the CATT cohort reported 27% of
311 eyes showing a loss of ≥ 1 line at 12 weeks went on to gain ≥ 1 line at 2 years.¹³ Thus, it may
312 be prudent to persist with anti-VEGF treatment even if the early response is poor in the
313 absence of effective alternative treatments.

314 Measuring the maximum rate of VA change between successive injections was a novel way
315 to assess the early response. We observed that approximately three quarters of eyes either
316 gained (54%) or lost (18%) more than 5 letters at least once between 2 successive injections,
317 with only 28% experiencing more gradual changes between successive injections. However,
318 the maximal rate of change and raw early VA change definitions provided somewhat similar
319 information, and the models using early VA change provided a better fit than the maximal
320 rate of change.

321 Lesion activity, or shorter time to lesion inactivation, may be another useful marker of early
322 treatment response. Eyes with a shorter time to lesion inactivity (3 or fewer injections) had
323 better vision than eyes requiring more than 3 injections, both at baseline (57.8 vs 53.9 letters
324 respectively) and at 3 years (65.3 vs. 59.6 letters respectively). They were also more likely to
325 have good vision (70 letters, 20/40) at 3 years although the change in VA at 3 years was not
326 significantly different ($P = 0.091$). A previous analysis of 12-month outcomes found eyes
327 with highly active lesions performed similarly to those with less active lesions,²⁷ however a
328 longer term analysis is warranted to clarify the relationship between highly active lesions and
329 visual outcomes.

330 We found that improvement in visual acuity up to the 4th injection of VEGF inhibitors was
331 the most robust clinical predictor of visual acuity 3 years after starting treatment. Previous
332 studies have also found greater predictive power between the 12 week change in VA with the
333 1 and 2 year outcomes compared with only using the baseline VA.¹³ This is probably because
334 the disease is still largely VEGF-driven in these cases with a good early response. Cases
335 which do not respond so well may be also be driven by other, less reversible, pathological
336 processes such as inflammation, fibrosis and macular atrophy.¹³

337 The present study has some limitations. Treatment schedules after the initial loading phase,
338 which might have influenced long-term outcomes,^{17,28} were at the discretion of the physician
339 and patient although most of the FRB! database practitioners use a treat and extend
340 regimen.^{29,30} It is however possible that patients with inferior initial responses may have
341 subsequently been less compliant or extended out by the physician and suffered inferior
342 outcomes as a result. Eyes with good VA at the 4th injection, tended to have better 3-year
343 outcomes, and also on average, received more injections. Overall, more injections are
344 associated with better visual acuity outcomes.²⁹ Still, eyes that were continued on monthly
345 injections after the 3 initial monthly injections due to persistent activity – and thus had a high
346 total number of injections - had worse outcomes at 3 years, possibly because their lesions
347 were more active. Anti-VEGF drug type was not considered in the present analysis because
348 previous studies have found no substantial difference between ranibizumab and aflibercept.³¹
349 Nor did we report switching rates as aflibercept was not yet available as a treatment option
350 for most of our follow-up period. We note that while switching treatments may be a possible
351 strategy when the early response is poor, there is currently little if any evidence that
352 switching anti-VEGF agents provides any obvious benefit.³²

353 High non-completion rates are common in observational studies, and this study was no
354 exception; 37% of eyes did not complete 3 years of follow-up during the study period.
355 Reasons for non-completion were reported for more than a third of the non-completers, with
356 most due to reasons that were not linked with efficacy. Around 40% of the eyes with a
357 recorded reason for non-completion were discontinued because further treatment was futile or
358 the patient declined further treatment. Patients were more likely to drop out if they
359 experienced early VA loss or their VA was less than 70 letters at the 4th injection. The
360 change in visual acuity at time of dropout between early response groups followed broadly
361 similar patterns to those observed in the completers, although the final vision at time of
362 dropout for the early response groups was, on average, 1-2 lines lower than their respective
363 completers.

364 In conclusion, the early response, particularly attainment of good vision and change in visual
365 acuity by the 4th injection, was more strongly associated with 3 year visual outcomes than
366 visual acuity at the time of starting treatment. As treatment protocols for nAMD generally
367 begin with 3 monthly injections, the response during this standardised period of treatment
368 may be useful to guide further treatment.³³

369

370 **Acknowledgements:** Fight Retinal Blindness! investigators: Auckland District Health Board,
371 New Zealand (Dr D Squirrell); Cairns Eye Surgery, Queensland (Dr A Field); Canberra
372 Hospital, Australian Capital Territory (Dr C Dayajeewa, Dr R Essex); Care Foresight, New
373 South Wales (Dr A Dunlop); Central Coast Eye Specialist, New South Wales (Dr S Young);
374 Centre for Eye Research Australia, Victoria (Professor R Guymer); Coastwide Eye Surgery,
375 New South Wales (Dr R Ferrier); Crest Eye Associates, New Zealand (Dr J Ah-Chan);
376 Doncaster Eye Center, Victoria (Dr L Chow); Dr Nadia Wittles Practice, South Australia (Dr
377 N Wittles); Dr. Phillip Windle, Queensland (Dr P Windle); Eye Associates, New South
378 Wales (Dr M Gillies, Dr A Hunt); Eye Surgeons Miranda, New South Wales (Dr A Hunt);
379 Eyemedics (Wayville), South Australia (Dr K Billings, Dr J Chen, Dr S Lake, Dr J Landers,
380 Dr M Perks, Dr R Phillips, Dr M Saha); Gladesville Eye Specialists, New South Wales (Dr S
381 Young); Hornsby Eye Specialists, New South Wales (Dr S Lal); Les Manning Practice,
382 Queensland (Dr L Manning); Lions Eye Institute, Western Australia (Professor I McAllister);
383 Marsden Eye Specialists, New South Wales (Dr J Arnold); Mona Vale Eye Centre, New
384 South Wales (Dr C Lim); Nepean Valley Eye Surgeons, New South Wales (Dr G Banerjee);

385 Retina Associates, New South Wales (Dr S Fraser-Bell, Dr A Fung, Professor A Hunyor, Dr
386 C Younan); Retina Consultants, New South Wales (Dr S Young); Retina Specialists, New
387 Zealand (Dr R Barnes, Dr A Vincent); Southern Eye Centre, Victoria (Dr D Louis); Specialist
388 Eye Group, Victoria (Dr L Chow); Strathfield Retina Clinic, New South Wales (Dr C Lim);
389 Sydney Eye Hospital, New South Wales (Dr J Wong); University Hospital Zurich,
390 Switzerland (Dr D Barthelmes); Victoria Parade Eye Consultants, Victoria (Dr M Daniell,
391 Professor R Guymer, Dr A Harper, Dr L Lim, Dr J O'Day)

392 **Figure Captions**

393 **Figure 1.** Predicted visual acuity (VA) over time from longitudinal generalised additive
394 models partitioned by A) whether VA was ≥ 70 letters at the 4th injection, B) absolute change
395 in VA at the 4th injection, C) length of the induction period and D) maximum rate of VA
396 change between successive injections. These models included data from completers and non-
397 completers.

398 **Figure 2.** Percentage of eyes partitioned by (A) VA change at the 4th injection and at 3 years,
399 and (B) VA change at the 4th injection, VA < 70 or ≥ 70 letters at the 4th injection (< 70 and
400 ≥ 70 respectively, labelled above bars), and VA < 70 or ≥ 70 letters at 3 years. Categories for
401 VA change included early loss (< 0 letter improvement), small gain (1-5 letter improvement)
402 and large gain (> 5 letter improvement). The number of eyes in each early VA change group
403 is shown above the bars.

404 **Figure 3.** Kaplan-Meier survival curves of time to non-completion partitioned by A) whether
405 VA was ≥ 70 letters at the 4th injection, B) absolute change in VA at the 4th injection, C)
406 length of the induction period and D) maximum rate of VA change between successive
407 injections

408

409 **References**

- 410 1. Rosenfeld PJ, Brown DM, Heier JS et al. Ranibizumab for neovascular age-related
411 macular degeneration. *N Engl J Med* 2006;355:1419-1431.
- 412 2. Boyer DS, Antoszyk AN, Awh CC et al. Subgroup analysis of the MARINA study of
413 ranibizumab in neovascular age-related macular degeneration. *Ophthalmology*
414 2007;114:246-252.
- 415 3. Rofagha S, Bhisitkul RB, Boyer DS et al. Seven-year outcomes in ranibizumab-
416 treated patients in ANCHOR, MARINA, and HORIZON. *Ophthalmology*
417 2013;120:2292-2299.
- 418 4. Finger RP, Wickremasinghe SS, Baird PN, Guymer RH. Predictors of anti-VEGF
419 treatment response in neovascular age-related macular degeneration. *Surv Ophthalmol*
420 2014;59:1-18.
- 421 5. Gillies MC, Campain A, Barthelmes D et al. Long-term outcomes of treatment of
422 neovascular age-related macular degeneration: data from an observational study.
423 *Ophthalmology* 2015;122:1837-1845.
- 424 6. Maguire MG, Martin DF, Ying G-s et al. Five-year outcomes with anti-vascular
425 endothelial growth factor treatment of neovascular age-related macular degeneration:
426 the Comparison of Age-Related Macular Degeneration Treatments Trials.
427 *Ophthalmology* 2016;123:1751-1761.
- 428 7. Gillies MC, Daien V, Nguyen V, Barthelmes D. Re: Comparison of Age-Related
429 Macular Degeneration Treatments Trials (CATT) Research Group, et al.: Five-year
430 outcomes with anti-vascular endothelial growth factor treatment of neovascular age-
431 related macular degeneration: The Comparison of Age-Related Macular Degeneration
432 Treatments Trials (*Ophthalmology* 2016;123:1751-1761). *Ophthalmology*
433 2017;124:e31-e32.
- 434 8. Kaiser PK, Brown DM, Zhang K et al. Ranibizumab for predominantly classic
435 neovascular age-related macular degeneration: subgroup analysis of first-year
436 ANCHOR results. *Am J Ophthalmol* 2007;144:850-857. e854.
- 437 9. Ying G-s, Huang J, Maguire MG et al. Baseline predictors for one-year visual
438 outcomes with ranibizumab or bevacizumab for neovascular age-related macular
439 degeneration. *Ophthalmology* 2013;120:122-129.
- 440 10. Gorin MB. Genetic insights into age-related macular degeneration: Controversies
441 addressing risk, causality, and therapeutics. *Mol Aspects Med* 2012;33:467-486.
- 442 11. Hagstrom SA, Ying G-s, Pauer GJT et al. Pharmacogenetics for Genes Associated
443 with Age-related Macular Degeneration in the Comparison of AMD Treatments Trials
444 (CATT). *Ophthalmology* 2013;120:593-599.
- 445 12. Hagstrom SA, Ying G, Pauer GT, et al. Vegfa and vegfr2 gene polymorphisms and
446 response to anti-vascular endothelial growth factor therapy: Comparison of age-
447 related macular degeneration treatments trials (catt). *JAMA Ophthalmology*
448 2014;132:521-527.
- 449 13. Ying G-s, Maguire MG, Daniel E et al. Association of Baseline Characteristics and
450 Early Vision Response with 2-Year Vision Outcomes in the Comparison of AMD
451 Treatments Trials (CATT). *Ophthalmology* 2015;122:2523-2531.e2521.
- 452 14. Von Elm E, Altman DG, Egger M et al. The Strengthening the Reporting of
453 Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting
454 observational studies. *Prev Med* 2007;45:247-251.
- 455 15. Rodrigues IA, Sprinkhuizen SM, Barthelmes D et al. Defining a minimum set of
456 standardized patient-centered outcome measures for macular degeneration. *Am J*
457 *Ophthalmol* 2016;168:1-12.

- 458 16. Gillies MC, Walton R, Liong J et al. Efficient capture of high-quality data on
459 outcomes of treatment for macular diseases: The Fight Retinal Blindness! Project.
460 Retina 2014;34:188-195.
- 461 17. Oubraham H, Cohen SY, Samimi S et al. Inject and extend dosing versus dosing as
462 needed: a comparative retrospective study of ranibizumab in exudative age-related
463 macular degeneration. Retina 2011;31:26-30.
- 464 18. Freund KB, Korobelnik J-F, Devenyi R et al. Treat-and-extend regimens with anti-
465 VEGF agents in retinal diseases: a literature review and consensus recommendations.
466 Retina 2015;35:1489-1506.
- 467 19. Essex RW, Nguyen V, Walton R et al. Treatment patterns and visual outcomes during
468 the maintenance phase of treat-and-extend therapy for age-related macular
469 degeneration. Ophthalmology 2016;123:2393-2400.
- 470 20. Hastie T, Tibshirani R. Generalized additive models. Statist Sci 1986;1:297-310.
- 471 21. Hastie TJ: Generalized additive models. In: *Statistical models in S*. edn.: Routledge;
472 2017: 249-307.
- 473 22. Nakagawa S, Schielzeth H. A general and simple method for obtaining R² from
474 generalized linear mixed-effects models. Methods Ecol Evol 2013;4:133-142.
- 475 23. R: A language and environment for statistical computing. Vienna, Austria: R
476 Foundation for Statistical Computing
- 477 24. Bates DM, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models
478 using lme4. J Stat Softw 2015;67:1-48.
- 479 25. Therneau T. coxme: Mixed effects Cox models. R package 2012.
- 480 26. Mehta H, Fraser-Bell S, Nguyen V et al. Short-term vision gains at 12 weeks correlate
481 with long-term vision gains at 2 years: results from the BEVORDEX randomised
482 clinical trial of bevacizumab versus dexamethasone implants for diabetic macular
483 oedema. Br J Ophthalmol 2017.
- 484 27. Barthelmes D, Walton R, Campain AE et al. Outcomes of persistently active
485 neovascular age-related macular degeneration treated with VEGF inhibitors:
486 observational study data. Br J Ophthalmol 2014;99:359-364.
- 487 28. Chin-Yee D, Eck T, Fowler S et al. A systematic review of as needed versus treat and
488 extend ranibizumab or bevacizumab treatment regimens for neovascular age-related
489 macular degeneration. Br J Ophthalmol 2016;100:914-917.
- 490 29. Arnold JJ, Campain A, Barthelmes D et al. Two-year outcomes of “treat and extend”
491 intravitreal therapy for neovascular age-related macular degeneration. Ophthalmology
492 2015;122:1212-1219.
- 493 30. Johnston RL, Carius H-J, Skelly A et al. A retrospective study of ranibizumab
494 treatment regimens for neovascular age-related macular degeneration (nAMD) in
495 Australia and the United Kingdom. Adv Ther 2017;34:703-712.
- 496 31. Gillies MC, Nguyen V, Daien V et al. Twelve-Month Outcomes of Ranibizumab vs.
497 Aflibercept for Neovascular Age-Related Macular Degeneration: Data from an
498 Observational Study. Ophthalmology 2016;123:2545-2553.
- 499 32. Barthelmes D, Campain A, Nguyen P et al. Effects of switching from ranibizumab to
500 aflibercept in eyes with exudative age-related macular degeneration. Br J Ophthalmol
501 2016;100:1640-1645.
- 502 33. Schmidt-Erfurth U, Bogunovic H, Sadeghipour A et al. Machine learning to analyze
503 the prognostic value of current imaging biomarkers in neovascular age-related
504 macular degeneration. Ophthalmology Retina 2018;2:24-30.
- 505

Table 1. Demographic characteristics of eyes grouped by their early response to treatment.

	Good Vision Achieved by 4th Injection		Good Vision Achieved by 4th Injection			Time to Inactive CNV			Maximum Rate of VA Change		
	Overall	VA <70 Letters	VA ≥70 Letters	Early Loss	Small Early Gain	Large Early Gain	Short	Long*	Rapid Loss	Gradual Change	Rapid Gain
Eyes	2051	1169	882	682	422	947	984	1067	368	568	1115
Patients	1828	1088	831	647	412	897	907	1002	352	550	1056
Females, %	61.5%	62.0%	60.8%	59.5%	62.8%	62.3%	62.2%	60.8%	61.4%	61.6%	61.4%
Baseline Age (SD)	79.6 (8.2)	80.7 (7.9)	78.3 (8.3)	79.7 (8.6)	79.4 (7.8)	79.8 (8)	80.7 (7.8)	78.7 (8.4)	79.4 (8.6)	79.2 (8.3)	80.0 (8.0)
Baseline VA (SD)	55.9 (18.7)	46.6 (17.8)	68.1 (11.2)	60.1 (19.1)	63.1 (14.4)	49.6 (18.1)	57.8 (18.4)	54.1 (18.8)	60.3 (16.4)	61.5 (19.2)	51.6 (18.0)
≥70 Letters, n (%)	572 (27.9%)	82 (7.0%)	490 (55.6%)	277 (40.6%)	189 (44.8%)	106 (11.2%)	318 (32.3%)	254 (23.8%)	136 (37.0%)	255 (44.9%)	181 (16.2%)
≤35 Letters, n (%)	308 (15.0%)	289 (24.7%)	19 (2.2%)	84 (12.3%)	23 (5.5%)	201 (21.2%)	124 (12.6%)	184 (17.2%)	36 (9.8%)	66 (11.6%)	206 (18.5%)
Baseline Lesion Size, Median μm (Q1, Q3)	2250 (1439, 3200)	2500 (1500, 3500)	1934 (1124, 2800)	2314 (1458, 3300)	2200 (1300, 3158)	2200 (1500, 3200)	2000 (1298, 3000)	2500 (1500, 3500)	2200 (1400, 3306)	2250 (1394, 3300)	2250 (1500, 3199)
Lesion Type, %											
<i>Occult</i>	56.6%	55.4%	58.2%	55.7%	64.2%	53.9%	57.5%	55.8%	60.1%	55.3%	56.1%
<i>Minimally Classic</i>	14.2%	14.5%	13.9%	14.5%	11.1%	15.4%	12.7%	15.7%	13.0%	13.4%	15.1%
<i>Predominantly Classic</i>	20.5%	21.6%	18.9%	21.1%	16.45%	21.9%	20.7%	20.2%	20.1%	20.1%	20.8%
<i>Other</i>	7.3%	7.4%	7.1%	7.3%	7.3%	7.3%	7.4%	7.2%	6.2%	9.0%	6.85%
<i>Not Recorded</i>	1.4%	1.0%	1.8%	1.3%	0.9%	1.6%	1.6%	1.1%	0.5%	2.3%	1.2%

* Includes 222 persistently active eyes whose lesion has yet to be graded as inactive by the end of completing 3 years of follow-up (69 eyes) or at their most recent visit if they did not complete 3 years of follow-up (153 eyes)

Table 2. Association between definitions of early response and achieving good vision (≥ 70 letters) at 3 years. Odds ratios and their respective p-values are presented only for categorical variables. Significant p-values are highlighted in bold.

Categorical Early Response Definitions	VA <70 Letters at 3 Years, n (%)	VA ≥ 70 Letters at 3 Years, n (%)	Odds Ratio for Achieving ≥ 70 Letters at 3 Years (95% CI)	P-value	Marginal R²	AIC
Overall	681 (53%)	608 (47%)	-			
Good VA at Baseline						
VA <70 Letters	574 (64%)	321 (36%)	1	<0.001	0.17	1606
VA ≥ 70 Letters	107 (27%)	287 (73%)	4.50 (3.30, 6.14)			
Good VA Achieved by 4th Injection						
VA <70 Letters	511 (77%)	149 (23%)	1	<0.001	0.30	1433
VA ≥ 70 Letters	170 (27%)	459 (73%)	9.78 (6.50, 14.70)			
Absolute Change in VA from Baseline at 4th Injection						
<i>Early Loss</i>	248 (62%)	152 (38%)	1	<0.001*	0.08	1711
<i>Small Early Gain</i>	135 (48%)	148 (52%)	1.75 (1.17, 2.61)			
<i>Large Early Gain</i>	298 (49%)	308 (51%)	1.76 (1.25, 2.45)			
Time to Inactive CNV						
<i>Short Induction</i>	300 (47%)	341 (53%)	1.59 (1.23, 2.07)	<0.001	0.07	1715
<i>Long Induction</i>	381 (59%)	267 (41%)	1			
Maximum Rate of VA Change						
<i>Rapid Loss</i>	141 (62%)	86 (38%)	1	0.011 †	0.07	1720
<i>Gradual Change</i>	165 (49%)	175 (51%)	1.70 (1.09, 2.64)			
<i>Rapid Gain</i>	375 (52%)	347 (48%)	1.55 (1.05, 2.30)			

Pairwise comparisons with Holm-Bonferroni adjustment for multiple comparisons:

* Early Loss vs. Small Early Gain (P = 0.002), Early Loss vs. Large Early Gain (P < 0.001), Small Early Gain vs. Large Early Gain (P = 0.986)

† Rapid Loss vs. Gradual Change ($P = 0.015$), Rapid Loss vs. Rapid Gain ($P = 0.018$), Gradual Change vs. Rapid Loss ($P = 0.523$)

ACCEPTED MANUSCRIPT

<i>Early Loss</i>	57.2 (22.2)	57.9 (55.5, 60.3)	< 0.001 *	-5.9 (-7.5, -4.3)	-5.6 (-7.2, -3.9)	< 0.001 **	0.23	10794
<i>Small Early Gain</i>	65.0 (17.2)	64.4 (61.7, 67.1)		0.7 (-0.9, 2.3)	0.7 (-1.3, 2.6)			
<i>Large Early Gain</i>	64.7 (17.6)	64.7 (62.6, 66.8)		12.8 (11.4, 14.1)	12.8 (11.4, 14.2)			
Time to Inactive CNV								
<i>Short Induction</i>	65.3 (17.9)	64.9 (62.9, 66.9)	< 0.001	5.3 (3.9, 6.6)	5.3 (3.7, 6.8)	0.145	0.02	11097
<i>Long Induction</i>	59.6 (20.3)	60.1 (58.1, 62.1)		3.4 (2.0, 4.9)	3.7 (2.2, 5.3)			
Maximum Rate of VA Change per 4 Weeks								
<i>Rapid Loss</i>	57.4 (20.7)	58.0 (55.1, 60.9)	< 0.001 †	-5.2 (-7.6, -2.8)	-4.7 (-7.0, -2.4)	< 0.001 ††	0.14	10937
<i>Gradual Change</i>	63.4 (19.9)	62.8 (60.2, 65.3)		-0.7 (-2.1, 0.7)	-0.9 (-2.8, 1.0)			
<i>Rapid Gain</i>	63.6 (18.4)	63.7 (61.6, 65.7)		9.7 (8.4, 11.0)	9.7 (8.4, 11.1)			

Pairwise comparisons with Holm-Bonferroni adjustment for multiple comparisons:

- * Early Loss vs. Small Early Gain (P < 0.001), Early Loss vs. Large Early Gain (P < 0.001), Small Early Gain vs. Large Early Gain (P = 0.826)
- ** Early Loss vs. Small Early Gain (P < 0.001), Early Loss vs. Large Early Gain (P < 0.001), Small Early Gain vs. Large Early Gain (P < 0.001)
- † Rapid Loss vs. Gradual Change (P = 0.006), Rapid Loss vs. Rapid Gain (P < 0.001), Gradual Change vs. Rapid Loss (P = 0.452)
- †† Rapid Loss vs. Gradual Change (P = 0.007), Rapid Loss vs. Rapid Gain (P < 0.001), Gradual Change vs. Rapid Loss (P < 0.001)


Table 4. Visual acuity, change in VA and the proportion of eyes with VA \geq 70 letters at time of dropout, non-completion rates and their association with different definitions of early response. Hazards ratios for non-completion and their respective p-values are presented only for categorical variables. Significant p-values are highlighted in bold.


Categorical Early Response Definition	VA at Time of Dropout (SD)	ΔVA at Time of Dropout (95% CI)	VA\geq70 Letters, n (%)	Non-completion Rate, n / N (%)	Hazards Ratio for non-completion (95% CI)	P-value
Overall	52.7 (24.5)	0.6 (-0.9, 2.0)	243 (32%)	762 / 2051 (37%)	-	-
Good VA at Baseline						
VA <70 Letters	46.9 (24.2)	2.0 (0.3, 3.8)	112 (19%)	584 / 1479 (39%)	1	0.003
VA \geq 70 Letters	71.6 (13.8)	-4.3 (6.3, -2.3)	131 (74%)	178 / 572 (31%)	0.77 (0.63, 0.90)	
Good VA by 4th Injection						
VA <70 Letters	43.0 (23.0)	-0.5 (-2.5, 1.4)	52 (10%)	509 / 1169 (44%)	1	<0.001
VA \geq 70 Letters	72.1 (13.4)	2.8 (0.9, 4.7)	191 (75%)	253 / 882 (29%)	0.61 (0.51, 0.71)	
Absolute Change in VA from Baseline at 4th Injection						
Early Loss	46.1 (26.2)	-9.7 (-11.8, -7.6)	78 (28%)	282 / 682 (41%)	1	0.016*
Small Early Gain	59.0 (23.4)	-1.8 (-4.4, 0.9)	61 (44%)	139 / 422 (33%)	0.75 (0.56, 0.93)	
Large Early Gain	55.6 (22.3)	10.0 (8.0, 12.1)	104 (30%)	341 / 947 (36%)	0.85 (0.69, 1.01)	
Time to Inactive CNV						
Short Induction	56.0 (24.0)	2.3 (0.3, 4.3)	129 (38%)	343 / 984 (35%)	0.90 (0.74, 1.05)	0.196
Long Induction	50.0 (24.7)	-0.9 (-2.9, 1.2)	114 (27%)	419 / 1067 (39%)	1	
Maximum Rate of VA Change						
Rapid Loss	44.2 (25.4)	-12.4 (-15.8, -9.0)	32 (23%)	141 / 368 (38%)	1	0.157
Gradual Change	55.2 (25.3)	-2.3 (-4.3, -0.3)	93 (41%)	228 / 568 (40%)	1.05 (0.78, 1.32)	
Rapid Gain	54.3 (23.1)	6.9 (4.9, 8.9)	118 (30%)	393 / 1115 (35%)	0.90 (0.69, 1.11)	


Pairwise comparisons with Holm-Bonferroni adjustment for multiple comparisons:

* Early Loss vs. Small Early Gain ($P = 0.018$), Early Loss vs. Large Early Gain ($P = 0.100$), Small Early Gain vs. Large Early Gain ($P = 0.206$)

ACCEPTED MANUSCRIPT







The early response to treatment for neovascular age-related macular degeneration is highly associated with treatment outcomes at 3 years and may provide a useful marker for guiding long-term treatment decisions.

ACCEPTED MANUSCRIPT