



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2005

**Epstein Barr–virus (EBV) associated T–cell clonopathy mimicking
lymphomatous meningitis**

Hermann, D M ; Fehr, J ; Bassetti, C L

DOI: <https://doi.org/10.1007/s00415-005-0744-9>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-155803>

Journal Article

Published Version

Originally published at:

Hermann, D M; Fehr, J; Bassetti, C L (2005). Epstein Barr–virus (EBV) associated T–cell clonopathy mimicking lymphomatous meningitis. *Journal of Neurology*, 252(6):736-737.

DOI: <https://doi.org/10.1007/s00415-005-0744-9>

D. M. Hermann
J. Fehr
C. L. Bassetti

Epstein Barr-virus (EBV) associated T-cell clonopathy mimicking lymphomatous meningitis

Received: 23 July 2004
Received in revised form:
17 November 2004
Accepted: 23 November 2004
Published online: 14 March 2005

Sirs: Epstein Barr-virus (EBV) exhibits strong B-cell tropism. In chronic EBV-infection, T-cells may also be affected, which often expand monoclonally and then may progress into T-cell lymphoma [2, 3]. We describe a young female with EBV-associated T-cell clonopathy presenting twice with monoclonal infiltration of cranial and spinal nerve roots, which have not converged into lymphoma by now after seven years.

In 1997, the 20-year-old patient developed fever ($\leq 40^{\circ}\text{C}$), fatigue, hepatosplenomegaly and mild lymphadenopathy with pancytopenia (Hb 8.3 g/dl, platelets 80000/ μl , neutrophils 500/ μl), mildly elevated CRP (26 mg/l), increased gammaglobulins (22 g/l), thymidin-kinase (59.2 U/l), soluble interleukin-2 receptor (sIL-2r) (10696 U/ml) and neopterin (56.9 ng/ml). A macrophage activation syndrome (characterized by elevated sIL-2r and neopterin) was suspected and a bone-marrow biopsy performed, revealing massive hemophagocytosis. The patient was splenectomized. Southern blots revealed monoclonal TCR β -rearrangement with singular restriction bands identical in bone-marrow, blood and spleen, PCR singular TCR γ -amplifications identical in blood

and spleen. Based on pediatric treatment schemes, aiming at macrophage inhibition, low-dose VP16-chemotherapy ($2 \times 150 \text{ mg/m}^2$) was initiated. Fever and inflammation markers declined.

Five weeks later, the patient developed fever, ear-pain and diplopia. Clinical examination revealed a mild meningism and right abducens nerve palsy. Three lumbar punctures showed mild pleocytosis (75/ μl , reactive T-cells) and elevated protein (1.63 g/l). Blood EBV-VCA IgG/IgM and EBV-EBNA IgG titres were increased. Cranial MRI exhibited meningeal contrast-enhancement along cranial nerves suggestive of lymphomatous meningitis. Intrathecal chemotherapy with methotrexate ($9 \times 15 \text{ mg}$) and AraC ($9 \times 40 \text{ mg}$) was started, followed by dexamethasone. CSF cell-counts decreased to 2/ μl , CSF protein to 0.49 g/l. Clinical symptoms and neurological abnormalities disappeared.

In 2002, hyposmia, hypoacusis and mild urinary disturbances were noted by the patient, followed by dizziness and gait disturbances three months later. On clinical examination, bilateral L3-S1 deficits with areflexia and positive Lasague's sign were found. Blood examinations showed normal cell-counts and CRP, reduced gammaglobulins (1.5 g/l), increased sIL-2r (1359 IU/ml) and neopterin (3.5 ng/ml), normal cyclin-D1 and positive EBV-PCR (125 DNA copies/ml). A mild CSF pleocytosis (36 cells/ μl , reactive T-cells) with elevated CSF protein (2.6 g/l) and positive CSF EBV-PCR were found. Both the blood and CSF still exhibited monoclonal TCR γ -rearrangement with identical clones as six years before. MRI revealed contrast-enhancement with dural broadening around cranial and spinal nerve roots (Figure). In the absence of lymphoma cells in three lumbar

puncture specimens, dexamethasone ($4 \times 8 \text{ mg}$) and intravenous immunoglobins ($3 \times 30 \text{ g}$) were started. Clinical symptoms rapidly improved. Today, fourteen months later, the polyradicular deficits have normalized. The patient complains about mild dizziness and (sensory) hearing difficulties.

In our patient, the recurrent cranial and spinal nerve root infiltrations reflect a steroid-responsive, EBV-related lymphoproliferative disorder. The absence of malignant cells in the blood and CSF, the negative tumour markers and clinical course allow the exclusion of the presence of a T-cell lymphoma. Neurological complications sometimes occur in acute and reactivated EBV, but rarely in chronic active EBV-infection [1]. Three patients with chronic active EBV-infection and encephalitis have been reported [1, 4, 5]. Two patients had fatal outcomes, all three patients exhibited brain parenchymal involvement on MRI. Thus, this is the first report of isolated meningeal infiltration in chronic active EBV-infection. In our patient, a gammaglobulin deficiency was detected during the second disease manifestation, whereas the first episode developed shortly after chemotherapy. Thus, immunodeficiency may have played a pathogenic role.

Monoclonal T-cell expansion following chronic EBV-infection is widely regarded as a guaranteed prelude to malignant lymphoma. In most cases, this is indeed clearly the case, although our patient shows that long-term survival is possible in a subset of patients. At present, our patient's long term prognosis remains unknown. In conclusion, meningeal monoclonal T-cell infiltration in the presence of chronic active EBV-infection should not mislead one to the diagnosis of T-cell lymphoma, unless a cytological diagnosis is




Fig. Gadolinium-enhanced T1-weighted MRI showing massive meningeal contrast enhancement (flashes) along cranial (A) and lumbar spinal (B) nerve roots

made. Clinical manifestations may well respond to steroids, allowing the avoidance of chemotherapy.

■ **Acknowledgement** We thank our Division of Neuroradiology (Director: Prof. A. Valavanis) for performing the MR studies.

References

- Häusler M, Ramaekers VT, Doenges M, Schweizer K, Ritter K, Schaade L (2002) Neurological complications of acute and persistent Epstein-Barr virus infection in paediatric patients. *J Med Virol* 68(2):253–263
- Kanegane H, Nomura K, Miyawaki T, Tosato G (2002) Biological aspects of Epstein-Barr virus (EBV)-infected lymphocytes in chronic active EBV infection and associated malignancies. *Crit Rev Oncol Hematol* 44(3):239–249
- Kasahara Y, Yachie A (2002) Cell type specific infection of Epstein-Barr virus (EBV) in EBV-associated hemophagocytic lymphohistiocytosis and chronic active EBV infection. *Crit Rev Oncol Hematol* 44(3):283–294
- Ohga S, Takada H, Honda K, Inamura T, Gondo K, Ohshima K, Yamamoto M, Hara T (1999) Central nervous system T-cell lymphoproliferative disorder in a patient with chronic active Epstein-Barr virus infection. *J Pediatr Hematol Oncol* 21(1):42–46
- Yamashita S, Murakami C, Izumi Y, Sawada H, Yamazaki Y, Yokota TA, Matsumoto N, Matsukura S (1998) Severe chronic active Epstein-Barr virus infection accompanied by virus-associated hemophagocytic syndrome, cerebellar ataxia and encephalitis. *Psychiatry Clin Neurosci* 52(4):449–452

Priv. Doz. Dr. D. M. Hermann (✉) ·
C. L. Bassetti
Dept. of Neurology
University Hospital Zurich (USZ)
Frauenklinikstr. 26
8091 Zurich, Switzerland
Tel.: +41-1/255-5526
Fax: +41-1/255-4507
E-Mail: dirk.hermann@usz.ch

J. Fehr
Dept. of Internal Medicine/Haematology
University Hospital Zurich (USZ)
Zurich, Switzerland