**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Largely flat latitudinal life history clines in the dung fly *Sepsis fulgens* across Europe (Diptera: Sepsidae)

Roy, Jeannine ; Blanckenhorn, Wolf U ; Rohner, Patrick T

Abstract: Clinal variation in body size and related life history traits is common and has stimulated the postulation of several eco-geographical rules. Whereas some clinal patterns are clearly adaptive, the causes of others remain obscure. We investigated intra-specific body size, development time and female fecundity (egg size and number) clines across 13 European populations of the dung fly *Sepsis fulgens* spanning 20° latitude from southern Italy to Estonia in a genetic common garden approach. Despite very short generation times (ca. 2 weeks at 24 °C), we found a converse Bergmann cline (smaller size at higher latitudes). As development time did not change with latitude (flat cline), integral growth rate thus likely declines towards the pole. At the same time, early fecundity, but not egg size, increased with latitude. Rather than being mediated by seasonal time constraints, the body size reduction in the northernmost flies from Estonia could suggest that these are marginal, edge populations, as when omitting them the body size cline became flat as well. Most of the other sepsid species investigated to date also show flat body size clines, a pattern that strikingly differs from *Drosophila*. We conclude that *S. fulgens* life history traits appear to be shaped by similar environmental pressures and selective mechanisms across Europe, be they adaptive or not. This reiterates the suggestion that body size clines can result as a secondary consequence of selection pressures shaping an entire life history syndrome, rendering them inconsistent and unpredictable in general.

DOI: <https://doi.org/10.1007/s00442-018-4166-7>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-157753>

Journal Article

Accepted Version

Originally published at:

Roy, Jeannine; Blanckenhorn, Wolf U; Rohner, Patrick T (2018). Largely flat latitudinal life history clines in the dung fly *Sepsis fulgens* across Europe (Diptera: Sepsidae). *Oecologia*, 187(3):851-862.

DOI: <https://doi.org/10.1007/s00442-018-4166-7>

Largely flat latitudinal life history clines in the dung fly *Sepsis fulgens* across Europe (Diptera: Sepsidae)

Jeannine Roy, Wolf U. Blanckenhorn*, Patrick T. Rohner

*Department of Evolutionary Biology & Environmental Studies, University of Zurich,
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

<wolf.blanckenhorn@ieu.uzh.ch>; <patrick.rohner@ieu.uzh.ch> j.roy@gmx.ch

Oecologia <https://doi.org/10.1007/s00442-018-4166-7>

Abstract (243 words)

Clinal variation in body size and related life history traits is common and has stimulated the postulation of several eco-geographical rules. Whereas some clinal patterns are clearly adaptive, the causes of others remain obscure. We investigated intraspecific body size, development time and female fecundity (egg size and number) clines across 13 European populations of the dung fly *Sepsis fulgens* spanning 20 degrees latitude from southern Italy to Estonia in a genetic common garden approach. Despite very short generation times (ca. 2 weeks at 24°C), we found a converse Bergmann cline (smaller size at higher latitudes). As development time did not change with latitude (flat cline), integral growth rate thus likely declines towards the pole. At the same time, early fecundity, but not egg size, increased with latitude. Rather than being mediated by seasonal time constraints, the body size reduction in the northernmost flies from Estonia could suggest that these are marginal, edge populations, as when omitting them the body size cline became flat as well. Most of the other sepsid species investigated to date also show flat body size clines, a pattern that strikingly differs from *Drosophila*. We conclude that *S. fulgens* life history traits appear to be shaped by similar environmental pressures and selective mechanisms across Europe, be they adaptive or not. This reiterates the suggestion that body size clines can result as a secondary consequence of selection pressures shaping an entire life history syndrome, rendering them inconsistent and unpredictable in general.

Keywords Body size, development time, egg size, fecundity, geographic differentiation, genetic differentiation, latitudinal cline, life history

Introduction

The study of large-scale geographic variation in life history traits, first and foremost body size, has a long tradition in evolutionary ecology (Partridge & Coyne 1997; Blackburn et al. 1999; Chown and Gaston 1999; Blanckenhorn and Demont 2004; Shelomi 2012). Body size of closely related species often increases with latitude, known as Bergmann's rule (Bergmann 1847), and corresponding intra-specific trends, sometimes called James' rule (James 1970), also commonly occur. Nevertheless, opposite, so-called converse Bergmann clines also exist (Mousseau 1997; Blanckenhorn and Demont 2004; Shelomi 2012). All these patterns are generally thought to be caused by systematic latitudinal climate variation, although the specific underlying environmental variables and causal mechanisms are frequently unclear. This is because body size is intimately entwined with other life history traits such as juvenile development time and growth rate, which ultimately co-determine final adult body size (Nylin and Gotthard 1998; Blanckenhorn 1999). Moreover, as body size affects most if not all life history and physiological traits of an organism in a concerted syndrome, several reproductive traits (egg size, egg number, etc.) also tend to exhibit geographic clines (e.g. Azevedo et al. 1996; Armbruster et al. 2001; Lardies et al. 2010). Though there are many exceptions, higher temperatures usually result in smaller body sizes in ectothermic organisms, an entirely plastic physiological response of unclear adaptive value (named temperature-size rule: Stevenson 1985; Atkinson 1994; van der Have & de Jong 1996; Van Voorhies 1996; Horne et al. 2015, 2018). While temperatures systematically decrease with latitude, season length decreases too, thus constraining an individual's growth period and hence final body size, producing adaptive converse Bergmann size clines especially in large species with long generation times, such as many beetles, grasshoppers and butterflies (smaller towards the poles: Blanckenhorn & Demont 2004; Nygren et al. 2008; Shelomi 2012). Analogous effects are exerted by altitude, albeit at much more condensed spatial scales, such that expected patterns

are often not found (Shelomi 2012; Klepsatel et al. 2014; Horne et al. 2018). It follows that several environmental factors (temperature, season length, etc.), more or less independently, act in conjunction on the entire life history syndrome of any particular species, resulting in diverse and potentially intermediate clinal patterns that are ultimately difficult to predict (Blanckenhorn & Demont 2004). More studies of geographic and clinal life history variation are therefore needed.

Although the literature by now boasts many studies on latitudinal clines for a great number of taxa, in insects work on the “model species” *Drosophila* tends to dominate (Shelomi 2012; Schilthuizen & Kellermann 2014; Flatt 2016; Horne et al. 2018). Such strong bias towards *Drosophila* cannot be representative from a biodiversity perspective, especially because Dipterans are very derived phylogenetically (see Misof et al. 2014). This occurs also because small insects, such as *Drosophila*, can be studied as genetic lineages under common garden conditions in the laboratory. Such work therefore tests for evolved, genetic clines, which from an evolutionary point of view is superior to merely phenotypic patterns derived from field-caught specimens, as commonly utilized in macro-ecology (Chown and Gaston 2010). In *Drosophila*, Bergmann clines are the rule within species (e.g. James et al. 1997; Huey et al. 2000; Shelomi 2012; Klepsatel et al. 2014; Fabian et al. 2015; Table 1), and apparently also across species (Rohner et al. 2018a). This can be expected in small species with short generation times, which consequently are multivoltine (multiple generations per year) over most of their distributional range (Blanckenhorn & Demont 2004). By contrast, larger insects such as beetles, grasshoppers or water bugs with typically long development times relative to the available season length often have univoltine life cycles (one generation per year; e.g. Masaki 1967; Blanckenhorn and Fairbairn, 1995; Mousseau 1997; Sota et al. 2000), and thus more likely exhibit converse Bergmann clines because they regularly experience seasonal time limitations forcing them to

abbreviate their development at the expense of a reduction in final body size (Blanckenhorn & Demont 2004).

Sepsid “black scavenger” or “dung” flies (Diptera: Sepsidae) are common and widespread worldwide, with many species in the tropics (Ang et al. 2013; see <http://sepsidnet-rmbr.nus.edu.sg/> for photographs). On average they are a bit larger than *Drosophila* (3-8 mm body length), and also multivoltine throughout most of their range. Sepsids generally depend on decaying organic material for reproduction and larval development, often breeding in livestock faeces (e.g. cow dung; Pont & Meier 2002), a common habitat in agricultural grasslands. In populating ephemeral habitats, sepsids are ecologically similar to *Drosophila* (Blanckenhorn 1999; Rohner et al. 2018a,b). Adult sepsids feed on dung (for protein) and nectar (for energy) and therefore, as *Drosophila*, are anautogenous income breeders, although males (but not females) likely produce their first batch of gametes already during the juvenile phase (Teuschl & Blanckenhorn 2007). Not least because they are easy to rear and have short generation times, sepsid flies are fast becoming model organisms in behaviour, ecology, and evolution (e.g. Teuschl & Blanckenhorn 2007; Puniamoorthy et al. 2009, 2012, 2014; Berger et al. 2013; Rohner et al. 2015; Esperk et al. 2016; Busso & Blanckenhorn, 2018). So far, the few studies of latitudinal variation in this group yield inconsistent evidence (summarized in Table 1). Puniamoorthy et al.’s. (2012) study suggests a Bergmann cline in European but not North American *Sepsis punctum* (see also Dmitriew & Blanckenhorn 2012), and Berger et al. (2013) show a cline in development rate in European *S. punctum* that is consistent with seasonal time constraints operating at higher latitudes (faster in northern Europe). No (i.e. flat) clines are evident in European *S. cynipsea* (Rohner et al. 2016) and *S. thoracica* (Busso & Blanckenhorn 2018).

Sepsis fulgens is one of the most ubiquitous, entirely black sepsid species in Europe about which almost nothing is known so far (Pont & Meier 2002; Fig. 1). We here examine clinal body size, development time and female fecundity (egg number and size) variation of 13 *Sepsis fulgens* populations across roughly 20 degrees latitude from southern Italy to Estonia. Similar to *Drosophila* studies, we took a genetic common garden approach by working with iso-female lines (i.e. the laboratory offspring of single field-caught females) to investigate evolved clines. We additionally assessed the extent of phenotypic plasticity in the above traits by rearing flies at a range of 2–4 laboratory temperatures, thus evaluating the temperature-size rule (Atkinson 1994; Horne et al. 2015, 2018). As this sepsid species is relatively small (body length ca. 4 mm), it features multiple generations per year (e.g. > 8 overlapping generations in lowland Switzerland (pers. obs.); and 2-3 generations per year even at higher latitude in Moscow, Russia: 55.45°N, 37.36°E; Pont & Meier, 2002; cf. Table 2), such that seasonal time limitations would generally not exert strong selection except in the last generation(s) before winter. We thus expected this species to show a Bergmann cline (larger towards the pole), and no countergradient pattern of accelerated development as found for the larger *S. punctum* by Berger et al. (2013; Blanckenhorn & Demont 2004). Consequently, following the expected body size pattern, we also expected either more or larger eggs in populations from higher latitudes.

Methods

Fly cultures and maintenance

Sepsis fulgens Meigen, 1826 (Diptera: Sepsidae) is a eurytopic species occurring throughout Europe, North Africa and the Middle East. The species is commonly found on cow, pig and horse dung, as well as on manure piles (Pont & Meier 2002). Contrary to many sepsids, the species is always entirely black (see http://sepsidnet-rmbr.nus.edu.sg/Sepsis_fulgens.html; Fig. 1).

We studied flies from 13 latitudinal field populations from southern Italy to Estonia (Fig. 1; Table 2 lists some standard climate characteristics of the collections sites). Live wild-caught females from these populations were collected and brought to our laboratory at different times mainly during the seasons 2013 & 2014. Cultures were subsequently kept as iso-female lines (i.e. the descendants of one wild-caught female) for multiple (> 15) generations in the laboratory (ca. 3-4 weeks per generation) before being used in our experiment in 2015. We had several (5–13) replicate families (i.e. iso-female lines) per population (Table 2), each housed in separate 1-litre plastic containers that were regularly supplied with fresh cow dung, sugar, and water ad libitum, and kept in a climate chamber at $24\pm 1^{\circ}\text{C}$, 60% relative humidity, and 13:11 h light:dark cycle. Assessing iso-female lines instead of wild-caught females minimises environmental variation influencing the phenotype, thus exposing mainly genetic variation, which remains best preserved in the laboratory by this culturing method.

Laboratory rearing experiments

Two rearing experiments were performed. The first rearing served to analyse body size and development time of both sexes, using 11 populations (cf. Fig. 1 and above): Lamezia, Padula, Terni, Transimeno, Maggia, Möschberg, Zürich, Bielefeld, Skelde, Rahinge and Pehka (Table 2). Iso-female lines were provided with fresh dung in their rearing containers, into which multiple females could lay eggs for a duration of up to 4 h. These egg masses were then split randomly among four environments differing in temperature only (5-100 eggs per mass culture dish). The larvae were subsequently allowed to develop in overabundant and uniform defrosted cow dung at 60% relative humidity and long photoperiods (≥ 13 h) at constant 12°C , 18°C , 24°C or 30°C in 50 ml glass vials. Vials were checked daily for offspring until no more individuals emerged. The

thorax length (distance between the protoniroleural lobe and the tip of the scutellum) of two randomly chosen offspring males and two offspring females was imaged and measured as an index of body size. The structural outline of the thorax best reflects overall body mass as it contains the important flight muscles.

The second rearing focused on female reproductive traits of 8 latitudinal populations: Lamezia, Petroia, Padula, Maggia, Möschberg, Zürich, Witterschlick and Rahinge (Fig. 1). Male and female flies from any given iso-female line, reared at constant 24°C, 60% relative humidity and 13 h photoperiod, were paired randomly and subsequently kept in one of two environments differing in temperature only (60% relative humidity and photoperiod ≥ 13 h); for each replicate iso-female line two pairs were kept in separate 50 ml glass vials with fresh dung and sugar at 18°C, and two pairs at 30°C. Every day the dung was checked for eggs. As soon as eggs were laid (= day of first reproduction), they were counted and five of them imaged, and the pair was provided with fresh dung for another day. In case the female laid additional eggs on the second day, we considered them part of the same clutch (as sepsids do not always lay discrete clutches within 24h). Thereafter females were sacrificed and measured. From this experiment we obtained data on adult age at first reproduction (days), the total number of laid eggs in the first 2 days (i.e. first clutch size, or early fecundity), egg size (length and width, from which egg volume was calculated using the formula for a spheroid: $(1/6)\pi \cdot \text{length} \cdot \text{width}^2$). For convenience, fore tibia length was used as an index of female body size; correlations among all structural length traits (including thorax) of *S. fulgens* exceed $r > 0.97$ in both males and females ($n > 40$ individuals).

All images (thorax, eggs, foreleg) were taken with a Leica MZ 12 stereomicroscope and a Leica DC490 digital camera. Measurements were performed using the program tpsDIG2 (© F. J. Rohlf 2010) and the program PAST (© O. Hammer 2003). All analyses were conducted with the

software IBM SPSS Statistics 23 (SPSS Inc.).

Results

Development time and body size (thorax length) were analysed with sex and rearing temperature as fixed factors and latitude as a continuous covariate with univariate ANOVAs (General Linear Models with underlying normal error distribution), using line (i.e. family) means (of typically two individuals per sex) as independent data points (and global error) because this is a genetic study and we were not specifically interested in within-line phenotypic variation. Development time did not change with latitude, i.e. the cline was flat ($F_{1,551} = 0.47, p > 0.3$; Fig. 2), with no significant interactions (e.g. temp*latitude: $F_{3,544} = 1.78, p = 0.149$; all other interactions $p > 0.2$). In contrast, body size of the flies clearly decreased with latitude ($F_{1,648} = 14.38, p < 0.001$; Fig. 2), again revealing no significant interactions (temp*latitude: $F_{3,651} = 1.16, p > 0.3$; all other interactions $p > 0.3$). Development time strongly increased as temperature decreased ($F_{3,551} = 477.83, p < 0.001$; Fig. 2), while body size was largest at 18°C, followed by 24°C, and smallest at both extreme temperatures of 30°C and 12°C ($F_{3,648} = 2.29, p = 0.067$, Fig. 2). As usual in insects, females are larger than males in this species ($F_{1,648} = 117.27, p < 0.001$). Sexual size dimorphism (calculated as the logarithm of the ratio between female and male thorax length) decreased from cold to warm temperatures ($F_{1,310} = 9.98, p = 0.002$); however, this pattern was only due to males being much smaller at 12°C, as sexual size dimorphism remained constant from 18°C to 30°C. When calculating an analogous index for the development time difference between the sexes, there was no difference between the sexes ($F_{1,201} = 0.08, p = 0.146$), and the development time difference did not vary strongly with temperature, although there was a trend of males taking overall longer to develop at cold temperatures ($F_{1,201} = 3.70, p = 0.056$; Fig. 2).

The female reproductive trait experiment was similarly analysed with temperature as fixed factor, and latitude, latitude² (to test for nonlinearity), and female size (tibia length) as continuous covariates, again using line means (of typically two females) as independent data points. Flies from all populations first reproduced after approximately the same amount of time, but oviposition occurred faster when warmer (18°C: 6.62 ± 0.073 (SE) days; 30°C: 4.18 ± 0.076 days), although this difference was not significant ($F_{1,76} = 2.48, p = 0.119$). Early fecundity (square-root-transformed number of eggs laid in the first 2 days) was unaffected by rearing temperature ($F_{1,76} = 0.11, p = 0.737$) but increased with body size ($F_{1,76} = 7.54, p = 0.008$) and latitude ($F_{1,76} = 13.30, p < 0.001$), and additionally showed a negative latitude² (i.e. hump-shaped, concave) effect ($F_{1,76} = 12.55, p = 0.001$), suggesting that the increase in fecundity with latitude could be asymptotic (Fig. 3). In contrast, cube-root-transformed egg volume was not affected by any of these factors (Figs. 3, 4). Consequently, no trade-off between the number and size of laid eggs was found.

Discussion

Despite *S. fulgens*' short generation time (2 weeks at 24°C, 3 weeks at 18°C; Fig. 2), we did not find the expected Bergmann size cline (or James' rule) in Europe, but rather a converse Bergmann cline (smaller size toward the north pole; cf. Blanckenhorn & Demont 2004). As development time did not change with latitude (Fig. 2), this implies that growth rate does not accelerate towards the pole, although our measurements are restricted to integral measurements of growth, which have been shown to sometimes differ from more precise growth rate measures (Tammaru and Esperk 2007; Rohner et al. 2017). Nevertheless, this result is opposite to what was found for the closely related but larger *S. punctum* (Berger et al. 2013), and also for the much larger yellow dung fly (Blanckenhorn et al. 2018). Rather than being mediated by seasonal time

constraints, which would imply an adaptive response (Roff 1980; Kivelä et al. 2011), the size decline in the northernmost flies from Estonia alternatively suggests that these could be marginal, edge populations in terms of environmental factors such as temperature, food, precipitation, or irradiation etc.; when omitting the two Estonian populations from the analysis (cf. Fig. 1), the European body size cline became flat as well. We therefore conclude that the clinal patterns found here are not necessarily adaptive, as body size, development time, fecundity and egg size are similar across all of Europe, hence presumably shaped by similar environmental pressures everywhere, or alternatively by stabilizing ecological selection relating to these flies' ephemeral food resource (dung). However, we have no direct evidence for the latter, and in the related dung fly *Sepsis cynipsea* overall balancing selection could not be found in an extensive study of one single Swiss population (Blanckenhorn 2007). Unfortunately, we could not obtain *S. fulgens* from more northern sites to verify whether these populations are indeed marginal and/or subject to seasonal time constraints there.

Clear body size clines also lack in the related *S. thoracica* and *S. cynipsea* in Europe (Rohner et al. 2016; Busso & Blanckenhorn 2018; Table 1). This situation for sepsids contrasts with that of various *Drosophila* species, the model insects also in evolutionary ecology, which generally show positive Bergmann (or James') clines within species that even have been demonstrated to re-evolve in relatively short time after invasion of a new continent (James et al. 1997; Huey et al. 2000; Zwaan et al. 2000; Klepsatel et al. 2014; Fabian et al. 2015; Flatt 2016; Table 1). Our common garden study implies that all the patterns found for *S. fulgens* (as well as the above-mentioned two other sepsid species) are genetic, i.e. evolved, as is also the case in the listed *Drosophila* studies. As sepsid flies are comparable to *Drosophila* in terms of nutrition ecology, life history, size, dispersal capacity and culturing methods (cf. Blanckenhorn 1999; Rohner et al. 2017, 2018a,b), this supports the overall conclusion that the evolution of clinal life

history patterns, and their underlying mechanisms, are complex and not easily predictable in general. In the end, therefore, we are left without being able to identify the mechanisms that mediate the cline differences between the two fly species summarized in Table 1 (cf. Blanckenhorn 1999).

Even though *S. fulgens* flies became smaller, they laid more but not larger eggs at higher latitudes, at least initially between 40 and 50 degrees latitude, beyond which early fecundity declined again in the potentially marginal Estonian populations (Fig. 3). This occurred despite the fact that within populations larger flies produced more eggs (Fig. 4), as is typical in insects (Honek 1993). Berger et al. (2008) showed that the realized fecundity of butterflies can be limited by cool weather, an effect that might select for increased size-specific early fecundity in insects at high latitudes. It could also be that northern flies invest more in female abdomens at the expense of thorax length, but we have no data on abdomen size, which is generally difficult to measure in insects because it is so flexible. Lacking sex by latitude interactions further suggest no sex-specific shifts in investment into the thorax relative to other body parts in females from high latitudes. Egg size also appears to be independent of body size in this species. Derived from Bergmann's rule and the temperature-size-rule (Stevenson 1985; Atkinson 1994; Partridge et al. 1994; van der Have & de Jong 1996; Van Voorhies 1996; Horne et al. 2015), our expectation was that the cooler northern climate would promote the evolution of larger eggs, a plastic response found e.g. in yellow dung flies (Blanckenhorn 2000; Blanckenhorn & Heyland 2004) and *D. melanogaster* (Azevedo et al. 1996), but this was not the case here. Our results also do not support the hypothesis that more extreme climates select for larger eggs providing offspring a competitive advantage during early development (Fox & Czesak 2000). That clutch size increases with latitude while egg size remains unaffected further implies the absence of a size-number trade-off, which is often expected (Smith & Fretwell 1974), perhaps due to seasonal time

constraints on this income breeder's life history at higher latitudes (Berger et al. 2008). Again, it seems that clines in reproductive parameters (fecundity and egg size) can turn out equally variable and unpredictable depending on species and environments (cf. Fox & Czesak 2000; e.g. Bauerfeind et al. 2018).

Despite the patterns documented here for *S. fulgens* being heritable, phenotypic plasticity in response to temperature was extensive for development time and body size (Fig. 2), albeit not for reproductive traits (Fig. 3). While development time lengthens strongly and expectedly as temperatures drop, body size did not strictly follow the temperature-size rule in this species (smaller when warmer: Atkinson 1994), as it was maximal at intermediate temperatures of 18°C and 24°C but lower at 12°C and 30°C (Fig. 2). For the related *S. cynipsea* the lower temperature threshold at which development ceases has been estimated to be roughly 9°C (Blanckenhorn 1999); 12°C appears to be a marginal temperature for *S. fulgens* as well, at which flies require 55 days to complete development to adulthood (Fig. 2). When reared at temperatures above this threshold, sexual size dimorphism did not vary with temperature, a common pattern in insects (Hirst et al. 2015). The increase in size dimorphism at 12°C further suggests that the sexes do differ in their temperature-size response at such marginal temperatures: males took longer time to emerge and emerged smaller, so they seem to do worse at cool temperatures. In contrast, rearing temperature of adult females did not affect fecundity here, as is often not the case and thus not expected, whereas egg size was, albeit non-significantly, reduced at 30°C relative to 18°C (Fig. 3), as expected on physiological grounds related to processes presumably mediating the temperature-size rule (Van Voorhies 1996; Atkinson & Sibly 1997; Fox & Czesak 2000; Blanckenhorn 2000; Fischer et al. 2003; Blanckenhorn & Heyland 2004; Garrad et al. 2016).

Flat latitudinal clines for body size, development time and/or growth rate in *S. fulgens* as demonstrated here do not necessarily imply that phenotypic clines based on field-caught specimens are also flat. As also shown here, all these traits are considerably plastic, such that lower average temperatures at high latitudes, often combined with low population densities and hence more abundant food per capita, could suffice to produce Bergmann clines (Blanckenhorn & Demont 2004). Unfortunately, we have no good data on field flies, as we concentrated on collecting live specimens. To convincingly demonstrate phenotypic (i.e. macro-ecological) clines based on field samples in this small and very plastic fly would require numerous specimens sampled systematically at all locations over the season. This is no trivial task. We thus cannot present a phenotypic field cline in addition to the genetic cline. Furthermore, while climate information for our sampling locations is available (Table 2), such data usually do not serve to single out the responsible environmental factors for a given cline, as the crucial parameters season length and temperature are typically highly correlated with latitude (and altitude) so that multiple regression approaches necessarily fail. Latitude thus remains the most easily assessable proxy (see Blanckenhorn & Busso 2018; Blanckenhorn et al. 2018).

In conclusion, latitudinal clines in body size, development time and reproductive traits in European *S. fulgens* populations are mostly flat, implying no systematic life history shifts across a wide range of climates. This may relate to these flies being relatively small and ubiquitous, because their substrate (livestock dung) is also ubiquitous in agricultural grasslands that are very common in Europe. Seasonal time constraints therefore probably do not exert strong selection shaping the life history of even the northernmost populations, where these flies still feature multiple generations per season (as e.g. in Moscow; op. cit.). As sepsid flies are widespread in the tropics (Pont & Meier 2002; Ang et al. 2013), northern European climes likely are secondary, marginal habitats for them, though not in terms of food (i.e. dung) availability. Despite being bad

fliers, sepsids are widespread such that gene flow should nevertheless be extensive, hampering local adaptation of populations (Busso & Blanckenhorn 2018). It remains puzzling that sepsids, being ecologically similar to drosophilids also in terms of dispersal capacity and thus gene flow between populations, show very different clinal life history patterns. So, this result reiterates the interpretation that clinal life history patterns are complex and not easily predictable in general, as temperature effects (presumably fostering positive Bergmann clines) and season length effects (fostering converse Bergmann clines) may cancel out, ultimately possibly producing the largely flat clines observed here (Blanckenhorn & Demont 2004).

Acknowledgements

We thank Martin Schäfer, Juan Pablo Busso, Anders Kjaersgaard, Nalini Puniamoorthy, Toomas Esperk, and Toomas Tammaru for collecting flies, and the entire Wolfpack for various support. This study was supported by grant no. SNF 31003A-143787 from the Swiss National Science Foundation.

330   **References** (65, 1606 words)

- 331   Ang YC, Puniamoorthy J, Pont AC, Bartak M, Blanckenhorn WU, Eberhard WG,
332   Puniamoorthy N, Silva V, Munari L, Meier R (2013) Sepsidnet: A plea for digital
333   reference collections and other science-based digitization initiatives in taxonomy.
334   Syst Entomol 38:637–644.
- 335   Armbruster P, Bradshaw WE, Ruegg K, Holzapfel CM (2001) Geographic variation
336   and the evolution of reproductive allocation in the pitcher-plant mosquito,
337   *Wyeomyia smithii*. Evolution 55:439-444.
- 338   Atkinson D (1994) Temperature and organism size a biological law for ectotherms?
339   Adv Ecol Res 25:1-58.
- 340   Atkinson D, Sibly RM (1997) Why are organisms usually bigger in colder
341   environments? Making sense of a life history puzzle. Trends Ecol Evol 12:235-
342   239.
- 343   Azevedo RBR, French V, Partridge L (1996) Thermal evolution of egg size in
344   *Drosophila melanogaster*. Evolution 50:2338-2345.
- 345   Bauerfeind SS, Schäfer MA, Berger D, Blanckenhorn WU, Fox CW (2018)
346   Replicated latitudinal clines in reproductive traits of European and North American
347   yellow dung flies. Oikos in press.
- 348   Berger D, Walters RJ, Gotthard K (2008) What limits insect fecundity? Body size-
349   and temperature-dependent egg maturation and oviposition in a butterfly. Funct
350   Ecol 22:523-529.
- 351   Berger D, Postma E, Blanckenhorn WU, Walters RJ (2013) Quantitative genetic
352   divergence and standing genetic (co)variance in thermal reaction norms along
353   latitude. Evolution 67:2385-2399.

354 Bergmann C (1847) Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer
 355 Grösse. Göttinger Studien 3:595–708.

356 Blackburn TM, Gaston KJ, Loder N (1999) Geographic gradients in body size: a
 357 clarification of Bergmann's rule. Diversity Distributions 5:165-174.

358 Blanckenhorn WU (1999) Different growth responses to food shortage and
 359 temperature in three insect species with similar life histories. Evol Ecol 13:395-
 360 409.

361 Blanckenhorn WU (2000) Temperature effects on egg size and their fitness
 362 consequences in the yellow dung fly. Evol Ecol 14:627-643.

363 Blanckenhorn WU (2007) Case studies of the differential equilibrium hypothesis of
 364 sexual size dimorphism in dung flies. In: Sex, Size and Gender Roles. Evolutionary
 365 Studies of Sexual Size Dimorphism (Fairbairn DJ, Blanckenhorn WU, Székely T,
 366 eds.), Oxford University Press, Oxford, pp. 106-114.

367 Blanckenhorn WU, Demont M (2004) Bergmann and converse Bergmann latitudinal
 368 clines in Arthropods: Two ends of a continuum? Integr Comp Biol 44:413–424.

369 Blanckenhorn WU, Fairbairn DJ (1995) Life history adaptation along a latitudinal
 370 cline in water striders. J Evol Biol 8:21-41.

371 Blanckenhorn WU, Heyland A (2004) The quantitative genetics of two life history
 372 trade-offs in the yellow dung fly in abundant and limited food environments. Evol
 373 Ecol 18:385-402.

374 Blanckenhorn WU, Bauerfeind SS, Berger D, Davidowitz G, Fox CW, Guillaume F,
 375 Nakamura S, Nishimura K, Sasaki H, Stillwell RC, Tachi T, Schäfer MA (2018)
 376 Season length—not temperature—drives life-history clines on three continents in a
 377 widespread dungfly. Ecography in press.

378 Busso JP, Blanckenhorn WU (2018) Climatic factors shaping plastic trade-offs in the
 379 polyphenic black scavenger fly *Sepsis thoracica* (Diptera: Sepsidae). J Biogeo
 380 45:593-603. DOI: 10.1111/jbi.13140.

381 Chown SL, Gaston KJ (1999) Exploring links between physiology and ecology at
 382 macro-scales: the role of respiratory metabolism in insects. Biol Rev 74:87-120.

383 Chown SL, Gaston KJ (2010) Body size variation in insects: a macroecological
 384 perspective. Biol Rev 85:139–169.

385 Dmitriew C, Blanckenhorn WU (2012) The role of sexual selection and conflict in
 386 mediating among-population variation in mating strategies and sexually dimorphic
 387 traits in the black scavenger fly *Sepsis punctum*. PlosOne 7:e49511.

388 Esperk T, Kjaersgaard A, Walters RJ, Berger D, Blanckenhorn WU (2016)
 389 Plastic and evolutionary responses to heat stress in a temperate dung fly:
 390 negative correlation between basal and induced heat tolerance? J Evol Biol
 391 29:900-915.

392 Fabian DK, Lack JB, Mathur V, Schloetterer C, Schmidt PS, Pool JE, Flatt T (2015)
 393 Spatially varying selection shapes life history clines among populations of
 394 *Drosophila melanogaster* from sub-Saharan Africa. J Evol Biol 28:826-840.

395 Fischer K, Brakefield PM, Zwaan BJ (2003) Plasticity in butterfly egg size: Why
 396 larger offspring at lower temperatures? Ecology 84:3138-3147.

397 Flatt T (2016) Genomics of clinal variation in *Drosophila*: disentangling the
 398 interactions of selection and demography. Mol Ecol 25:1023-1026.

399 Fox CW, Czesak ME (2000) Evolutionary ecology of progeny size in arthropods. Ann
 400 Rev Entomol 45:341-369.

401 Garrad R, Booth DT, Furlong MJ (2016) The effect of rearing temperature on
 402 development, body size, energetics and fecundity of the diamondback moth. Bull
 403 Entomol Res 106:175–181.

404 Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005). Very high resolution
 405 interpolated climate surfaces for global land areas. Int J Climatol 25: 1965-1978.

406 Hirst AG, Horne CR, Atkinson D (2015) Equal temperature–size responses of the
 407 sexes are widespread within arthropod species. Proc R Soc B 282:20152475.
 408 <http://dx.doi.org/10.1098/rspb.2015.2475>

409 Honek A (1993) Intraspecific variation in body size and fecundity in insects: a general
 410 relationship. Oikos 66:483-492.

411 Horne CR, Hirst AG, Atkinson D (2015) Temperature-size responses match
 412 latitudinal size clines in arthropods, revealing critical differences between aquatic
 413 and terrestrial species. Ecol Lett 18:327-335.

414 Horne CR, Hirst AG, Atkinson D (2018) Insect temperature-body size trends common
 415 to laboratory, latitudinal and seasonal gradients are not found across altitudes.
 416 Funct Ecol in press.

417 Huey RB, Gilchrist GW, Carlson ML, Berrigan D, Serra L (2000) Rapid evolution of
 418 a geographic cline in size in an introduced fly. Science 287: 308–309.

419 James AC, Azevedo RBR, Partridge L (1997) Genetic and environmental responses to
 420 temperature of *Drosophila melanogaster* from a latitudinal cline. Genetics
 421 146:881-890.

422 James FC (1970) Geographic size variation in birds and its relationship to climate.
 423 Ecology 51:365–390.

424 Kivelä SM, Välimäki P, Carrasco D, Mäenpää MI, Oksanen J (2011) Latitudinal
425 insect body size clines revisited: a critical evaluation of the saw-tooth model. J
426 Anim Ecol 80:1184-95.

427 Klepsatel P, Galikova M, Huber CD, Flatt T (2014) Similarities and differences in
428 altitudinal versus latitudinal variation for morphological traits in *Drosophila*
429 *melanogaster*. Evolution 68:1385–1398.

430 Lardies MA, Arias MB, Bacigalupe LD (2010) Phenotypic covariance matrix in life-
431 history traits along a latitudinal gradient: a study case in a geographically
432 widespread crab on the coast of Chile. Mar Ecol Prog Ser 412:179-187.

433 Masaki S (1967) Geographic variation and climatic adaptation in a field cricket.
434 Evolution 21:725-741.

435 Misof B, et al. (2014) Phylogenomics resolves the timing and pattern of insect
436 evolution. Science 346:763-767.

437 Mousseau TA (1997) Ectotherms follow the converse to Bergmann's rule. Evolution
438 51:630–632.

439 Nygren GH, Bergström A, Nylin S (2008) Latitudinal body size clines in the butterfly
440 *Polyommatus icarus* are shaped by gene-environment interactions. J Insect Science
441 8:47.

442 Nylin S, Gotthard K (1998) Plasticity in life history traits. Ann Rev Entomol 43:63-
443 83.

444 Partridge L, Coyne JA (1997) Bergmann's rule in ectotherms: is it adaptive?
445 Evolution 51:632-635.

446 Partridge L, Barrie B, Fowler K, French V (1994) Evolution and development of body
447 size and cell size in *Drosophila melanogaster* in response to temperature.

448 Evolution 48:1269-1276.

449 Pont AC, Meier R (2002) The Sepsidae (Diptera) of Europe. Fauna Entomol Scand
450 37:1–221.

451 Puniamoorthy,N, Rifqi M, Tan D, Meier R (2009) From kissing to belly stridulation:
452 Comparative analysis reveals surprising diversity, rapid evolution, and much
453 homoplasy in the mating behavior of 27 Species of sepsid flies (Diptera: Sepsidae).
454 J Evol Biol 22:2146-2156.

455 Puniamoorthy N, Schäfer MA, Blanckenhorn WU (2012) Sexual selection accounts
456 for geographic reversal of sexual size dimorphism in the dung fly, *Sepsis punctum*
457 (Diptera: Sepsidae). Evolution 66:2117-2126.

458 Puniamoorthy N, Schäfer MA, Römbke J, Meier R, Blanckenhorn WU (2014)
459 Ivermectin sensitivity is an ancient trait affecting all ecdysozoa but shows
460 phylogenetic clustering among sepsid flies. Evol Appl 7:548-554.

461 Roff D (1980) Optimizing development time in a seasonal environment: the ‘ups and
462 downs’ of clinal variation. Oecologia 45:202–208.

463 Rohner PT, Bächli G, Pollini Paltrinieri L, Duelli P, Obrist MK, Jochmann R,
464 Blanckenhorn WU (2015) Distribution, diversity gradients and Rapoport’s
465 elevational rule in the black scavenger flies of the Swiss Alps (Diptera: Sepsidae).
466 Insect Conserv Diversity 4:367-376.

467 Rohner PT, Blanckenhorn WU, Puniamoorthy N (2016) Sexual selection on male size drives
468 the evolution of male-biased sexual size dimorphism via the prolongation of male
469 development. Evolution 70:1189-1199.

470 Rohner PT, Blanckenhorn WU, Schäfer MA (2017) Critical weight mediates sex-
 471 specific body size plasticity and sexual dimorphism in the yellow dung fly
 472 *Scathophaga stercoraria* (Diptera: Scatophagidae). *Evol Dev* 19:147-156.
 473 Rohner PT, Pitnick S, Blanckenhorn WU, Snook RR, Bächli G, Lüpold S (2018a)
 474 Global macroecology of size, sexual size dimorphism, dispersal and range size in
 475 fruit flies (Diptera: Drosophilidae). *Ecography* 41: in press.
 476 Rohner PT, Teder T, Esperk T, Lüpold S, Blanckenhorn WU (2018b) The evolution
 477 of male-biased sexual size dimorphism is associated with increased body size
 478 plasticity in males. *Funct Ecol* 32:581-591. DOI: 10.1111/1365-2435.13004
 479 Sota T, Takami Y, Kubota K, Ujiie M, Ishikawa R (2000) Interspecific body size
 480 differentiation in species assemblages of the carabid subgenus *Ohomopterus* in
 481 Japan. *Pop Ecol* 42:279–291.
 482 Schäfer MA, Berger D, Rohner PT, Kjærsgaard A, Bauerfeind SS, Guillaume F, Fox
 483 CW, Blanckenhorn WU (2018) Geographic clines in wing morphology relate to
 484 colonization history in New but not Old World populations of the yellow dung fly.
 485 *Evolution* submitted.
 486 Schilthuizen M, Kellermann V (2014) Contemporary climate change and terrestrial
 487 invertebrates: evolutionary versus plastic changes. *Evol Appl* 7:56-67.
 488 Shelomi M (2012) Where are we now? Bergmann's rule sensu lato in insects. *Am Nat*
 489 180:511-519.
 490 Smith C C, Fretwell SD (1974) The optimal balance between size and number of
 491 offspring. *Am Nat* 108:499-506.

492 Stevenson RD (1985) Body Size and Limits to the Daily Range of Body Temperature
493 in Terrestrial Ectotherms. Am Nat 125:102-117.

494 Teuschl Y, Blanckenhorn WU (2007) The reluctant fly: What makes *Sepsis cynipsea*
495 females willing to copulate? Anim Behav 73:85-97.

496 Tammaru T, Esperk T (2007) Growth allometry of immature insects: larvae do not
497 grow exponentially. Funct Ecol 21:1099-1105.

498 van der Have TM, de Jong G (1996) Adult size in ectotherms: Temperature effects on
499 growth and differentiation. J Theoret Biol 18:329-340.

500 Van Voorhies WA (1996) Bergmann size clines: A simple explanation for their
501 occurrence in ectotherms. Evolution 50:1259-1264.

502 Zwaan BJ, Azevedo RBR, James AC, Van't Land J, Partridge L (2000) Cellular basis
503 of wing size variation in *Drosophila melanogaster*: a comparison of latitudinal
504 clines on two continents. Heredity 84:338-347.

Figure Captions

Fig. 1 Map indicating sampling sites of our 13 European *Sepsis fulgens* populations.

510 **Fig. 2** Population mean $\pm$ SE egg-to-adult development time (top) and body size
 511 (thorax length; bottom) for females (left) and males (right) of 11 latitudinal *Sepsis*
 512 *fulgens* populations at four constant laboratory temperatures (12°C, 18°C, 24°C,
 513 30°C).

514

515

516 **Fig. 3** Line mean number of eggs laid (i.e. first clutch size; top) and size of eggs
 517 (bottom) for 8 latitudinal populations of lab-reared *Sepsis fulgens* at 18°C or 30°C.

520 **Fig. 4** Line mean number of laid eggs (top) and size of eggs (bottom) of *Sepsis*
521 *fulgens* females from 8 latitudinal populations lab-reared at 18°C (left) or 30°C (right)
522 as a function of body size (fore tibia length).

525 **Table 1.** Comparison of clinal studies of Drosophilidae and Sepsidae (Diptera). All slopes refer to wing length (mm/deg. latitude; rescaled from
526 other structural traits for sepsids). Slopes that differ significantly from 0 are in bold.
527

Sepsidae	Sex*	Geographic range	Temp	Lat. range	Slope	SE	95% CI low	95% CI hi	no. Pops	Reference
<i>Sepsis cynipsea</i>	M & F	Europe	24	43 – 60	0.00008	0.00050	-0.00091	0.00106	7	Rohner et al. 2016
<i>Sepsis fulgens</i>	M & F	Europe	18	38 – 60	-0.00849	0.00502	-0.01833	0.00134	11	this study
<i>Sepsis neocynipsea</i>	M & F	North America	24	33 – 44	0.00324	0.00469	-0.00596	0.01243	6	Rohner et al. 2016
<i>Sepsis punctum</i>	M & F	Europe	18	43 – 60	0.01471	0.00493	0.00506	0.02437	7	Berger et al. unpublished
	M & F	North America	18	34 – 46	0.00144	0.00629	-0.01088	0.01376	6	Berger et al. unpublished
<i>Sepsis thoracica</i>	M & F	Europe	18	38 – 60	0.00239	0.01787	-0.03264	0.03741	15	Busso & Blanckenhorn 201
Drosophilidae										
<i>Drosophila kikkawai</i>	F	Indian subcontinent	24	7 – 31	0.01868	0.00156	0.01562	0.02174	9	Karan et al. 1998
<i>Drosophila melanogaster</i>	F	Eurasia	25	38 – 58	0.00210	–	–	–	16	Imasheva et al. 1994
<i>Drosophila serrata</i>	F	East Coast Australia	25	14 – 34	0.11400	0.00760	0.09910	0.12890	19	Hallas et al. 2002
	M		25		0.06900	0.01550	0.03862	0.09938		
<i>Drosophila simulans</i>	F	Eastern Australia	25	25 – 42	0.00337	0.00100	0.00141	0.00533	10	Arthur et al. 2008
	M		25		0.00370	0.00100	0.00174	0.00566		
<i>Drosophila subobscura</i>	F	Europe	20	37 – 56	0.00200	0.00060	0.00082	0.00318	11	Calboli et al. 2003
	M		20		0.00250	0.00060	0.00132	0.00368		
	F	North America	20	35 – 50	0.00190	0.00070	0.00053	0.00327	11	
	M		20		0.00040	0.00060	-0.00078	0.00158		
	F	South America	20	29 – 45	0.00210	0.00070	0.00073	0.00347	10	
	M		20		0.00150	0.00070	0.00013	0.00287		
<i>Zaprionus indianus</i>	F	Indian subcontinent	24	9 – 31	0.01190	0.00230	0.00739	0.01641	10	Karan et al. 2000
	M		24		0.01120	0.00420	0.00297	0.01943		

528 * M & F: average slope for both sexes

529 **Table 2.** Characteristics and climate conditions at the sampling locations of our 13 populations.
530

Population	Collection (YYYY.M)	No. lines	Latitude	Longitude	Altitude (m)	Temperature* (°C)		Mean annual range (bio7)	Mean of warmest quarter (bio10)	Annual precipitation (mm)
						Winter length ^{\$} (d)	Mean annual (bio1)			
Lamezia (I)	2013.4	13	38.92	16.25	244	15	17.6	22.0	24.8	864
Padula (I)	2013.4	13	40.22	15.65	720	128	13.2	22.6	20.6	764
Terni (I)	2012.9	7	42.57	12.62	132	119	14.6	28.1	22.7	853
Lake Trasimeno (I)	2012.9	5	43.13	12.1	258	155	13.3	27.3	21.4	792
Petroia (I)	2014.8	6	43.25	12.55	513	157	12.2	26.0	20.5	936
Maggia (CH)	2013.5	4	46.25	8.7	372	172	10.8	26.6	19.0	1240
Möschberg (CH)	2013.4	11	46.91	7.63	518	200	7.5	25.2	15.4	1067
Zürich (CH)	2013.6	8	47.34	8.55	489	189	9.5	25.8	17.6	1077
Witterschlick (D)	2014.7	6	50.69	7.02	96	186	9.6	23.6	16.9	773
Bielefeld (D)	2013.6	12	52.03	8.53	105	196	8.8	23.1	16.2	833
Skelde (DK)	2013.8	5	54.85	9.73	11	215	8.3	21.5	15.7	750
Rahinge (Est)	2014.6	7	58.37	26.7	72	238	4.9	31.3	15.9	600
Pehka (Est)	2013.5	12	59.48	26.37	57	240	4.7	31.1	15.7	618

531
532 ^{\$} Mean number of days below 10°C (assumed developmental threshold)

533 * Bioclimatic variables 1, 7, 10 from: <http://worldclim.org>; Hijmans et al. (2005)

534

535 **Supplementary Table S1:** Sex- and temperature-specific latitudinal slope estimates for development time, thorax length, clutch size and egg
536 volume.
537

	Temp	Dev. time			Thorax length			Clutch size			Egg volume		
		slope	SE	<i>r</i>	slope	SE	<i>r</i>	slope	SE	<i>r</i>	slope	SE	<i>r</i>
Females	12	0.039	0.042	0.109	-0.002	0.001	-0.155						
	18	-0.001	0.024	-0.005	-0.002	0.001	-0.153	0.572	0.607	0.149	0	0	-0.024
	24	0.013	0.016	0.093	-0.002	0.001	-0.216						
	30	-0.023	0.011	-0.241	0	0.001	-0.036	0.21	0.614	0.055	0	0	-0.067
Males	12	0.035	0.043	0.095	-0.001	0.001	-0.081						
	18	0.002	0.024	0.013	-0.001	0.001	-0.172						
	24	0.011	0.017	0.071	-0.002	0.001	-0.261						
	30	-0.023	0.011	-0.243	-0.001	0.001	-0.082						

538