



Year: 2018

Serotypes and virulence profiles of Shiga toxin-producing *Escherichia coli* strains isolated during 2017 from human infections in Switzerland

Nüesch-Inderbinen, Magdalena ; Morach, Marina ; Cernela, Nicole ; Althaus, Denise ; Jost, Marianne ; Mäusezahl, Mirjam ; Bloomberg, Guido ; Stephan, R

Abstract: Since 2015, the Swiss Federal Office of Public Health registered an increase of notifications of STEC, probably due to the adoption of culture independent stx screening tests in diagnostic laboratories. This study aimed to identify the serotypes and virulence genes of 120 STEC isolated from human clinical stx positive specimens during 2017 in order to estimate any changes in serotype distribution and toxin profiles of STEC compared to the time span 2010-2014. Culturing of STEC from stool samples was achieved using the streak plate technique on MacConkey agar. We performed O and H serotyping by PCR and by micro array. Virulence genes were identified and subtyped using molecular methods, including stx1 and stx2 subtypes, and the intimin encoding gene, eae. STEC were recovered from 27.5% of the stx positive samples. STEC O157:H7 accounted for 7.5% of all isolates, and STEC O80:H2, O91:H10/H14/H21, O103:H2/H11, and O26:H11 accounted for 36.9% of the non-O157 strains. Forty-five isolates with stx1 variants, 47 with stx2 variants and 28 isolates with both stx1 and stx2 variants were identified. Forty (33.3% of all isolates) carried the subtypes associated with high pathogenic potential, stx2a, stx2c, or stx2d. The eae gene for intimin was detected in 54 strains (45% of all strains). Compared to 2010-2014, our data show that the proportion of the so called "top five" serogroups, STEC O26, O111, O103, and O157 declined from 53.7% to 28.3% in 2017. The proportion of isolates with stx2a, stx2c, or stx2d decreased from 50.5% to 33.3%. We also observed an increase of STEC harbouring the low pathogenic subtypes stx2b and stx2e from 12.6% to 29.2%, and of eae negative STEC from 29.5% in 2010-2014 to 55% in 2017. Simultaneously, there was a sharp increase of the patients' median age from 24 years to 46.5 years. Clinical manifestations in the patients included abdominal pain without diarrhea (22.3%), diarrhea (77.7%), and the haemolytic-uremic syndrome (HUS) (7.4%). Our data show that a greater number and a wider range of STEC serotypes are detected by culture-independent testing, with implications for public health services.

DOI: <https://doi.org/10.1016/j.ijmm.2018.06.011>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-168119>

Journal Article

Accepted Version

Originally published at:

Nüesch-Inderbinen, Magdalena; Morach, Marina; Cernela, Nicole; Althaus, Denise; Jost, Marianne; Mäusezahl, Mirjam; Bloomberg, Guido; Stephan, R (2018). Serotypes and virulence profiles of Shiga toxin-producing *Escherichia coli* strains isolated during 2017 from human infections in Switzerland. *International Journal of Medical Microbiology* : IJMM, 308(7):933-939.

DOI: <https://doi.org/10.1016/j.ijmm.2018.06.011>

1 **Serotypes and virulence profiles of Shiga toxin-producing *Escherichia coli* strains**
2 **isolated during 2017 from human infections in Switzerland**

3

4 Magdalena Nüesch-Inderbinnen¹, Marina Morach¹, Nicole Cernela¹, Denise Althaus¹,
5 Marianne Jost², Mirjam Mäusezahl², Guido Bloomberg¹, Roger Stephan¹

6

7

8 ¹Swiss National Centre for Enteropathogenic Bacteria and *Listeria* (NENT), Institute for
9 Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Switzerland

10

11 ²Swiss Federal Office of Public Health, Division Communicable Diseases

12

13

14

15 *Corresponding author:

16 Roger Stephan, Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of
17 Zurich, Winterthurerstrasse 272, CH-8057 Zurich, Switzerland.

18 Phone +41 44 635 86 51, Fax +41 44 635 89 08, e-mail stephanr@fsafety.uzh.ch

19

20 Abstract

21 Since 2015, the Swiss Federal Office of Public Health registered an increase of notifications
22 of STEC, probably due to the adoption of culture independent *stx* screening tests in
23 diagnostic laboratories. This study aimed to identify the serotypes and virulence genes of 120
24 STEC isolated from human clinical *stx* positive specimens during 2017 in order to estimate
25 any changes in serotype distribution and toxin profiles of STEC compared to the time span
26 2010–2014. Culturing of STEC from stool samples was achieved using the streak plate
27 technique on MacConkey agar. We performed O and H serotyping by PCR and by micro
28 array. Virulence genes were identified and subtyped using molecular methods, including *stx1*
29 and *stx2* subtypes, and the intimin encoding gene, *eae*. STEC were recovered from 27.5% of
30 the *stx* positive samples. STEC O157:H7 accounted for 7.5% of all isolates, and STEC
31 O80:H2, O91:H10/H14/H21, O103:H2/H11, and O26:H11 accounted for 36.9% of the non-
32 O157 strains. Forty-five isolates with *stx1* variants, 47 with *stx2* variants and 28 isolates with
33 both *stx1* and *stx2* variants were identified. Forty (33.3% of all isolates) carried the subtypes
34 associated with high pathogenic potential, *stx2a*, *stx2c*, or *stx2d*. The *eae* gene for intimin
35 was detected in 54 strains (45% of all strains). Compared to 2010–2014, our data show that
36 the proportion of the so called "top five" serogroups, STEC O26, O111, O103, and O157
37 declined from 53.7% to 28.3% in 2017. The proportion of isolates with *stx2a*, *stx2c*, or *stx2d*
38 decreased from 50.5% to 33.3%. We also observed an increase of STEC harbouring the low
39 pathogenic subtypes *stx2b* and *stx2e* from 12.6% to 29.2%, and of *eae* negative STEC from
40 29.5% in 2010–2014 to 55% in 2017. Simultaneously, there was a sharp increase of the
41 patients' median age from 24 years to 46.5 years. Clinical manifestations in the patients
42 included abdominal pain without diarrhea (22.3%), diarrhea (77.7%), and the haemolytic-
43 uremic syndrome (HUS) (7.4%). Our data show that a greater number and a wider range of

44 STEC serotypes are detected by culture-independent testing, with implications for public
45 health services.

46

47

48 **Keywords**

49 STEC, human, multiplex PCR, culture, serotypes

50 **1. Introduction**

51 Shiga toxin (Stx)-producing *Escherichia coli* (STEC) are etiological agents of outbreaks and
52 of sporadic cases of human gastrointestinal illnesses which may include non-bloody or
53 bloody diarrhea, haemorrhagic colitis (HC), and the haemolytic uremic syndrome (HUS)
54 (Karch et al., 2005). STEC are characterized by the production of one or more Stx, which
55 consist of two groups designated Stx1 (consisting of the three variants Stx1a, Stx1c and
56 Stx1d) and Stx2 (composed of seven distinct variants Stx2a, Stx2b, Stx2c, Stx2d, Stx2e,
57 Stx2f, and Stx2g). STEC associated with severe disease tend to feature variants Stx2a, Stx2c
58 and Stx2d, whereas STEC producing Stx2b and Stx2e are linked to mild clinical symptoms or
59 asymptomatic fecal carriage (Stephan and Hoelzle, 2000; Friedrich et al., 2002; Fuller et al.,
60 2011). Virulence may further be increased by the presence of intimin, the product of the *eae*
61 gene, which mediates attaching and effacing lesions on gastrointestinal epithelial cells (Kaper
62 et al., 2004). STEC belonging to the serogroups O157, O26, O103, O111, and O145
63 constitute the so called “top five” serogroups of human pathogenic STEC in the EU and
64 Switzerland, and are considered, together with a few others such as O91 and O113, important
65 serogroups in public health (EFSA, 2017).

66 Human infection with STEC is a notifiable disease in Switzerland. Notification of confirmed
67 cases to the federal office for public health (FOPH) is based on the isolation of STEC from
68 faeces, or on the detection of *stx1* and/or *stx2* in faeces or from a clinical isolate of *E. coli*.

69 The number of notifications of STEC in Switzerland has been increasing since 2015, possibly
70 due to the introduction of PCR based, increasingly sensitive *stx* screening tests in laboratory
71 testing practices. However, such culture-independent testing (CIDT) generally does not yield
72 an isolate, and positive results are not always culture confirmed. From the public health point
73 of view, the advantages of rapid and broad range pathogen detection are therefore challenged
74 by possible loss of strain subtyping with consequent disruption of monitoring trends in

75 serotype or Stx distribution (Cronquist et al., 2012). Recognition of the trends in serotypes
76 and toxin profiles of STEC is however of great importance in order to estimate their potential
77 for causing disease and to anticipate epidemiological changes. The aim of this study was to
78 gain epidemiological and serotyping information on STEC isolated during 2017. Therefore,
79 from May to December 2017, the FOPH requested all diagnostic laboratories to forward
80 clinical materials that tested positive for *stx* by CIDT to the Swiss National Reference Centre
81 for Enteropathogenic Bacteria and *Listeria* (NENT) for culture and further strain
82 characterization. The isolates were analysed with regard to their serotypes, *stx* subtypes and
83 presence of the *eae* gene. The results were compared with earlier data from Switzerland
84 investigated over the 5-year period 2010-2014 (Fierz et al., 2017).

85

86 **2. Material and Methods**

87 **2.1. Sample collection**

88 Human stool samples that tested positive for *stx* using multiplex molecular methods were
89 submitted to the NENT from May to December 2017 from clinical diagnostic laboratories
90 distributed nationwide.

91 From a total of 457 submitted specimens, 436 were included for analysis after the elimination
92 of repeat specimens (i.e., specimens obtained from the same patient). Of the 436 patients,
93 data of age and gender were known for 431 individuals. Thereof, 240 (55.7%) were from
94 female and 191 (44.3 %) were from male patients. The median age was 46.5 years (range 0–
95 99 years). Forty-six (10.7%) were isolated from patients ≤ 5 years of age.

96

97

98 **2.2. Strain isolation**

99 Specimens were cultured on MacConkey agar using the streak plate technique. From each
100 plate, six individual colonies, if possible of different morphology, were picked und
101 subcultured on sheep blood agar (Difco™ Columbia Blood Agar Base EH; Becton Dickinson
102 AG, Allschwil, Switzerland). Isolates that were confirmed to possess *stx* (*stx1*
103 *and/or stx2*) by real-time PCR (LightCycler R 2.0 Instrument, Roche Diagnostics
104 Corporation, Indianapolis, IN, USA) (EURL, 2013a) were selected for further analysis. From
105 plates yielding more than one *stx* positive colony, one isolate was randomly chosen for
106 subsequent characterization. Proportions of STEC in stool samples were defined as the
107 numbers of *stx* positive colonies among six *E. coli* colonies.

108

109 **2.3. Serotyping**

110 Strains were examined by PCR for the presence of genes associated with 14 selected
111 serogroups including the top-five serogroups, namely O26, O45, O55, O80, O91, O103,
112 O104, O111, O113, O121, O128, O145, O146, and O157 (Perelle et al., 2004; EURL, 2013a;
113 EURL, 2014; Soysal et al., 2016). Strains were tested for the presence of flagellar genes
114 related to H2, H4, H7, H8, H10, H11, H19, H21, H25, and H28 (Mora et al., 2012; EURL,
115 2013b; Beutin et al., 2015; Alonso et al., 2017). Strains belonging to other O groups and H
116 types were serotyped using the Alere™ *E. coli* SeroGeno typing AS-1 kit (Alere
117 Technologies, Jena, Germany).

118

119 **2.4. Virulence markers**

120 The identification of *stx1* subtypes (*stx1a*, *stx1c*, *stx1d*) and *stx2* subtypes (*stx2a*, *stx2b*, *stx2c*,
121 *stx2d*, *stx2e*, *stx2g*) was carried out by conventional PCR amplification (Scheutz et al., 2012).
122 Screening of the strains for *eae* was performed by real-time PCR according to the guidelines
123 of the European Union Reference laboratory (EURL, 2013a).

124

125 **3. Results**

126 **3.1. Recovery rate**

127 Out of a total of 436 human fecal specimens that tested positive by multiplex PCR for the
128 presence of *stx*, 120 samples yielded an STEC isolate for further characterisation, amounting
129 to a recovery rate of 27.5%.

130 Categorising the 120 samples into those with high numbers of *stx* positive colonies (five or
131 six positive colonies per sample), those with intermediate numbers (three or four positive
132 colonies) and those with low numbers (one or two colonies), resulted in 46 (38.3%) stool
133 samples with a high proportion of STEC colonies, 35 (29.2%) with an intermediate
134 proportion, and 39 (32.5%) stool samples with a low proportion of STEC, respectively
135 Strains exhibiting high or low colony numbers are listed in Table 1.

136

137 **3.2. Serological diversity**

138 Twenty-five different O-serogroups were identified among the 120 STEC isolates, in addition
139 to 17 O-non-typeable (Ont) serogroups, and 4 ambiguous results. Eighteen different H-types
140 were determined, including two non-typeable H-types and two ambiguous results. An
141 overview of the serotypes is given in Table 1.

142 Among the 120 isolates, 9 (7.5%) were O157:H7, and 111 (92.5%) were non-O157 STEC
143 strains. Together with STEC O157, the top five serogroups were represented by STEC O103
144 (n=11), O145 (n=6), and O26 (n=8), amounting to 28.3% of the isolates. No STEC O111
145 were detected. Isolates belonging to O80:H2 (n=11), and O91:H10/H14/H21 (n=11)
146 accounted each for 9.2% of the isolates, respectively. Other serotypes included
147 O174:H2/H8/H21 (n=7; 5.8%), and O146:H21 (n=6; 5%) (Table 1). Other serotypes were
148 represented by four or less STEC isolates (Table 1).

149

150 3.3. Distribution of virulence genes among the serotypes

151 Of the 120 STEC strains, 45 (37.5%) carried *stx1* genes only: *stx1a* (n = 35), *stx1c* (n = 9)
152 and *stx1d* (n=1). Forty-seven strains (39.1%) carried *stx2* genes only: *stx2a* (n = 14), *stx2b* (n
153 = 14), *stx2c* (n = 5), *stx2d* (n = 12), and *stx2e* (n = 2). Twenty-eight (23.3%) harboured
154 combinations of *stx1* and *stx2* genes. Forty (33.3% of all isolates) carried the subtypes
155 associated with high pathogenic potential, *stx2a*, *stx2c*, or *stx2d* (Table 1). The
156 majority thereof (n = 25/62.5% of the 40 strains) were associated with O80:H2 (n=11),
157 O157:H7 (n = 9) and O145:H28 (n = 5).

158 Thirty-five (29.2%) isolates harboured the low pathogenic subtypes *stx2b* and *stx2e* and were
159 mainly associated to the serogroup O146 and Ont serogroups.

160 The *eae* gene encoding intimin was detected in 54 strains (45% of all strains). Thereof, 31
161 (57.4% of the *eae* positive strains) were associated with *stx2a*, *stx2c*, or *stx2d*, and one (4.3%)
162 with *stx2b*. The remaining 22 (40.7%) of the *eae* positive isolates carried *stx1a* alone (Table
163 1).

164 The majority (25 strains, 80.6%) of the *eae* positive strains harbouring *stx2* subtypes
165 belonged to STEC O80:H2, O145:H28, and O157:H7. By contrast, of the 66 (55% of all
166 strains) that tested negative for *eae*, nine (13.6% of all *eae* negative strains) harboured *stx2a*,
167 *stx2c*, or *stx2d*, while 34 (51.5%) were associated with *stx2b* or *stx2e*. The remaining 23
168 (34.8%) carried *stx1a* *stx1c* or *stx1d* alone (Table 1).

169 The distribution of serogroups and genotypes compared to earlier data from 2010-2014 (Fierz
170 et al., 2017) is illustrated in Figure 1.

171

172 **3.4. Relationship between STEC type, age of patients, clinical symptoms, and**
173 **proportions of *stx* positive *E. coli* in stool samples**

174
175 Patients were classified into four groups, according to their age at the time of sampling. Age
176 group 1 consisted of infants and children ≤ 5 years of age (n=14), group 2 contained children
177 and young adults between 6 and 17 years (n=16). Group 3 consisted of adult patients between
178 18 and 60 years (n=56), and group 4 of patients >60 years (n=33). For one patient, the age
179 was unknown (Table 1). Clinical data were provided for 94 (78.3%) of the patients.

180 Abdominal pain without diarrhea (AP) was reported for 21 (22.3%) of the patients. The
181 majority (77.7%) suffered from diarrhea (D). HUS was present in 7 (7.4%) of the patients, six
182 thereof with D, and one with AP. Two further patients (2.1%) had presumptive HUS with
183 acute kidney failure (AKF). Nineteen (20.2%) patients were hospitalized (two patients with
184 AP only and 17 patients with D).

185 The distribution of STEC serotypes and of *stx2* and *eae* genes among the patients' age groups
186 is listed in Table 1. STEC belonging to the top five serogroups were found in 71.2% of the
187 STEC infected children from age group 1 and in 56.3% from age group 2. Infections due to
188 the top five STEC serotypes were less frequent among age group 3 (16.1% of the patients),
189 and age group 4 (18.2% of the patients). By contrast, of the two most prevalent serogroups
190 from this study, STEC O80 was isolated more frequently from patients in age group 4 (15.2%
191 of the patients) than from patients in groups 1 (7.1%), group 2 (6.3%) and group 3 (7.1%),
192 respectively. STEC O91 was detected only among isolates from patients of age groups 3 and
193 4, accounting for 16.1% and 6% of the infections, respectively. STEC harbouring
194 *stx2a/stx2d/stx2c* were frequent among patients from age groups 1, 2 and 4 (50%, 43.8%, and
195 42.4% respectively), and least frequent among patients from age group 3 (19.6%). STEC
196 containing *stx2b/stx2e* was more frequent among patients from age groups 3 and 4 (33.9%
197 and 33.3%, respectively), compared to patients from age groups 1 and 2 (7.1% and 25%,

198 respectively). Similarly, *eae* positive STEC were observed at higher rates among patients
199 from groups 1 and 2 (85.7% and 62.5%, respectively) than among those from groups 3 and 4
200 (30.4% and 42.2%, respectively). By contrast, *eae* negative STEC were remarkably less
201 frequent in isolates from patients of age groups 1 and 2 (14.3% and 37.5%, respectively) than
202 in those from group 3 and 4 (69.6% and 57.6%, respectively). The distribution of STEC
203 serogroups and of *stx2* and *eae* genes among the patients' age groups is illustrated in Figure
204 2A and B, respectively.

205 STEC serotypes and virulence genes associated with patients with abdominal pain only,
206 diarrhea, HUS, and with patients that were hospitalised are listed in Table 1. STEC belonging
207 to the top five serogroups were found in 14.3% of 21 patients with AP, in 30.1 % of 73
208 patients with D and in 42.9% of 7 patients with HUS (Table 1). The top five serogroups were
209 furthermore associated with 31.6% of 19 hospitalised patients. STEC O80 was isolated from
210 4.8% of patients with AP and 12.3% of patients with D. Moreover, STEC O80:H2 was
211 associated with one case of HUS, two cases of AKF and isolated from 21% of hospitalised
212 patients (Table 1). STEC O91 was detected at similar rates among patients with AP and D
213 (14.3% and 9.6%, respectively), in one HUS case and in 15.8% of hospitalised patients.

214 STEC harbouring *stx2a/stx2d/stx2c* were less frequent among patients with AP (14.3%), than
215 among isolates from patients with D (41%) and HUS patients (100%), and were found in
216 68.4% of hospitalised patients. Similarly, *eae* positive STEC were observed at a lower rate in
217 patients with AP (23.8%) than among patients with D (47.9%) or HUS (85.7%), and
218 hospitalised patients (63.2%) (Table 1). By contrast, STEC carrying *stx2b/stx2e* were more
219 frequent among patients with AP (57.2%) than among patients with D (21.9%), and absent
220 among HUS patients.

221 Finally, *eae* negative STEC were accountable for 76.2% of patients with AP only, and for
222 52.1% and 14.3% of patients with D and HUS, respectively. Furthermore, *eae* negative STEC
223 were recovered from 36.8% of hospitalised patients.

224 The distribution of STEC serogroups and of *stx2* variants and *eae* genes among patients with
225 AP, D, HUS, and hospitalized patients is illustrated in Figure 2C and D, respectively.

226 The proportions of STEC among *E. coli* isolated from the patients' stool samples varied
227 according to serotypes and virulence genes. The majority of the STEC O145, STEC O157,
228 and STEC O91 isolates (66.7%, 55.6%, and 54.5%, respectively) were found in high numbers
229 in the stool samples, whereas high proportions were less frequent for serogroups O26 (50%)
230 and O103 (36.4%), and remarkably less (28.1%), for isolates belonging to serogroups other
231 than the top five, O80 or O91 (Figure 2E). STEC isolates harbouring *stx2a/ stx2c/ stx2d* were
232 observed more frequently in higher proportions than those harbouring *stx2b/stx2e* (50% and
233 34.3%, respectively). Similarly, *eae* positive STEC were found more frequently in high
234 proportions among *E. coli* from patients' stool samples than *eae* negative STEC (44.4% and
235 33.3%, respectively, Figure 2F).

236 Among patients with AP, the proportion of STEC in stool was high for 33.3% and low for
237 47.6%. Among patients with D and HUS, proportions were high for 37% and 42.9%,
238 respectively, and low for 32.9% and 28.6% (Table 1).

239

240 **4. Discussion**

241

242 Since 2015, the Swiss FOPH has registered an increase of notifications of STEC related
243 infections compared to previous years, with the increasing use of *stx* screening tests driving
244 this trend (Hächler and Stephan, 2015). In order to estimate any changes in serotype

245 distribution and toxin profiles of STEC, this study aimed to identify the serotypes and
246 virulence genes of 120 STEC isolated from human clinical specimens during 2017.
247 The five most common serogroups were O157, O103, O26, O91 and O80, with *E. coli*
248 O157:H7 accounting for 7.5% of the STEC strains. By comparison, during 2000–2009,
249 30.6% of the STEC strains isolated from humans in Switzerland were *E. coli* O157:H7, and
250 during 2010–2014 this rate further decreased to 19% (Käppeli et al., 2011; Fierz et al., 2017).
251 Thus, as reported for other countries in the EU, the proportion of STEC O157 isolated from
252 humans continues to decrease in Switzerland (EFSA, 2017). Similarly, the proportion STEC
253 O145 which belongs to the top five serogroups in Europe (EFSA, 2017), decreased from
254 12.6% during 2010–2014 to 5.4% in 2017 (Fierz et al., 2017). By contrast, with regard to the
255 serogroups that do not belong to the top five, we observed an increase of STEC O80 from
256 6.3% to 9.2% and of STEC O91 from 3.2% to 9.2%, compared to 2010–2014 (Fierz et al.,
257 2017). In particular, the increase of STEC O80:H2 is noteworthy. This hypervirulent,
258 multidrug resistant serotype harbours genetic characteristics of a hybrid STEC/ extraintestinal
259 pathogenic *E. coli* (ExPEC) pathotype and has recently emerged in France and Switzerland
260 associated with severe disease including bacteremia and HUS (Soysal et al., 2016; Fierz et
261 al., 2017; Nüesch-Inderbinen et al., 2018).

262 Compared to the study period 2010-2014, we also observed a remarkable decrease in the
263 percentage of STEC harbouring *stx2* variants that are associated with severe disease (i.e.,
264 *stx2a*, *stx2c*, *stx2d*) from 50.5% to 33.3%, and of *eae* positive STEC from 70.5% to 45%.
265 Simultaneously, we noted a sharp increase from 12.6% to 29.2%, of STEC associated with
266 *stx2b* and *stx2e*, variants that are linked to mild clinical symptoms and asymptomatic carriage
267 (Stephan and Hoelzle, 2000; Fuller et al., 2011), and to *eae* negative STEC from 29.5% to
268 55%. Correlating to this development, the patients' age median increased from 24 years to

269 46.5 years, and the proportion of patients ≤ 5 years dropped from 43.2% in 2010–2014 to
270 10.7% in 2017.

271 The present study analysed STEC cultured stool samples that were positive for *stx* by PCR
272 screening. The relatively low success rate of 27% for isolation of STEC by culture compared
273 with other studies 72%-96.5% (Friedrich et al., 2002; Tunsjø et al., 2015) may be explained
274 by the loss of *stx* genes over time due to transport conditions or time delay to culture in the
275 reference laboratory. A further explanation is that the concentration of STEC in some of the
276 stool samples may have been too low to allow detection of isolates. Alternatively, the low
277 recovery may have been caused by the use of PCR to directly target *stx* genes in faeces. Free
278 *stx*-carrying bacteriophages have been detected at a prevalence of 62% in fecal samples of
279 healthy people and may thus be responsible for some of the *stx* positive stool samples
280 (Martinez-Castillo et al., 2013; Urdahl et al., 2013). Therefore, caution should be employed
281 when diagnosing disease on the basis of *stx* positive results obtained directly from faecal
282 samples.

283 A definite attribution of the STEC isolates from this study to disease was precluded by the
284 lack of sufficient information on co-detection of pathogens other than STEC by multiplex
285 PCR based CIDT. Overall, it cannot be excluded that a certain number of the isolated STEC
286 were not the actual etiological agent of disease. Indeed, ten (8.3%) of the STEC strains
287 identified in this study belong to serotypes detected previously in faecal samples of healthy
288 humans including O8:H25, O128:H10, O146:H21, O146:H28, and O174:H8 (Stephan and
289 Untermann, 1999; Stephan and Schumacher, 2001; Urdahl et al., 2013).

290 Our data are supportive of previous studies that established an association of *eae* negative
291 STEC with severe clinical symptoms including HUS (Beutin et al., 2004). However, further
292 studies are needed in order to correlate the presence of infrequently detected STEC serotypes

293 and *stx2b/stx2e* harbouring STEC with the severity of disease and, ultimately, to distinguish
294 public health relevant infections from the non-relevant.
295 Conclusively, in spite of the increased notification of STEC during 2017, there was no
296 evidence for an outbreak situation. Our data are suggestive of an ongoing trend towards a
297 wider spectrum of serologically different STEC types most likely due to the introduction of
298 PCR based, increasingly sensitive *stx* screening tests in laboratory testing practices. This
299 study emphasises the importance of combining molecular methods of detection with classical
300 culture techniques to enable the detection of emerging STEC serotypes or outbreak situations.
301 Moreover, recognition of the trends in serotypes and toxin profiles of STEC is of great
302 importance in order to estimate epidemiological changes.

303

304 **Funding**

305 This work was partly supported by the Swiss Federal Office of Public Health, Division
306 Communicable Diseases.

307

308 **Conflicts of interest**

309 None to declare.

310

311 **Acknowledgements**

312

313 The authors thank all contributing diagnostic laboratories.

314 **References**

- 315 Alonso, C. A., Mora, A., Díaz, D., Blanco, M., González-Barrio, D., Ruiz-Fons, F., Simón,
316 C., Blanco, J., Torres, C., 2017. Occurrence and characterization of *stx* and/or *eae*-
317 positive *Escherichia coli* isolated from wildlife, including a typical EPEC strain from a
318 wild boar. *Vet. Microbiol.* 207, 69-73. doi=10.1016/j.vetmic.2017.05.028.
- 319 Beutin, L., Krause, G., Zimmermann, S., Kaulfuss, S., Gleier, K., 2004. Characterization of
320 Shiga toxin-producing *Escherichia coli* strains isolated from human patients in Germany
321 over a 3-year period. *J. Clin. Microbiol.* 42, 1099-1108.
- 322 Beutin, L., Delannoy, S., Fach, P., 2015. Genetic Diversity of the *fliC* genes encoding the
323 flagellar antigen H19 of *Escherichia coli* and application to the specific identification of
324 enterohemorrhagic *E. coli* O121:H19. *Appl. Environ. Microbiol.* 81, 4224-4230.
325 DOI=10.1128/AEM.00591-15.
- 326 Cronquist, A. B., Mody, R. K., Atkinson, R., Besser, J., Tobin D'Angelo, M., Hurd, S.,
327 Robinson, T., Nicholson, C., Mahon, B. E., 2012. Impacts of culture-independent
328 diagnostic practices on public health surveillance for bacterial enteric pathogens. *Clin.*
329 *Infect. Dis.* 54 Suppl 5, S432-439. DOI=10.1093/cid/cis267.
- 330 EFSA (European Food Safety Authority) and ECDC (European Centre for Disease
331 Prevention and Control), 2017. The European Union summary report on trends and
332 sources of zoonoses, zoonotic agents and food-borne outbreaks in 2016. *EFSA J.*
333 15(12):5077, 228 pp. <https://doi.org/10.2903/j.efsa.2017.5077>.
- 334 European Union Reference Laboratory (EURL), 2013a. Identification and
335 characterization of verocytotoxin-producing *Escherichia coli* (VTEC) by real
336 time PCR amplification of the main virulence genes and the genes associated
337 with the serogroups mainly associated with severe human infections. EU-RL

338 VTEC_Method_02_Rev 0. Available online at:
339 http://old.iss.it/binary/vtec/cont/EU_RL_VTEC_Method_02_Rev_0.pdf
340 European Union Reference Laboratory (EURL), 2013b. Detection and
341 identification of verotoxin-producing *Escherichia coli* (VTEC) O104:H4 in
342 food by real time PCR. EU-RL VTEC_Method_04_Rev 1. Available
343 online at: http://old.iss.it/binary/vtec/cont/EU_RL_VTEC_Method_04_Rev_1.pdf.
344 European Union Reference Laboratory (EURL), 2014. Identification of the
345 VTEC serogroups mainly associated with human infections by conventional
346 PCR amplification of O-associated genes. EU-RL VTEC_Method_03_Rev 01.
347 Available online at:
348 http://old.iss.it/binary/vtec/cont/EU_RL_VTEC_Method_03_Rev_1.pdf.
349 Fierz, L., Cernela, N., Hauser, E., Nüesch-Inderbilen, M., Stephan, R. 2017. Characteristics
350 of Shigatoxin-producing *Escherichia coli* strains isolated during 2010-2014 from human
351 infections in Switzerland. *Front. Microbiol.* 8, 1471. DOI=10.3389/fmicb.2017.01471.
352 Friedrich, A. W., Bielaszewska, M., Zhang, W. L., Pulz, M., Kuczius, T., Ammon, A., Karch,
353 H. 2002. *Escherichia coli* harboring Shiga toxin 2 gene variants: frequency and
354 association with clinical symptoms. *J. Infect. Dis.* 185, 74-84. DOI=10.1086/338115.
355 Fuller, C. A., Pellino, C. A., Flagler, M. J., Strasser, J. E., Weiss, A. A., 2011. Shiga toxin
356 subtypes display dramatic differences in potency. *Infect. Immun.* 79, 3, 1329-1337.
357 doi=10.1128/IAI.01182-10.
358 Hächler, H., Stephan, R., 2015. Auffälliger Anstieg der Meldezahlen enterohämorrhagischer
359 *E. coli*-Infektionen über die letzten Monate in der Schweiz: Einfluss neuer Multiplex
360 PCR-Methoden in der Primär-Diagnostik? *Bull. Swiss FOPH*, 52, p987-989.
361 Kaper, J. B., Nataro, J. P., Mobley, H. L., 2004. Pathogenic *Escherichia coli*. *Nat. Rev.*
362 *Microbiol.* 2, 123-140. DOI=10.1038/nrmicro818.

363 Käppeli, U., Hächler, H., Giezendanner, N., Cheasty, T., Stephan, R., 2011. Shiga toxin-
364 producing *Escherichia coli* O157 associated with human infections in Switzerland, 2000-
365 2009. *Epidemiol. Infect.* 139, 1097-1104. DOI=10.1017/S0950268810002190.

366 Karch, H., Tarr, P. I., Bielaszewska, M., 2005. Enterohaemorrhagic *Escherichia coli* in
367 human medicine. *Int. J. Med. Microbiol.* 295, 405-418.
368 DOI=10.1016/j.ijmm.2005.06.009.

369 Martinez-Castillo, A., Quirós, P., Navarro, F., Miró, E., Muniesa, M. 2013. Shiga toxin 2-
370 encoding bacteriophages in human fecal samples from healthy individuals. *Appl.*
371 *Environ. Microbiol.* 79, 4862-4868. DOI=10.1128/AEM.01158-13.

372 Mora, A., López, C., Dhabi, G., López-Beceiro, A. M., Fidalgo, L. E., Díaz, E. A., Martínez-
373 Carrasco, C., Mamani, R., Herrera, A., Blanco, J. E., Blanco, M., Blanco, J., 2012.
374 Seropathotypes, phylogroups, Stx subtypes, and intimin types of wildlife-carried, Shiga
375 toxin-producing *Escherichia coli* strains with the same characteristics as human-
376 pathogenic isolates. *Appl. Environ. Microbiol.* 78, 2578-2585.
377 DOI=10.1128/AEM.07520-11.

378 Nüesch-Inderbinen, M., Cernela, N., Wüthrich, D., Egli, A., Stephan, R. 2018. Genetic
379 characterization of Shiga toxin producing *Escherichia coli* belonging to the emerging
380 hybrid pathotype O80:H2 isolated from humans 2010-2017. *Int. J. Med. Microbiol.* 308,
381 534-538. DOI: 10.1016/j.ijmm.2018.05.007

382 Perelle, S., Dilasser, F., Grout, J., Fach, P., 2004. Detection by 5'-nuclease PCR of Shiga-
383 toxin producing *Escherichia coli* O26, O55, O91, O103, O111, O113, O145 and
384 O157:H7, associated with the world's most frequent clinical cases. *Mol. Cell Probes.* 18,
385 185-192. DOI=10.1016/j.mcp.2003.12.004.

386 Scheutz, F., Teel, L. D., Beutin, L., Piérard, D., Buvens, G., Karch, H., Mellmann, A.,
387 Caprioli, A., Tozzoli, R., Morabito, S., Strockbine, N. A., Melton-Celsa, A. R., Sanchez,

388 M., Persson, S., O'Brien, A. D., 2012. Multicenter evaluation of a sequence-based
389 protocol for subtyping Shiga toxins and standardizing Stx nomenclature. J. Clin.
390 Microbiol. 50, 2951-2963. DOI=10.1128/JCM.00860-12.

391 Soysal, N., Mariani-Kurkdjian, P., Smail, Y., Liguori, S., Gouali, M., Loukiadis, E., Fach, P.,
392 Bruyand, M., Blanco, J., Bidet, P., Bonacorsi, S., 2016. Enterohemorrhagic *Escherichia*
393 *coli* hybrid pathotype O80:H2 as a new therapeutic challenge. Emerg. Infect. Dis. 22,
394 1604-1612. doi=10.3201/eid2209.160304

395 Stephan, R., Hoelzle, L. E., 2000. Characterization of Shiga toxin type 2 variant B-subunit in
396 *Escherichia coli* strains from asymptomatic human carriers by PCR-RFLP. Lett. Appl.
397 Microbiol. 31, 139-142.

398 Stephan, R., Schumacher, S. 2001. Resistance patterns of non-O157 Shiga toxin-producing
399 *Escherichia coli* (STEC) strains isolated from animals, food and asymptomatic human
400 carriers in Switzerland. Lett. Appl. Microbiol. 32, 114-117. DOI=10.1046/j.1472-
401 765x.2001.00867.x.

402 Stephan, R., Untermann, F., 1999. Virulence factors and phenotypical traits of verotoxin-
403 producing *Escherichia coli* strains isolated from asymptomatic human carriers. J. Clin.
404 Microbiol. 37, 1570-1572.

405 Tunsjø, H. S., Kvissel, A. K., Follin-Arbelet, B., Brotnov, B. M., Ranheim, T. E., Leegaard,
406 T. M. 2015. Suitability of *stx*-PCR directly from fecal samples in clinical diagnostics of
407 STEC. APMIS. 123, 872-878. DOI=10.1111/apm.12428.

408 Urdahl, A. M., Solheim, H. T., Vold, L., Hasseltvedt, V., Wasteson, Y. 2013. Shiga toxin-
409 encoding genes (*stx* genes) in human faecal samples. APMIS. 121, 202-210.
410 DOI=10.1111/j.1600-0463.2012.02957.x.

411
412
413

414 **Figure legends**

415 Figure 1: Comparative distribution of serogroups and genotypes of STEC isolated during
416 2010-2014 and in 2017 from humans.

417 A: Percentage of the “top five” STEC serogroups and other selected STEC serogroups
418 isolated during 2010-2014 and in 2017.

419 B: Percentage of STEC harbouring *stx1* and *stx2* subtypes, and percentage of *eae* positive and
420 *eae* negative STEC isolated during 2010-2014 and in 2017.

421 Figure 2: Relationship between STEC type, patients’ age, clinical symptoms, and proportions
422 of STEC types among *E. coli* in stool samples.

423 A: Percentage of patients by age group infected with isolates belonging to the top five STEC
424 serogroups (O26, O103, O111, O145, or O157), STEC O80, STEC O91, or other STEC
425 serogroups.

426 B: Percentage of patients by age group infected with STEC harbouring *stx2* virulence genes
427 associated with high pathogenicity (*stx2a/stx2c/stx2d*), *stx2* genes associated with milder
428 symptoms (*stx2b/stx2e*), *eae* positive (*eae* +) STEC, and *eae* negative (*eae* -) STEC.

429 C: Distribution of STEC serogroups isolated among patients with abdominal pain only (AP),
430 diarrhea (D) or the hemorrhagic uremic syndrome (HUS), and from hospitalised patients (H).

431 D: Distribution of STEC harbouring *stx2* virulence genes, and percent *eae* + and *eae* - STEC
432 among patients with AP, D or HUS and among hospitalised patients.

433 E: Percentage of selected STEC serogroups present in high or low proportions among *E. coli*
434 isolated from human stool samples. Stool samples with five or six *stx* positive colonies out of
435 six *E. coli* colonies were classified as high. Samples with one or two STEC per six *E. coli*
436 colonies were defined as low proportion.

437 F: Percentage of STEC harbouring *stx2* subtypes, and percentage of *eae* positive and *eae*
438 negative STEC present in high or low proportions among *E. coli* isolated from human stool
439 samples.

440

441

442

443