



Year: 2019

The Effect of Vitamin C on Clinical Outcome in Critically Ill Patients: A Systematic Review With Meta-Analysis of Randomized Controlled Trials

Putzu, Alessandro ; Daems, Anne-Marie ; Lopez-Delgado, Juan Carlos ; Giordano, Vito Federico ; Landoni, Giovanni

Abstract: **OBJECTIVES** The effects of vitamin C administration on clinical outcome in critically ill patients remain controversial. **DATA SOURCES** Online databases were searched up to October 1, 2018. **STUDY SELECTION** We included randomized controlled trials on the use of vitamin C (any regimen) in adult critically ill patients versus placebo or no therapy. **DATA EXTRACTION** Risk ratio for dichotomous outcome and standardized mean difference for continuous outcome with 95% CI were calculated using random-effects model meta-analysis. **DATA SYNTHESIS** Forty-four randomized studies, 16 performed in ICU setting (2,857 patients) and 28 in cardiac surgery (3,598 patients), published between 1995 and 2018, were included in the analysis. In ICU patients, vitamin C administration was not associated with a difference in mortality (risk ratio, 0.90; 95% CI, 0.74-1.10; $p = 0.31$), acute kidney injury, ICU or hospital length of stay compared with control. In cardiac surgery, vitamin C was associated to a reduction in postoperative atrial fibrillation (risk ratio, 0.64; 95% CI, 0.52-0.78; $p < 0.0001$), ICU stay (standardized mean difference, -0.28 d; 95% CI, -0.43 to -0.13 d; $p = 0.0003$), and hospital stay (standardized mean difference, -0.30 d; 95% CI, -0.49 to -0.10 d; $p = 0.002$). Furthermore, no differences in postoperative mortality, acute kidney injury, stroke, and ventricular arrhythmia were found. **CONCLUSIONS** In a mixed population of ICU patients, vitamin C administration is associated with no significant effect on survival, length of ICU or hospital stay. In cardiac surgery, beneficial effects on postoperative atrial fibrillation, ICU or hospital length of stay remain unclear. However, the quality and quantity of evidence is still insufficient to draw firm conclusions, not supporting neither discouraging the systematic administration of vitamin C in these populations. Vitamin C remains an attractive intervention for future investigations aimed to improve clinical outcome.

DOI: <https://doi.org/10.1097/CCM.0000000000003700>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-180606>

Journal Article

Published Version

Originally published at:

Putzu, Alessandro; Daems, Anne-Marie; Lopez-Delgado, Juan Carlos; Giordano, Vito Federico; Landoni, Giovanni (2019). The Effect of Vitamin C on Clinical Outcome in Critically Ill Patients: A Systematic Review With Meta-Analysis of Randomized Controlled Trials. *Critical Care Medicine*, 47(6):774-783.

DOI: <https://doi.org/10.1097/CCM.0000000000003700>

The Effect of Vitamin C on Clinical Outcome in Critically Ill Patients: A Systematic Review With Meta-Analysis of Randomized Controlled Trials

Alessandro Putzu, MD¹; Anne-Marie Daems, MD¹; Juan Carlos Lopez-Delgado, MD, PhD^{2,3}; Vito Federico Giordano, MD⁴; Giovanni Landoni, MD^{4,5}

Objectives: The effects of vitamin C administration on clinical outcome in critically ill patients remain controversial.

Data Sources: Online databases were searched up to October 1, 2018.

Study Selection: We included randomized controlled trials on the use of vitamin C (any regimen) in adult critically ill patients versus placebo or no therapy.

Data Extraction: Risk ratio for dichotomous outcome and standardized mean difference for continuous outcome with 95% CI were calculated using random-effects model meta-analysis.

Data Synthesis: Forty-four randomized studies, 16 performed in ICU setting (2,857 patients) and 28 in cardiac surgery (3,598 patients), published between 1995 and 2018, were included in the analysis. In ICU patients, vitamin C administration was not associated with a difference in mortality (risk ratio, 0.90; 95% CI, 0.74–1.10; $p = 0.31$), acute kidney injury, ICU or hospital length of stay compared with control. In cardiac surgery, vitamin C was associated to a reduction in postoperative atrial fibrillation (risk ratio, 0.64; 95% CI, 0.52–0.78; $p < 0.0001$), ICU stay (standardized mean difference, -0.28 d; 95% CI, -0.43 to -0.13 d; $p = 0.0003$), and hospital stay (standardized mean difference,

-0.30 d; 95% CI, -0.49 to -0.10 d; $p = 0.002$). Furthermore, no differences in postoperative mortality, acute kidney injury, stroke, and ventricular arrhythmia were found.

Conclusions: In a mixed population of ICU patients, vitamin C administration is associated with no significant effect on survival, length of ICU or hospital stay. In cardiac surgery, beneficial effects on postoperative atrial fibrillation, ICU or hospital length of stay remain unclear. However, the quality and quantity of evidence is still insufficient to draw firm conclusions, not supporting neither discouraging the systematic administration of vitamin C in these populations. Vitamin C remains an attractive intervention for future investigations aimed to improve clinical outcome. (*Crit Care Med* 2019; XX:00–00)

Key Words: ascorbic acid; cardiac surgery; critical care; intensive care; perioperative medicine; vitamin C

¹Division of Anesthesiology, Department of Anesthesiology, Pharmacology & Intensive Care Medicine, Geneva University Hospitals, Geneva, Switzerland.

²Intensive Care Department, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Barcelona, Spain.

³Institut d'Investigació Biomèdica Bellvitge (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain.

⁴Department of Anesthesia and Intensive Care, IRCCS San Raffaele Scientific Institute, Milan, Italy.

⁵Vita-Salute San Raffaele University, Milan, Italy.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website (<http://journals.lww.com/ccmjournal>).

The authors have disclosed that they do not have any potential conflicts of interest.

For information regarding this article, E-mail: landoni.giovanni@hsr.it

Copyright © 2019 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

DOI: 10.1097/CCM.0000000000003700

this micronutrient, and supra-physiologic doses of vitamin C could have further beneficial effects (7–9). Side effects are rare, but reported in the literature (1, 4, 10). Nevertheless, despite there is a rationale in favor of the use of vitamin C, the clinical effectiveness of vitamin C supplementation is still unclear. In this systematic review with meta-analysis, meta-regression, and trial sequential analysis (TSA), we aimed to assess the effect of vitamin C administration on major clinical outcome in ICU or cardiac surgery patients.

MATERIALS AND METHODS

The present systematic review was conducted in compliance with Cochrane methodology (11), Preferred Reporting Items Systematic Reviews and Meta-Analysis (PRISMA) guidelines (12), and according to a pre-published protocol on the international prospective register of systematic reviews (PROSPERO, CRD42017069790). A complete PRISMA checklist is provided in **eTable 1** (Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>) (13).

Search Strategy

Two investigators (A.P.,V.F.G.) independently searched PubMed, EMBASE, and the Cochrane Central Register of clinical trials (last updated October 1, 2018) for appropriate articles. The full PubMed search strategy is available in the supplement (**eMethods 1**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>) and it includes all the most common vitamin C definitions employed in the literature. Abstracts from recent international conferences were searched for additional relevant studies. In addition, we hand scanned references of retrieved articles and pertinent reviews of the literature. No language restriction was enforced.

Study Selection

References obtained from searches were first independently examined at an abstract level by two investigators (A.P., V.F.G.) and then, if potentially relevant, collected as complete articles. Eligible studies met the following PICOS criteria: 1) Population: adult ICU or cardiac surgery patients; 2) Intervention: any type of vitamin C formulation and regimen; 3) Comparison intervention: placebo or no therapy; 4) Outcome: mortality at longest follow-up available; and 5) Study design: randomized controlled trials (RCTs). There were no restrictions on vitamin C formulation and dose or time of administration. The exclusion criteria were overlapping populations and pediatric studies. Two investigators (A.P., A.M.D.) independently assessed selected studies for the final analysis, with eventual divergences finally resolved by consensus. If the article did not report data on primary or secondary outcomes, the corresponding author was contacted for further information.

Data Abstraction and Study Characteristics

Three authors (A.P., A.M.D., V.F.G.) extracted relevant information from each selected study and entered them into predefined databases; divergences were resolved by consensus with a fourth

author (G.L.). We collected potential sources of significant clinical heterogeneity, such as study design, sample size, clinical setting, inclusion and exclusion criteria, vitamin C regimen and association with other drugs, control treatment, as well as primary and secondary endpoints. We evaluated the possible authors' conflicts of interests and the funding of each study.

The primary endpoint of the present review was mortality at the longest follow-up available. Secondary outcomes included acute kidney injury (AKI), supraventricular arrhythmia, ventricular arrhythmia, stroke, length of ICU stay, and length of hospital stay. The outcomes were reported as per-trial definition. We applied the intention-to-treat methods whenever possible.

Internal Validity and Risk of Bias Assessment

The internal validity of each trial included was independently evaluated by two authors (A.P., A.M.D.) for bias according to The Cochrane Collaboration methods (11) (**eMethods 2**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>). Eventual divergences were resolved by consensus. We rated the potential risk of bias by applying a rating of “low,” “high,” or “unclear” to each study.

Data Analysis and Synthesis

For dichotomous variables, we calculated the odds ratio (OR) with a 95% CI and in case of common events, defined as the frequency of the event occurring in the control group more than or equal to 10%, we calculated the risk ratio (RR) with a 95% CI. For continuous outcome, we calculated standardized mean difference (sMD) with 95% CI. The *p* value for the comparison between the groups was calculated, and a *p* value of less than or equal to 0.05 was considered as statistically significant. Heterogeneity was assessed using the chi-square test and the *I*² statistic. We post hoc decided to include only ICU and cardiac surgery patients, presenting the results separately using a random-effects model meta-analysis. Publication bias was assessed by visually inspecting funnel plots if 10 or more trials are included in the meta-analysis (11). We performed various sensitivity, subgroup, and meta-regression analyses (**eMethods 3**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>). Meta-analysis was performed using Review Manager (RevMan, Version 5.3. Copenhagen, Denmark: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014).

TSA is a methodology that could quantify the statistical reliability of data in a cumulative meta-analysis adjusting significance levels for the risks of random errors due sparse data and repetitive testing on accumulating data. The TSA combines an information size calculation (equal to the cumulated sample sizes of all included trials), with a threshold for a statistically significant effect and a threshold for futility of the intervention, both adjusted to better control of type I and type II errors. Conclusions made using TSA show the potential to be more consistent than those using traditional meta-analysis methodology (14–16). Therefore, we performed a post hoc random-effects TSA for mortality with the intent of maintaining an overall 5% risk of type I error and a 10% risk of type II error, at a power of 90% (14–16). We assumed a relative risk

reduction of 15%, and we derived the control event proportion from the actual control dataset. The resulting required information size was further diversity (D2)-adjusted. We used the TSA software (TSA Viewer [Computer program], Version 0.9.5.5 Beta, Copenhagen Trial Unit, Centre for Clinical Intervention Research, Rigshospitalet, Copenhagen, Denmark, 2016).

RESULTS

The selection process is summarized in **Figure 1**. A total of 1,894 studies were identified and, after exclusion of ineligible studies, 44 studies ($n = 6,455$) were included in the meta-analysis (8–10, 17–61). Major exclusions are reported in **eTable 2** (Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

Thirty-four studies were single-center, one trial was two-centers, three trials were three-centers, and one trial was multicenter. In 10 cases, we received further information from the

authors (21, 23, 31, 38, 39, 50, 53, 60–62) (**eTable 3**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

Three trials were considered to carry a low risk of bias in all bias domains (20, 45, 58), five trials were at unclear risk of bias (9, 22–24, 40), and 36 trials were at high risk of bias (**eFigs. 1** and **2**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

ICU Patients

Sixteen studies (2,857 patients) administered vitamin C supplements in critically ill ICU patients: six in a general ICU population (2,425 patients), six in a septic population (260 patients), three in patients with severe acute pancreatitis (135 patients), and one in patients with major burns (37 patients) (**Table 1**; and **eTable 4**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

The administration of vitamin C in critically ill patients was associated with no difference in mortality at the longest follow-up available (400 of 1,453 patients [28%] in the vitamin C group vs 411 of 1,404

[29%] in the control group; RR, 0.90; 95% CI, 0.74–1.10; $p = 0.31$) (**Fig. 2**). Funnel plot was symmetric, not suggesting publication bias (**eFig. 3**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>). Similar results were found at subgroup analyses, in particular when assessing trials giving only vitamin C, IV/enteral vitamin C, and high-dose IV vitamin C (**Fig. 3**; and **eTable 5**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

TSA was inconclusive, since 19.40% of the information size was accrued and the cumulative curve did not reach the area of futility (**Fig. 4**), suggesting the need of further trials for a firm conclusion on the effects of vitamin C therapy.

Vitamin C administration was associated with no significant reduction in length of ICU stay in critically ill patients (**eFig. 4**, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

No significant difference in AKI between vitamin C and control was found (RR, 1.03; 95% CI, 0.84–1.25; $p = 0.78$; TSA inconclusive) and in




Figure 1. Study flow diagram. PRISMA = Preferred Reporting Items Systematic Reviews and Meta-Analysis.

TABLE 1. Characteristics of the ICU Trials Included in the Analysis

References	Journal	Country	Setting	No. of Patients	Experimental Intervention	Vitamin C Regimen	Control
ICU patients							
Bansal et al (19)	<i>Saudi J Gastroenterol</i>	India	Acute pancreatitis	39	Vitamin C, vitamin E, vitamin A	Vitamin C 1,000 mg IV qd for 14 d	No treatment
Berger et al (20)	<i>Crit Care</i>	Switzerland	Mixed ICU	200	Vitamin C, vitamin B1, selenium, zinc	Vitamin C 1,100 mg IV bid on first 2 d and qd until day 5	Placebo
Crimi et al (24)	<i>Anesth Analg</i>	Italy	Mixed ICU	216	Vitamin C, vitamin E	Vitamin C 500 mg E qd for 10 d	Placebo
Ferrón-Celma et al (30)	<i>J Surg Res</i>	Spain	Sepsis/septic shock	20	Vitamin C	Vitamin C 450 mg IV qd for 6 d	Placebo
Fowler et al (7)	<i>J Transl Med</i>	United States	Sepsis/septic shock	24	Vitamin C	Vitamin C 12.5 or 50 mg/kg IV qid for 4 d	Placebo
Galley et al (31)	<i>Free Radic Biol Med</i>	United Kingdom	Sepsis/septic shock	30	Vitamin C, vitamin E, NAC	Vitamin C 1,000 mg IV qd for 1 d	Placebo
Heyland et al (58)	<i>N Engl J Med</i>	United States, Canada, Europe	Mixed ICU	1,218	Vitamin C, vitamin E, zinc, beta-carotene, selenium, \pm glutamine	Vitamin C 1,500 mg E qd for 28 d or until ICU discharge	Placebo
Howe et al (9)	<i>Am J Crit Care</i>	United States	Mixed ICU	72	Vitamin C, vitamin E, \pm NAC	Vitamin C 1,000 mg E qd for 28 d or until ICU discharge	Placebo
Nabil Habib et al (54)	<i>Int J Microbiol Adv Immunol</i>	Egypt	Sepsis/septic shock	100	Vitamin C	Vitamin C 1,500 mg IV qid until ICU discharge	No treatment
Nathens et al (8)	<i>Ann Surg</i>	United States	Mixed ICU	595	Vitamin C, vitamin E	Vitamin C 1,000 mg IV three times a day until ICU discharge	No treatment
Palli et al (33)	<i>Crit Care</i>	Greece	Mixed ICU	124	Vitamin C, NAC	Vitamin C 2,000 mg IV 2 hr before and at 10 hr and 18 hr following the infusion of contrast agent	Placebo
Sateesh et al (44)	<i>Trop Gastroenterol</i>	India	Acute pancreatitis	53	Vitamin C, NAC, multivitaminic compound	Vitamin C 500 mg E qd for 5 d	No treatment
Schneider et al (55)	<i>Clin Nutr</i>	Germany	Sepsis/septic shock	58	Vitamin C, vitamin E, glutamine, beta-carotene, selenium, zinc, glycine, tributyrin	Vitamin C 1,500 mg E qd	Placebo
Siriwardena et al (45)	<i>Gut</i>	United Kingdom	Acute pancreatitis	43	Vitamin C, NAC, selenium	Vitamin C 2,000 mg IV over 24 hr for the first 2 d and 1,000 mg IV over 24 hr for the following 5 d	Placebo
Tanaka et al (47)	<i>Arch Surg</i>	Japan	Burn patients	37	Vitamin C	Vitamin C 66 mg/kg/hr IV for 24 hr	No treatment
Zabet et al (49)	<i>J Res Pharm Pract</i>	Iran	Sepsis/septic shock	28	Vitamin C	Vitamin C 25 mg/kg enteral qid for 3 d	Placebo

E = enteral, NAC = N-acetylcysteine, qd = once a day, qid = four times a day.




Figure 2. Mortality in ICU patients. Forest plot for mortality at longest follow-up available in patients randomized to vitamin C (any regimen) or control. *df* = degrees of freedom, M-H = Mantel-Haenszel.

other secondary outcomes (eFig. 5 and eTable 4, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

Cardiac Surgery

Twenty-eight trials administered vitamin C supplements in cardiac surgery (3,598 patients): 21 in coronary artery bypass grafting surgery (2,465 patients) and seven in a mixed cardiac surgery population (1,133 patients) (eTables 6 and 7, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

Mortality rate was low and similar between vitamin C and control group (15 of 968 patients [1.55%] in the vitamin C group vs 17 of 927 [1.83%] in the control group; RR, 1.00; 95% CI, 0.48–2.08; *p* = 1.00; TSA inconclusive)

(eFig. 6, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

Vitamin C was associated with lower supraventricular arrhythmias versus control in cardiac surgery patients (RR, 0.64; 95% CI, 0.52–0.78; *p* < 0.0001). Atrial fibrillation was the main reported supraventricular arrhythmia. The analysis was limited by the high statistical heterogeneity (*I*² = 57%; *p*_{heterogeneity} = 0.0005) and significant asymmetry of funnel plot, suggesting publication bias; most of the heterogeneity was explained by Europe/United States trials, which found no difference in postoperative supraventricular arrhythmias (RR, 0.92; 95% CI, 0.76–1.11; *p* = 0.38) in comparison to an important significant effect in other countries (RR, 0.49; 95% CI, 0.39–0.61; *p* < 0.00001)




Figure 3. Trial sequential analysis for mortality at longest follow-up available in a mixed ICU population. Trial sequential analysis of the included trials (black square fill icons) shows that the cumulative Z curve did not cross nor the traditional boundary nor the trial sequential monitoring boundary for futility and did not reach the required information size ($n = 14,725$), suggesting the need for more trials to establish firm conclusions about vitamin C effects in critically ill patients. X-axis: the number of patients randomized; y-axis: the cumulative z score; horizontal dotted lines: conventional boundaries (upper for benefit, z score = 1.96, lower for harm, z score = -1.96, two-sided, $p = 0.05$); oblique lines with diamond icons: trial sequential monitoring boundaries; oblique line with square fill icons: Z curve; and vertical line with diamonds: required information size.

($p_{\text{subgroup}} < 0.0001$) (eFigs. 7 and 8, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

Vitamin C administration was associated to a small reduction in postoperative length of ICU stay (sMD, -0.28 d; 95% CI, -0.43 to -0.13 d; $p = 0.0003$) and hospital stay (sMD, -0.30 d; 95% CI, -0.49 to -0.10 d; $p = 0.002$). The results were limited by high statistical heterogeneity ($I^2 > 62\%$; $p_{\text{heterogeneity}} < 0.001$) and possible publication bias. Subgroup analysis including only Europe/United States trials found no effects of vitamin C on length of ICU or hospital stay (eFigs. 9-11 and eTable 8, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

No significant differences in ventricular arrhythmia (OR, 0.79; 95% CI, 0.33-1.88; $p = 0.59$; TSA inconclusive), AKI (RR, 1.18; 95% CI, 0.74-1.89; $p = 0.48$; TSA inconclusive), and stroke (OR, 0.47; 95% CI, 0.13-1.69; $p = 0.25$; TSA inconclusive) were found (eFig. 6, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

Vitamin C Regimens

The results were consistent at subgroup, sensitivity, and meta-regression analyses. No effects of dose, length of therapy, route of administration, and combination with other antioxidants

were suggested on any outcome assessed in ICU patients. In cardiac surgery, meta-regression suggested possible beneficial effects of longer supplementation with low-dose vitamin C on supraventricular arrhythmia (eTables 5, 8 and 9, and eMethods 1, Supplemental Digital Content 1, <http://links.lww.com/CCM/E416>).

DISCUSSION

This meta-analysis of RCTs showed that vitamin C supplementation, alone or in association with other antioxidant drugs, is associated with no effects on survival and ICU or hospital stay in a mixed population of ICU patients. In cardiac surgery, the observed beneficial effects on postoperative atrial fibrillation and ICU/hospital length of stay remain unclear, due to the presence of publication bias and subgroup effects according to geographical area. The quality and quantity of randomized evidence is still insufficient to draw firm conclusions on clinical effects of vitamin C, not supporting neither discouraging the systematic administration in critically ill patients.

Cardiac surgery and ICU patients represent two different populations, often managed by same medical teams. Oxidative stress plays a central role in the pathophysiology of




Figure 4. Main subgroup analyses in a mixed ICU population. Forest plot for mortality at longest follow-up available in patients randomized to receive: only vitamin C, enteral vitamin C (any regimen), IV vitamin C (any regimen), or high-dose IV vitamin C (> 5,000 mg a day). *df* = degrees of freedom, M-H = Mantel-Haenszel.

inflammatory syndrome in both populations. A high proportion of surgical ICU patients suffer from vitamin C deficiency (60) and, in addition, the intense inflammatory response can increase metabolic turnover of vitamin C resulting in a severe deficiency (61). Analogously, the use of cardiopulmonary bypass in cardiac surgery can also induce oxidative stress leading to vitamin C depletion (62). An imbalance in the production of oxidizing species, such as reactive oxygen species, superoxide anions, and other radicals, leads to an increased oxidative stress,

which causes an increased endothelial permeability with an impairment of microcirculatory flow, an inadequate cell-mediated immunity, followed by an activation of apoptotic mitochondrial mechanisms that may contribute to cell death and organ failure (4, 5). Hypothetically, vitamin C can counterbalance oxidizing species, providing a preventing role in the oxidative stress development and restoring neutrophil function (4, 5). Some randomized trials included in the present meta-analysis have shown an improvement in the inflammatory profile of ICU patients

receiving vitamin C, confirming its potential immunomodulatory activity (7, 8, 24). Indeed, two of these studies showed also an improvement in organ dysfunction and failure (7, 8), as also suggested by a recent nonrandomized trial (63). Despite vitamin C might exert various positive effects, our analysis failed to find any benefit in terms of mortality.

Human cells have vitamin C transporters coupled with sodium transport and their activity is limited by blood level of vitamin C; kidneys excrete the excess of vitamin C and its blood levels remain strictly controlled (64). Thus, higher doses of vitamin C administered orally (> 400 mg) can saturate the intestinal absorption system, achieving plasma concentrations that may not be enough to scavenge an excess of oxidizing species (64, 65). In this scenario, IV infusion route should be the preferred in order to achieve higher plasma concentrations in order to scavenge oxidizing species, protecting from their undesirable effects (66). For this purpose, doses higher than the usually recommended daily intake could be recommended especially if we consider that, given the hydrophilic chemical structure of vitamin C, its plasma concentrations may be affected by the higher distribution volume due to the positive fluid balance that critically ill patients frequently suffer from (67). A dosage of 2–3 g per day might be needed to restore normal levels of vitamin C in critically ill patients (68). However, our study failed to find any subgroup effects in patients randomized to receive IV or high-dose ascorbic acid.

Higher levels of oxidizing species can cause tubular injury and glomerular edema, leading to AKI (69). Recent findings from a retrospective before-after study suggest that the early use of IV vitamin C, together with corticosteroids and thiamine, may be effective in preventing progressive organ dysfunction including AKI and reducing the mortality of septic shock (63). However, the external validity of this retrospective study is limited, mainly due to the nonrandomized, single-centered design, the small sample size, and the presence of baseline imbalances and other limitations (70). Our analysis did not show any effect in terms of renal failure, even if AKI can be a rare complication of vitamin C therapy and represents so an outcome of interest, it was underreported in the included trials. Results from cardiac surgery, another setting where AKI prevention is crucial, were inconclusive too.

In cardiac surgery, vitamin C administration has emerged as a strategy of postoperative atrial fibrillation prevention (6). The substantial difference in atrial fibrillation results from Europe/United States in comparison to other countries, together with substantial publication bias (6, 71), rise doubts on the external validity of the positive aggregate results of our meta-analysis on atrial fibrillation in cardiac surgery. Some epidemiologic evidence suggested possible larger treatment effects in less developed countries (72). Nevertheless, differences in population characteristics, lifestyle factors including nutrition, treatment response, and healthcare systems exist and, at least partially, could explain the large difference in effect estimate. The prevention of atrial fibrillation, a potential source of embolic stroke, together with the evidence that oxidizing species enhance secondary brain damage by means

of increased permeability of blood-brain barrier generating edema in the acute stage of stroke (73, 74), leave a role for vitamin C. Once again, our results do not firmly support this hypothesis.

This is the first comprehensive meta-analysis systematically assessing the effects of vitamin C supplementation on important clinical outcomes in critically ill patients. With 44 trials and more than 6,000 patients, this is the largest meta-analysis on this topic performed to date. We focused on some important clinical outcomes, such as mortality, and we performed a systematic review of several databases, aiming to reduce the possibility of missing minor publications. Traditional limitations of aggregate patient data meta-analyses are present, including different patients baseline characteristics, clinical subsettings, and different vitamin C regimens. In particular, differences in routes of administration (enteral vs IV), dosage (low- vs high-dose), total duration of supplementation, and combined treatment with other antioxidants drugs, increase the inconsistency of the aggregate analysis. However, we performed several subgroup analyses to explore the possible effects of different vitamin C regimens, with inconclusive results. Most of the eligible trials had an high risk of bias and publication bias is probable (6). In ICU trials, mortality and length of ICU or hospital stay were the most reported outcomes and several other outcomes of interests, such as organ failure or quality-of-life endpoints, were modestly and heterogeneously reported. On the contrary, atrial fibrillation was the most reported outcome in cardiac surgery while mortality was low, making the aggregate analysis underpowered for this latter outcome.

Adverse effects following vitamin C administration are limited (1, 4) and are mainly related to oxalate kidney stones formation and oxalate nephropathy (10). However, most of the included trials did not systematically assess adverse effects, and a systematic evaluation before extensive implementation of vitamin C therapy is warranted.

These issues, together with inconclusive results from TSA, suggest that the actual randomized evidence on vitamin C supplementation is inconsistent and the current use of vitamin C as adjunctive therapy cannot be recommended or discouraged neither in a mixed population of ICU patients nor in cardiac surgery.

CONCLUSIONS

Vitamin C administration is not associated with a reduction in mortality in a mixed ICU population. In cardiac surgery, the observed beneficial effects on postoperative atrial fibrillation and ICU/hospital length of stay remain unclear, due to the presence of publication bias and subgroup effects according to geographical area. The quality and quantity of evidence is still insufficient to draw firm conclusions on the possible effect of vitamin C on clinical outcome and aggregate randomized evidence does not support neither discourage the systematic administration of vitamin C in ICU or cardiac surgery. Vitamin C remains an attractive intervention for future investigations aimed at reducing mortality and morbidity.

REFERENCES

- Amrein K, Oudemans-van Straaten HM, Berger MM: Vitamin therapy in critically ill patients: Focus on thiamine, vitamin C, and vitamin D. *Intensive Care Med* 2018; 44:1940–1944
- Koekkoek WA, van Zanten AR: Antioxidant vitamins and trace elements in critical illness. *Nutr Clin Pract* 2016; 31:457–474
- Berger MM, Oudemans-van Straaten HM: Vitamin C supplementation in the critically ill patient. *Curr Opin Clin Nutr Metab Care* 2015; 18:193–201
- Spoelstra-de Man AME, Elbers PWG, Oudemans-van Straaten HM: Making sense of early high-dose intravenous vitamin C in ischemia/reperfusion injury. *Crit Care* 2018; 22:70
- Marik PE: Vitamin C for the treatment of sepsis: The scientific rationale. *Pharmacol Ther* 2018; 189:63–70
- Hemilä H, Suonsyrjä T: Vitamin C for preventing atrial fibrillation in high risk patients: A systematic review and meta-analysis. *BMC Cardiovasc Disord* 2017; 17:49
- Fowler AA 3rd, Syed AA, Knowlson S, et al; Medical Respiratory Intensive Care Unit Nursing: Phase I safety trial of intravenous ascorbic acid in patients with severe sepsis. *J Transl Med* 2014; 12:32
- Nathens AB, Neff MJ, Jurkovich GJ, et al: Randomized, prospective trial of antioxidant supplementation in critically ill surgical patients. *Ann Surg* 2002; 236:814–822
- Howe KP, Clochesy JM, Goldstein LS, et al: Mechanical ventilation antioxidant trial. *Am J Crit Care* 2015; 24:440–445
- Colliou E, Mari A, Delas A, et al: Oxalate nephropathy following vitamin C intake within intensive care unit. *Clin Nephrol* 2017; 88:354–358
- Higgins JP, Green S (Eds): *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from <http://handbook.cochrane.org>
- Liberati A, Altman DG, Tetzlaff J, et al: The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: Explanation and elaboration. *BMJ* 2009; 339:b2700
- Moher D, Liberati A, Tetzlaff J, et al; PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: The PRISMA Statement. *PLoS Med* 2009; 6:e1000097
- Brok J, Thorlund K, Wetterslev J, et al: Apparently conclusive meta-analyses may be inconclusive—trial sequential analysis adjustment of random error risk due to repetitive testing of accumulating data in apparently conclusive neonatal meta-analyses. *Int J Epidemiol* 2009; 38:287–298
- Thorlund K, Devereaux PJ, Wetterslev J, et al: Can trial sequential monitoring boundaries reduce spurious inferences from meta-analyses? *Int J Epidemiol* 2009; 38:276–286
- Thorlund K, Engström J, Wetterslev J, et al: User manual for trial sequential analysis (TSA). Copenhagen Trial Unit, Centre for Clinical Intervention Research, Copenhagen, Denmark. User Manual Seq Anal (TSA) Copenhagen Trial Unit, Cent Clin Interv Res Copenhagen, Denmark 2011:1–115. Available at: www.ctu.dk/tsa. Accessed May 15, 2018
- Bjordahl PM, Helmer SD, Gosnell DJ, et al: Perioperative supplementation with ascorbic acid does not prevent atrial fibrillation in coronary artery bypass graft patients. *Am J Surg* 2012; 204:862–867; discussion 867
- Bailey DM, Raman S, McEneny J, et al: Vitamin C prophylaxis promotes oxidative lipid damage during surgical ischemia-reperfusion. *Free Radic Biol Med* 2006; 40:591–600
- Bansal D, Bhalla A, Bhasin DK, et al: Safety and efficacy of vitamin-based antioxidant therapy in patients with severe acute pancreatitis: A randomized controlled trial. *Saudi J Gastroenterol* 2011; 17:174–179
- Berger MM, Soguel L, Shenkin A, et al: Influence of early antioxidant supplements on clinical evolution and organ function in critically ill cardiac surgery, major trauma, and subarachnoid hemorrhage patients. *Crit Care* 2008; 12:R101
- Braga M, Bissolati M, Rocchetti S, et al: Oral preoperative antioxidants in pancreatic surgery: A double-blind, randomized, clinical trial. *Nutrition* 2012; 28:160–164
- Castillo R, Rodrigo R, Perez F, et al: Antioxidant therapy reduces oxidative and inflammatory tissue damage in patients subjected to cardiac surgery with extracorporeal circulation. *Basic Clin Pharmacol Toxicol* 2011; 108:256–262
- Colby JA, Chen WT, Baker WL, et al: Effect of ascorbic acid on inflammatory markers after cardiothoracic surgery. *Am J Health Syst Pharm* 2011; 68:1632–1639
- Crimi E, Liguori A, Condorelli M, et al: The beneficial effects of antioxidant supplementation in enteral feeding in critically ill patients: A prospective, randomized, double-blind, placebo-controlled trial. *Anesth Analg* 2004; 99:857–863, table of contents
- Das D, Sen C, Goswami A: Effect of Vitamin C on adrenal suppression by etomidate induction in patients undergoing cardiac surgery: A randomized controlled trial. *Ann Card Anaesth* 2016; 19:410–417
- Dehghani MR, Majidi N, Rahmani A, et al: Effect of oral vitamin C on atrial fibrillation development after isolated coronary artery bypass grafting surgery: A prospective randomized clinical trial. *Cardiol J* 2014; 21:492–499
- Healy RM, Day D, Van Gorder C, et al: Ascorbic acid utilization for atrial-fibrillation prophylaxis post coronary-artery-bypass graft and valve replacement surgeries: An interim analysis of a prospective, randomized study [abstract manuscript]. American College of Clinical Pharmacy (ACCP) 2010 Annual Meeting, Austin, TX, October 17–20, 2010
- Duffy MJ, O’Kane CM, Stevenson M, et al: A randomized clinical trial of ascorbic acid in open abdominal aortic aneurysm repair. *Intensive Care Med Exp* 2015; 3:50
- Eslami M, Badkoubeh RS, Mousavi M, et al: Oral ascorbic acid in combination with beta-blockers is more effective than beta-blockers alone in the prevention of atrial fibrillation after coronary artery bypass grafting. *Tex Heart Inst J* 2007; 34:268–274
- Ferrón-Celma I, Mansilla A, Hassan L, et al: Effect of vitamin C administration on neutrophil apoptosis in septic patients after abdominal surgery. *J Surg Res* 2009; 153:224–230
- Galley HF, Howdle PD, Walker BE, et al: The effects of intravenous antioxidants in patients with septic shock. *Free Radic Biol Med* 1997; 23:768–774
- Akbarzadeh M, Eftekhari MH, Shafa M, et al: Effects of a new metabolic conditioning supplement on perioperative metabolic stress and clinical outcomes: A randomized, placebo-controlled trial. *Iran Red Crescent Med J* 2016; 18:e26207
- Palli E, Makris D, Papanikolaou J, et al: The impact of N-acetylcysteine and ascorbic acid in contrast-induced nephropathy in critical care patients: An open-label randomized controlled study. *Crit Care* 2017; 21:269
- Papoulidis P, Ananiadou O, Chalvatzoulis E, et al: The role of ascorbic acid in the prevention of atrial fibrillation after elective on-pump myocardial revascularization surgery: A single-center experience—a pilot study. *Interact Cardiovasc Thorac Surg* 2011; 12:121–124
- Petersen F, Rodrigo R, Richter M, et al: The effects of polyunsaturated fatty acids and antioxidant vitamins on atrial oxidative stress, nitrotyrosine residues, and connexins following extracorporeal circulation in patients undergoing cardiac surgery. *Mol Cell Biochem* 2017; 433:27–40
- Polymeropoulos E, Bagos P, Papadimitriou M, et al: Vitamin C for the prevention of postoperative atrial fibrillation after cardiac surgery: A meta-analysis. *Adv Pharm Bull* 2016; 6:243–250
- Pourmatroud E, Hormozi L, Hemadi M, et al: Intravenous ascorbic acid (vitamin C) administration in myomectomy: A prospective, randomized, clinical trial. *Arch Gynecol Obstet* 2012; 285:111–115
- Rebrova Tlu, Shipulin VM, Afanas’ev SA, et al: The experience of the application of ascorbic acid as antioxidant after coronary artery surgery with use of cardiopulmonary bypass. *Kardiologija* 2012; 52:73–76
- Rodrigo R, Gutiérrez R, Fernández R, et al: Ageing improves the antioxidant response against postoperative atrial fibrillation: A randomized controlled trial. *Interact Cardiovasc Thorac Surg* 2012; 15:209–214
- Rodrigo R, Korantzopoulos P, Cereceda M, et al: A randomized controlled trial to prevent post-operative atrial fibrillation by antioxidant reinforcement. *J Am Coll Cardiol* 2013; 62:1457–1465
- Sadeghpour A, Alizadehasl A, Kyavar M, et al: Impact of vitamin C supplementation on post-cardiac surgery ICU and hospital length of stay. *Anesth Pain Med* 2015; 5:e25337
- Safaei N, Babaei H, Azarfarin R, et al: Comparative effect of grape seed extract (*Vitis vinifera*) and ascorbic acid in oxidative stress in-

- duced by on-pump coronary artery bypass surgery. *Ann Card Anaesth* 2017; 20:45–51
43. Samadikhah J, Golzari SE, Sabermarouf B, et al: Efficacy of combination therapy of statin and vitamin C in comparison with statin in the prevention of post-CABG atrial fibrillation. *Adv Pharm Bull* 2014; 4:97–100
 44. Sateesh J, Bhardwaj P, Singh N, et al: Effect of antioxidant therapy on hospital stay and complications in patients with early acute pancreatitis: A randomised controlled trial. *Trop Gastroenterol* 2009; 30:201–206
 45. Siriwardena AK, Mason JM, Balachandra S, et al: Randomised, double blind, placebo controlled trial of intravenous antioxidant (n-acetylcysteine, selenium, vitamin C) therapy in severe acute pancreatitis. *Gut* 2007; 56:1439–1444
 46. Sisto T, Paajanen H, Metsä-Ketelä T, et al: Pretreatment with antioxidants and allopurinol diminishes cardiac onset events in coronary artery bypass grafting. *Ann Thorac Surg* 1995; 59:1519–1523
 47. Tanaka H, Matsuda T, Miyagantani Y, et al: Reduction of resuscitation fluid volumes in severely burned patients using ascorbic acid administration: A randomized, prospective study. *Arch Surg* 2000; 135:326–331
 48. Van Wagoner DR, Palumbo R, Li J, et al: Supplemental vitamin C did not reduce the incidence of atrial arrhythmia following cardiac bypass surgery [abstract manuscript]. Heart Rhythm Society; 2003. Available at: <http://www.mv.helsinki.fi/home/hemila/CAF/vanWagoner2003.pdf>. Accessed May 12, 2018
 49. Zabet MH, Mohammadi M, Ramezani M, et al: Effect of high-dose ascorbic acid on vasopressor's requirement in septic shock. *J Res Pharm Pract* 2016; 5:94–100
 50. Sarzaeem M, Shayan N: Effect of vitamin C in prevention of atrial fibrillation after coronary artery bypass graft: A double-blind randomized clinical trial. *Tehran Univ Med J* 2014; 71:787–793
 51. Antonic M: Effect of ascorbic acid on postoperative acute kidney injury in coronary artery bypass graft patients: A pilot study. *Heart Surg Forum* 2017; 20:E214–E218
 52. Obied HY: 2. Role of ascorbic acid & statin in reduction of the incidence of the atrial fibrillation in patients under B-blocker and undergoing coronary artery bypass graft operation in early post-operative period. *J Saudi Hear Assoc* 2013; 27:299–300
 53. Jouybar R, Kabgani H, Kamalipour H, et al: The perioperative effect of ascorbic acid on inflammatory response in coronary artery bypass graft surgery; a randomized controlled trial coronary artery bypass graft surgery. *Iran Cardiovasc Res J* 2012; 6:13–17
 54. Nabil Habib T, Ahmed I: Early adjuvant intravenous vitamin C treatment in septic shock may resolve the vasopressor dependence. *Int J Microbiol Adv Immunol* 2017; 5:77–81
 55. Schneider A, Markowski A, Momma M, et al: Tolerability and efficacy of a low-volume enteral supplement containing key nutrients in the critically ill. *Clin Nutr* 2011; 30:599–603
 56. Stanger O, Aigner I, Schimetta W, et al: Antioxidant supplementation attenuates oxidative stress in patients undergoing coronary artery bypass graft surgery. *Tohoku J Exp Med* 2014; 232:145–154
 57. Moludi J, Keshavarz S, Pakzad R, et al: Effect of vitamin C supplementation in the prevention of atrial fibrillation. *Tehran Univ Med J* 2016; 73:791–797
 58. Heyland D, Muscedere J, Wischmeyer PE, et al; Canadian Critical Care Trials Group: A randomized trial of glutamine and antioxidants in critically ill patients. *N Engl J Med* 2013; 368:1489–1497
 59. Heyland DK, Tougas G, King D, et al: Impaired gastric emptying in mechanically ventilated, critically ill patients. *Intensive Care Med* 1996; 22:1339–1344
 60. Luo M, Fernandez-Estivariz C, Jones DP, et al: Depletion of plasma antioxidants in surgical intensive care unit patients requiring parenteral feeding: Effects of parenteral nutrition with or without alanyl-glutamine dipeptide supplementation. *Nutrition* 2008; 24:37–44
 61. Doll S, Ricou B: Severe vitamin C deficiency in a critically ill adult: A case report. *Eur J Clin Nutr* 2013; 67:881–882
 62. Rodemeister S, Duquesne M, Adolph M, et al: Massive and long-lasting decrease in vitamin C plasma levels as a consequence of extracorporeal circulation. *Nutrition* 2014; 30:673–678
 63. Marik PE, Khangoora V, Rivera R, et al: Hydrocortisone, vitamin C, and thiamine for the treatment of severe sepsis and septic shock: A retrospective before-after study. *Chest* 2017; 151:1229–1238
 64. Rodrigo R, Guichard C, Charles R: Clinical pharmacology and therapeutic use of antioxidant vitamins. *Fundam Clin Pharmacol* 2007; 21:111–127
 65. Graumlich JF, Ludden TM, Conry-Cantilena C, et al: Pharmacokinetic model of ascorbic acid in healthy male volunteers during depletion and repletion. *Pharm Res* 1997; 14:1133–1139
 66. Jackson TS, Xu A, Vita JA, et al: Ascorbate prevents the interaction of superoxide and nitric oxide only at very high physiological concentrations. *Circ Res* 1998; 83:916–922
 67. Blanchard J, Tozer TN, Rowland M: Pharmacokinetic perspectives on megadoses of ascorbic acid. *Am J Clin Nutr* 1997; 66:1165–1171
 68. Long CL, Maull KI, Krishnan RS, et al: Ascorbic acid dynamics in the seriously ill and injured. *J Surg Res* 2003; 109:144–148
 69. Dennis JM, Witting PK: Protective role for antioxidants in acute kidney disease. *Nutrients* 2017; 9:718
 70. Kalil AC, Johnson DW, Cawcutt KA: Vitamin C is not ready for prime time in sepsis but a solution is close. *Chest* 2017; 152:676
 71. Hemilä H: Publication bias in meta-analysis of ascorbic acid for postoperative atrial fibrillation. *Am J Health Syst Pharm* 2017; 74:372–373
 72. Panagiotou OA, Contopoulos-Ioannidis DG, Ioannidis JP: Comparative effect sizes in randomised trials from less developed and more developed countries: Meta-epidemiological assessment. *BMJ* 2013; 346:f707
 73. Chen GC, Lu DB, Pang Z, et al: Vitamin C intake, circulating vitamin C and risk of stroke: A meta-analysis of prospective studies. *J Am Heart Assoc* 2013; 2:e000329
 74. Jin R, Liu L, Zhang S, et al: Role of inflammation and its mediators in acute ischemic stroke. *J Cardiovasc Transl Res* 2013; 6:834–851