



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Habitat selection of an old-growth forest specialist in managed forests

Ettwein, Antonia ; Pasinelli, Gilberto ; Korner, Pius ; Lanz, Michael

DOI: <https://doi.org/10.17011/conference/eccb2018/107273>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-182725>

Conference or Workshop Item

Published Version

Originally published at:

Ettwein, Antonia; Pasinelli, Gilberto; Korner, Pius; Lanz, Michael (2018). Habitat selection of an old-growth forest specialist in managed forests. In: 5th European Congress of Conservation Biology, Jyväskylä, Finland, 12 June 2018 - 15 June 2018, University of Jyväskylä.

DOI: <https://doi.org/10.17011/conference/eccb2018/107273>

1 **Habitat selection of an old-growth forest specialist in managed forests**

2 Antonia Ettwein: Swiss Ornithological Institute, Sempach

3 Pius Korner: Swiss Ornithological Institute, Sempach

4 Michael Lanz: Swiss Ornithological Institute, Sempach

5 Thibault Lachat: School of Agricultural, Forest and Food Sciences HAFL, Zollikofen & Swiss Federal Institute for Forest,
6 Snow and Landscape Research WSL, Birmensdorf

7 Hanna Kokko: Department of Evolutionary Biology and Environmental Studies, University of Zurich

8 Gilberto Pasinelli: Swiss Ornithological Institute, Sempach & Department of Evolutionary Biology and Environmental
9 Studies, University of Zurich

10

11

12 Corresponding author: Antonia Ettwein: Seerose 1, 6204 Sempach, Switzerland, antonia.ettwein@vogelwarte.ch

13

14

15 Short title: Habitat selection of the white-backed woodpecker

16

1 **Abstract**

2 Old-growth forest specialists are among the species most affected by commercial forestry. However, it is often
3 unclear whether such species can persist and what their habitat needs are in managed forests. We investigated
4 habitat selection of one such old-growth forest specialist, the white-backed woodpecker (*Dendrocopos leucotos*), a
5 species highly dependent on dead wood and typically found in primeval forests. Our aim was to understand factors
6 affecting occupancy probability in managed forests in Central Europe, based on detection/non-detection data in 62
7 squares of 1 km² in 2015 and 2016. We used occupancy models to compare *a priori* expectations about the
8 relationships between occupancy and habitat characteristics at two spatial scales while accounting for imperfect
9 detection. Occupancy was best explained by a proxy for food availability at a large (1 km²) scale and increased with
10 the abundance of emergence holes produced by saproxylic beetles on standing and lying dead wood. Furthermore,
11 occupancy was positively related to the mean DBH (diameter at breast height) of live trees and standing dead wood
12 at a small scale (0.25 km² with high amounts of dead wood). Detection probability was negatively related to time of
13 day, date, and number of accessible survey points, and positively related to the number of observers. Our results
14 demonstrate that detailed knowledge about a species' foraging ecology is important for its effective conservation as
15 surrogate criteria such as dead wood availability might not reflect the key factors required. For white-backed
16 woodpeckers, it is important that the available dead wood is sufficiently colonized by saproxylic beetles, and for the
17 conservation of the species, the habitat requirements of saproxylic beetles thus have to be taken into account as
18 well.

19
20 **Keywords:** occupancy modeling, habitat selection, white-backed woodpecker, *Dendrocopos leucotos*, habitat
21 specialist, forest management

22

1 Introduction

2 A long history of human land-use has altered forests in many parts of the world. In Central Europe, most forests are
3 characterized by homogeneous tree species composition, vertical stratification, and age structure, as well as by low
4 amounts of habitat trees (live or dead trees providing ecological niches, Bütler et al., 2013) and dead wood
5 (Kuuluvainen et al., 1996; Commarmot et al., 2005; Merino et al., 2007; Müller, Hothorn, & Pretzsch, 2007; Brumelis
6 et al., 2011) As a consequence, wildlife communities have also changed – often at the expense of specialist species
7 which are sensitive to changes in their environment because of their narrow ecological niche and other
8 characteristics, such as low population size, population growth rate, competitive ability, or dispersal ability (Grove,
9 2002; Henle et al., 2004; Devictor, Julliard, & Jiguet, 2008; Clavel, Julliard, & Devictor, 2010). Old-growth forest
10 specialists are among the species most affected by commercial forestry (Bengtsson et al., 2000; Fraixedas, Lindén, &
11 Lehtikoinen, 2015) in which short rotation periods usually inhibit the occurrence of late successional stages. The
12 decline of many old-growth species (Fraixedas, Lindén, & Lehtikoinen, 2015) highlights the need to increase our
13 understanding of these species' biological requirements to develop effective measures for their conservation. This
14 particularly applies if old-growth species occur in managed forests. While knowledge of the factors affecting habitat
15 selection is important for the conservation of any species, it is crucial for the conservation of specialized species in
16 managed habitats, as such habitats are prone to becoming unsuitable for the specialists if key resources are
17 removed.

18 We examined habitat selection of the white-backed woodpecker (*Dendrocopos leucotos leucotos*) in managed
19 forests in the Eastern Alps (western Austria, eastern Switzerland, Liechtenstein) during the mating and breeding
20 season (end of February to mid-May). The white-backed woodpecker is a resident bird species found in old-growth
21 and mature deciduous and mixed forests (Aulén, 1988; Hogstad & Stenberg, 1994). As it is considered a species with
22 high demands in terms of its ecological requirements (Scherzinger, 1982), the white-backed woodpecker has been
23 proposed as an umbrella species for communities associated with old deciduous-dominated forests rich in dead
24 wood (Roberge, Mikusiński, & Svensson, 2008) and an indicator of forest biodiversity (Mikusiński, Gromadzki, &
25 Chylarecki, 2001). Its preference for forests rich in dead wood is due to its foraging ecology: the white-backed

1 woodpecker is a food specialist feeding mainly on the larvae of saproxylic beetles (beetles that are dependent on
2 dead or dying wood, wood inhabiting fungi, or the presence of other saproxylics (Speight, 1989)), where it prefers
3 large larvae such as those of *Cerambycidae* (Aulén, 1988; Glutz von Blotzheim & Bauer, 1994). Due to intensive forest
4 management, many European populations have massively declined (Virkkala et al., 1993; Carlson, 2000; Czeszczewik
5 & Walankiewicz, 2006). As a consequence, the species is considered threatened in the red lists of several European
6 countries and is listed in the Annex I of the European Union's Birds Directive.

7 Contrary to the generally rather negative population trends in Europe, the species has colonized western Austria,
8 eastern Switzerland, and Liechtenstein since the 1970s (Mollet, Zbinden, & Schmid, 2009). Forests in western
9 Austria, eastern Switzerland, and Liechtenstein are mostly managed, while most previous studies on habitat
10 selection or habitat use of the white-backed woodpecker have been conducted in regions containing large areas of
11 forest not commercially used or even relicts of primeval forest (Scherzinger, 1982; Hogstad & Stenberg, 1994; Frank,
12 2002; Czeszczewik, 2009). The expansion of the species' range to regions dominated by managed forests raises the
13 question as to which factors enable the occurrence of an old-growth habitat specialist in this type of landscape. We
14 built a set of candidate models, each of which representing a hypothesis to examine whether forest structure
15 (hypothesis (1) "Old mixed and deciduous forests"), volume or diameter of dead wood (hypotheses (2) "Dead wood
16 volume" and (3) "Large pieces of dead wood"), nest site availability (hypothesis (4) "Abundance of suitable nesting
17 trees"), saproxylic beetle abundance (hypothesis (5) "Food availability"), or topography (hypotheses (6-8) "South-
18 exposed slopes in intermediate elevation") best explain white-backed woodpecker occupancy (see Table 1 for
19 candidate models and *a priori* expectations). Each candidate model was evaluated at two spatial scales (1 km² and
20 0.25 km²). The key aim was to understand the habitat needs of this old-growth forest specialist to develop measures
21 for the conservation of the species in managed forests.

22 **Methods**

23 **Study area**

24 The study presented here was conducted in the Eastern Alps in western Austria (province Vorarlberg), eastern
25 Switzerland (cantons Grisons and St. Gallen), and the Principality of Liechtenstein. The study region is characterized

1 by human settlements and agricultural land in the valley bottoms and less intensive land use in the mountainous
2 area, which is mostly covered with forest up to the timber line at approximately 1600 – 1800 m. The dominating tree
3 species are Norway spruce (*Picea abies*), silver fir (*Abies alba*), and European beech (*Fagus sylvatica*). The natural
4 vegetation are beech-dominated forests in the submontane belt, mixed forests in the montane belt, and pure
5 coniferous forests in the subalpine belt (Ellenberg & Strutt, 2009). Forests easily accessible from the valley bottoms
6 are typically intensively managed (except for protection forests), whereas forests at remote or inaccessible locations
7 in the mountains tend to be used less intensively or sometimes not at all for timber production.

8 Within the study region, we selected 62 squares of 1 km² (Fig. 1) in which we recorded detection/non-detection of
9 white-backed woodpeckers, habitat structure, and a measure of saproxylic beetle abundance. The white-backed
10 woodpecker occurs at low density in the study area, thus to avoid having a vast majority of squares not occupied,
11 half of the 62 squares included locations at which the species had been observed since 1975 (based on data from the
12 Swiss Ornithological Institute, inatura Dornbirn, and BirdLife Vorarlberg). The rest of the squares were placed at a
13 distance of two to five kilometers to the next square with a previous observation, but without any previously known
14 white-backed woodpecker observations in the square itself. This distance was chosen because it was large enough to
15 avoid attracting individuals from nearby squares to an unoccupied square during the white-backed woodpecker
16 surveys on the one hand; on the other hand, the distance was small enough to ensure that all squares could have
17 potentially been reached by white-backed woodpeckers (i.e., the unoccupied squares were not isolated from
18 occupied ones). We selected only squares without previous observations which could potentially contain suitable
19 white-backed woodpecker habitat: squares without previous observations had a forest cover at least as high as
20 squares with observations ($\geq 45\%$) and were situated at the same elevation (635 – 1520 m above sea level), resulting
21 in a mean (means are reported with \pm SD throughout the whole text) forest cover of $75.0 \pm 15.3 \%$ for squares with
22 and $75.1 \pm 17.3 \%$ for squares without observations, and a mean elevation of 958 ± 242 m and 1074 ± 239 m,
23 respectively. It is important to note that the non-random selection of squares on the one hand led to a higher
24 occupancy probability in our results than generally found in the study area, and probably weakened the effect of
25 forest area and elevation in our models on the other hand. Our approach was, however, necessary to obtain enough

1 white-backed woodpecker observations to be able to model occupancy in the first place, and to proceed beyond
2 results which are already well-known (e.g., that the species occupies forested areas).

3 **White-backed woodpecker surveys**

4 Data on the detection/non-detection history of each square were collected with up to two replicate surveys in 2015
5 and 2016 (i.e., with a total of four surveys per square for most squares). White-backed woodpeckers were searched
6 for at four predefined survey points per square (Fig. 1). Each survey point was situated at a distance of 300 m from
7 the edges of the square to ensure that only white-backed woodpeckers within the square were recorded. At each
8 survey point, one or two field workers used playbacks of white-backed woodpeckers to stimulate responses of
9 individuals. We used a series of drumming (available in Schulze et al., 2003), two different calls (the first call can be
10 found in Schulze et al., 2003, the second one in Stübing & Bergmann, 2006), and the imitation of foraging pecking
11 sounds by knocking on a tree with a stick in a manner that mimics (to the human ear at least) the typical behavior of
12 woodpeckers. The sounds were played back/produced for a minute each in a random order. After one minute of
13 playback/knocking, we waited for a reaction for three minutes. If no woodpecker responded, the next one minute
14 series was played back and responses observed during the following three minutes, etc. If no woodpecker was
15 detected after a total of 16 minutes of playback, knocking, and waiting, the field workers proceeded to the next
16 point. On the other hand, as soon as a white-backed woodpecker reacted, the playback was stopped to minimize
17 disturbance, and the species was considered to be present at that point. The white-backed woodpeckers usually
18 responded by approaching the playback tape while calling, or by drumming at a nearby tree. In some cases, the
19 woodpeckers did not approach the playback but were drumming in some distance. The species can be distinguished
20 from other woodpecker species by its drumming as well as its calls, and both visual and acoustic detections were
21 recorded.

22 When the species had been detected at one of the first points, the other survey points within a square were also
23 checked for white-backed woodpeckers in 2015, but not in 2016, because preliminary analyses of the 2015 data
24 suggested that the playback influenced the location of response by the woodpeckers within a square (Ettwein, 2016).

1 Consequently, the species was considered detected in a square if a white-backed woodpecker responded at one of
2 the four survey points per square.

3 In 2015, all squares were surveyed twice between the beginning of March and mid-May. In 2016, not all of the 62
4 squares could be surveyed twice due to high amounts of snow in April, leading to time constraints in May: 48
5 squares were surveyed twice, 11 randomly selected squares once, and three squares were not surveyed at all.

6 **Habitat measurements**

7 Habitat structure was mapped in the summer of 2015 in two circular sample plots of 500 m² (radius of 12.62 m) per
8 survey point (i.e., eight sample plots per square). The centers of these plots were located 50 m to the west and to
9 the east of the survey point (Fig. 1). We mapped live trees, snags (standing dead wood with a height > 1.30 m),
10 stumps (standing dead wood with a height ≤ 1.30 m), and lying dead wood (logs, pieces of branches). We measured
11 diameter at breast height (DBH) of all live trees with a DBH ≥ 12 cm, recorded tree species and whether dead
12 branches with a diameter ≥ 10 cm were present or not. Snags and stumps were mapped when their diameter
13 exceeded 5 cm, and height and diameter (DBH for snags, diameter in the middle for stumps) were recorded. We
14 used the formula for a cylinder to calculate the volume of snags and stumps. Lying dead wood was recorded along
15 three transects with a method used in the Swiss National Forest Inventory when the diameter at the intersection
16 with the transect was ≥ 7 cm (for details see Böhl & Brändli (2007) and Keller (2011)).

17 To estimate the abundance of saproxylic beetles (a measure of relative food availability), we counted all emergence
18 holes of saproxylic beetles within a 20 cm wide strip around each mapped piece of dead wood (for logs lying on the
19 ground, the inaccessible part was excluded). The strip was situated at the same place at which the diameter was
20 measured. The number of emergence holes within this strip was then used to calculate the density of holes per m²
21 on the surface of the piece of dead wood, and the mean density of holes on standing and lying dead wood was taken
22 as the relative abundance of saproxylic beetles per sample plot.

23 Aspect [°] was determined with a compass, elevation above sea level with a hand-held GPS device (Garmin eTrex
24 20, Olathe, Kansas, USA), and slope inclination with a hypsometer (Haglöf Vertex IV, Långsele, Sweden).

1 **Statistical analyses**

2 We analyzed the detection/non-detection data of the white-backed woodpecker surveys using occupancy models to
3 account for imperfect detection (MacKenzie et al., 2002). To relate the occupancy and detection probabilities to
4 predictors, we used the logit-link-function. For intercepts and slopes we used normal priors with mean zero and
5 standard deviation 1.5 and 5, respectively. For a simple prior sensitivity analysis, we rerun the best models using a
6 narrower prior (SD 2) for the slopes of the occupancy part of the model, which yielded the same model ranking. Also,
7 effect plots changed very little using the narrower prior (see Appendix I). Models were fitted in a Bayesian
8 framework using Hamiltonian Monte Carlo as implemented in the open-source software Stan via R (R Core Team,
9 2018; R-package *rstan*, Stan Development Team 2018). This technique allows for sampling from the posterior
10 distribution of the model parameters, hence, with enough samples, the posterior distribution can be approximated.
11 The posterior distribution can be used to predict detection and occupancy probability for any combination of the
12 predictors (as e.g. done for the effect plots we present), or to report the estimate of a parameter together with a
13 95% credible interval (median and 95%-interquartile range of the marginal posterior distribution).

14 Model convergence was confirmed using the \hat{R} statistics and diagnostic plots (package *shinystan*, Gabry, 2018). We
15 used 2000 after 8000 warm-up samples from each of four Markovian chains. To ease model convergence, all linear
16 predictors were centered and scaled to one standard deviation. Highly correlated covariates ($|r| > 0.7$, see Appendix
17 II for a correlation matrix) were not used in the same model. For quadratic effects, we used orthogonal polynomials.
18 For model-ranking, we used the package *loo* (Vehtari et al., 2019) which produces a leave-one-out cross-validation
19 score LooCVS for each model (a measure that is approximated by the well-known Akaike's Information Criterion AIC).

20 Initially, we attempted to fit dynamic occupancy models (MacKenzie et al., 2003) which estimate first-year
21 occupancy, a yearly colonization and extinction probability, and detection probability (package *unmarked*, Fiske &
22 Chandler (2011), function *colext*) or single-season occupancy models using a frequentist approach (and AIC model
23 ranking). However, many models did not converge. Similarly, occupancy models with square as random factor could
24 not be fitted successfully in the Stan framework as the detection/non-detection data from only two years apparently
25 did not contain enough information to fit such complex models. Hence, we treated each square-year combination as

1 a separate square, resulting in a sample size of 121 (2 x 62 minus the three squares that were not surveyed in 2016)
2 instead of 62 squares. We are aware of the pseudoreplication in our analyses by not including the square as random
3 effect in the models. However, using data from one year only yielded the same best models, and effect plots based
4 on the data from single years (Appendix III) show similar effects.

5 We are also aware of the risk of having single false positive errors (individuals that had been attracted into
6 unoccupied squares due to the use of playbacks) in our data, which would violate an assumption of occupancy
7 models (Royle & Link, 2006); even so, we feel confident that our results are reliable, and false positives should be
8 rare as the range of the playbacks which depended on terrain and background noise was usually less than 300 m.

9 *Model selection procedure*

10 As we had a large number of covariates, we used a multistep process to build models following Kéry & Royle (2015)
11 (Fig. 2, Table 2, Table 3). At a given step, we compared models based on LooCVS and used the model with the lowest
12 value to proceed to the next step.

13 The model selection procedure was conducted as follows. In step 1, we compared models containing covariates
14 describing detection probability p while holding ψ constant: In step 1.1, year and the following survey-specific
15 predictors were included (see Table 2): time, date, number of observers, and number of accessible survey points (2
16 to 4 as at least two survey points were accessible in all squares; 31 out of the 248 points were inaccessible due to
17 steep terrain, barriers such as rock ledges, or high amounts of snow). Time and date were tested as both linear and
18 quadratic effects. Then, in step 1.2, we considered models adding *a priori* selected habitat covariates on p (see Table
19 2). Note that p was modeled at the 1 km² scale only.

20 In step 2, we selected covariates describing occupancy probability ψ by comparing a set of *a priori* models (Table 1)
21 and including the covariates selected in step 1 for p . To evaluate whether the mean habitat structure of the whole
22 square or the habitat structure of the supposedly most suitable part of the square explained occupancy best, we
23 included habitat data at two spatial scales in our analyses and ran each model twice (once for each spatial scale): in
24 step 2.1a), each covariate in the occupancy part of the models was represented by the mean across the eight habitat

1 sample plots per square (1 km² scale). In step 2.1b), the values of the quarter with the highest amount of dead wood
2 was used for each covariate (0.25 km² scale), an approach that ignores the other three quarters of each square. For
3 b), values for the covariates were calculated as means of the two respective habitat sample plots. In step 2.2, we
4 included interactions (see Table 3). Interactions were not included in the *a priori* models due to the high number of
5 covariates for ψ but all possible two-way interactions were evaluated in step 2.2 with the covariates of the best
6 model from step 2.1.

7 **Results**

8 White-backed woodpeckers were found in 20 out of 62 squares in 2015 and in 18 out of 59 squares in 2016. In 10
9 squares, white-backed woodpeckers were observed in both years. In eight squares, the species was found in 2016
10 but not in 2015; in 10 squares, it was found in 2015 but not in 2016.

11 When modeling detection probability, the best-supported model included time of day, date, number of observers,
12 and the number of accessible survey points. Detection probability decreased with time, date, and number of
13 accessible survey points, and increased with the number of observers (Table 4, Fig. 3).

14 When modeling occupancy probability, the model including the covariates abundance of beetle emergence holes on
15 standing and lying dead wood (representing the food availability hypothesis) had the lowest LooCVS value at both
16 spatial scales, with the model at the 1 km² scale having a remarkably lower LooCVS value than all other models at
17 both spatial scales (Δ LooCVS to the next model at the 1 km² scale = 16.2, Δ LooCVS to the top-ranked model at the
18 0.25 km² scale = 10.6, Table 3). Occupancy was positively related to the abundances of beetle emergence holes on
19 standing and lying dead wood at each spatial scale. In general, models describing forest structure yielded lower
20 LooCVSs than models describing topography. Two covariates not included in the top-ranked model, mean DBH of live
21 trees and of snags, were also strongly related to occupancy (95% credible interval did not include 0) at the 0.25 km²
22 scale (Table 5, Fig. 4).

23

24 **Discussion**

1 By using occupancy modeling with habitat data at two spatial scales, we demonstrated that food availability at a
2 large scale (1 km²) appears to be more important for occupancy probability of the white-backed woodpecker in a
3 landscape dominated by managed forests than forest structure characteristics or geographical aspects (food
4 availability hypothesis supported). While most approaches for the conservation of old-growth forest species focus on
5 quantitative targets of structural elements such as the volume of dead wood, our results indicate that detailed
6 knowledge about how a species uses these structural elements is required for its effective conservation. Here, the
7 preference for saproxylic beetle larvae turned out to be the key to understand why a proxy for food availability
8 explained occupancy much better than more general habitat factors.

9

10 *Habitat factors related to occupancy probability*

11 At both spatial scales, the model including the abundances of saproxylic beetles on lying and standing dead wood
12 performed best; however, the LooCVS of this model was much lower at the 1 km² scale than at the 0.25 km² scale,
13 indicating that saproxylic beetles should ideally be available in high densities over a large area. Consistent with this
14 result, the white-backed woodpecker is considered a food specialist with relatively large area requirements (e.g.,
15 Scherzinger, 1982). Both the observations of foraging individuals and of individuals feeding their nestlings have
16 revealed that the larvae of saproxylic beetles, particularly large larvae such as those of *Cerambycidae*, make up the
17 majority of the species' prey (Aulén, 1988; Hogstad & Stenberg, 1997).

18 Contrary to our expectations, volume of dead wood, the habitat of saproxylic beetles, was not included in well-
19 supported models, which could have two reasons. The first might be related to our approach for processing the field
20 data for dead wood for the models. At the 1 km² scale, we calculated the mean volume of standing and lying dead
21 wood from the eight sample plots. Because of the often high heterogeneity of forests within the squares, this
22 approach probably failed to discern forest stands relevant to white-backed woodpeckers. At the 0.25 km² scale, we
23 used only the quarter with the highest amount of dead wood in squares with and without white-backed woodpecker
24 observations. This approach resulted in dead wood volumes comparable to those found in old forest reserves or
25 primeval forests (e.g., Christensen et al., 2005; Drößler, 2006; Holeksa et al., 2007; Motta et al., 2010) and white-

1 backed woodpecker habitats elsewhere (Frank, 2002; Bühler, 2009; Czeszczewik, 2009) in squares with and without
2 white-backed woodpecker observations (Table 1). Focusing on the 0.25 km² with the highest dead wood volume may
3 thus have removed the potential effect of this covariate in the models. Second, high amounts of dead wood are not
4 necessarily equal to high abundances of saproxylic beetles. Several studies have shown that other factors such as the
5 diameter, decomposition, or type (standing, lying, stump) of dead wood can play a more important role for
6 saproxylic beetle abundance (Redolfi DeZan, Battisti, & Carpaneto, 2014; Procházka & Schlaghamerský, 2019) and
7 diversity (see review by Lassauce et al., 2011) than dead wood volume alone, and the correlation between dead
8 wood volume and saproxylic beetle abundance was also weak in our study system (Appendix II). Besides the
9 characteristics of dead wood, the spatial and temporal habitat continuity may influence saproxylic beetle
10 communities (Sverdrup-Thygeson, Gustafsson, & Kouki, 2014). Consequently, dead wood varies in quality as foraging
11 substrate for woodpeckers (Löhmus, Kinks, & Soon, 2010). Although the high importance of dead wood for white-
12 backed woodpeckers has consistently been pointed out, it is important to keep in mind that dead wood is mainly
13 necessary because it contains saproxylic beetle larvae and its quality is probably at least as important as its quantity.

14 Previous studies on the habitat requirements of this species have revealed a preference for stands with high
15 amounts of lying deadwood (Czeszczewik, 2009), a high number of dying trees and a low density of forest roads
16 (Kajtoch, Figarski, & Pelka, 2013), and for old stands with a high density of dead and dying trees (Hogstad &
17 Stenberg, 1994). In contrast, covariates describing forest structure are not included in our top-ranked model.

18 However, none of the studies above has considered food availability in their analyses. The few studies on food
19 specialist woodpeckers, in which this important covariate was included in habitat models, have also highlighted its
20 high importance compared to forest structure characteristics. For instance, Bonnot, Millspaugh, & Rumble (2009)
21 found that availability and distribution of food explained territory selection of black-backed woodpeckers (*Picoides*
22 *arcticus*) better than nest site availability. Similarly, American three-toed woodpecker (*Picoides dorsalis*) occupancy
23 was positively related to the number of trees infested by bark beetles, whereas no relationship to the density of old
24 snags or the diameter of healthy or infested trees was found (Kelly et al., 2019).

1 That our models describing food availability received overwhelming support compared with the other models does
2 not imply that other factors do not affect occupancy. Accordingly, the covariates mean DBH of live trees and snags
3 also showed strong positive relationships to occupancy at the 0.25 km² scale (Fig. 4, Table 5). A high mean DBH of
4 live trees represents stands with large old trees which are often considered as habitat trees because they might offer
5 various microhabitats such as crown dead wood or bark injuries to arthropods (Bütler et al., 2013; Larrieu et al.,
6 2018), and may therefore be important for foraging. Snags are also known to be important as foraging substrate for
7 white-backed woodpeckers, with all diameter classes being used (Scherzinger, 1982; Aulén, 1988; Aulén & Lundberg,
8 1991; Frank & Hochebner, 2001; Stenberg & Hogstad, 2004; Bühler, 2009). The positive relationship between
9 occupancy and mean DBH of snags might be due to the higher availability of large larvae (preferred by the white-
10 backed woodpecker) in forest stands with large-diameter dead wood: the mean body size of saproxylic beetles has
11 been shown to increase with the diameter of dead wood (Brin et al., 2011) and the availability of large snags
12 (Janssen et al., 2017).

13 *Detection probability*

14 An important aspect of our study was the explicit modeling of the presence/absence measurement error. Consistent
15 with the known general decrease in acoustic activity of birds over the day (Bibby et al., 2000), detection probability
16 decreased with time of day, even though we used playbacks to stimulate responses of woodpeckers. Detection
17 probability was also negatively related to date. White-backed woodpeckers are known to have the highest drumming
18 activity during the pre-breeding season from the end of February to the beginning of April as well as shortly before
19 the chicks fledge at the end of May/beginning of June, while they behave inconspicuously during the breeding and
20 early nestling periods (Scherzinger, 1982). As we finished our surveys in mid-May, the linear relationship between
21 detection probability and date met our expectations. Contrary to our expectation, detection probability decreased
22 when more survey points were accessible within the square. This result might be explained by the fact that most of
23 the skipped survey points were inaccessible due to the steepness of the terrain and thus probably situated in forest
24 stands with low management intensity. Lastly, detection probability was slightly higher when two observers
25 conducted the survey, which might be due to a higher chance of hearing drumming individuals that did not approach

1 the playback/survey point. However, the credible interval included zero, indicating that the effect is relatively
2 unimportant.

3 It is generally accepted that accounting for detection probability provides more reliable and less biased estimates of
4 the biological parameters related to the occupancy of a site by a species (Kéry et al. 2013). Our study emphasizes
5 that detection probability should also be accounted for when using playbacks.

6 **Conservation implications**

7 The occurrence of the white-backed woodpecker in western Austria, eastern Switzerland, and Liechtenstein in
8 managed forests shows that this old-growth specialist is not restricted to primeval forests or large conservation
9 areas without (ongoing) forest management. However, it is important to note that the management intensity of the
10 forests considered in the present study is generally rather low, as shown by the relatively high amounts of dead
11 wood: the mean volume of dead wood was approximately 55 m³/ha in both squares with and without white-backed
12 woodpecker observations (Table 1), which is much higher than the mean dead wood volumes reported for all
13 European countries, where the values range between 8 m³/ha in North Europe to 20 m³/ha in Central-West Europe
14 (Forest Europe, 2015). In addition, patches with amounts of dead wood comparable to those found in primeval
15 forests (on average 79 m³/ha of lying and 26 m³/ha of standing dead wood) were found in our study area. Occupancy
16 was positively related to the mean DBH of live trees and standing dead wood in these patches, indicating that large-
17 diameter trees and snags in stands rich in dead wood should be present and retained for the conservation of the
18 species.

19 More important than covariates describing forest structure, however, was food availability at a large (1 km²) scale.
20 This result is particularly interesting in the context of approaches currently applied for biodiversity conservation in
21 forests. Forest management often focuses on reaching thresholds in dead wood quantity although it is well-known
22 that the diversity of saproxylic species at the stand level is also linked to dead wood diversity (Lassauce et al., 2011).
23 Likewise, present recommendations for the conservation of specialized woodpecker species such as the white-
24 backed and three-toed woodpecker (*Picoides tridactylus*) are mainly based on quantitative targets regarding dead
25 wood (e.g., Bütler et al., 2004; Kajtoch, Figarski, & Pelka, 2013). Although factors such as volume of dead wood or

1 number of habitat trees per hectare are valuable criteria for forest management guidelines because they are
2 relatively easy to assess and implement in the field, quantitative targets alone might not be the appropriate criteria
3 for the conservation of dead wood specialists in managed forests. For example, dead wood in managed forests is
4 often composed of a high proportion of stumps and small-diameter dead wood compared to old-growth forests
5 (Keren & Diaci, 2018). Even when high amounts of dead wood with these characteristics are retained, only a certain
6 proportion of the saproxylic species will benefit from this measure, whereas the habitat might remain unsuitable for
7 species preferring large-diameter snags or logs (Gossner et al. 2013). Similarly, white-backed woodpeckers are
8 unlikely to occur where dead wood is available but not sufficiently colonized by saproxylic beetles. Thus, we
9 recommend that the existing knowledge about saproxylic species and dead wood management (see e.g. Vítková et
10 al., 2018), particularly the promotion of large-diameter dead wood, be taken into account when implementing
11 conservation measures in managed forests. For developing conservation measures specifically directed at white-
12 backed woodpeckers in temperate managed forest, however, more knowledge about their preferred prey species
13 and foraging substrates is needed.

14

15 **Acknowledgements**

16 We thank Jérôme Guélat and Gabriele Hilke Peter for providing advice and support with GIS work, Ueli Bühler and
17 Georg Willi for stimulating discussions on the white-backed woodpecker, Gianna Petendi, Marlies Sperandio, Sandra
18 Melcher, Johanna Kronberger and Carolyn Nabholz for help during field work, and inatura Erlebnis Naturschau
19 GmbH, Dornbirn, for financial support. Inatura and BirdLife Vorarlberg provided us with data on previous
20 observations of the study species, for which we are very grateful.

21

22 **References**

23 Aulén, G. (1988). Ecology and distribution of the white-backed woodpecker *Dendrocopos leucotos* in Sweden
24 (Dissertation). Swedish University of Agricultural Sciences, Uppsala.

- 1 Aulén, G., & Lundberg, A. (1991). Sexual dimorphism and patterns of territory use by the white-backed woodpecker
2 *Dendrocopos leucotos*. *Ornis Scand. Scand. J. Ornithol.* **22**, 60–64.
- 3 Bengtsson, J., Nilsson, S. G., Franc, A., & Menozzi, P. (2000). Biodiversity, disturbances, ecosystem function and
4 management of European forests. *For. Ecol. Manag.* **132**, 39–50.
- 5 Bibby, C. J., Burgess, N. D., Hill, D. A., & Mustoe, S. (2000). Bird census techniques. 2nd ed. London: Academic Press.
- 6 Böhl, J., & Brändli, U.-B. (2007). Deadwood volume assessment in the third Swiss National Forest Inventory: methods
7 and first results. *Eur. J. For. Res.* **126**, 449–457.
- 8 Bonnot, T., Millsbaugh, J., & Rumble, M. (2009). Multi-scale nest-site selection by black-backed woodpeckers in
9 outbreaks of mountain pine beetles. *For. Ecol. Manag.* **259**, 220–228.
- 10 Brin, A., Bouget, C., Brustel, H., & Jactel, H. (2011). Diameter of downed woody debris does matter for saproxylic
11 beetle assemblages in temperate oak and pine forests. *J. Insect Conserv.* **15**, 653–669.
- 12 Brumelis, G., Jonsson, B. G., Kouki, J., Kuuluvainen, T., & Shorohova, E. (2011). Forest naturalness in northern
13 Europe : perspectives on processes, structures and species diversity. *Silva Fenn.* **45**, 807–821.
- 14 Bühler, U. (2009). Totholz – existenziell für den Weissrückenspecht in Nordbünden | Dead wood – a vital necessity
15 for the white-backed woodpecker in the Grisons. *Schweiz. Z. Forstwes.* **160**, 210–217.
- 16 Bütler, R, Angelstam, P., Ekelund, P., & Schlaepfer, R. (2004). Dead wood threshold values for the three-toed
17 woodpecker presence in boreal and sub-Alpine forest. *Biol. Conserv.* **119**, 305–318.
- 18 Bütler, Rita, Lachat, T., Larrieu, L., & Paillet, Y. (2013). Habitat trees: key elements for forest biodiversity. In
19 Integrative approaches as an opportunity for the conservation of forest biodiversity. Kraus D., Krumm F.
20 (eds). pp. 84–91. Freiburg: European Forest Institute.
- 21 Carlson, A. (2000). The effect of habitat loss on a deciduous forest specialist species: the white-backed woodpecker
22 (*Dendrocopos leucotos*). *For. Ecol. Manag.* **131**, 215–221.
- 23 Christensen, M., Hahn, K., Mountford, E. P., Ódor, P., Standovár, T., Rozenbergar, D., Diaci, J., Wijdeven, S., Meyer,
24 P., Winter, S., & Vrska, T. (2005). Dead wood in European beech (*Fagus sylvatica*) forest reserves. *For. Ecol.*
25 *Manag.* **210**, 267–282.

- 1 Clavel, J., Julliard, R., & Devictor, V. (2010). Worldwide decline of specialist species: toward a global functional
2 homogenization? *Front Ecol Env.* **8**, 222–228.
- 3 Commarmot, B., Bachofen, H., Bundziak, Y., Bürgi, A., Ramp, B., Shparyk, Y., Sukhariuk, D., Viter, R., & Zingg, A.
4 (2005). Structures of virgin and managed beech forests in Uholka (Ukraine) and Sihlwald (Switzerland): a
5 comparative study. *For. Snow Landsc. Res.* **79**, 45–56.
- 6 Czeszczewik, D. (2009). Marginal differences between random plots and plots used by foraging white-backed
7 woodpeckers demonstrates supreme primeval quality of the Białowieża National Park, Poland. *Ornis Fenn.*
8 **86**, 30–37.
- 9 Czeszczewik, D., & Walankiewicz, W. (2006). Logging affects the white-backed woodpecker *Dendrocopos leucotos*
10 distribution in the Białowieża Forest. *Ann. Zool. Fenn.* **43**, 221–227.
- 11 Devictor, V., Julliard, R., & Jiguet, F. (2008). Distribution of specialist and generalist species along spatial gradients of
12 habitat disturbance and fragmentation. *Oikos* **117**, 507–514.
- 13 Drößler, L. (2006). Struktur und Dynamik von zwei Buchenurwäldern in der Slowakei. Dissertation, Georg-August-
14 Universität Göttingen, Göttingen.
- 15 Ellenberg, H., & Strutt, G. K. (2009). Vegetation ecology of Central Europe. 4th ed. Cambridge: Cambridge University
16 Press.
- 17 Ettwein, A. (2016). Habitatwahl des Weißrückenspechts (*Dendrocopos leucotos*) in Vorarlberg, der Ostschweiz und
18 dem Fürstentum Liechtenstein. Master's thesis, Universität für Bodenkultur Wien, Wien.
- 19 Fiske, I. J., & Chandler, R. B. (2011). Unmarked: an R package for fitting hierarchical models of wildlife occurrence and
20 abundance. *J. Stat. Softw.* **43**, 1–23.
- 21 Forest Europe. (2015). State of Europe's Forests 2015. Ministerial Conference on the Protection of Forests in Europe.
- 22 Fraixedas, S., Lindén, A., & Lehtikoinen, A. (2015). Population trends of common breeding forest birds in southern
23 Finland are consistent with trends in forest management and climate change. *Ornis Fenn.* **92**, 187–203.
- 24 Frank, G. (2002). Brutzeitliche Einnischung des Weissrückenspechtes *Dendrocopos leucotos* im Vergleich zum
25 Buntspecht *Dendrocopos major* in montanen Mischwäldern der nordlichen Kalkalpen. *Vogelwelt* **123**, 225–
26 239.

- 1 Frank, G., & Hochebner, T. (2001). Erfassung der Spechte - insbesondere des Weißrückenspechtes *Picoides leucotos* -
2 im Rahmen des LIFE-Projektes Wildnisgebiet Dürrenstein. in: LIFE- Projekt Wildnisgebiet Dürrenstein. Amt
3 der Niederösterreichischen Landesregierung (Ed.). pp. 116–148. St. Pölten: Amt der Niederösterreichischen
4 Landesregierung.
- 5 Gabry, J. (2018). shinystan: interactive visual and numerical diagnostics and posterior analysis for bayesian models. R
6 package version 2.5.0. <https://CRAN.R-project.org/package=shinystan>.
- 7 Glutz von Blotzheim, U. N., & Bauer, K. (1994). Handbuch der Vögel Mitteleuropas. Band 9: *Columbiformes* -
8 *Piciformes*. 2nd ed. Wiesbaden: Aula-Verlag.
- 9 Grove, S. J. (2002). Saproxylic insect ecology and the sustainable management of forests. *Annu. Rev. Ecol. Syst.* **33**, 1–
10 23.
- 11 Henle, K., Davies, K. F., Kleyer, M., Margules, C., & Settele, J. (2004). Predictors of species sensitivity to
12 fragmentation. *Biodivers. Conserv.* **13**, 207–251.
- 13 Hogstad, O., & Stenberg, I. (1994). Habitat selection of a viable population of white-backed woodpeckers
14 *Dendrocopos leucotos*. *Fauna Nor. Ser. C Cinclus* **17**, 75–94.
- 15 Hogstad, O., & Stenberg, I. (1997). Breeding success, nestling diet and parental care in the white-backed woodpecker
16 *Dendrocopos leucotos*. *J. Für Ornithol.* **138**, 25–38.
- 17 Holeksa, J., Saniga, M., Szwagrzyk, J., Dziedzic, T., Ferenc, S., & Wodka, M. (2007). Altitudinal variability of stand
18 structure and regeneration in the subalpine spruce forests of the Pol'ana biosphere reserve, Central
19 Slovakia. *Eur. J. For. Res.* **126**, 303–313.
- 20 Janssen, P., Fuhr, M., Cateau, E., Nusillard, B., & Bouget, C. (2017). Forest continuity acts congruently with stand
21 maturity in structuring the functional composition of saproxylic beetles. *Biol. Conserv.* **205**, 1–10.
- 22 Kajtoch, L., Figarski, T., & Pelka, J. (2013). The role of forest structural elements in determining the occurrence of two
23 specialist woodpecker species in the Carpathians, Poland. *Ornis Fenn.* **90**, 23–40.
- 24 Keller, M. (2011). Swiss National Forest Inventory. Manual of the field survey 2004 - 2007. p. 269. Birmensdorf: Swiss
25 Federal Research Institute WSL.

- 1 Kelly, J. J., Latif, Q. S., Saab, V. A., & Veblen, T. T. (2019). Spruce beetle outbreaks guide American three-toed
2 woodpecker *Picoides dorsalis* occupancy patterns in subalpine forests. *Ibis* **161**, 172–183.
- 3 Keren, S., & Diaci, J. (2018). Comparing the quantity and structure of deadwood in selection managed and old-
4 growth forests in south-east Europe. *Forests* **9**, 76.
- 5 Kéry, M., & Royle, J. A. (2016). Applied hierarchical modeling in ecology: analysis of distribution, abundance and
6 species richness in R and BUGS. Volume 1: Prelude and static models. 1st ed. Cambridge: Academic Press.
- 7 Kuuluvainen, T., Penttinen, A., Leinonen, K., & Nygren, M. (1996). Statistical opportunities for comparing stand
8 structural heterogeneity in managed and primeval forests: an example from boreal spruce forest in
9 southern Finland. *Silva Fenn.* **30**, 315–328.
- 10 Larrieu, L., Paillet, Y., Winter, S., Bütler, R., Kraus, D., Krumm, F., Lachat, T., Michel, A. K., Regnery, B., &
11 Vandekerckhove, K. (2018). Tree related microhabitats in temperate and Mediterranean European forests: a
12 hierarchical typology for inventory standardization. *Ecol. Indic.* **84**, 194–207.
- 13 Lassauce, A., Paillet, Y., Jactel, H., & Bouget, C. (2011). Deadwood as a surrogate for forest biodiversity: meta-analysis
14 of correlations between deadwood volume and species richness of saproxylic organisms. *Ecol. Indic.* **11**,
15 1027–1039.
- 16 Lõhmus, A., Kinks, R., & Soon, M. (2010). The importance of dead-wood supply for woodpeckers in Estonia. *Balt. For.*
17 **16**, 76–86.
- 18 MacKenzie, D. I., Nichols, J. D., Hines, J. E., Knutson, M. G., & Franklin, A. B. (2003). Estimating site occupancy,
19 colonization, and local extinction when a species is detected imperfectly. *Ecology* **84**, 2200–2207.
- 20 MacKenzie, D. I., Nichols, J. D., Lachman, G. B., Droege, S., Royle, J. A., & Langtimm, C. A. (2002). Estimating site
21 occupancy rates when detection probabilities are less than one. *Ecology* **83**, 2248–2255.
- 22 Merino, A., Real, C., Álvarez-González, J. G., & Rodríguez-Gutián, M. A. (2007). Forest structure and C stocks in
23 natural *Fagus sylvatica* forest in southern Europe: the effects of past management. *For. Ecol. Manag.* **250**,
24 206–214.
- 25 Mikusiński, G., Gromadzki, M., & Chylarecki, P. (2001). Woodpeckers as indicators of forest bird diversity. *Conserv.*
26 *Biol.* **15**, 208–217.

- 1 Mollet, P., Zbinden, N., & Schmid, H. (2009). Steigende Bestandszahlen bei Spechten und anderen Vogelwarten dank
2 Zunahme von Totholz? *Schweiz. Z. Forstwes.* **160**, 334–340.
- 3 Motta, R., Berretti, R., Castagneri, D., Lingua, E., Nola, P., & Vacchiano, G. (2010). Stand and coarse woody debris
4 dynamics in subalpine Norway spruce forests withdrawn from regular management. *Ann. For. Sci.* **67**, 803–
5 803.
- 6 Müller, J., Hothorn, T., & Pretzsch, H. (2007). Long-term effects of logging intensity on structures, birds, saproxylic
7 beetles and wood-inhabiting fungi in stands of European beech *Fagus sylvatica* L. *For. Ecol. Manag.* **242**,
8 297–305.
- 9 Procházka, J., & Schlaghamerský, J. (2019). Does dead wood volume affect saproxylic beetles in montane beech-fir
10 forests of Central Europe? *J. Insect Conserv.* **23**, 157–173.
- 11 Redolfi DeZan, L., Battisti, C., & Carpaneto, G. M. (2014). Bird and beetle assemblages in relict beech forests of
12 central Italy: A multi-taxa approach to assess the importance of dead wood in biodiversity conservation.
13 *Community Ecol.* **15**, 235–245.
- 14 Roberge, J.-M., Mikusiński, G., & Svensson, S. (2008). The white-backed woodpecker: umbrella species for forest
15 conservation planning? *Biodivers. Conserv.* **17**, 2479–2494.
- 16 Royle, J. A., & Link, W. A. (2006). Generalized site occupancy models allowing for false positive and false negative
17 errors. *Ecology* **87**, 835–841.
- 18 Scherzinger, W. (1982). Die Spechte im Nationalpark Bayerischer Wald. Vol. Heft 9. Grafenau:
19 Nationalparkverwaltung Bayerischer Wald.
- 20 Schulze, A., Roché, J. C., Chappuis, C., Dingler, K. H., & Werle, A. (2003). Die Vogelstimmen Europas, Nordafrikas und
21 Vorderasiens. Germering: Ample Edition.
- 22 Speight, M. C. D. (1989). Saproxylic invertebrates and their conservation. p. 78. Strasbourg, France: Council of
23 Europe.
- 24 Stan Development Team. (2018). RStan: the R interface to Stan. R package version 2.18.2. <http://mc-stan.org/>.
- 25 Stenberg, I., & Hogstad, O. (2004). Sexual dimorphism in relation to winter foraging in the white-backed woodpecker
26 (*Dendrocopos leucotos*). *J. Ornithol.* **145**, 321–326.

- 1 Stübing, S., & Bergmann, H. H. (2006). Methodenstandards zur Erfassung der Brutvögel Deutschlands:
2 Klangattrappen. 1st ed. Radolfzell: Verein Sächsischer Ornithologen.
- 3 Sverdrup-Thygeson, A., Gustafsson, L., & Kouki, J. (2014). Spatial and temporal scales relevant for conservation of
4 dead-wood associated species: current status and perspectives. *Biodivers. Conserv.* **23**, 513–535.
- 5 Vehtari, A., Gabry, J., Yao, Y., & Gelman, A. (2019). Efficient leave-one-out cross-validation and WAIC for Bayesian
6 models." R package version 2.1.0, <URL: <https://CRAN.R-project.org/package=loo>>.
- 7 Virkkala, R., Alanko, T., Laine, T., & Tiainen, J. (1993). Population contraction of the white-backed woodpecker
8 *Dendrocopos leucotos* in Finland as a consequence of habitat alteration. *Biol. Conserv.* **66**, 47–53.
- 9 Vítková, L., Bače, R., Kjučukov, P., & Svoboda, M. (2018). Deadwood management in Central European forests: Key
10 considerations for practical implementation. *For. Ecol. Manag.* **429**, 394–405.
- 11
- 12


1 **Figures**

2 Figure 1 Overview of study area and study design. Detection/non-detection of white-backed woodpeckers was
3 recorded in 62 1-km²-squares at four survey points (cross) per square. Habitat structure and saproxylic beetle
4 abundance were mapped in eight circular sample plots (dots) with a size of 500 m² each per square.

5 Figure 2 Model selection procedure for modeling white-backed woodpecker detection (p) and occupancy
6 probabilities (ψ).


7 Figure 3 Relationships between white-backed woodpecker occupancy probability within surveyed squares at the 1
8 km² scale and abundance of beetle emergence holes on standing (a) and lying (b) deadwood (note the logarithmic x-
9 axis), and between detection probability and (c) time of day and (d) day of year based on the top-ranked model from
10 step 2.2 (Table 4). The solid lines are fitted values, calculated with the values for other covariates set to their means.
11 Dotted line = 95% credible interval, circles = raw data. n = 121.

12 Figure 4 Relationships between white-backed woodpecker occupancy probability within surveyed squares at the 0.25
13 km² scale and (a) mean diameter at breast height (DBH) of live trees and (b) mean DBH of standing dead wood. The
14 solid lines are fitted values, calculated with the values for other covariates set to their means. Dotted line = 95%
15 credible interval, circles = raw data. n = 121.



1


2 Figure 1



1

2 Figure 2



3



1

2 Figure 3

3



1

2 Figure 4

1 **Tables**

2 Table 1 Covariates used for modeling white-backed woodpecker occupancy (ψ) and detection (p) probabilities according to
 3 different hypotheses (*a priori* models). Each hypothesis was represented by one model containing the corresponding listed
 4 covariates. Predicted (pred.) effects are the *a priori* expectations: + = positive correlation between parameter and covariate was
 5 predicted, - = negative correlation predicted, x = no prediction, but correlation could be present, ^= quadratic effect predicted.
 6 Mean \pm SD of the covariates are reported for squares with and for squares without white-backed woodpecker (WBW)
 7 observations at the 1 km² scale as well as for the 0.25 km² with the highest amount of dead wood.

Hypothesis	Covariate name	Description	pred. effect	1km ² square		0.25 km ² with the most dw			
				p	$\psi/\epsilon/\gamma$	without WBW	with WBW	without WBW	with WBW
(1) Old mixed deciduous forests	dbh	mean DBH of live trees (cm)	+	+/-/+	32.3 \pm 5.0	33.2 \pm 4.2	29.9 \pm 6.1	34.1 \pm 7.0	
	forest	area (ha) of forest per 1km ² square/quarter of the square		+/-/+	76.6 \pm 12.8	77.2 \pm 14.4	19.3 \pm 4.1	20.4 \pm 4.5	
	deadbr	number of live trees with dead branches/ha		+/-/+	12.9 \pm 11.9	20.7 \pm 19.6	9.0 \pm 12.9	17.1 \pm 24.3	
	dec	proportion of deciduous trees	+	+/-/+	41.5 \pm 28.4	45.7 \pm 20.0	40.8 \pm 31.6	46.2 \pm 24.2	
(2) Dead wood volume	vollog	volume of lying dead wood (m ³)		+/-/+	36.4 \pm 33.7	38.2 \pm 40.5	69.3 \pm 70.5	79.1 \pm 82.2	
	volsnag	volume of snags (m ³)		+/-/+	19.4 \pm 10.9	16.5 \pm 7.5	34.0 \pm 29.8	26.1 \pm 17.5	
(3) Large pieces of dead wood	dlog	mean diameter of lying dead wood		+/-/+	15.4 \pm 6.1	15.2 \pm 5.1	16.9 \pm 12.8	19.5 \pm 11.8	
	dsnag	mean DBH of snags		+/-/+	18.5 \pm 8.5	19.0 \pm 9.1	18.9 \pm 17.6	20.7 \pm 13.6	
(4) Abundance of suitable nesting trees	snag30	number of snags with a DBH > 30 cm		+/-/+	4.2 \pm 4.6	4.6 \pm 3.5	6.6 \pm 9.4	6.8 \pm 6.7	
(5) Food availability	beetlel	number of beetle emergence holes in lying dead wood per m ²		+/-/+	457 \pm 582	599 \pm 545	346 \pm 357	580 \pm 539	
	beetles	number of beetle emergence holes in snags per m ²		+/-/+	254 \pm 136	291 \pm 101	252 \pm 177	274 \pm 183	
(6-8) South-exposed, steep slopes in intermediate elevation	northness	aspect computed as $\cos(\text{aspect}[\text{°}] * \pi / 180)$		-/+/-	-0.2 \pm 0.52	-0.28 \pm 0.48	-0.21 \pm 0.71	-0.36 \pm 0.72	
	eastness	aspect computed as $\sin(\text{aspect}[\text{°}] * \pi / 180)$		x	-0.09 \pm 0.58	-0.04 \pm 0.57	-0.06 \pm 0.69	-0.1 \pm 0.61	
	slope	slope (%), linear and quadratic effects		x	+/-/+	54.8 \pm 16.5	53.0 \pm 11.9	57.5 \pm 20.6	55.2 \pm 24.9
	elev	elevation above sea level (m), linear and quadratic effects		^/+/^	1008 \pm 260	978 \pm 189	999 \pm 269	998 \pm 229	
Survey-specific covariates									
	time	time of day, linear and quadratic effects		-					
	date	day of year, linear and quadratic effects		-					
	accessible	number of accessible survey points (2, 3, 4)		+					
	obs	number of observers during white-backed woodpecker survey (1, 2)		x					
	year	year of white-backed woodpecker survey (2015, 2016)		x					

1 Table 2 Results of model selection for modeling white-backed woodpecker detection (p) probability. Model selection was
 2 performed based on the leave-one-out cross-validation score (LooCVS). np = number of estimated parameters, Δ LooCVS =
 3 difference between a model's LooCVS and the LooCVS of the top-ranked model of the respective section. Models printed bold
 4 were selected for further modeling. Abbreviations of covariate names are given in Table 1.

Model	np	LooCVS	Δ LooCVS
Step 1.1: Including survey-specific covariates			
$\psi(.)p(\text{time+date+obs+accessible})$	5	220.2	0
$\psi(.)p(\text{time+date+accessible})$	5	220.6	0.4
$\psi(.)p(\text{time+date})$	4	222	1.8
$\psi(.)p(\text{time+date+obs})$	5	222.9	2.7
$\psi(.)p(\text{time+date+year})$	5	223.2	3
$\psi(.)p(\text{time})$	3	224.3	4.1
$\psi(.)p(\text{time+time}^2)$	4	225.6	5.4
$\psi(.)p(\text{date})$	3	229.1	8.9
$\psi(.)p(\text{accessible})$	3	229.4	9.2
$\psi(.)p(.)$	2	230.2	10
$\psi(.)p(\text{accessible+obs})$	4	231.2	11
$\psi(.)p(\text{year})$	3	231.8	11.6
$\psi(.)p(\text{obs})$	3	231.8	11.6
$\psi(.)p(\text{date+date}^2)$	4	231.8	11.6
$\psi(.)p(\text{obs+accessible+year})$	5	232.8	12.6
Step 1.2 : Including habitat covariates			
$\psi(.)p(\text{time+date+obs+accessible})$	6	220.2	0
$\psi(.)p(\text{time+date+obs+accessible+dec})$	7	222	1.8
$\psi(.)p(\text{time+date+obs+accessible+dbh})$	7	223.4	3.2
$\psi(.)p(\text{time+date+obs+accessible+forest})$	7	223.4	3.2
$\psi(.)p(\text{time+date+obs+accessible+slope})$	7	226.5	6.3

5

6

1 Table 3 Results of model selection for modeling white-backed woodpecker occupancy (ψ) probability. Model selection was
 2 performed based on the leave-one-out cross-validation score (LooCVS). np = number of estimated parameters, Δ LooCVS =
 3 difference between a model's LooCVS and the LooCVS of the top-ranked model of the respective section. Models printed bold
 4 were selected for further modeling. Abbreviations of covariate names are given in Table 1.

Model	np	LooCVS	Δ LooCVS
Step 2.1a: Testing the <i>a priori</i> hypotheses at the 1 km² scale			
$\psi(\text{beetlel+beetles})p(\text{time+date+obs+accessible})$	8	202.4	0
$\psi(\text{snag30})p(\text{time+date+obs+accessible})$	7	218.6	16.2
$\psi(.)p(\text{time+date+obs+accessible})$	6	220	17.6
$\psi(\text{vollog+volsnag})p(\text{time+date+obs+accessible})$	8	221.2	18.8
$\psi(\text{dbh+forest+deadbr+dec})p(\text{time+date+obs+accessible})$	10	223.3	20.9
$\psi(\text{dlog+dsnag})p(\text{time+date+obs+accessible})$	8	223.3	20.9
$\psi(\text{northness+eastness+slope+slope}^2\text{+elev})p(\text{time+date+obs+accessible})$	11	224.7	22.3
$\psi(\text{northness+eastness+slope+elev+elev}^2\text{)p}(\text{time+date+obs+accessible})$	11	227.1	24.7
$\psi(\text{northness+eastness+slope+elev})p(\text{time+date+obs+accessible})$	10	227.7	25.3
Step 2.1b: Testing the <i>a priori</i> hypotheses at the 0.25 km² scale			
$\psi(\text{beetlel+beetles})p(\text{time+date+obs+accessible})$	8	213	0
$\psi(\text{dbh+forest+deadbr+dec})p(\text{time+date+obs+accessible})$	10	216.5	3.5
$\psi(\text{dlog+dsnag})p(\text{time+date+obs+accessible})$	8	219.5	6.5
$\psi(.)p(\text{time+date+obs+accessible})$	6	220	7
$\psi(\text{snag30})p(\text{time+date+obs+accessible})$	7	220.1	7.1
$\psi(\text{vollog+volsnag})p(\text{time+date+obs+accessible})$	8	225.1	12.1
$\psi(\text{northness+eastness+slope+elev})p(\text{time+date+obs+accessible})$	10	226.1	13.1
$\psi(\text{northness+eastness+slope+slope}^2\text{+elev})p(\text{time+date+obs+accessible})$	9	226.4	13.4
$\psi(\text{northness+eastness+slope+elev+elev}^2\text{)p}(\text{time+date+obs+accessible})$	11	228.9	15.9
Step 2.2: Testing the effect of year			
$\psi(\text{beetlel+beetles})p(\text{time+date+obs+accessible})$	8	202.4	0
$\psi(\text{beetlel+beetles+year})p(\text{time+date+obs+accessible})$	9	204.6	2.2
Step 2.3: Including interactions			
$\psi(\text{beetlel+beetles})p(\text{time+date+obs+accessible})$	8	202.4	0
$\psi(\text{beetlel+beetles+beetlel:beetlex})p(\text{time+date+obs+accessible})$	9	215.7	13.3

5

6

1 Table 4 Medians and credible intervals (2.5 % and 97.5 % quantiles) of the marginal posterior distribution of the covariates
 2 included in each *a priori* occupancy model attempting to predict white-backed woodpecker occupancy and detection probability
 3 at the 1 km² scale. Models are ordered according to their leave-one-out cross-validation score, from lowest on the left to highest
 4 on the right. Explanations of covariate names and hypothesis (i.e., *a priori* model) numbers are given in Table 1.

Covariate	Hypothesis							
	(5)	(4)	(2)	(1)	(3)	(6)	(7)	(8)
occupancy probability								
intercept	1.24 [0.17;3.10]	1.01 [-0.08;2.98]	1.97 [0.23;4.40]	1.27 [-0.05;3.86]	1.41 [0.06;3.69]	2.85 [0.55;5.16]	2.98 [1.02;5.15]	2.97 [0.86;5.32]
dbh	-	-	-	0.49 [-6.45;7.21]	-	-	-	-
forest	-	-	-	0.35 [-6.47;7.25]	-	-	-	-
deadbr	-	-	-	0.56 [-0.48;1.56]	-	-	-	-
dec	-	-	-	0.36 [-1.42;2.22]	-	-	-	-
vollog	-	-	-0.31 [-1.74;0.99]	-	-	-	-	-
volsnag	-	-	1.18 [-1.64;3.41]	-	-	-	-	-
dlog	-	-	-	-	-0.13 [-1.38;1.16]	-	-	-
dsnag	-	-	-	-	0.95 [-0.56;2.98]	-	-	-
snag30	-	0.98 [-0.01;2.21]	-	-	-	-	-	-
beetle	1.76 [0.54;3.51]	-	-	-	-	-	-	-
beetles	2.61 [1.12;4.92]	-	-	-	-	-	-	-
northness	-	-	-	-	-	-0.70 [-2.87;1.40]	0.00 [-2.20;1.82]	0.07 [-2.28;1.97]
eastness	-	-	-	-	-	1.45 [-0.71;3.71]	0.81 [-1.41;3.20]	0.52 [-2.29;3.47]
slope	-	-	-	-	-	1.53 [-0.65;3.60]	0.71 [-2.26;2.75]	0.15 [-2.25;2.32]
slope_q	-	-	-	-	-	-	-	-1.31 [-3.50;0.88]
elev	-	-	-	-	-	-1.48 [-4.43;1.26]	0.09 [-3.78;3.17]	0.50 [-3.28;3.49]
elev_q	-	-	-	-	-	-1.62 [-3.31;-0.11]	-	-
detection probability (only given for the top-ranked model)								
intercept	-1.59 [-2.75;-0.47]							
time	-0.73 [-1.22;-0.31]							
date	-0.47 [-0.99;0.00]							
obs	0.44 [-0.27;1.18]							
accessible	-0.72 [-1.26;-0.24]							

5

6


1 Table 2 Medians and credible intervals (2.5 % and 97.5 % quantiles) of the marginal posterior distribution of the covariates
 2 included in each *a priori* occupancy model attempting to predict white-backed woodpecker occupancy and detection probability
 3 at the 0.25 km² scale. Models are ordered according to their leave-one-out cross-validation score, from lowest on the left to
 4 highest on the right. Explanations of covariate names and hypothesis (i.e., *a priori* model) numbers are given in Table 1.

Covariate	Hypothesis							
	(5)	(1)	(3)	(4)	(2)	(6)	(7)	(8)
occupancy probability								
intercept	1.03 [-0.08;3.09]	0.89 [-0.23;3.19]	0.74 [-0.20;2.78]	1.10 [-0.06;3.33]	1.32 [-0.02;4.08]	2.74 [0.72;5.09]	2.53 [0.70;4.88]	2.62 [0.96;4.91]
dbh	-	1.42 [0.41;3.00]	-	-	-	-	-	-
forest	-	0.76 [-0.35;2.03]	-	-	-	-	-	-
deadbr	-	0.82 [-0.21;2.40]	-	-	-	-	-	-
dec	-	0.77 [-0.42;2.41]	-	-	-	-	-	-
vollog	-	-	-	-	0.09 [-1.43;1.28]	-	-	-
volsnag	-	-	-	-	0.31 [-2.65;1.58]	-	-	-
dlog	-	-	0.45 [-0.73;1.48]	-	-	-	-	-
dsnag	-	-	0.94 [0.22;1.97]	-	-	-	-	-
snag30	-	-	-	0.86 [-0.15;2.86]	-	-	-	-
beetlel	1.27 [0.41;2.56]	-	-	-	-	-	-	-
beetles	1.22 [0.27;2.75]	-	-	-	-	-	-	-
northness	-	-	-	-	-	0.67 [-1.33;2.73]	0.62 [-1.47;2.73]	-0.04 [-1.76;2.10]
eastness	-	-	-	-	-	-0.73 [-3.41;1.92]	-0.50 [-2.70;1.97]	-0.14 [-2.21;2.31]
slope	-	-	-	-	-	0.20 [-1.34;1.72]	0.02 [-1.37;1.21]	1.12 [-1.11;3.26]
slope_q	-	-	-	-	-	-	-	2.16 [-0.16;5.36]
elev	-	-	-	-	-	1.13 [-1.69;4.18]	1.31 [-1.92;3.39]	1.03 [-2.13;3.35]
elev_q	-	-	-	-	-	-0.44 [-2.09;1.98]	-	-
detection probability (only given for the top-ranked model)								
intercept	-1.63 [-2.81;-0.52]							
time	-0.73 [-1.22;-0.31]							
date	-0.56 [-1.08;-0.09]							
obs	0.40 [-0.28;1.15]							
accessible	-0.58 [-1.10;-0.14]							

5

6

1 Appendix I Fitted values and 95% credible intervals of relationships between white-backed woodpecker occupancy probability
2 and abundance of beetle emergence holes on (a) standing and (b) lying deadwood (note the logarithmic x-axis)
3 modeled with two different priors (solid lines = used prior, dotted lines = alternative prior). Fitted values were calculated with the values for other
4 covariates set to their means. Circles = raw data. The plots are based on the top-ranked model from step 2.2 (Table 4). n = 121




- 1 Appendix II Correlations between the covariates used for occupancy modeling (Pearson correlation coefficient). Values above the
- 2 diagonal represent correlations at the 0.25 km² scale, values below the diagonal represent correlations at the 1 km² scale.
- 3 Covariates are explained in Table 1.

	dbh	forest	deadbr	dec	vollog	volsnag	dlog	dsnag	snag30	beetle	beetles	northness	eastness	slope	elev
dbh	1.00	-0.10	0.28	-0.32	0.22	0.15	0.18	0.13	0.16	-0.03	0.37	0.06	0.33	0.26	0.19
forest	-0.07	1.00	-0.19	-0.01	0.01	-0.15	0.02	-0.01	-0.15	0.00	-0.20	-0.08	-0.13	-0.01	0.03
deadbr	0.27	-0.04	1.00	0.10	0.17	0.12	0.22	0.20	0.15	-0.08	0.15	0.00	0.11	0.09	-0.08
dec	-0.19	-0.03	0.21	1.00	0.12	-0.19	0.11	-0.05	0.01	0.01	-0.24	-0.06	-0.14	0.23	-0.40
vollog	-0.01	-0.22	0.07	0.36	1.00	-0.06	0.78	0.24	0.17	-0.11	0.10	0.11	0.08	0.09	0.09
volsnag	0.28	-0.10	0.18	-0.04	0.20	1.00	-0.01	0.21	0.56	0.19	0.41	-0.26	-0.08	0.08	0.16
dlog	0.01	-0.16	0.10	0.17	0.75	0.19	1.00	0.17	0.31	0.03	0.08	0.15	0.08	0.01	0.14
dsnag	0.30	-0.07	0.12	0.22	0.25	0.54	0.14	1.00	0.47	-0.08	0.31	-0.10	-0.05	0.12	0.24
snag30	0.27	-0.12	0.23	0.21	0.37	0.65	0.34	0.70	1.00	0.21	0.39	-0.20	-0.19	0.28	0.19
beetle	-0.13	-0.06	-0.07	-0.05	-0.10	0.23	0.11	0.13	0.36	1.00	-0.06	-0.20	0.17	0.03	0.39
beetles	0.08	0.06	0.46	0.16	0.20	0.26	0.18	0.26	0.37	0.16	1.00	-0.07	-0.06	0.16	0.13
northness	-0.10	-0.07	-0.08	-0.16	0.09	-0.20	0.03	-0.26	-0.43	0.05	-0.05	1.00	-0.12	0.08	-0.13
eastness	0.21	-0.19	-0.07	0.01	0.04	0.06	0.05	0.16	0.11	0.03	0.05	-0.28	1.00	-0.02	0.24
slope	0.13	-0.18	0.20	0.16	0.11	0.16	0.16	0.23	0.20	-0.03	-0.08	-0.11	-0.07	1.00	0.11
elev	0.21	-0.14	-0.04	-0.41	-0.09	0.19	0.28	0.08	0.06	-0.16	0.04	-0.13	0.27	0.18	1.00

4

5

1 Appendix III: Relationships between white-backed woodpecker occupancy probability and abundance of beetle emergence holes
 2 on standing and lying deadwood (note the logarithmic x-axis) modeled separately for the years 2015 and 2016. The solid lines are
 3 fitted values, calculated with the values for other covariates set to their means. Dotted line = 95% credible interval, circles = raw
 4 data. All plots are based on the top-ranked model from step 2.2 (Table 4). n = 62 in 2015, n = 59 in 2016.



5

6