**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

COVID-19: A series of important recent clinical and laboratory reports in immunology and pathogenesis of SARS-CoV-2 infection and care of allergy patients

Pfaar, Oliver ; Torres, Maria J ; Akdis, Cezmi A

Abstract: The “coronavirus disease 2019 (COVID-19)” outbreak was first reported in December 2019 (China). Since then, this disease has rapidly spread across the globe and in March 2020 the World Health Organization (WHO) declared the COVID-19 pandemic.¹ Since the outbreak was first announced, our journal has extensively focused on the clinical features, outcomes, diagnosis, immunology, and pathogenesis of COVID-19 and its infectious agent severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).

DOI: <https://doi.org/10.1111/all.14472>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-195080>

Journal Article

Accepted Version

Originally published at:

Pfaar, Oliver; Torres, Maria J; Akdis, Cezmi A (2021). COVID-19: A series of important recent clinical and laboratory reports in immunology and pathogenesis of SARS-CoV-2 infection and care of allergy patients. *Allergy*, 76(3):622-625.

DOI: <https://doi.org/10.1111/all.14472>

PROF. OLIVER PFAAR (Orcid ID : 0000-0003-4374-9639)

PROF. CEZMI AKDIS (Orcid ID : 0000-0001-8020-019X)

Article type : Editorial

COVID-19: A series of important recent clinical and laboratory reports in immunology and pathogenesis of SARS-CoV-2 infection and care of allergy patients

Oliver Pfaar¹, Maria J. Torres² and Cezmi A. Akdis³

¹ Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Germany.

² Allergy Clinical Unit, Hospital Regional Universitario de Málaga-UMA-ARADyAL, Málaga, Spain.

³ Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich, Davos, Switzerland.

Corresponding author:

Prof. Dr. med. Oliver Pfaar

Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy,
University Hospital Marburg, Philipps-Universität Marburg,
Marburg, Germany
Baldingerstr
D-35043 Marburg
Germany

E-mail: oliver@pfaar.org

Telephone: +49 6421 58 66479

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ALL.14472](https://doi.org/10.1111/ALL.14472)

This article is protected by copyright. All rights reserved

The “*coronavirus disease 2019* (COVID-19)” outbreak was first reported in December 2019 (China). Since then, this disease has rapidly spread across the globe and in March 2020 the World Health Organization (WHO) declared the COVID-19 pandemic.¹ Since the outbreak was first announced, our journal has extensively focused on the clinical features, outcomes, diagnosis, immunology, and pathogenesis of COVID-19 and its infectious agent severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). We published our first COVID-19 article on 19 February, focusing for the first time on the clinical characteristics of 140 cases of human-to-human coronavirus transmission without any links to the Huanan Wet Market.² Hypertension and diabetes were mentioned as risk factors and there was no increased prevalence in allergic patients. This early study reported that the main symptoms at hospital admission were fever (91.7%), cough (75.0%), fatigue (75.0%), gastrointestinal symptoms (39.6%), and dyspnea (36.7%). Lymphopenia and eosinopenia were also reported as important signs and biomarkers for monitoring and severity of the patients.² The prevalent eosinopenia in COVID-19 patients and the possible anti-viral role of eosinophils were further discussed in several following publications in *Allergy*.^{3,4} Our second COVID-19 paper brought attention to the wide range of clinical manifestations of this disease, from asymptomatic cases to patients with mild and severe

symptoms, with or without pneumonia as well as with only diarrhea.⁵ Patients with common allergic diseases did not develop distinct symptoms and severe courses. Cases with pre-existing chronic obstructive pulmonary disease or complicated with a secondary bacterial pneumonia were severe. Another article, timely appearing in our journal, alerted the scientific community that even in experienced hands there was a 14.1% false negative polymerase chain reaction (PCR) diagnosis in COVID-19 cases and were later diagnosed positive after repeated tests.⁶ A pediatric article was also published extensively analyzing 182 cases and it was reported that children with COVID-19 showed a mild clinical course.⁷ Patients with pneumonia had a higher proportion of fever and cough and increased inflammatory biomarkers compared to those without pneumonia. There were 43 allergic patients in this series and there was no significant difference between allergic and non-allergic COVID-19 children in disease incidence, clinical features, laboratory, and immunological findings. Allergy was not a risk factor for disease and severity of SARS-CoV-2 infection and did not significantly influence the disease course of COVID-19 in children.⁷

The immunology of COVID-19 was extensively reviewed in two articles from leading experts with a comprehensive discussion of the tip of the iceberg in COVID-19 epidemiology, anti-viral response, antibody response to SARS-CoV-2, acute phase reactants, cytokine storm, and pathogenesis of tissue injury and severity.^{8,9}

Two studies timely reported the role of possible trained immunity in countries with a Bacillus Calmette-Guérin (BCG) vaccination programme and a relatively low COVID-19 prevalence and mortality rate.^{10,11} In an extensive RNA sequencing analyses of SARS-CoV-2 receptor and their molecular partners revealed that ACE2 and TMPRSS2 were coexpressed at the epithelial sites of the lung and skin, whereas CD147 (BSG), cyclophilins (PPIA and PPIB), CD26 (DPP4) and related molecules were expressed in both, epithelium and in immune cells.¹²

Allergists, respiratory physicians, pediatricians, and other health care providers treating patients with allergic diseases are frequently in contact with patients potentially infected with SARS-CoV-2. Practical considerations and recommendations given by experts in the field of allergic diseases can provide useful recommendations for clinical daily work. Since the beginning

of this current pandemic, our journal has disseminated clinical reports,^{2,3,5,6,13} statements on the urgent need for accuracy in designing and reporting clinical trials in COVID-19,¹⁴ preventive measures,^{10,11,15} and Position Statements elaborated by experts in the field in close collaboration with the European Academy of Allergy and Clinical Immunology (EAACI) and its task force “Allergy and Its Impact on Asthma (ARIA)”.¹⁶⁻²⁸ (keynote information in table 1). A compendium answering 150 frequently encountered questions regarding COVID-19 and allergic diseases has been recently published by experts in their respective area (Figure 1).²⁹ In addition, readers can put further questions regarding this “living” compendium electronically to the authors and their answers will be available through a new category in the journal’s webpage.³⁰ Besides, EAACI in collaboration with ARIA, has provided recommendations on operational plans and practical procedures for ensuring optimal standards in the daily clinical care of patients with allergic diseases, whilst ensuring the safety of patients and healthcare workers.²³

Table 1: Examples of recently published recommendations, statements and Position Papers of the EAACI

Clinical Topic	Keynote message(s)	Title and Reference
Asthma	<ul style="list-style-type: none"> Prescribed inhaled (or oral) corticosteroids as controller medication should be further applied to patients with asthma as cessation may lead to asthma exacerbations. 	ARIA-EAACI statement on Asthma and COVID-19. ¹⁶
Allergic rhinitis	<ul style="list-style-type: none"> Intranasal corticosteroids (including nasal sprays) can be further applied in COVID-19 patients at the recommended dosages and a cessation or interruption is not recommended. Furthermore, a cessation or interruption of allergen immunotherapy (AIT) for both the subcutaneous route (SCIT) or sublingual route (SLIT) is not recommended in asymptomatic patients without suspicion for SARS-CoV-2 infection and/or contact to SARS-CoV-2 positive individuals or in other conditions as 	<p>corticosteroids in allergic rhinitis in COVID-19 infected patients: An ARIA-EAACI statement</p> <p>Intranasal corticosteroids in allergic rhinitis in COVID-19 infected patients: An ARIA-EAACI statement.²¹</p> <p>Handling of allergen immunotherapy in the COVID-19 pandemic: An ARIA-EAACI statement.²²</p>

	outlined.	
Biological-Treatment	<ul style="list-style-type: none"> · Biologicals should be applied in type 2 inflammations (asthma, atopic dermatitis, chronic rhinosinusitis with nasal polyps) via self-application in non-infected patients · In patient with an active SARS-CoV-2 infection treatment with biologicals should be paused and re-initiated after recovery and SARS-CoV-2 negativity of patients. 	Considerations on biologicals for patients with allergic disease in times of the COVID-19 pandemic: an EAACI Statement. ²⁸
Children	<ul style="list-style-type: none"> · Less severe courses of COVID-19 in the pediatric population have been reported. · Similar to the adult age group, severe forms of asthma and immunodeficiencies are reported to be risk factors. · A sufficient control of symptoms in children with allergies, asthma and immunodeficiency in accordance with current guidelines is key in the current pandemic. 	Managing childhood allergies and immunodeficiencies during respiratory virus epidemics - The 2020 COVID-19 pandemic: A statement from the EAACI-section on pediatrics. ²⁵
Drug allergy	<ul style="list-style-type: none"> · Drugs being used in different phases of COVID-19 disease seem to cause rare but potentially severe drug hypersensitivity reactions (DHRs). · Most of these DHRs affect the skin and are non-immediate. · The most important differential diagnosis of these DHRs is COVID-19 related exanthems. 	Diagnosis and management of the drug hypersensitivity reactions in coronavirus disease drug hypersensitivity. ²⁷
Ocular allergy	<ul style="list-style-type: none"> · An international survey revealed that experts in the field recommend that during the ongoing pandemic therapy strategies in line with current guidelines should be followed. · However, particular caution for the use of corticosteroids and immunomodulators in COVID-19 infected patients is expressed. 	Managing ocular allergy in the time of COVID-19. ¹⁹
Organization of an allergy clinic	<ul style="list-style-type: none"> · Recommendations on operational plans and procedures to maintain high standards in the daily clinical care of allergic patients whilst ensuring necessary safety in the current 	COVID-19 pandemic: Practical considerations on the organization of an allergy clinic - an EAACI/ARIA Position Paper. ²³

	COVID-19 pandemic should be followed. · Nine different scenarios and topics are discussed.	
--	---	--

Even though there has been a decrease in both infection rates and prevalence of COVID-19 in some parts of the world, in some countries the spread of COVID-19 has not been adequately controlled and the numbers are still rising. Epidemiologists have warned of possible subsequent waves until an effective vaccination becomes available. Until then, recommendations for different aspects of patient care are strictly needed, which underlines the importance of the COVID-19 series of publications in this journal. The recommendations provided in these articles will inform allergists and healthcare providers on the management and treatment of patients during this pandemic. It should be noted that these statements are based on current evidence and medical knowledge and there is a paucity of prospective clinical trials regarding some of the information provided in these documents. These clinical statements need to be regularly updated with incoming new clinical data.

Figure 1. Picture representing the wide range of clinical aspects involved in the pathogenesis of COVID-19.

References

1. Coronavirus disease (COVID-2019) situation reports.
<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>, Accessed on 16 Apr 2020.
2. Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy*. 2020.
3. Jesenak M, Banovcin P, Diamant Z. COVID-19, chronic inflammatory respiratory diseases and eosinophils - Observations from reported clinical case series. *Allergy*. 2020.
4. Zhang M, Guogang X, Fengming D, al. e. The role of peripheral blood eosinophil counts in COVID-19 patients. *Allergy*. 2020 (in press).

5. Dong X, Cao YY, Lu XX, et al. Eleven faces of coronavirus disease 2019. *Allergy*. 2020.
6. Zhang JJ, Cao YY, Dong X, et al. Distinct characteristics of COVID-19 patients with initial rRT-PCR-positive and rRT-PCR-negative results for SARS-CoV-2. *Allergy*. 2020.
7. Du H, Dong X, Zhang JJ, et al. Clinical characteristics of 182 pediatric COVID-19 patients with different severities and allergic status. *Allergy*. 2020.
8. Sokolowska M, Agache I, Akdis CA. Immune response in Covid-19: mechanisms, clinical outcome, diagnostics and perspectives – an EAACI report. *Allergy*. 2020 (in press).
9. Azkur AK, Akdis M, Azkur D, et al. Immune response to SARS-CoV-2 and mechanisms of immunopathological changes in COVID-19. *Allergy*. 2020.
10. Gursel M, Gursel I. Is global BCG vaccination-induced trained immunity relevant to the progression of SARS-CoV-2 pandemic? *Allergy*. 2020.
11. Ozdemir C, Kucuksezer UC, Tamay ZU. Is BCG vaccination affecting the spread and severity of COVID-19? *Allergy*. 2020.
12. Radzikowska U, Ding M, Tan G, et al. Distribution of ACE2, CD147, CD26 and other SARS-CoV-2 associated molecules in tissues and immune cells in health and in asthma, COPD, obesity, hypertension, and COVID-19 risk factors. *Allergy*. 2020.
13. Cai Q, Huang D, Ou P, et al. COVID-19 in a designated infectious diseases hospital outside Hubei Province, China. *Allergy*. 2020.
14. Bonini S, Maltese G. COVID-19 Clinical trials: quality matters more than quantity. *Allergy*. 2020.
15. Sotgiu G, Gerli AG, Centanni S, et al. Advanced forecasting of SARS-CoV-2-related deaths in Italy, Germany, Spain, and New York State. *Allergy*. 2020.
16. Bousquet J, Jutel M, al. ACe. ARIA-EAACI statement on Asthma and COVID-19. *Allergy* 2020 (submitted).
17. Carli G, Cecchi L, Stebbing J, Parronchi P, Farsi A. Is asthma protective against COVID-19? *Allergy*. 2020.
18. Johnston SL. Asthma and COVID-19: is asthma a risk factor for severe outcomes? *Allergy* 2020 (in press).
19. Leonardi A, Fauquert JL, Doan S, et al. Managing ocular allergy in the time of COVID-19. *Allergy*. 2020.

20. Licari A, Votto M, Brambilla I, et al. Allergy and asthma in children and adolescents during the COVID outbreak: What we know and how we could prevent allergy and asthma flares. *Allergy*. 2020.
21. Bousquet J, Akdis C, Jutel M, et al. Intranasal corticosteroids in allergic rhinitis in COVID-19 infected patients: An ARIA-EAACI statement. *Allergy*. 2020.
22. Klimek L, Jutel M, Akdis C, et al. Handling of allergen immunotherapy in the COVID-19 pandemic: An ARIA-EAACI statement. *Allergy*. 2020.
23. Pfaar O, Klimek L, Jutel M, et al. COVID-19 pandemic: Practical considerations on the organization of an allergy clinic - an EAACI/ARIA Position Paper. *Allergy*. 2020.
24. Klimek L, Jutel M, Bousquet J, al. e. Treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in the COVID-19 pandemic – An EAACI Position Paper. *Allergy* 2020 (submitted).
25. Brough HA, Kalayci O, Sediva A, et al. Managing childhood allergies and immunodeficiencies during respiratory virus epidemics - The 2020 COVID-19 pandemic: A statement from the EAACI-section on pediatrics. *Pediatr Allergy Immunol*. 2020.
26. Untersmayr E, al. e. Immunology of COVID-19: mechanisms, clinical outcome, diagnostics and perspectives – a report of the European Academy of Allergy and Clinical Immunology (EAACI). *Allergy*. 2020.
27. Gelincik A, Brockow K, Celik GE, et al. Diagnosis and management of the drug hypersensitivity reactions in Coronavirus disease 19. *Allergy*. 2020.
28. Vultaggio A, Agache I, Akdis CA, et al. Considerations on Biologicals for Patients with allergic disease in times of the COVID-19 pandemic: an EAACI Statement. *Allergy*. 2020.
29. Riggioni C, Comberiati P, Giovannini M, et al. A compendium answering 150 questions on COVID-19 and SARS-CoV-2. *Allergy*. 2020.
30. Allergy homepage. [https://onlinelibrary.wiley.com/doi/toc/10.1111/\(ISSN\)1398-9995.COVID-19](https://onlinelibrary.wiley.com/doi/toc/10.1111/(ISSN)1398-9995.COVID-19). Accessed on 16 Jun 2020.

Disclosure of conflict of interests:

Dr. Pfaar reports grants and/or personal fees from ALK-Abelló, Allergopharma, Stallergenes Greer, HAL Allergy Holding B.V./HAL Allergie GmbH, Bencard Allergie GmbH/Allergy Therapeutics, Lofarma, Biomay, Circassia, ASIT Biotech Tools S.A., Laboratorios LETI/LETI Pharma, MEDA Pharma/MYLAN, Anergis S.A., Mobile Chamber Experts (a GA2LEN Partner), Indoor Biotechnologies, GSK, Astellas Pharma Global, EUFOREA, ROXALL, NOVARTIS, SANOFI AVENTIS, Med Update Europe GmbH, and streamedup! GmbH. Dr. Torres has no conflict of interest in relation to this work. Dr. Akdis reports grants from Allergopharma, Idorsia, Swiss National Science Foundation, Christine Kühne-Center for Allergy Research and Education, European Commission's Horizon's 2020 Framework Programme, Cure, Novartis Research Institutes, AstraZeneca, SciBase, GSK, and other from Sanofi and Regeneron.

This article is protected by copyright. All rights reserved