**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

**Delayed infection of a pelvic lymphocele following robotic radical
prostatectomy and pelvic lymphadenectomy: two cases**

Tremp, M ; Sulser, T ; Seifert, H H

DOI: <https://doi.org/10.1159/000251192>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-20482>

Journal Article

Accepted Version

Originally published at:

Tremp, M; Sulser, T; Seifert, H H (2009). Delayed infection of a pelvic lymphocele following robotic radical prostatectomy and pelvic lymphadenectomy: two cases. *Urologia internationalis*, 83(4):479-481.

DOI: <https://doi.org/10.1159/000251192>

Delayed infection of a pelvic lymphocele following robotic radical prostatectomy and pelvic lymphadenectomy: two cases

Mathias Tremp, MD¹, Tullio Sulser, MD, Hans-Helge Seifert, MD

¹ Corresponding author.

Department of Urology, University Hospital

Frauenklinikstrasse 10, 8091 Zürich, Switzerland

Tel. +41 44 255 54 07

e-mail: mathias.tremp@usz.ch

Key words: Infection, Lymphocele, Laparoscopy, Robotic surgery, Prostatic neoplasm

Abstract

Pelvic lymphocele is an infrequent complication of pelvic surgery, usually presenting shortly after surgery. We report two cases with a delayed infected pelvic lymphocele presenting after transperitoneal pelvic lymphadenectomy and robotic radical prostatectomy for adenocarcinoma of the prostate.

These cases illustrate that late infection of pelvic lymphoceles may occur following radical prostatectomy and pelvic lymphadenectomy. The practicing urologist should be aware of this possibility and look for an infected lymphocele in postoperative pelvic lymphadenectomy patients presenting with fever and leukocytosis of uncertain etiology, regardless of the time elapsed since surgery.

To date, there is a paucity of data in the literature on robotic-assisted laparoscopic resection of a lymphocele after radical prostatectomy. The minimally invasive technique can be considered as a possible alternative to lymphocele percutaneous drainage. It is effective, results in minimal patient morbidity and allows for a rapid recovery.

Introduction

Pelvic lymphocele has long been recognized as an infrequent [1] complication following pelvic lymphadenectomy for adenocarcinoma of the prostate. The incidence of symptomatic lymphocele following simultaneous pelvic lymph node dissection (PLND) is between 3 and 14% depending on the extent of lymph node dissection and the operating surgeon.[1-3] Lymphocele development is a problem for the patient when it leads to sequelae relevant to health. In addition to secondary infection of the lymphocele, these sequelae comprise mainly thromboembolic events due to compression of the pelvic vessels. Another relevant consequence of lymphoceles is the significantly higher incidence of re-intervention [3] We report on the development and treatment of two patients with delayed infected lymphoceles after a transperitoneal robotic-assisted laparoscopic radical prostatectomy and extended pelvic lymph node dissection.

Case 1

A 70-year-old caucasian evaluated with transrectal ultrasonography and prostate biopsy for an elevated prostate-specific antigen level of 4,3 µg/l was diagnosed with Gleason score 3 + 4 adenocarcinoma of the prostate, clinical stage T1c. Following an uncomplicated transperitoneal non-nerve-sparing robotic radical prostatectomy with pelvic lymphadenectomy, the pathologic examination revealed pT3b and all 15 removed lymph nodes were negative for metastasis. No drains were placed during the operation. Patient follow-up was performed every 3 months.

The patient remained in a state of good health until 20 months postoperatively, when he began experiencing symptoms of malaise, intermittent fever of 39°C, and progressive right lower abdominal pain.

On physical examination, his rectal examination revealed no evidence of recurrent cancer, and there were no fluctuant or tender areas. Laboratory data revealed a white blood cell count of

17.4 x 10³/mm³ (normal, 3 to 9.6 x 10³/mm³), hematocrit of 35.6% (normal, 40 to 50%), and normal serum electrolyte values, including a creatinine of 78 µmol/l (normal 62 – 106 µmol/l). Urinalysis was unremarkable. Blood cultures and urine culture showed no bacterial growth.

Given his continued febrile state, an abdominal and pelvic computed tomography (CT) scan was obtained (Fig. 1), revealing a 8.5 x 6.7cm low attenuation lesion over the right iliopsoas muscle with a thick capsule and infiltration of the surrounding soft tissue, consistent with an infected lymphocele. Because of its low morbidity and feasibility, we decided to perform an ultrasound-guided percutaneous drainage of the fluid collection with aspiration of a purulent fluid. Culture of the fluid showed profuse gram positive bacteria (*S. aureus*), sensible to amoxicillin/clavulan acid. An 8.5 F pigtail catheter was placed for drainage and the patient began a course of amoxicillin/clavulan acid. He immediately defervesced, his white blood cell count became normal to 6.06 x 10³/mm³ and he was discharged home after 9 days on oral amoxicillin/clavulan acid. His drainage gradually decreased and repeat sonogram showed no significant fluid collection. The drain was removed uneventfully after 12 days. A CT-scan after 4 weeks showed an almost complete resolution of the lymphocele (1 x 4.5 cm) (Figure 2) so robotic-assisted laparoscopic resection of the lymphocele was not necessary.

Case 2

In August 2006, a 79-year-old patient with a prostate-specific antigen level of 1.96 µg/l and a Gleason score 3 + 4 adenocarcinoma of the prostate underwent a transperitoneal robotic radical prostatectomy and pelvic lymphadenectomy. Pathologic examination revealed pT3a and all 5 removed lymph nodes were negative for metastases. In April 2007, the patient began experiencing symptoms of progressive left lower abdominal pain, pelvic fullness, abdominal distension with edema and paresthesia on his abdomen and left leg. Rectal examination revealed no evidence of recurrent cancer. He had no fever; laboratory data revealed a white

blood cell count of $9.49 \times 10^3/\text{mm}^3$, hematocrit 32.8%, and normal serum electrolyte values, including a creatinine of 77 $\mu\text{mol/l}$. Urinalysis was unremarkable. An abdominal ultrasound was obtained, revealing a 5.1 x 5.7 x 5.2 cm low attenuation lesion with a thick capsule in the paravesical fossa and infiltration of the surrounding soft tissue. No hydronephrosis was detected. However, in the CT-scan metastasis was not excludable. Bone scintigraphy showed no bone metastasis.

An attempt at percutaneous ultrasound-guided drainage was not diagnostic. Therefore, we decided to treat the patient with a robotic-assisted laparoscopic resection of the lesion. Intraoperatively and pathologically, the lesion has been confirmed to be a lymphocele. Culture of the fluid showed profuse gram positive bacteria (*S. aureus*), sensible to amoxicillin/clavulan acid. The patient recovered uneventfully after the operation and his symptoms dissolved rapidly.

Comment

The cases presented are unusual because the patients' pelvic lymphoceles became infected more than 6 months after transperitoneal robotic radical prostatectomy and pelvic lymphadenectomy. A careful review of the literature revealed only 3 other reported cases of delayed infected lymphocele after surgery[4-6] Two of the reported cases had an infected lymphocele more than one year after surgery.[5,6] In all cases, the patients presented with chills and fever of unknown etiology. In our cases, the patients had an existing lymphocele seeded with a known skin commensal, *S. aureus*. However, the patients had no skin lesions or disruptions to explain the entry of *S. aureus* into his lymphatic system.

Pepper et al. showed the rate of symptomatic lymphocele formation was low after retropubic radical prostatectomy, with an overall incidence of 3.5%.[7] In addition, Solberg et al. showed that laparoscopic lymph node dissection was associated with a statistically significant lower

frequency of lymphocele formation (37%) compared to open pelvic lymph node dissection (61%).[8]

In their series of 99 patients after robotic-assisted laparoscopic extended pelvic lymph node dissection for prostate cancer, Feicke et al. reported symptomatic lymphocele in 5 patients (5%), although they placed no drains during the operation. However, lymphoceles needed to be drained percutaneously only in two patients (2%).[9] The median number of lymph nodes harvested was 19 (range: 8–53).[9] Pelvic drain placement after prostatectomy has been discussed not only to prevent lymphocele formation but also to prevent urinoma formation and postoperative hematoma. Two recently published articles addressed the issue of pelvic drainage after prostatectomy, concluding that pelvic drainage can be omitted in up to 90% of robotic assisted prostatectomies.[10,11] Feicke et al. assessed the integrity of the vesicourethral anastomosis intraoperatively in all patients of their series with a bladder filling with 50-100 ml saline. As no leakage was observed, they chose not to place a pelvic drain.[9] Indications on placement of a pelvic drain are rectal injury, urinary bladder injury, fish mouth bladder neck deformity, tension at the urethrovesical anastomosis, non-water-tight anastomosis, inadequate homeostasis or a large median lobe.[10,11]

The decision to treat postoperative lymphoceles has traditionally depended on associated symptoms and the patient's overall clinical status. Specifically, asymptomatic pelvic lymphoceles without mass effect on nearby organs warrant no intervention.

Aspiration as definitive treatment has been discouraged because rapid reaccumulation usually occurs.[12] Finally, in patients with giant lymphoceles not amenable to external drainage, laparoscopic unroofing or open surgical drainage of infected lymphocele has been described.[3]

Conclusion

Our cases illustrate that late infection of asymptomatic pelvic lymphoceles can occur. Robotic-assisted laparoscopic resection is a minimally invasive technique and can be considered as a possible alternative to lymphocele percutaneous drainage. It is effective, results in minimal patient morbidity and allows for a rapid recovery.

References

- 1 Novicki DE, Ferrigni RG: Non-malignant processes of the lymphatic system of clinical importance to the urologist. AUA Update Series VI 1987;22
- 2 Stolzenburg JU, Wasserscheid J, Rabenalt R, Do M, Schwalenberg T, McNeill A, Constantinides C, Kallidonis P, Ganzer R, Liatsikos E: Reduction in incidence of lymphocele following extraperitoneal radical prostatectomy and pelvic lymph node dissection by bilateral peritoneal fenestration. World J Urol 2008;26:581-586.
- 3 Musch M, Klevecka V, Roggenbuck U, Kroepfl D: Complications of pelvic lymphadenectomy in 1,380 patients undergoing radical retropubic prostatectomy between 1993 and 2006. The Journal of Urology 2008;179:923-929.
- 4 Nakano M, Miwa K, Kanimoto Y: A case of delayed infection of a pelvic lymphocele following radical prostatectomy 4 months after operation. Hinyokika Kiyo, 2003 Jul; 49(7):419-21
- 5 Spring DB, Schroeder D, Babu S, Agee R, Gooding GA: Ultrasonic evaluation of lymphocele formation after staging lymphadenectomy for prostatic carcinoma. Radiology 1981;141:479-483.
- 6 Carbone JM, Nadler RB, Bullock AD, Basler JW: Delayed infection of a pelvic lymphocele following pelvic lymphadenectomy. Urology 1996;47:140-142.
- 7 Pepper RJ, Pati J, Kaisary AV: The incidence and treatment of lymphoceles after radical retropubic prostatectomy. BJU Int 2005;95:772-775.
- 8 Solberg A, Angelsen A, Bergan U, Haugen OA, Viset T, Klepp O: Frequency of lymphoceles after open and laparoscopic pelvic lymph node dissection in patients with prostate cancer. Scand J Urol Nephrol 2003;37:218-221.
- 9 Feicke A, Baumgartner M, Talimi S, Schmid DM, Seifert H-H, Muntener M, Fatzer M, Sulser T, Strebel RT: Robotic-assisted laparoscopic extended pelvic lymph node dissection for prostate cancer: Surgical technique and experience with the first 99 cases. European Urology 2009;55:876-884.
- 10 Sharma S, Kim HL, Mohler JL: Routine pelvic drainage not required after open or robotic radical prostatectomy. Urology 2007;69:330-333.
- 11 Araki M, Manoharan M, Vyas S, Nieder AM, Soloway MS: A pelvic drain can often be avoided after radical retropubic prostatectomy--an update in 552 cases. European Urology 2006;50:1241-1247.
- 12 Kay R, Fuchs E, Barry JM: Management of postoperative pelvic lymphoceles. Urology 1980;15:345-347.

Figure 1 – Computed tomography of patient one shows a thick-walled right pelvic obturator fossa lymphocele (arrow), 20 months after robotic radical prostatectomy and pelvic lymphadenectomy

**Figure 2 – CT of patient one shows almost complete resolution of lymphocele (arrow)
after 4 weeks**