



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Cardiovascular and Antiobesity Effects of Resveratrol Mediated through the Gut Microbiota

Bird, Julia K ; Raederstorff, Daniel ; Weber, Peter ; Steinert, Robert E

DOI: <https://doi.org/10.3945/an.117.016568>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-207547>

Journal Article

Published Version

Originally published at:

Bird, Julia K; Raederstorff, Daniel; Weber, Peter; Steinert, Robert E (2017). Cardiovascular and Antiobesity Effects of Resveratrol Mediated through the Gut Microbiota. *Advances in Nutrition*, 8(6):839-849.

DOI: <https://doi.org/10.3945/an.117.016568>

Cardiovascular and Antiobesity Effects of Resveratrol Mediated through the Gut Microbiota

Julia K Bird,¹ Daniel Raederstorff,¹ Peter Weber,¹ and Robert E Steinert^{1,2}

¹Human Nutrition and Health, DSM Nutritional Products, Basel, Switzerland, and ²Department of Surgery, Division of Visceral and Transplantation Surgery, University Hospital Zürich, Zürich, Switzerland

ABSTRACT

Encouraging scientific research into the health effects of dietary bioactive resveratrol has been confounded by its rapid first-pass metabolism, which leads to low in vivo bioavailability. Preliminary studies have shown that resveratrol can modulate gut microbiota composition, undergo biotransformation to active metabolites via the intestinal microbiota, or affect gut barrier function. In rodents, resveratrol can modify the relative Bacteroidetes:Firmicutes ratio and reverse the gut microbial dysbiosis caused by a high-fat diet. By upregulating the expression of genes involved in maintaining tight junctions between intestinal cells, resveratrol contributes to gut barrier integrity. The composition of the gut microbiome and rapid metabolism of resveratrol determines the production of resveratrol metabolites, which are found at greater concentrations in humans after ingestion than their parent molecule and can have similar biological effects. Resveratrol may affect cardiovascular risk factors such as elevated blood cholesterol or trimethylamine *N*-oxide concentrations. Modulating the composition of the gut microbiota by resveratrol may affect central energy metabolism and modify concentrations of satiety hormones to produce antiobesity effects. Encouraging research from animal models could be tested in humans. *Adv Nutr* 2017;8:839–49.

Keywords: resveratrol, microbiome, obesity, cardiovascular disease, inflammation, gut barrier integrity

Introduction

Resveratrol is a plant-derived stilbenoid with antifungal and antibacterial properties. It is found in the human diet from various foods such as grapes and the wine produced therefrom, berries such as cranberries and red currants, and peanut skins (1). Interest in the health-promoting properties of resveratrol has been greatly influenced by experiments that have shown an increase in the lifespan of simple organisms by 1) stimulating the activity of enzymes such as AMP-activated protein kinase and the sirtuins that are also activated by calorie restriction (2) and 2) putatively affecting metabolic pathways that modulate molecular damage (3). Interestingly, resveratrol has also been shown to protect against metabolic disturbances induced by a high-fat diet and shows potential in reversing the negative effects of the metabolic syndrome in rodent models (4, 5). A series of

landmark studies has also shown strong cancer chemopreventive activity through the inhibition of cyclooxygenase and anti-inflammatory actions (6–8). The latter has been postulated as the mechanism whereby moderate wine consumption is associated with cardioprotection and other beneficial effects on health, commonly known as the French paradox, a term that refers to the relatively low incidence of coronary artery disease in France despite a high consumption of saturated fats (9, 10).

Resveratrol intakes from the diet are low compared with other polyphenols (11). A study that pooled resveratrol intakes from France, Germany, Greece, and Italy estimated median intakes of resveratrol and its main glucoside to be 100 $\mu\text{g}/\text{d}$ (12), which is similar to median intakes estimated in an older population in the Netherlands [123 $\mu\text{g}/\text{d}$ (13)] and somewhat lower than was found in another older population located in a wine-growing municipality in Chianti, Italy [500 $\mu\text{g}/\text{d}$ (14)]. In addition to these low dietary intakes, resveratrol is poorly bioavailable, either when imbibed as red wine (15) or consumed as food or a dietary supplement (16). In fact, when administered orally, only ~1–8% of free resveratrol is found in serum samples, whereas 25% is excreted without absorption, and the rest (generally

The authors reported no funding received for this study.

Author disclosures: JKB, DR, PW, and RES are employees of DSM Nutritional Products, a supplier of nutritional ingredients, including resveratrol, for use in food, dietary supplements, pharmaceutical applications, and animal feed.

Address correspondence to JKB (e-mail: julia.bird@dsm.com).

Abbreviations used: EGCG, epigallocatechin gallate; GLP-1, glucagon-like peptide 1; HFS, high-fat and high-sucrose; mTOR, mechanistic target of rapamycin; TMAO, trimethylamine *N*-oxide.

>70%) is extensively metabolized by the liver, intestinal tract, and gut microbiota (16). A schematic description of the metabolism and disposition of resveratrol is shown in **Figure 1**. Hence, plasma concentrations of free resveratrol are considerably lower than its metabolites, which calls into question whether results obtained from in vitro experiments with the use of resveratrol can be used to elucidate the mechanisms behind improvements in human health.

With the increased interest in the role of the gut microbiome in the maintenance of human health and the pathogenesis of many diseases, several mechanisms have been proposed by which resveratrol can exert its health effects through modulating the composition of the gut microbiota (17, 18). In addition, direct effects of resveratrol and its metabolites on gut integrity and barrier function have been suggested (17). Broad-spectrum antibacterial activity of resveratrol has been found in several clinically important bacterial species, including opportunistic pathogens of the digestive tract: *Escherichia coli*, *Enterococcus faecalis*, and *Salmonella enterica* subsp. *enterica* (19). The bactericidal effect of resveratrol is, however, differential, with much lower activity seen against commensal bacterial species such as *Lactobacillus* spp. (20). Resveratrol may thus be able to affect the composition of bacterial species in the gut to one that is conducive to human health. In addition, whereas high intersubject variability of resveratrol absorption, metabolism, and excretion determines the pharmacokinetic characteristics of resveratrol (21), the gut microbiota that is capable of metabolizing resveratrol further modulates the fate and physiologic effects of resveratrol. Preliminary studies have shown that gut microorganisms affect the pattern of resveratrol metabolites produced; for example, the Coriobacteriaceae family of Actinobacteria is capable of metabolizing resveratrol to dihydroresveratrol, as may *Bacteroides* spp. (21). The complex metabolic activity of microbes in the large intestine may thus be responsible for the variation in resveratrol metabolite concentrations between individuals and ultimately for its effects on clinically important endpoints, because potent antioxidant and proliferative effects similar to those found with resveratrol have also been seen with its major metabolites (22).

The aim of this review is to briefly summarize the current evidence for the cardiovascular and antiobesity effects of resveratrol that are mediated through the gut microbiome. As a basis for our review, we searched the PubMed database with the use of the keywords “resveratrol gut microbiota” to identify primary research articles that have investigated the effect of resveratrol on the gut microbiome, and we support these articles with relevant background citations. We focus on pre-clinical and clinical evidence and discuss the mechanisms by which an interaction between resveratrol and the gut microbiome may affect clinically relevant outcomes.

Current Status of Knowledge

Metabolism and disposition of resveratrol in humans

Poor water solubility is likely responsible for the low oral bioavailability of resveratrol in vivo (23). Walle et al. (24)

showed that >70% of an oral dose of 25 mg was absorbed, but only 1–8% of total resveratrol in plasma was found as free resveratrol because of the rapid metabolism of resveratrol, mainly to glucuronides and sulfate conjugates (24–26). There is a trend to a higher proportion of free resveratrol in plasma as doses increase (25, 26). The recovery of ingested resveratrol has been shown to be highly variable: 55–85% of the 25-mg oral dose in Walle et al. (24) was recovered in the urine and 0–40% in the feces. Interestingly, ~10% of an intravenous dose was also recovered in the feces, indicating that resveratrol metabolites in the circulation are excreted back into the intestines (24), presumably via biliary secretion (23). In vitro fecal fermentation experiments have confirmed that 18.3 mg resveratrol/L, a much higher concentration than obtainable by the diet, can be almost completely degraded to metabolites by the gut microbiome of healthy volunteers within 24 h, with the main metabolites, including lunularin, dihydroresveratrol, and 3,4-dihydroxy-trans-stilbene, produced by the gut (21).

Proposed mechanisms of action of resveratrol on the gut microbiome

Modulation of the composition of the gut microbiota

Resveratrol has the potential to affect the growth of species within the gut microbiota and alter its composition, as shown by changes caused by resveratrol to bacterial monocultures or to the relative abundance of particular microorganisms within complex communities (**Table 1**). In vitro bacterial growth has been shown to be slowed by the addition of resveratrol at concentrations of 50 mg/L for most of a panel of 43 human- and animal-origin bacteria, whereas growth was stimulated for one strain of *Lactobacillus acidophilus* (27). In the same series of experiments, the motility of the human pathogen *Proteus mirabilis* was inhibited, which potentially curtailed its swarming behavior, and 25% of the strains were able to transform resveratrol to several metabolites, including resveratrolsides, piceid, and dihydroresveratrol (27). The doses used in this study were several orders of magnitude higher than normal dietary intakes; however, the exposure of the gut microbiome to the concentration used is possible with dietary supplementation. In an in vitro fermentation model, a stilbene extract from grapes containing resveratrol affected microbial metabolism and composition. The stilbene extract containing 0.76 and 1.0 g resveratrol/L, a pharmacologic dose, decreased the production of NH_4^+ and SCFAs; Enterobacteriaceae numbers increased, whereas *Bacteroides* decreased, and there was no apparent effect on lactobacilli or clostridia (28). Taken together, the in vitro studies indicate that resveratrol has a broad antibacterial activity at concentrations that can be reached with dietary supplements. Moreover, it appears that stilbenes inhibit bacteria involved in saccharolytic and proteolytic activities, whereas gram-negative species are less sensitive to the antimicrobial effects of stilbenes.

In vivo work in experimental mouse and rat models has also demonstrated shifts in microbial populations in




FIGURE 1 The metabolism and disposition of RES. RES is mainly absorbed in the small intestine after oral intake. It is then quickly and extensively metabolized to RES-GLU and RES-SUL forms in the intestine and the liver. The RES-GLU and RES-SUL forms are excreted by the kidney in the urine. In the human body, sulfatases and glucuronidases can convert the conjugated forms of RES back to the free RES form. Enterohepatic recycling also occurs by biliary excretion and intestinal reabsorption. Moreover, in the intestine, unabsorbed RES is converted to microbial metabolites—for example, DHR—which can also be further metabolized to DHR-GLU and DHR-SUL derivatives.

Finally, the unabsorbed RES and its metabolites are excreted in the feces. DHR, dihydroresveratrol; GLU, glucuronide; RES, resveratrol; SUL, sulfate.

response to resveratrol supplementation (Table 1). A study conducted in outbred mice showed that 4 g resveratrol/kg caused gut microbiota remodeling with an increase in the relative abundance of *Bacteroides*, *Lactobacillus*, *Bifidobacterium*, and *Akkermansia*, and decreases in *Prevotella*, Ruminococcaceae, *Anaerotruncus*, *Alistipes*, *Helicobacter*, and Peptococcaceae (29). Moreover, at the phylum level, resveratrol substantially increased Bacteroidetes:Firmicutes ratios in both C57BL/6J and apoE^{-/-} mice. Similarly, in mice with surgically induced heart failure, 4 g resveratrol/kg increased the relative abundance of *Parabacteroides*, *Bilophila*, and *Akkermansia*, and decreased the relative abundance of Lachnospiraceae; moreover, the Bacteroidetes:Firmicutes ratio increased (30). In several studies in mice, the combined effect of an obesogenic diet and resveratrol has been investigated. The combination of a high-fat diet and resveratrol on the Bacteroidetes:Firmicutes ratio [decreases that are associated with obesity development (38, 39) and occur in response to a high-fat diet] was intermediate between the high-fat diet and the control diet in one study: 200 mg resveratrol · kg⁻¹ · d⁻¹ could partly mitigate the effects of a high-fat diet on this ratio (31). Substantial changes in the composition of the gut microbiota upon 60 mg resveratrol · kg⁻¹ · d⁻¹ supplementation were also observed in mice fed a high-fat diet compared with a control diet, and resveratrol normalized changes in the microbiota induced by the high-fat diet, although in this case resveratrol inhibited the growth of *Bacteroides vulgatus* as well as *Parabacteroides johnsonii* and *Alistipes putredinis* to the extent that these species were no longer detectable by denaturing gradient gel electrophoresis after supplementation (32). In one study in mice with the use of a 2 × 2 factorial design [standard diet, standard diet and resveratrol, high-fat and high-sucrose

(HFS) diet, and HFS diet and resveratrol], 4 g resveratrol/kg affected the gut microbiome only in mice fed the obesogenic diet; resveratrol had no effect in mice fed the standard diet (34). In particular, resveratrol treatment increased the Bacteroidetes:Firmicutes ratio, decreased the relative abundance of Turicibacteraceae, *Moryella*, Lachnospiraceae, and *Akkermansia*, and increased *Bacteroides* and *Parabacteroides* in the HFS diet group only (34). In yet another mouse study, similar results were found: 200 mg resveratrol · kg⁻¹ · d⁻¹ had marked effects on the composition of the gut microbiota and ameliorated changes caused by a high-fat diet (33). The increase in *Lactococcus*, *Clostridium*, *Oscillibacter*, *Hydrogenoaerobacterium*, and *Flavonifractor* induced by a high-fat diet was partially attenuated by resveratrol treatment in this study.

These results were also confirmed in experiments in rats, in which 1 mg resveratrol · kg⁻¹ · d⁻¹ increased *Lactobacillus* and *Bifidobacterium* while diminishing the increase of Enterobacteriaceae upon dextran sulfate sodium-induced colitis (35). On the other hand, a study in rats fed an HFS diet showed no phylum-level changes caused by 15 mg resveratrol · kg⁻¹ · d⁻¹, but Gracilibacteraceae (Firmicutes) concentrations were lower than in HFS-only rats, and *Parabacteroides* and various *Clostridium* spp. as well as *Gracilibacter thermotolerans* and *Parabacteroides distasonis* were also suppressed by resveratrol compared with the control diet (36).

Some preliminary research on changes in urinary metabolites derived from the gut microbiota after resveratrol supplementation has been conducted in humans (40). Specifically, concentrations of tryptamine and tyramine from amino acid metabolism were lower after 1000 mg resveratrol/d supplementation in 24 middle-aged men with metabolic syndrome. Lower concentrations of tryptamine may reflect

TABLE 1 Effect of resveratrol on gut microbial populations in *in vitro* and in rodent models¹

Model/system	Treatment	Effect relative to control									
		Lactic acid bacteria	Bifidobacteria	Enterococcus faecalis	Bacteroides spp.	Akkermansia spp.	Prevotella spp.	Ruminococcaceae	Firmicutes	Reference	
In vitro monoculture panel	Control: liquid culture medium	↑ ²	—	↓	↓	—	—	↓	↓	—	27
	Intervention: 50 mg resveratrol/L	—	—	—	—	—	—	—	—	—	—
In vitro mixed-culture fermentation model	Control: liquid culture medium	↔	—	—	↓	—	—	—	—	↔	28
	Intervention 1: 0.76 g resveratrol/L (stilbenes: 1.5 g/L)	↔	—	—	↓	—	—	—	—	↔	—
	Intervention 2: 1.0 g resveratrol/L (stilbenes: 2 g/L)	↔	—	—	↓	—	—	—	—	↔	—
	Control: standard diet	↑	↑	—	↑	↑	↓	↓	↓	↓	29
Mouse	Intervention 1: 4 g resveratrol/kg in C57BL/6J mice	↑	↑	—	↑	↑	↓	↓	↓	↓	—
	Intervention 2: 4 g resveratrol/kg in apoE ^{-/-} mice	↑	↑	—	↑	↑	↓	↓	↓	↓	—
Mouse	Control: standard diet	—	—	—	↑	↑	—	—	—	—	30
	Intervention: 450 mg resveratrol · kg ⁻¹ · d ⁻¹	—	—	—	↑	↑	—	—	—	—	—
Mouse	Control: standard diet	↓	↓↓	↑	↓	—	—	—	—	—	31
	Intervention 1: high-fat diet	↔	—	↔	↔	—	—	—	—	—	—
Mouse	Intervention 2: resveratrol and high-fat diet (200 mg · kg ⁻¹ · d ⁻¹)	—	—	—	↓	—	—	—	—	—	32
	Control: high-fat diet	—	—	—	↓	—	—	—	—	—	—
Mouse	Intervention: resveratrol and high-fat diet (60 mg · kg ⁻¹ · d ⁻¹)	↔	—	—	—	—	—	—	—	↔	33
	Control 1: standard diet	—	—	—	—	—	—	—	—	—	—
Mouse	Intervention 1: 200 mg resveratrol · kg ⁻¹ · d ⁻¹	↓	—	—	—	—	—	—	—	↓	—
	Control 2: high-fat diet	—	—	—	—	—	—	—	—	—	—
Mouse	Intervention 2: resveratrol and high-fat diet (200 mg · kg ⁻¹ · d ⁻¹)	—	—	—	↔	↔	↔	↔	↔	↔	34
	Control 1: standard diet	—	—	—	↔	↔	↔	↔	↔	↔	—
Mouse	Intervention 1: 4 g resveratrol/kg	—	—	—	↔	↔	↔	↔	↔	↔	—
	Control 2: HFS diet	—	—	—	↑	↑	—	—	—	—	—
Rat	Intervention 3: HFS and resveratrol (diet containing 4 g/kg)	—	—	—	—	—	—	—	—	—	—
	Control: standard diet	—	—	—	—	—	—	—	—	—	35
Rat	Intervention 1: DSS-induced colitis model	↑	↑	↑	—	—	—	—	—	—	—
	Intervention 2: DSS-induced colitis model and 1 mg resveratrol · kg ⁻¹ · d ⁻¹	↑	↑	↔	—	—	—	—	—	—	—

(Continued)

TABLE 1 (Continued)

Model/system	Treatment	Effect relative to control										Reference
		Lactic acid bacteria	Bifidobacteria	Enterococcus faecalis	Bacteroides spp.	Akkermansia spp.	Prevotella spp.	Ruminococcaceae	Firmicutes			
Rat	Control: high-fat sucrose diet Intervention: HFS and resveratrol (diet containing 15 mg · kg ⁻¹ · d ⁻¹)	↔	—	—	—	↔	↔	↔	↔	↓		36
Human	Control: placebo (microcrystalline cellulose) Intervention: 282 mg EGCG and 80 mg resveratrol/d	—	—	—	↓ ³	↔	—	—	↔			37

¹ DSS, dextran sulfate sodium; EGCG, epigallocatechin gallate; HFS, high-fat and high-sucrose; ↑, significant increase; ↓, significant decrease; ↔, no significant difference compared with control.

² Several strains were tested with divergent results.

³ Men only.

a relative decrease in Firmicutes and therefore a reduction in the functional ability of the gut microbiome to produce tryptophan decarboxylases (41). Likewise, resveratrol treatment may inhibit the increase in *E. faecalis* occurring during a high-fat diet or in inflammatory states in rodent models, thereby reducing the activity of tyrosine decarboxylase in producing tyramine (42). Although the composition of the gut microbiome was not measured directly in the human study, substantial changes in the production of microbiome-associated metabolites mirrored trends found in animal studies in which resveratrol was found to modulate the gut microbiota (40).

The effect of a resveratrol supplement in combination with the dietary polyphenol epigallocatechin gallate (EGCG) on the gut microbiota has been investigated in a single clinical trial conducted in 37 healthy untrained overweight and obese adults (37). Dietary supplements containing 141 mg EGCG and 40 mg resveratrol were consumed twice daily for 12 wk. At baseline, there were significant differences in the microbiome of men compared with women, which indicates that sex should be controlled for in future human studies in this area. The intervention substantially reduced the abundance of Bacteroidetes in men but not in women. The abundance of other bacterial phyla and species, including Firmicutes, Actinobacteria, γ -Proteobacteria, *Akkermansia muciniphila* (phylum Verrucomicrobiae), sulfate-reducing bacteria, acetogenic bacteria, and the archeon *Methanobrevibacter smithii* was unchanged. Because resveratrol and EGCG were tested together, however, it is difficult to speculate on the role each of these compounds would have had if dosed separately.

In conclusion, there is good evidence that based on both in vitro and rodent studies, resveratrol directly modifies the composition of the gut microbiota either by inhibiting the growth of individual microbial species or causing population shifts. In some experiments, resveratrol appeared to reverse dysbiosis arising from an obesogenic diet, for example by increasing the Bacteroidetes:Firmicutes ratio. However, the most robust results have been limited to studies in rodents, and their clinical significance for humans is therefore unclear. In most rodent studies, doses used were considerably higher than the 450-mg acceptable daily intake established for humans, and although there was good concordance in the pharmacologic response between species (43), doses used in animal models cannot be directly translated to humans. Further experiments establishing efficacy at appropriate doses in humans are warranted.

Beneficial effects of resveratrol metabolites produced by the gut microbiota. Resveratrol undergoes rapid and extensive glucuronidation and sulfation reactions in the host liver and duodenum (44, 45). Moreover, the gut microbiota can metabolize resveratrol, as found in both in vitro fermentation experiments with human fecal cultures and controlled intervention studies with healthy volunteers, although with pronounced interindividual differences (21). One study of 73 men and women at high cardiovascular disease

risk found 21 different urinary resveratrol metabolites, including glucuronide and sulfate conjugates of dihydroresveratrol formed by the intestinal microbiota after the chronic consumption of 272 mL red wine or dealcoholized red wine/d for 4 wk, as determined with the use of ultra-HPLC–mass spectrometry (46). There was no difference in resveratrol bioavailability because of the alcohol content of the wine. In a secondary study by the same research group in 10 healthy men who consumed either grape extract or red wine, the metabolic fate of resveratrol was examined (47). Seventeen metabolites were identified in plasma and urine. The dietary supplement formulation released resveratrol more slowly into the digestive tract than wine and showed a delayed absorption, leading to 2- to 4-fold higher urinary concentrations of microbial metabolites (47).

Dihydroresveratrol, 3,4'-dihydroxy-*trans*-stilbene, and 3,4'-dihydroxybibenzyl (lunularin) have been identified both in vitro and in vivo as the main microbiota-derived metabolites formed from resveratrol (48). However, resveratrol metabolism by the gut microbiota has shown pronounced interindividual differences, both in terms of the proportion of ingested resveratrol metabolized and the pattern of metabolites formed (21). Two strains out of a battery of 31 intestinal-origin bacterial strains, *Slackia equolifaciens* and *Adlercreutzia equolifaciens*, were identified as producers of dihydroresveratrol (21); interestingly, these strains are also capable of bioconverting other plant-derived polyphenols such as isoflavones and various catechins (49, 50). Dihydroresveratrol plasma concentrations have been found, however, to vary considerably between individuals after resveratrol supplementation, although one study found that it was produced in all 12 volunteers after daily supplementation with 150 mg resveratrol (51).

Although microbiota-derived resveratrol metabolites have not been researched as intensively as resveratrol, several studies point to their biological effects. The NO synthase inhibition activity of dihydroresveratrol is similar to resveratrol itself (52), and it displays comparable antioxidant activity and effects on cell proliferation and cycling (24, 53). Interestingly, the glucose-lowering drug metformin appears to affect dihydroresveratrol formation: there is an apparent shift toward higher concentrations of dihydroresveratrol with an increasing dose of metformin without a change in the circulating concentrations of resveratrol (54), suggesting the production of other metabolites is reduced. Taken with other results that have shown that metformin attenuated favorable effects from resveratrol supplementation on glucose control and the blood lipid profile (55), the production of metabolites other than dihydroresveratrol by the gut microbiota may be responsible for its cardiovascular benefits.

The resveratrol metabolite lunularin, also found in plants such as the common cow wheat and common celery, has bioactive properties, including reducing the secretion of proinflammatory mediators after an LPS challenge (56). Moreover, it showed moderately cytotoxic effects against

liver carcinoma cells and mildly antibacterial activity against the gram-negative bacterium *Pseudomonas aeruginosa* (57). Likewise, several studies have identified in vitro effects of 3,4'-dihydroxy-*trans*-stilbene: chemopreventive effects via the nuclear factor erythroid-2–related factor 2 pathway (58) were found, it was more effective than resveratrol in ameliorating insulin resistance (59), and it appeared to have cytogenetic activity comparable to resveratrol (60). The current limited research base shows that similar to resveratrol, its gut-derived metabolites have anti-inflammatory properties, may be mildly chemopreventive, and could improve glucose control or insulin sensitivity.

Improved gut barrier function. Resveratrol may influence the crosstalk between the gut microbiome and the immune system via its effects on barrier function and integrity. The integrity of proteins in the junction of cells is important in maintaining the intestinal barrier function, and tight junction protein upregulation has been associated with improved intestinal cell integrity (61). Several studies have shown that resveratrol can improve cell junction integrity by upregulating the expression of intestinal tight junction proteins such as tight junction protein 2 (36) and occludin (62). Moreover, the increase in gut permeability caused by the common mycotoxin deoxynivalenol could be arrested by resveratrol, which has been shown to promote the assembly of tight junction protein claudin 4 (63). Considering the importance of the gut barrier function for immunity and in the pathophysiology of several disease processes (“leaky gut”), the role of resveratrol in contributing to the maintenance of an intact intestinal barrier clearly warrants further investigation.

Health effects of resveratrol mediated through the gut microbiota

Research into the health effects of resveratrol or its metabolites that are mediated by the gut microbiota are limited to experimental animal models. The focus of the research is on cardiovascular effects and the metabolic syndrome, although these areas overlap into immunity and other areas of health.

Cardiovascular disease. A major contributor to the progression of cardiovascular disease is the process of atherosclerosis. Trimethylamine-*N*-oxide (TMAO) is a metabolite produced by a 2-step process whereby dietary L-carnitine, choline, and lecithin are converted to trimethylamine by the gut microbiota, and host flavin-containing monoamine oxidases are responsible for the conversion to TMAO, which is implicated in the development of atherosclerosis (64). In a 5-y follow-up cohort study, a higher plasma TMAO concentration was associated with a 4-fold increase in mortality risk in patients with stable coronary artery disease (65). Circulating TMAO concentrations have been shown to predict increases in carotid intima-media thickness, an early marker of atherosclerosis, in healthy overweight individuals with a family history of

type 2 diabetes participating in a recent longitudinal cross-sectional study (66).

Bacteria from several families commonly found in the gut are capable of producing trimethylamine, although strains within one species may differ in their ability to produce trimethylamine (67). Overall, studies in rodents have indicated that resveratrol reduces the increase in bacteria of Firmicutes (25, 26, 32, 38, 39), which partially modulates the catabolism of choline and carnitine to trimethylamine (68). In a series of experiments with the use of both standard and atherosclerosis-prone mice, resveratrol treatment (0.4%) inhibited the synthesis of trimethylamine and its further conversion to TMAO through gut microbiota remodeling, which was associated with an increase in the Bacteroides:Firmicutes ratio, and potentially through the inhibition of microbial trimethylamine lyase activity (29, 69). Moreover, increases in the proportion of *Lactobacillus* and *Bifidobacterium* appeared to increase the activity of bile salt hydrolase, which increased bile acid excretion and thus lowered cholesterol by increased endogenous bile acid production via repression of the enterohepatic farnesoid X receptor fibroblast growth factor 15 (29). In atherosclerosis-prone apoE^{-/-} mice, 4 g resveratrol/kg attenuated atherosclerosis both in the presence and absence of dietary choline. Moreover, the improvement in atherosclerosis induced by resveratrol in choline-fed mice was inhibited by antibiotic treatment. The latter results suggest that mechanisms other than the inhibition of the gut microbial trimethylamine production are involved in the antiatherosclerotic effects of resveratrol in this murine model of atherosclerosis (29). Thus, beyond shifting the microbiome's composition toward a healthier pattern, it could be modulated by resveratrol to specifically target the production of secondary metabolites (70).

Obesity. There is a complex relation between obesity, diet, and the composition and functionality of the gut microbiome (18, 71). Studies in mice have suggested that resveratrol can influence this relation in several ways, either by directly modulating the gut microbiome to one associated with a healthy weight (31, 34) or by changing the expression and activity of genes involved in weight maintenance, such as fasting-induced adipose factor or the mechanistic target of rapamycin (mTOR), known regulators of peripheral lipid and glucose metabolism as well as central energy metabolism (31, 33, 72). Changes in gene expression related to inflammation, apoptosis, and mitochondrial FA oxidation by resveratrol may also mitigate metabolic disturbances caused by obesity (35).

One study involved C57BL/6N mice in a 2 × 2 factorial design, with resveratrol and the control compared with a standard diet and an HFS diet (45% of calories from fat, 17% of calories from sucrose) (34). Resveratrol treatment (4 g/kg) reduced fat mass in mice fed the HFS diet only, and there was a nonsignificant decrease in body weight in both resveratrol-treated groups. Resveratrol also improved glucose homeostasis in mice fed the obesogenic diet. Fecal microbiota

transplantations from mice fed resveratrol or a control diet were used to determine whether these changes could arise from changes in the gut microbiome that were seen. Glucose homeostasis (glucose AUC) improved considerably in obese mice that received fecal microbiota transplantations from resveratrol-treated mice, but there was no change from the baseline in mice receiving a fecal microbiota transplantation from mice fed the control diet, which suggests that the glucose homeostasis improvements were caused by changes in the gut microbiota or metabolites remaining in the feces rather than resveratrol itself.

In a similar study conducted in mice fed with a high-fat diet (50% of calories from fat), a high-fat diet with 200 mg resveratrol · kg⁻¹ · d⁻¹, or a control diet (10% of calories from fat), resveratrol prevented weight gain caused by the high-fat diet (31). Several changes in the composition of the gut microbiota were noted; most importantly, it prevented a high-fat diet-induced gut microbiota dysbiosis, namely by increasing the Bacteroidetes:Firmicutes ratio. Bacteria from Firmicutes, which are relatively more prevalent in the obese (71), produce greater amounts of energy from dietary fiber than the other major gut bacterial phylum, Bacteroidetes (18), by increasing the production of SCFAs (73). Thus, the available energy content of the diet is augmented by increasing the amount of harvestable energy available from otherwise indigestible carbohydrates (74). In addition, a greater abundance of Bacteroidetes in humans may also increase postprandial fat oxidation (37). However, the relation is complex. SCFA production also stimulates the secretion of the satiety-related hormones leptin, glucagon-like peptide 1 (GLP-1), and pancreatic peptide YY₃₋₃₆, which are known to reduce ad libitum eating and thus lower body weight (75–79). In addition, the expression of fasting-induced adipose factor, which normally has to be suppressed in ileal cells to enable the microbiota-induced deposition of TGs in adipocytes (80), was promoted in both the control and high-fat resveratrol diets, suggesting that resveratrol affected how gut microbes enabled lipid transport in the body (31).

Resveratrol may have marked effects on the signaling activity of mTOR (33), a central regulator of energy storage and consumption (81). A study performed in mice exposed to a high-fat or control diet found that high-fat diet-related microbial dysbiosis correlated with increased body weight and biomarkers of the metabolic syndrome such as elevated blood glucose concentrations and insulin resistance. The specific inhibition of the mTOR1 signaling pathway, which reduces fat deposition and impairs adipocyte maintenance (82), with 200 mg resveratrol · kg⁻¹ · d⁻¹ may have caused the reductions in body weight and improvements in markers of glucose homeostasis in the resveratrol-treated group (33). Moreover, the activation of the mTOR2 signaling pathway followed by mTOR1 inhibition with resveratrol has been speculated to directly suppress the growth of obesity-associated gut microorganisms such as *Lactococcus*, *Clostridium* cluster XI, *Oscillibacter*, and *Hydrogenoanaerobacterium* (33).

Impaired GLP-1 signaling is also thought to play a role in obesity and type 2 diabetes development (83). Modifications of the gut microbiota that are derived from the diet may affect secreted GLP-1 concentrations and the consequences of the metabolic syndrome (84). In mice, gut microbiota composition changes were concurrent with increases in circulating GLP-1 concentrations in resveratrol-treated mice maintained on a high-fat diet compared with the control and high-fat diets. These changes were associated with improved glucose control, specifically reducing the glucose AUC compared with mice that consumed a high-fat diet, corresponding to an increase in insulin secretion (32). In addition, this study showed that resveratrol induced beneficial changes in markers of inflammation: TNF- α was suppressed in many tissues, whereas anti-inflammatory cytokine IL-10 was enhanced (32). In contrast, however, 500 mg resveratrol 2 times/d in a pilot study in humans had no effect on GLP-1 secretion, glycemic control, gastric emptying, body weight, or energy intake (85).

Future Research Directions

The current state of the research on resveratrol and its effects on human health via the microbiota has several limitations that hamper its development. The composition of the gut microbiome is complex, with a common set of microbial species present in all individuals, although the relative proportions may vary widely within a much larger set of bacteria, fungi, and archaea that make up the human gut microbiome (86). Commonly used methods to analyze gut microbial composition include isolating bacteria on selective media (35), quantitative real-time PCR, fluorescence in situ hybridization combined with flow cytometry (31), and the main metagenomic approaches, 16S ribosomal RNA-based amplicon sequencing and whole shotgun sequencing (21). In the literature that describes the effects of resveratrol, gut microbiota diversity was investigated according to taxonomic classification levels that varied from strain to phylum. Clear inconsistencies in how specific compositional changes have been compared exist between studies. The translatability of results between research groups is reduced by both the use of different methods of characterizing the gut microbiota and the taxonomic level used to compare microbial composition between groups. The use of standard methodologies and reference genomes could be beneficial in comparing research results with the effect of resveratrol on the composition of the gut microbiota (87–89). Moreover, the gut microbiota is affected by the composition of the diet, and variations in the composition of nondefined animal diets used add a potential source of random error to experimental work. Purified diets could improve reproducibility between animal experiments.

An inherent difficulty in conducting research into the effects of dietary components via the modulation of the gut microbiota is that microbiota-modulating effects may be extraneous to direct biological effects—or result from them. Gnotobiotic rodent models offer the opportunity to show that biological effects are mediated through effects on the

microbiome. Therefore, germ-free models should be further used in resveratrol research. For example, a landmark paper recently showed that improvements in insulin sensitivity from resistant starch occurred independent of the gut microbiome with the use of a germ-free mouse model (90).

A few studies have reported encouraging health effects in animals; however, human studies that link resveratrol's health effects with its modulation of the human microbiome are sparse. The most reproducible outcomes of resveratrol supplementation in rodent models appear to involve the attenuation of the negative health effects of a high-fat diet. Considering the effect of certain metabolite-producing types and links to the composition of the gut, pre- and probiotics should be examined to determine whether they can amplify the effects of resveratrol. Moreover, the results showing that resveratrol can improve gut barrier functioning have applications that overlap with cardiovascular health and obesity, such as immunity and gastrointestinal health. The use of *in vitro* techniques, particularly at physiologic rather than pharmacologic doses, would be beneficial in identifying relevant mechanisms by which resveratrol affects the microbiome. Further research should aim to determine whether modulating the gut microbiota with the use of resveratrol could modify metabolic factors associated with obesity.

In conclusion, resveratrol is capable of affecting the composition of the gut microbiome. It can preferentially slow the growth of certain microbes, which results in a more favorable microbial profile, particularly under obesity-promoting dietary conditions. In addition, resveratrol metabolites produced by gut bacteria have distinct biological effects: it may be possible to steer the production of these metabolites through the concurrent use of pre- or probiotics. Resveratrol or its metabolites can also influence the production of appetite hormones and metabolic byproducts of certain dietary components and the integrity of the intestinal epithelium, which may have antiobesity or cardioprotective effects. Further elucidation of the biological effects of resveratrol mediated via the gut microbiota in humans is warranted.

Acknowledgments

We thank Sophie Fehlbaum for carefully proofreading the manuscript, particularly in regard to the correct nomenclature of microorganisms. The authors' responsibilities were as follows—PW and RES: developed the concept for the review; JKB: wrote the original paper with contributions from RES and had primary responsibility for the final content; DR and RES: contributed substantially to revisions; and all authors: read and approved the final manuscript.

References

1. Rothwell JA, Perez-Jimenez J, Neveu V, Medina-Remón A, M'Hiri N, García-Lobato P, Manach C, Knox C, Eisner R, Wishart DS, et al. Phenol-Explorer 3.0: a major update of the Phenol-Explorer database to incorporate data on the effects of food processing on polyphenol content. *Database* (Oxford) 2013;2013:bat070.

2. Most J, Tosti V, Redman LM, Fontana L. Calorie restriction in humans: an update. *Ageing Res Rev* 2017;39:36–45.
3. Howitz KT, Bitterman KJ, Cohen HY, Lamming DW, Lavu S, Wood JG, Zipkin RE, Chung P, Kisielewski A, Zhang LL, et al. Small molecule activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. *Nature* 2003;425:191–6.
4. Poulsen MM, Jorgensen JO, Jessen N, Richelsen B, Pedersen SB. Resveratrol in metabolic health: an overview of the current evidence and perspectives. *Ann N Y Acad Sci* 2013;1290:74–82.
5. Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, Prabhu VV, Allard JS, Lopez-Lluch G, Lewis K, et al. Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* 2006;444:337–42.
6. Jang M, Cai L, Udeani GO, Slowing KV, Thomas CF, Beecher CW, Fong HH, Farnsworth NR, Kinghorn AD, Mehta RG, et al. Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* 1997;275:218–20.
7. Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov* 2006;5:493–506.
8. Surh YJ, Chun KS, Cha HH, Han SS, Keum YS, Park KK, Lee SS. Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF-kappa B activation. *Mutat Res* 2001;480–481:243–68.
9. Vidavalur R, Otani H, Singal PK, Maulik N. Significance of wine and resveratrol in cardiovascular disease: French paradox revisited. *Exp Clin Cardiol* 2006;11:217–25.
10. Renaud S, de Lorgeril M. Wine, alcohol, platelets, and the French paradox for coronary heart disease. *Lancet* 1992;339:1523–6.
11. Zamora-Ros R, Knaze V, Rothwell JA, Hemon B, Moskal A, Overvad K, Tjonneland A, Kyro C, Fagherazzi G, Boutron-Ruault MC, et al. Dietary polyphenol intake in Europe: the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Eur J Nutr* 2016;55:1359–75.
12. Zamora-Ros R, Rothwell JA, Achaintre D, Ferrari P, Boutron-Ruault MC, Mancini FR, Affret A, Kuhn T, Katzke V, Boeing H, et al. Evaluation of urinary resveratrol as a biomarker of dietary resveratrol intake in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Br J Nutr* 2017;117:1596–602.
13. Siedlinski M, Boer JM, Smit HA, Postma DS, Boezen HM. Dietary factors and lung function in the general population: wine and resveratrol intake. *Eur Respir J* 2012;39:385–91.
14. Rabassa M, Zamora-Ros R, Urpi-Sarda M, Bandinelli S, Ferrucci L, Andres-Lacueva C, Cherubini A. Association of habitual dietary resveratrol exposure with the development of frailty in older age: the Invecchiare in Chianti study. *Am J Clin Nutr* 2015;102:1534–42.
15. Vitaglione P, Sforza S, Galaverna G, Ghidini C, Caporaso N, Vescovi PP, Fogliano V, Marchelli R. Bioavailability of trans-resveratrol from red wine in humans. *Mol Nutr Food Res* 2005;49:495–504.
16. Walle T. Bioavailability of resveratrol. *Ann N Y Acad Sci* 2011;1215:9–15.
17. Chen F, Wen Q, Jiang J, Li HL, Tan YF, Li YH, Zeng NK. Could the gut microbiota reconcile the oral bioavailability conundrum of traditional herbs? *J Ethnopharmacol* 2016;179:253–64.
18. Komaroff AL. The microbiome and risk for obesity and diabetes. *JAMA* 2017;317:355–356.
19. Paulo L, Ferreira S, Gallardo E, Queiroz JA, Domingues F. Antimicrobial activity and effects of resveratrol on human pathogenic bacteria. *World J Microbiol Biotechnol* 2010;26:1533–8.
20. Nawrocki EM, Bedell HW, Humphreys TL. Resveratrol is cidal to both classes of *Haemophilus ducreyi*. *Int J Antimicrob Agents* 2013;41:477–9.
21. Bode LM, Bunzel D, Huch M, Cho GS, Ruhland D, Bunzel M, Bub A, Franz CMAP, Kulling SE. In vivo and in vitro metabolism of trans-resveratrol by human gut microbiota. *Am J Clin Nutr* 2013;97:295–309.
22. Storniolo CE, Moreno JJ. Resveratrol metabolites have an anti-proliferative effect on intestinal epithelial cancer cells. *Food Chem* 2012;134:1385–91.
23. Delmas D, Lin HY. Role of membrane dynamics processes and exogenous molecules in cellular resveratrol uptake: consequences in bioavailability and activities. *Mol Nutr Food Res* 2011;55:1142–53.
24. Walle T, Hsieh F, DeLegge MH, Oatis JE Jr., Walle UK. High absorption but very low bioavailability of oral resveratrol in humans. *Drug Metab Dispos* 2004;32:1377–82.
25. Boocock DJ, Faust GE, Patel KR, Schinas AM, Brown VA, Ducharme MP, Booth TD, Crowell JA, Perloff M, Gescher AJ, et al. Phase I dose escalation pharmacokinetic study in healthy volunteers of resveratrol, a potential cancer chemopreventive agent. *Cancer Epidemiol Biomarkers Prev* 2007;16:1246–52.
26. Brown VA, Patel KR, Viskaduraki M, Crowell JA, Perloff M, Booth TD, Vasilinin G, Sen A, Schinas AM, Piccirilli G, et al. Repeat dose study of the cancer chemopreventive agent resveratrol in healthy volunteers: safety, pharmacokinetics, and effect on the insulin-like growth factor axis. *Cancer Res* 2010;70:9003–11.
27. Jung CM, Heinze TM, Schnackenberg LK, Mullis LB, Elkins SA, Elkins CA, Steele RS, Sutherland JB. Interaction of dietary resveratrol with animal-associated bacteria. *FEMS Microbiol Lett* 2009;297:266–73.
28. Giuliani C, Marzorati M, Innocenti M, Vilchez-Vargas R, Vital M, Pieper DH, Van de Wiele T, Mulinacci N. Dietary supplement based on stilbenes: a focus on gut microbial metabolism by the in vitro simulator M-SHIME(R). *Food Funct* 2016;7:4564–75.
29. Chen ML, Yi L, Zhang Y, Zhou X, Ran L, Yang J, Zhu JD, Zhang QY, Mi MT. Resveratrol attenuates trimethylamine-N-oxide (TMAO)-induced atherosclerosis by regulating TMAO synthesis and bile acid metabolism via remodeling of the gut microbiota. *MBio* 2016;7:e02210–15.
30. Sung MM, Byrne NJ, Robertson IM, Kim TT, Samokhvalov V, Levasseur J, Soltys CM, Fung D, Tyreman N, Denou E, et al. Resveratrol improves exercise performance and skeletal muscle oxidative capacity in heart failure. *Am J Physiol Heart Circ Physiol* 2017;312:H842–53.
31. Qiao Y, Sun J, Xia S, Tang X, Shi Y, Le G. Effects of resveratrol on gut microbiota and fat storage in a mouse model with high-fat-induced obesity. *Food Funct* 2014;5:1241–9.
32. Dao TMA, Waget A, Klopp P, Serino M, Vachoux C, Pechere L, Drucker DJ, Champion S, Barthélemy S, Barra Y, et al. Resveratrol increases glucose induced GLP-1 secretion in mice: a mechanism which contributes to the glycemic control. *PLoS One* 2011;6.
33. Jung MJ, Lee J, Shin NR, Kim MS, Hyun DW, Yun JH, Kim PS, Whon TW, Bae JW. Chronic repression of mTOR complex 2 induces changes in the gut microbiota of diet-induced obese mice. *Sci Rep* 2016;6.
34. Sung MM, Kim TT, Denou E, Soltys CM, Hamza SM, Byrne NJ, Masson G, Park H, Wishart DS, Madsen KL, et al. Improved glucose homeostasis in obese mice treated with resveratrol is associated with alterations in the gut microbiome. *Diabetes* 2017;66:418–25.
35. Larrosa M, Yañez-Gascón MJ, Selma MV, González-Sarrías A, Toti S, Cerón JJ, Tomás-Barberán F, Dolara P, Espín JC. Effect of a low dose of dietary resveratrol on colon microbiota, inflammation and tissue damage in a DSS-induced colitis rat model. *J Agric Food Chem* 2009;57:2211–20.
36. Etxeberria U, Arias N, Boqué N, Macarulla MT, Portillo MP, Martínez JA, Milagro FL. Reshaping faecal gut microbiota composition by the intake of trans-resveratrol and quercetin in high-fat sucrose diet-fed rats. *J Nutr Biochem* 2015;26:651–60.
37. Most J, Penders J, Lucchesi M, Goossens GH, Blaak EE. Gut microbiota composition in relation to the metabolic response to 12-week combined polyphenol supplementation in overweight men and women. *Eur J Clin Nutr* 2017 Jun 7 (Epub ahead of print; DOI: 10.1038/ejcn.2017.89).
38. Bervoets L, Van Hoorenbeeck K, Kortleven I, Van Noten C, Hens N, Vael C, Goossens H, Desager KN, Vankerckhoven V. Differences in gut microbiota composition between obese and lean children: a cross-sectional study. *Gut Pathog* 2013;5:10.
39. Dugas LR, Fuller M, Gilbert J, Layden BT. The obese gut microbiome across the epidemiologic transition. *Emerg Themes Epidemiol* 2016;13:2.

40. Korsholm AS, Kjaer TN, Ornstrup MJ, Pedersen SB. Comprehensive metabolomic analysis in blood, urine, fat, and muscle in men with metabolic syndrome: a randomized, placebo-controlled clinical trial on the effects of resveratrol after four months' treatment. *Int J Mol Sci* 2017;18.
41. Williams BB, Van Benschoten AH, Cimermancic P, Donia MS, Zimmermann M, Taketani M, Ishihara A, Kashyap PC, Fraser JS, Fischbach MA. Discovery and characterization of gut microbiota decarboxylases that can produce the neurotransmitter tryptamine. *Cell Host Microbe* 2014;16:495–503.
42. Pessione E, Pessione A, Lamberti C, Coisson DJ, Riedel K, Mazzoli R, Bonetta S, Eberl L, Giunta C. First evidence of a membrane-bound, tyramine and beta-phenylethylamine producing, tyrosine decarboxylase in *Enterococcus faecalis*: a two-dimensional electrophoresis proteomic study. *Proteomics* 2009;9:2695–710.
43. Edwards JA, Beck M, Riegger C, Bausch J. Safety of resveratrol with examples for high purity, trans-resveratrol, resVida(RR). *Ann N Y Acad Sci* 2011;1215:131–7.
44. Wenzel E, Somoza V. Metabolism and bioavailability of trans-resveratrol. *Mol Nutr Food Res* 2005;49:472–81.
45. Wu B, Basu S, Meng S, Wang X, Hu M. Regioselective sulfation and glucuronidation of phenolics: insights into the structural basis. *Curr Drug Metab* 2011;12:900–16.
46. Rotches-Ribalta M, Urpi-Sarda M, Llorach R, Boto-Ordóñez M, Jauregui O, Chiva-Blanch G, Perez-García L, Jaeger W, Guillen M, Corella D, et al. Gut and microbial resveratrol metabolite profiling after moderate long-term consumption of red wine versus dealcoholized red wine in humans by an optimized ultra-high-pressure liquid chromatography tandem mass spectrometry method. *J Chromatogr A* 2012;1265:105–13.
47. Rotches-Ribalta M, Andres-Lacueva C, Estruch R, Escribano E, Urpi-Sarda M. Pharmacokinetics of resveratrol metabolic profile in healthy humans after moderate consumption of red wine and grape extract tablets. *Pharmacol Res* 2012;66:375–82.
48. Juan ME, Alfaras I, Planas JM. Determination of dihydroresveratrol in rat plasma by HPLC. *J Agric Food Chem* 2010;58:7472–5.
49. Jin JS, Kitahara M, Sakamoto M, Hattori M, Benno Y. *Slackia equolifaciens* sp. nov., a human intestinal bacterium capable of producing equol. *Int J Syst Evol Microbiol* 2010;60:1721–4.
50. Takagaki A, Nanjo F. Bioconversion of (–)-epicatechin, (+)-epicatechin, (–)-catechin, and (+)-catechin by (–)-epigallocatechin-metabolizing bacteria. *Biol Pharm Bull* 2015;38:789–94.
51. van der Made SM, Plat J, Mensink RP. Resveratrol does not influence metabolic risk markers related to cardiovascular health in overweight and slightly obese subjects: a randomized, placebo-controlled crossover trial. *PLoS One* 2015;10:e0118393.
52. Kageura T, Matsuda H, Morikawa T, Toguchida I, Harima S, Oda M, Yoshikawa M. Inhibitors from rhubarb on lipopolysaccharide-induced nitric oxide production in macrophages: structural requirements of stilbenes for the activity. *Bioorg Med Chem* 2001;9:1887–93.
53. Stivala LA, Savio M, Carafoli F, Perucca P, Bianchi L, Maga G, Forti L, Pagnoni UM, Albini A, Prosperi E, et al. Specific structural determinants are responsible for the antioxidant activity and the cell cycle effects of resveratrol. *J Biol Chem* 2001;276:22586–94.
54. Timmers S, de Ligt M, Phielix E, van de Weijer T, Hansen J, Moonen-Kornips E, Schaart G, Kunz I, Hesselink MK, Schrauwen-Hinderling VB, et al. Resveratrol as add-on therapy in subjects with well-controlled type 2 diabetes: a randomized controlled trial. *Diabetes Care* 2016;39:2211–7.
55. Movahed A, Nabipour I, Lieben Louis X, Thandapilly SJ, Yu L, Kalantarhormozi M, Rekabpour SJ, Netticadan T. Antihyperglycemic effects of short term resveratrol supplementation in type 2 diabetic patients. *Evid Based Complement Alternat Med* 2013;2013:851267.
56. Vogl S, Atanasov AG, Binder M, Bulusu M, Zehl M, Fakhrudin N, Heiss EH, Picker P, Wawrosch C, Saukel J, et al. The herbal drug *melampyrum pratense* L. (Koch): isolation and identification of its bioactive compounds targeting mediators of inflammation. *Evid Based Complement Alternat Med* 2013;2013:395316.
57. Lu ZQ, Fan PH, Ji M, Lou HX. Terpenoids and bisbibenzyls from Chinese liverworts *Conocephalum conicum* and *Dumortiera hirsuta*. *J Asian Nat Prod Res* 2006;8:187–92.
58. Lin D, Dai F, Sun LD, Zhou B. Toward an understanding of the role of a catechol moiety in cancer chemoprevention: the case of copper- and o-quinone-dependent Nrf2 activation by a catechol-type resveratrol analog. *Mol Nutr Food Res* 2015;59:2395–406.
59. Ito-Nagahata T, Kurihara C, Hasebe M, Ishii A, Yamashita K, Iwabuchi M, Sonoda M, Fukuhara K, Sawada R, Matsuoka A, et al. Stilbene analogs of resveratrol improve insulin resistance through activation of AMPK. *Biosci Biotechnol Biochem* 2013;77:1229–35.
60. Matsuoka A, Takeshita K, Furuta A, Ozaki M, Fukuhara K, Miyata N. The 4'-hydroxy group is responsible for the in vitro cytogenetic activity of resveratrol. *Mutat Res* 2002;521:29–35.
61. Hsieh CY, Osaka T, Moriyama E, Date Y, Kikuchi J, Tsuneda S. Strengthening of the intestinal epithelial tight junction by *Bifidobacterium bifidum*. *Physiol Rep* 2015;3.
62. Wang N, Han Q, Wang G, Ma WP, Wang J, Wu WX, Guo Y, Liu L, Jiang XY, Xie XL, et al. Resveratrol protects oxidative stress-induced intestinal epithelial barrier dysfunction by upregulating heme oxygenase-1 expression. *Dig Dis Sci* 2016;61:2522–34.
63. Ling KH, Wan ML, El-Nezami H, Wang M. Protective capacity of resveratrol, a natural polyphenolic compound, against deoxyvalenol-induced intestinal barrier dysfunction and bacterial translocation. *Chem Res Toxicol* 2016;29:823–33.
64. Velasquez MT, Ramezani A, Manal A, Raj DS. Trimethylamine N-oxide: the good, the bad and the unknown. *Toxins (Basel)* 2016;8.
65. Senthong V, Wang Z, Li XS, Fan Y, Wu Y, Tang WH, Hazen SL. Intestinal microbiota-generated metabolite trimethylamine-N-oxide and 5-year mortality risk in stable coronary artery disease: the contributory role of intestinal microbiota in a COURAGE-like patient cohort. *J Am Heart Assoc* 2016;5.
66. Randrianarisoa E, Lehn-Stefan A, Wang X, Hoene M, Peter A, Heinzmann SS, Zhao X, Konigsrainer I, Konigsrainer A, Balletshofer B, et al. Relationship of serum trimethylamine N-oxide (TMAO) levels with early atherosclerosis in humans. *Sci Rep* 2016;6:26745.
67. Romano KA, Vivas EL, Amador-Noguez D, Rey FE. Intestinal microbiota composition modulates choline bioavailability from diet and accumulation of the proatherogenic metabolite trimethylamine-N-oxide. *MBio* 2015;6:e02481.
68. Wilson A, McLean C, Kim RB. Trimethylamine-N-oxide: a link between the gut microbiome, bile acid metabolism, and atherosclerosis. *Curr Opin Lipidol* 2016;27:148–54.
69. Wang Z, Roberts AB, Buffa JA, Levison BS, Zhu W, Org E, Gu X, Huang Y, Zamanian-Daryoush M, Culley MK, et al. Non-lethal inhibition of gut microbial trimethylamine production for the treatment of atherosclerosis. *Cell* 2015;163:1585–95.
70. Brown JM, Hazen SL. Targeting of microbe-derived metabolites to improve human health: the next frontier for drug discovery. *J Biol Chem* 2017;292:8560–8.
71. Clarke SF, Murphy EF, Nilaweera K, Ross PR, Shanahan F, O'Toole PW, Cotter PD. The gut microbiota and its relationship to diet and obesity: new insights. *Gut Microbes* 2012;3:186–202.
72. Kim HK, Youn BS, Shin MS, Namkoong C, Park KH, Baik JH, Kim JB, Park JY, Lee KU, Kim YB, et al. Hypothalamic Angptl4/Fiaf is a novel regulator of food intake and body weight. *Diabetes* 2010;59:2772–80.
73. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 2006;444:1027–31.
74. DiBaise JK, Zhang H, Crowell MD, Krajmalnik-Brown R, Decker GA, Rittmann BE. Gut microbiota and its possible relationship with obesity. *Mayo Clin Proc* 2008;83:460–9.
75. Reimann F. Molecular mechanisms underlying nutrient detection by incretin-secreting cells. *Int Dairy J* 2010;20:236–42.
76. Samuel BS, Shaito A, Motoike T, Rey FE, Backhed F, Manchester JK, Hammer RE, Williams SC, Crowley J, Yanagisawa M, et al. Effects of the gut microbiota on host adiposity are modulated by the short-chain fatty-acid binding G protein-coupled receptor, Gpr41. *Proc Natl Acad Sci USA* 2008;105:16767–72.

77. Tazoe H, Otomo Y, Kaji I, Tanaka R, Karaki SI, Kuwahara A. Roles of short-chain fatty acids receptors, GPR41 and GPR43 on colonic functions. *J Physiol Pharmacol* 2008;59(Suppl 2):251–62.
78. Tolhurst G, Heffron H, Lam YS, Parker HE, Habib AM, Diakogiannaki E, Cameron J, Grosse J, Reimann F, Gribble FM. Short-chain fatty acids stimulate glucagon-like peptide-1 secretion via the G-protein-coupled receptor FFAR2. *Diabetes* 2012;61:364–71.
79. Xiong Y, Miyamoto N, Shibata K, Valasek MA, Motoike T, Kedzierski RM, Yanagisawa M. Short-chain fatty acids stimulate leptin production in adipocytes through the G protein-coupled receptor GPR41. *Proc Natl Acad Sci USA* 2004;101:1045–50.
80. Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, Nagy A, Semenkovich CF, Gordon JI. The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci USA* 2004;101:15718–23.
81. Woods SC, Seeley RJ, Cota D. Regulation of food intake through hypothalamic signaling networks involving mTOR. *Annu Rev Nutr* 2008;28:295–311.
82. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell* 2012;149:274–93.
83. Steinert RE, Feinle-Bisset C, Asarian L, Horowitz M, Beglinger C, Geary N. Ghrelin, CCK, GLP-1, and PYY(3–36): secretory controls and physiological roles in eating and glycemia in health, obesity, and after RYGB. *Physiol Rev* 2017;97:411–63.
84. Everard A, Cani PD. Gut microbiota and GLP-1. *Rev Endocr Metab Disord* 2014;15:189–96.
85. Thazhath SS, Wu T, Bound MJ, Checklin HL, Standfield S, Jones KL, Horowitz M, Rayner CK. Administration of resveratrol for 5 wk has no effect on glucagon-like peptide 1 secretion, gastric emptying, or glycaemic control in type 2 diabetes: a randomized controlled trial. *Am J Clin Nutr* 2016;103:66–70.
86. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N, Levenez F, Yamada T, et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 2010;464:59–65.
87. Nelson KE, Weinstock GM, Highlander SK, Worley KC, Creasy HH, Wortman JR, Rusch DB, Mitreva M, Sodergren E, et al. A catalog of reference genomes from the human microbiome. *Science* 2010;328:994–9.
88. Sinha R, Abnet CC, White O, Knight R, Huttenhower C. The microbiome quality control project: baseline study design and future directions. *Genome Biol* 2015;16:276.
89. Li J, Jia H, Cai X, Zhong H, Feng Q, Sunagawa S, Arumugam M, Kultima JR, Prifti E, Nielsen T, et al. An integrated catalog of reference genes in the human gut microbiome. *Nat Biotechnol* 2014;32:834–41.
90. Bindels LB, Segura Munoz RR, Gomes-Neto JC, Mutemberezi V, Martinez I, Salazar N, Cody EA, Quintero-Villegas MI, Kittana H, de Los Reyes-Gavilan CG, et al. Resistant starch can improve insulin sensitivity independently of the gut microbiota. *Microbiome* 2017;5:12.