**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

Tuning Tetrazole Photochemistry for Protein Ligation and Molecular Imaging

Fay, Rachael ; Holland, Jason P

Abstract: Photochemistry provides a wide range of alternative reagents that hold potential for use in bimolecular functionalisation of proteins. Here, we report the synthesis and characterisation of metal ion binding chelates derivatised with disubstituted tetrazoles for the photoradiochemical labelling of monoclonal antibodies (mAbs). The photophysical properties of tetrazoles featuring extended aromatic systems and auxochromic substituents to tune excitation toward longer wavelengths (365 and 395 nm) were studied. Two photoactivatable chelates based on desferrioxamine B (DFO) and the aza-macrocyclic NODAGA were functionalised with a tetrazole and developed for protein labelling with ⁸⁹Zr, ⁶⁴Cu and ⁶⁸Ga radionuclides. DFO-tetrazole (1) was assessed by direct conjugation to formulated trastuzumab and subsequent radiolabelling with ⁸⁹Zr. Radiochemical studies and cellular-based binding assays demonstrated that the radiotracer remained stable in vitro retained high immunoreactivity. Positron emission tomography (PET) imaging and biodistribution studies were used to measure the tumour specific uptake and pharmacokinetic profile in mice bearing SK-OV-3 xenografts. Experiments demonstrate that tetrazole-based photochemistry is a viable approach for the light-induced synthesis of PET radiotracers.

DOI: <https://doi.org/10.1002/chem.202100061>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-213028>

Journal Article

Accepted Version

Originally published at:

Fay, Rachael; Holland, Jason P (2021). Tuning Tetrazole Photochemistry for Protein Ligation and Molecular Imaging. *Chemistry - A European Journal*, 27(15):4893-4897.

DOI: <https://doi.org/10.1002/chem.202100061>

Tuning tetrazole photochemistry for protein ligation and molecular imaging

Rachael Fay^[a] and Jason P. Holland^{*[a]}

Abstract: Photochemistry provides a wide range of alternative reagents that hold potential for use in bimolecular functionalisation of proteins. Here, we report the synthesis and characterisation of metal ion binding chelates derivatised with disubstituted tetrazoles for the photoradiochemical labelling of monoclonal antibodies (mAbs). The photophysical properties of tetrazoles featuring extended aromatic systems and auxochromic substituents to tune excitation toward longer wavelengths (365 and 395 nm) were studied. Two photoactivatable chelates based on desferrioxamine B (DFO) and the *aza*-macrocycle NODAGA were functionalised with a tetrazole and developed for protein labelling with ⁸⁹Zr, ⁶⁴Cu and ⁶⁸Ga radionuclides. DFO-tetrazole (**1**) was assessed by direct conjugation to formulated trastuzumab and subsequent radiolabelling with ⁸⁹Zr. Radiochemical studies and cellular-based assays binding demonstrated that the radiotracer remained stable *in vitro* retained high immunoreactivity. Positron emission tomography (PET) imaging and biodistribution studies were used to measure the tumour specific uptake and pharmacokinetic profile in mice bearing SK-OV-3 xenografts. Experiments demonstrate that tetrazole-based photochemistry is a viable approach for the light-induced synthesis of PET radiotracers.

The photochemical reactivity of tetrazoles was first reported by Huisgen *et al.* in 1967.^[1] Light-induced activation of diaryl-tetrazoles populates an electronic excited state that undergoes rapid loss of N₂(g), to form a highly reactive nitrile imine. This nitrile imine intermediate can behave as a 1,3-dipole and participate in cycloaddition reactions^[2] or as an electrophile, reacting with native nucleophiles on protein.^[3] In 2008, the group of Qing Lin reported the use of tetrazoles for labelling of post-translationally modified proteins *via* photoinduced [3+2] cycloadditions with alkene substituents.^[4] This remarkable and fast reaction was described as a 'photo-click' process and is fluorogenic, generating emissive pyrazoline products.^[5] Subsequent work used genetically encoded artificial amino acids bearing alkenes^[6] and cyclopropanes^[7] for site-specific, photo-induced labelling of pre-modified proteins *in vitro* and *in vivo*. One of the attractive features of tetrazole-alkene photochemistry is that reactivity can be modified to increase coupling rates *via* structural and frontier orbital control on the tetrazole^[8–11] or the alkene partners.^[12,13] However, a drawback of tetrazole photochemistry is the need to use short wavelength light (~302 nm) to form the nitrile imine which is potentially damaging to proteins like mAbs.^[14,15]

The photo-click reaction between tetrazoles and alkenes was reported to be bioorthogonal.^[5,16] However, Huisgen *et al.* noted the reactivity of thermally-induced nitrile imines with thiophenol in 1961.^[3] More recently, the bioorthogonality of the 'photo-click' chemistry has been questioned. Detailed experimental evidence confirmed that the electrophilic nitrile imine produced *via* photochemical activation from tetrazoles reacts with nucleophilic amino acid residues in proteins.^[17,18] Li *et al.* demonstrated the photochemical induced reactivity of tetrazoles toward a range of carboxylic acids including pent-4-enoic acid, AcOH, free alanine and proteins labelled at Glu residues.^[17] Whilst the authors also reported reactivity toward a range of biologically relevant nucleophiles such as thiols, amines and alcohols, they noted that at physiological pH, carboxylic acids reacted preferentially with the nitrile imine, even outcompeting the cycloaddition process. Similarly, Zhao *et al.* reported the photo-induced reactivity of tetrazoles with propionic acid and a protected Asp derivative.^[18] Competition experiments with amino acids containing phenol, thiol, and amino functional groups, reduced the yield of the reaction between their model tetrazole and Boc-Asp-OTfBu, indicating that other nucleophiles can also react with the nitrile imine.^[18] Following these studies, the photo-induced reactivity of tetrazoles with thiols (mercaptoethanol) and amino acids (Trp, Pro, His, Ser, Asn, Tyr, Asn) was reported.^[19–21]

Until recently, the combination of radiochemistry and photochemistry has received minimal attention. Early work established that compounds radiolabelled with ^{99m}Tc, ¹⁸F, ¹⁸⁸Re and ¹¹¹In radionuclides could be functionalised with substituted (perfluoro-, nitro-) and non-substituted aryl azide (ArN₃) for photochemically induced labelling of different bioactive molecules.^[22–31] Sun *et al.* used the photo-induced cycloaddition process and labelled an alkene-functionalised, tumour-specific peptide with a modified tetrazole which allowed ⁶⁸Ga and ⁶⁴Cu radiolabelling for PET and optical imaging in a glioblastoma model.^[32] Our group has also explored the use of photoactivatable chelates bearing an aryl azide to make viable ⁶⁸Ga and ⁸⁹Zr PET radiotracers *via* direct functionalisation of mAbs.^[33–36]

Here, we demonstrate that the photophysical properties of tetrazoles can be tuned through synthetic control over the arylene substituents to facilitate the rapid and direct, bimolecular protein labelling at wavelengths in the UVA region of the spectrum. Two-step photochemical conjugation and radiolabelling produced ⁸⁹Zr-labelled trastuzumab for PET imaging of human epidermal growth-factor receptor 2 (HER2/*neu*) expression in tumours.

[a] R. Fay, Prof. Dr J. P. Holland*
University of Zurich, Department of Chemistry, Winterthurerstrasse
190, Zurich, CH-8057, Switzerland
Correspondence to:
E-mail: jason.holland@chem.uzh.ch
Tel: +41.44.63.53.990
Website: www.hollandlab.org

Supporting information is available

Tetrazole compounds **7**, **9**, **10**, and **12** (Figure 1) were synthesised by reacting phenylsulfonylhydrazones with arylene diazonium salts (generated *in situ*) and isolated in moderate yields (23–42%).^[37] Compound **8** was synthesised from compound **9** by nucleophilic aromatic substitution of the bromo-substituent by using NaSMe. Compound **13** was synthesised in two-steps involving isolation and deprotection of the corresponding OBn derivative (**14**). Efforts to synthesise the anthracene analogue (**11**) were not successful. Full experimental details and characterisation data are given in the supporting information (Schemes S1–S8, Figures S1–S36)

Figure 1. General synthetic scheme to yield long-wavelength tetrazole analogues.

With six tetrazole scaffolds in hand, we measured their photophysical and photochemical properties by using electronic absorption spectroscopy (UV/vis) analytical high-performance liquid chromatography (HPLC) to analyse the formation of photolysis products. (Figure 2, Figures S37–S38, and Table S1). Tetrazoles **10** and **12** displayed no photoactivity when irradiated at 365 nm. For compound **9**, minor decomposition was observed under 365 nm irradiation but this was attributed to homolytic cleavage of the bromo-substituent.^[38] Whilst the naphthalene derivative (**12**) was not photo-reactive at 365 nm, naphthylene analogues functionalised with auxochromic groups (OMe, SMe, OH) exhibited bathochromic shifts that allowed photolysis of the tetrazole to occur at longer wavelengths. In general, changes in the absorption spectra for compounds **7**, **8** and **13** followed the Woodward-Fieser-Kuhn rules where the methyl-mercaptan substituent (**9**) induced the greatest shift in λ_{max} to 332 nm (+42 nm with respect to the unsubstituted naphthalene-derivative (**12**), λ_{max} = 290 nm; Figure 2A).^[39,40] Kinetic studies on the photoactivation revealed that the rates of photolysis increased with increasing molecular absorption coefficient (Table S1). Irradiation of compound **7** (ϵ_{365} = 298 M⁻¹ cm⁻¹) with light at 365 nm induced a rapid first-order decomposition with an observed rate constant, $k_{\text{obs}}(365\text{nm})$ = 0.44 ± 0.12 min⁻¹. Although compound **8** displayed the largest red-shift and fastest rate of photolysis, thioethers are susceptible to oxidation *in vivo*.^[41] Therefore, compounds were selected for further applications in protein ligation.

Figure 2. Photophysical and photochemical data on tetrazoles **7**, **9** and **13**. A) Electronic absorption spectra recorded in DMSO. B) HPLC chromatograms before and after photolysis of compound **7**. C) Kinetic plots showing the relative change in the concentration of compound **7** versus time during extended photolysis at 365 and 395 nm.

To apply compound **7** in the photoradiochemical synthesis of ⁸⁹Zr-labelled mAbs, we synthesised the desferrioxamine-tetrazole derivative, compound **1** (Scheme S8 and Figures S31–S36). DFO is a hexadentate chelate which is used in clinical PET with ⁸⁹Zr-mAbs to coordinate the ⁸⁹Zr⁴⁺ metal ion (half-life, $t_{1/2}$ = 78.41 h).^[42] In addition, to increase radiochemical scope, we also synthesised a NODAGA-based tetrazole (**4**) for ⁶⁸Ga and ⁶⁴Cu radiochemistry (Scheme S7 and Figures S29–S31). Full details of synthesis and radiosynthesis of [^{nat/68}Ga]Ga-**4**, [^{nat/64}Cu]Cu-**4** and [^{nat/68}Ga]Ga-**1** are presented in the supporting information (Figures S39–S41). Details on the photochemical conjugation of compound **4** to human serum albumin (HSA) and dinutuximab (a chimeric mAb against the disialoganglioside GD2 in neuroblastoma) and radiolabelling with ⁶⁸Ga and ⁶⁴Cu, respectively, are also shown (Figures S42–S45).

The DFO analogue (compound **1**) was synthesised from (**7**) in 4-steps with an overall yield of 31%. Briefly, compound **7** was saponified and a Boc protected PEG linker was coupled to yield compound **3** (Scheme S8). Deprotection of the Boc group revealed the free amine (**2**) which was then coupled with DFO-succinate via amide bond formation to yield the desired compound **1**. Photophysical experiments with (**1**) indicated that the compound retained the photoactivity seen for the tetrazole core (ϵ_{365} = 312 M⁻¹ cm⁻¹, $k_{\text{obs}}(365\text{nm})$ = 1.88 ± 1.09 min⁻¹; ϵ_{395} = 200 M⁻¹ cm⁻¹, $k_{\text{obs}}(395\text{nm})$ = 0.19 ± 0.09 min⁻¹, Figure S38). The calculated photochemical quantum yields (ϕ_p) for (**1**), which is a measure of the efficiency of light-induced activation, were 2.2 ± 0.4% and 0.23 ± 0.04% at 365 and 395 nm, respectively. Complexation of ^{nat}Zr⁴⁺ with (**1**) was achieved by using standard methods and the ^{nat}Zr-**1** complex was characterised by HRMS and HPLC. The radioactive complex, ⁸⁹Zr-**1**⁺ was prepared at 23 °C in ≤10 minutes and was characterised by using radio-iTLC, and by radio-HPLC which gave a single peak in the chromatogram (Figure 3). Co-injection of ⁸⁹Zr-**1**⁺ with the non-radioactive complex measured by electronic absorption showed coincident retention times in HPLC.

Figure 3. Chromatographic data on the complexation of ^{nat}⁸⁹Zr⁴⁺ by compound **1**. A) Radio-iTLC data ⁸⁹Zr-**1**⁺ before (black) and after (blue) irradiation and a control (red, [⁸⁹Zr][Zr(DTPA)]³⁻). B) Analytical HPLC chromatograms for **1** (blue), ^{nat}Zr-**1**⁺ (green) and ⁸⁹Zr-**1**⁺ (black).

Next, we performed the photochemical conjugation between compound **1** and trastuzumab (formulated as HerceptinTM; 10-to-

1 chelate-to-mAb ratio, pH7) by irradiating reactions at either 365 or 395 nm for 10 min at room temperature (Figure 4A). Aliquots of the crude mixtures were retained for analysis and the DFO-mAb conjugates were isolated from small-molecule impurities by preparative size-exclusion chromatography (SEC). Samples of the crude photoconjugation mixtures were radiolabelled with $^{89}\text{Zr}^{4+}$ giving decay-corrected radiochemical yields (RCYs) of $27.9 \pm 10.5\%$, ($n=3$) and $34.8 \pm 3.0\%$ ($n=2$), at 365 and 395 nm, respectively (measured by manual size-exclusion chromatography with PD-10 columns; Figures S46). For a more accurate quantitative analysis we also measured the radiolabelled mixtures by radio-SEC which showed that 15% of activity was associated with trastuzumab (Figure 4, panel B, blue trace). Combined with the initial 10-to-1 stoichiometric ratio of reagents, the isolated DFO-mAb conjugate used in future studies had an average of 1.5 accessible DFO ligands per mAb. For *in vitro* and *in vivo* experiments, the purified DFO-Np(OMe)-Tz-trastuzumab conjugate was radiolabelled with an excess of ^{89}Zr at pH 8.5 and purified by SEC to give [^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab in sterile PBS (pH 7.4) with a RCY of 39.8%, a radiochemical purity (RCP) >95%, and a molar activity of $A_m = 4.01 \text{ MBq nmol}^{-1}$ of mAb (Figure 4B).

Figure 4. Photochemical conjugation and ^{89}Zr -radiolabelling of DFO-Np(OMe)-Tz-trastuzumab. A) Schematic showing the structure of compound 1 and the photochemical conjugation reaction with trastuzumab. B) SEC chromatograms showing the crude (blue) and purified (black) samples of the ^{89}Zr -mAb, the electronic absorption (280 nm) profile of trastuzumab, and the elution of [^{89}Zr]Zr(DTPA) $^{-}$ as a control. C) Cellular binding assays showing the specific binding of [^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab to SK-OV-3 ovarian cancer cells (HER2/*neu*-positive). Control study: n.s. = non-specific binding (achieved by pre-blocking with a 1025-fold excess of non-labelled trastuzumab).

The stability of [^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab was assessed in PBS and human serum (Figure S47–S48). Data indicated that the radiotracer was stable for up to 24 h at 37 °C in serum with the main peak in radio-SEC still associated with the ^{89}Zr -mAb. In sterile PBS, the radiotracer was stable for 3 days with no change in RCP. Cellular binding studies using SK-OV-3 cells were performed to assess the biochemical integrity of the radiotracer after the photoconjugation and radiolabelling. Binding curve analysis indicated that the [^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab sample had an immunoreactive fraction of $82 \pm 4\%$, and retained specificity toward HER2/*neu* (Figure 4C).^[43]

PET imaging and biodistribution studies were performed to measure the pharmacokinetics, tumour specificity and stability of [^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab in athymic nude mice bearing SK-OV-3 xenografts (Figure 5, Figures S50–S55, Table S2-S3). All animals received the same amount of activity ($0.53 \pm 0.02 \text{ MBq}$) but three different mass doses of ^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab. Group 1 received the highest molar activity dose ($20 \mu\text{g}$, $A_m = 4.01 \text{ MBq nmol}^{-1}$), group 2 received a 13-fold increase in protein dose ($A_m = 0.30 \text{ MBq nmol}^{-1}$) and group 3 received a full blocking dose (72-fold excess, $A_m = 0.06 \text{ MBq nmol}^{-1}$). SK-OV-3 tumours display extremely high expression of the HER2/*neu* protein and as consequence, no differences were observed in the PET images and biodistribution data between groups 1 and 2. For example, biodistribution data revealed that 72 h post-administration the tumour associated activity was $45.8 \pm 14.0 \text{ \%ID g}^{-1}$ for group 1 and $47.0 \pm 7.4 \text{ \%ID g}^{-1}$ for group 2. In contrast, the tumour uptake in the blocking group was reduced by ~67% to $15.5 \pm 5.2 \text{ \%ID g}^{-1}$, indicating specific tumour uptake. The tumour-to-tissue contrast was high although activity in the bone was observed in all three groups ($11.8 \pm 4.0 \text{ \%ID g}^{-1}$, $10.1 \pm 5.7 \text{ \%ID g}^{-1}$, $9.1 \pm 3.9 \text{ \%ID g}^{-1}$ for groups 1, 2, and 3, respectively). Bone uptake was in line with previous on ^{89}Zr -trastuzumab.^[44]

Figure 5. PET imaging and biodistribution data for [^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab in mice bearing SK-OV-3 tumours. A) Coronal and axial PET images shown through the tumour centre for the three different study groups. T = tumour, H = heart, L = liver. B) Biodistribution data recorded at 72 h post-administration.

Whole-body excretion studies were also used to measure the effective half-life of [^{89}Zr]ZrDFO-Np(OMe)-Tz-trastuzumab in each mouse (Figure S56). Measurements indicated that no significant difference in pharmacokinetics was observed between animals in three different dose groups with $t_{1/2}(\text{eff})$ equal to $29.8 \pm 0.8 \text{ h}$, $35.5 \pm 0.6 \text{ h}$, and $35.2 \pm 7.4 \text{ h}$, respectively. Values were also consistent with prior measurements using ^{89}Zr -trastuzumab

prepared via simultaneous photoradiolabelling with an aryl azide species.^[45]

In conclusion, the experiments provide irrefutable evidence that a direct photochemical conjugation occurs between the light-induced nitrile imine and the protein. The nature of the conjugate bonds formed remain uncertain but others have reported convincing evidence that carboxylate, amine and sulfhydryl nucleophiles from amino acids including Glu, Asp, Lys and Cys can attack the nitrile imine intermediate (Scheme S9).^[17–21] Tetrazole photochemistry can be adapted for labelling mAbs in the presence of standard formulation buffers (containing high concentrations of trehalose, polysorbate, ascorbic acid, histidine, etc). Some drawbacks of the existing tetrazole compound **1** species include the low water solubility and the lack of chemo- and regioselective functionalisation. Further experiments are underway to address these limitations. This work expands the scope of photoradiolabelling and provides an encouraging benchmark demonstrating that ⁸⁹Zr-mAbs can be produced using tetrazole photochemistry.

Acknowledgements

J.P.H. thanks the Swiss National Science Foundation (SNSF Professorship PP00P2_163683 and PP00P2_190093), the European Research Council (ERC-2015-StG, NanoSCAN-676904; ERC-2020-CoG, PhotoPHARMA, 101001734), the Swiss Cancer League (Krebsliga Schweiz; KLS-4257-08-2017), and the University of Zurich (UZH) for financial support. We thank all members of the Radiochemistry and Imaging Science group at UZH.

Keywords: tetrazoles • photochemistry • radiochemistry • antibodies • positron emission tomography

- [1] J. S. Clovis, A. Eckell, R. Huisgen, R. Sustmann, *Chem. Ber.* **1967**, *100*, 60–70.
- [2] J. P. Holland, M. Gut, S. Klingler, R. Fay, A. Guillou, *Chem. Eur. J.* **2020**, *26*, 33–48.
- [3] R. Huisgen, J. Sauer, M. Seidel, *Chem. Ber.* **1961**, *94*, 2503–2509.
- [4] W. Song, Y. Wang, J. Qu, M. M. Madden, Q. Lin, *Angew. Chemie - Int. Ed.* **2008**, *47*, 2832–2835.
- [5] C. P. Ramil, Q. Lin, *Curr. Opin. Chem. Biol.* **2014**, *21*, 89–95.
- [6] W. Song, Y. Wang, J. Qu, Q. Lin, *J. Am. Chem. Soc.* **2008**, *130*, 9654–9655.
- [7] Z. Yu, Y. Pan, Z. Wang, J. Wang, Q. Lin, *Angew. Chemie - Int. Ed.* **2012**, *51*, 10600–10604.
- [8] Y. Wang, W. J. Hu, W. Song, R. K. V. Lim, Q. Lin, *Org. Lett.* **2008**, *10*, 3725–3728.
- [9] Y. Wang, W. Song, W. J. Hu, Q. Lin, *Angew. Chemie - Int. Ed.* **2009**, *48*, 5330–5333.
- [10] R. K. V. Lim, Q. Lin, *Chem. Commun.* **2010**, *46*, 7993–7995.
- [11] Z. Yu, L. Y. Ho, Z. Wang, Q. Lin, *Bioorganic Med. Chem. Lett.* **2011**, *21*, 5033–5036.
- [12] Z. Yu, Q. Lin, *J. Am. Chem. Soc.* **2014**, *136*, 4153–4156.
- [13] X. Shang, R. Lai, X. Song, H. Li, W. Niu, J. Guo, *Bioconjugate Chem.* **2017**, *28*, 2859–2864.
- [14] H. Durchschlag, C. Fochler, B. Feser, S. Hausmann, T. Seroneit, M. Swientek, E. Swoboda, A. Winklmair, C. Wlček, P. Zipper, *Radiat. Phys. Chem.* **1996**, *47*, 501–505.
- [15] D. I. Pattison, A. S. Rahmanto, M. J. Davies, *Photochem. Photobiol. Sci.* **2012**, *11*, 38–53.
- [16] R. K. V. Lim, Q. Lin, *Acc. Chem. Res.* **2011**, *44*, 828–830.
- [17] Z. Li, L. Qian, L. Li, J. C. Bernhammer, H. V. Huynh, J.-S. Lee, S. Q. Yao, *Angew. Chemie Int. Ed.* **2016**, *55*, 2002–2006.
- [18] S. Zhao, J. Dai, M. Hu, C. Liu, R. Meng, X. Liu, C. Wang, T. Luo, *Chem. Commun.* **2016**, *52*, 4702–4705.
- [19] W. Siti, A. K. Khan, H.-P. M. De Hoog, B. Liedberg, M. Nallani, *Org. Biomol. Chem.* **2015**, *13*, 3202.
- [20] W. Feng, L. Li, C. Yang, A. Welle, O. Trapp, P. A. Levkin, *Angew. Chemie - Int. Ed.* **2015**, *54*, 8732–8735.
- [21] Y. Zhang, W. Liu, Z. Zhao, *Molecules* **2013**, *19*, 306–315.
- [22] R. S. Pandurangi, S. R. Karra, K. V. Katti, R. R. Kuntz, W. A. Volkert, *J. Org. Chem.* **1997**, *62*, 2798–2807.
- [23] R. S. Pandurangi, P. Lusiak, R. R. Kuntz, W. A. Volkert, J. Rogowski, M. S. Platz, *J. Org. Chem.* **1998**, *63*, 9019–9030.
- [24] R. Rajagopalan, R. R. Kuntz, U. Sharma, W. A. Volkert, R. S. Pandurangi, *J. Org. Chem.* **2002**, *67*, 6748–6757.
- [25] C. W. Lange, H. F. Vanbrocklin, S. E. Taylor, S. E. Taylor, *J. Label. Compd. Radiopharm. J Label Compd Radiopharm* **2002**, *45*, 257–268.
- [26] K. Hashizume, N. Hashimoto, Y. Miyake, *J. Org. Chem.* **1995**, *60*, 6680–6681.
- [27] M. Nishikawa, T. Nakano, T. Okabe, N. Hamaguchi, Y. Yamasaki, Y. Takakura, F. Yamashita, M. Hashida, *Bioconjug. Chem.* **2003**, *14*, 955–961.
- [28] M. A. Stalteri, S. J. Mather, *Eur. J. Nucl. Med.* **1996**, *23*, 178–187.
- [29] T. R. Sykes, T. K. Woo, R. P. Baum, P. Qi, A. A. Noujaim, *J. Nucl. Med.* **1995**, *36*, 1913–22.
- [30] T. R. Sykes, V. V. Somayaji, S. Bier, T. K. Woo, C. S. Kwok, V. Snieckus, A. A. Noujaim, *Appl. Radiat. Isot.* **1997**, *48*, 899–906.
- [31] H. J. Wester, K. Hamacher, G. Stöcklin, *Nucl. Med. Biol.* **1996**, *23*, 365–72.
- [32] L. Sun, J. Ding, W. Xing, Y. Gai, J. Sheng, D. Zeng, *Bioconjug. Chem.* **2016**, *27*, 1200–1204.
- [33] M. Patra, L. S. Eichenberger, G. Fischer, J. P. Holland, *Angew. Chemie - Int. Ed.* **2019**, *58*, 1928–1933.
- [34] R. Fay, M. Gut, J. P. Holland, *Bioconjug. Chem.* **2019**, *30*, 1814–1820.
- [35] L. S. Eichenberger, M. Patra, J. P. Holland, *Chem. Commun.* **2019**, *55*, 2257–2260.
- [36] S. Klingler, R. Fay, J. P. Holland, *J. Nucl. Med.* **2020**, *61*, 1072–1078.
- [37] S. Ito, Y. Tanaka, A. Kakehi, K. Kondo, *Bull. Chem. Soc. Jpn.* **1976**, *49*, 1920–1923.
- [38] Y. Zakon, L. Halicz, F. Gelman, *Environ. Sci. Technol.* **2013**, *47*, 14147–14153.
- [39] R. B. Woodward, *J. Am. Chem. Soc.* **1941**, *63*, 1123–1126.
- [40] L. F. Fieser, M. Fieser, S. Rajagopalan, *J. Org. Chem.* **1948**, *13*, 800–806.
- [41] K. A. Usmani, E. D. Karoly, E. Hodgson, R. L. Rose, *Drug Metab. Dispos.* **2004**, *32*, 333 LP-339.
- [42] D. J. Vugts, C. Klaver, C. Sewing, A. J. Poot, K. Adamzek, S. Huegeli, C. Mari, G. W. M. Visser, I. E. Valverde, G. Gasser, T. L. Mindt, G. A. M. S. van Dongen, *Eur. J. Nucl. Med. Mol. Imaging* **2017**, *44*, 286–295.
- [43] T. Lindmo, E. Boven, F. Cuttitta, J. Fedorko, P. A. Bunn, *J. Immunol. Methods* **1984**, *72*, 77–89.
- [44] J. P. Holland, E. Caldas-Lopes, V. Divilov, V. a Longo, T. Taldone, D. Zatorska, G. Chiosis, J. S. Lewis, *PLoS One* **2010**, *5*, e8859.
- [45] M. Patra, S. Klingler, L. S. Eichenberger, J. Holland, *iScience* **2019**, *13*, 416–431.

*Rachael Fay and Jason P. Holland**

Page No. – Page No.

Tuning tetrazole photochemistry for protein ligation and molecular imaging

Tetrazole-based photolabelling of antibodies for PET. Tuning the photophysical properties of di-aryl tetrazole scaffolds can induce a bathochromic shift in the photoactivation window that facilitates protein ligation at long wavelengths. Photoactivatable metal binding chelates functionalised with a tetrazole are developed for applications in light-induced photoradiolabelling of monoclonal antibodies for positron emission tomography.