

COHOMOTOPY INVARIANTS AND THE UNIVERSAL COHOMOTOPY INVARIANT JUMP FORMULA

CHRISTIAN OKONEK, ANDREI TELEMAN

ABSTRACT. Starting from ideas of Furuta, we develop a general formalism for the construction of cohomotopy invariants associated with a certain class of S^1 -equivariant non-linear maps between Hilbert bundles. Applied to the Seiberg-Witten map, this formalism yields a new class of cohomotopy Seiberg-Witten invariants which have clear functorial properties with respect to diffeomorphism of 4-manifolds. For 4-manifolds with $b_1 = 0$ and $b_+ > 1$ our invariants are equivalent to the Bauer-Furuta invariants, but they are finer in general. We study fundamental properties of the new invariants in a very general framework. In particular we prove a universal cohomotopy invariant jump formula and a multiplicative property. The formalism applies to other gauge theoretical problems, e.g. to the theory of gauge theoretical (Hamiltonian) Gromov-Witten invariants.

CONTENTS

1. Introduction	2
1.1. Motivation	2
1.2. Summary of results	4
2. Cohomotopy groups associated with elements in $K(B)$	6
2.1. Definition of ${}_{S^1}\alpha_B^*(X, Y)$	6
2.2. The computation of ${}_{S^1}\alpha^k(B_+, V^+)$	8
2.3. The groups $\alpha^*(x)$ associated with an element $x \in K(B)$	10
2.4. The S^1 -equivariant J -map and the description of $\alpha^*(x)$	14
2.5. Stabilization	17
2.6. The cohomotopy Euler class of an element in $K(B)$	17
3. Cohomotopy invariants associated with certain non-linear maps between Hilbert bundles	18
3.1. The cylinder construction	18
3.2. General properties of the invariant $\{\mu\}$	19
3.3. A class of non-linear maps between Hilbert bundles	22
3.4. The Seiberg-Witten map in dimension 4	23
3.5. Finite dimensional approximation	25
3.6. Compatibility properties	28
4. Fundamental properties of the cohomotopy invariants	30
4.1. The Hurewicz image of the cohomotopy invariant	30
4.2. Cohomotopy invariant jump formulae	35
4.3. A product formula and a vanishing theorem	42
5. Appendix	45
5.1. Inductive limits of functors	45
5.2. Bundle maps between pointed sphere bundles	49

1. INTRODUCTION

1.1. Motivation. The goal of this article is to develop a general formalism for the construction of cohomotopy invariants associated with a certain class of non-linear maps between Hilbert bundles. The main example we have in mind is the Seiberg-Witten map, but the formalism applies to other interesting classes of maps related to gauge theoretical problems as well.

The idea to define stable-homotopy Seiberg-Witten invariants is due to Furuta, who first used “finite dimensional approximations” of the monopole map in his work on the 11/8 conjecture [Fu1], and then introduced a class of refined Seiberg-Witten invariants (called “stable homotopy version of the Seiberg-Witten invariants”) in a geometric, non-formalized way in [Fu2]. In this preprint Furuta acknowledges independent work by Bauer [B3].

According to Furuta, the new invariants belong to a certain inductive limit of sets of homotopy classes of maps associated with “finite dimensional approximations” of the Seiberg-Witten map. The structure and the functorial properties of this inductive limit (with respect to diffeomorphisms between 4-manifolds) have not been worked out in this article. A precise version of the new invariants has been introduced later by Bauer-Furuta in [BF]: the Bauer-Furuta classes belong to certain stable cohomotopy groups associated with a presentation (E, F) of the K-theory element $\text{ind}(\not{D})$ defined by a fixed $Spin^c$ -structure. This element $\text{ind}(\not{D})$ belongs to the K-theory group $K(B)$, where $B = H^1(X; \mathbb{R})/H^1(X; \mathbb{Z})$ is the Picard group of the base manifold X .

In this article we propose a different construction of cohomotopy invariants which has the following advantages: Our construction yields a larger class of invariants, which in general are finer than the Bauer-Furuta classes, and have better functorial properties with respect to diffeomorphisms. In order to explain the advantages of the new formalism in a non-technical way, we consider again the Seiberg-Witten case.

It is well known that the Seiberg-Witten map μ can be regarded as an S^1 -equivariant bundle map between Hilbert bundles over the torus B (see [BF] and section 3.4 of this article). We first choose the perturbing form in the second Seiberg-Witten equation in the “bad way”, i.e. such that the equations have reducible solutions (solutions with trivial spinor component); we make this “bad choice” even in the case $b_+(X) > 1$! In “classical” Seiberg-Witten theory one perturbs the second Seiberg-Witten equation using a nontrivial self-dual harmonic form $\kappa \in i\mathbb{H}^+ \setminus \{0\}$, and gets a new map μ_κ which defines a moduli space which does not contain reductions. Instead of a constant perturbation κ , we consider a map $\kappa : B \rightarrow i\mathbb{H}^+ \setminus \{0\}$, and perturb the Seiberg-Witten map μ (regarded as bundle map over B) using this map. The associated invariant will depend on the homotopy class $[\kappa] \in [B, S(i\mathbb{H}^+)]$. This leads to the following questions:

- (1) Does one obtain new invariants in this way?
- (2) If so, does one have a *universal cohomotopy invariant jump formula*, i.e. a formula which describes the jump of the cohomotopy invariant when one passes from one homotopy class to another?

- (3) Use again constant perturbation forms κ , but let κ vary in the sphere $S(i\mathbb{H}^+)$ and regard the obtained map $\tilde{\mu}$ as an S^1 -equivariant bundle map over the larger basis $B \times S(i\mathbb{H}^+)$. Does this universal perturbation $\tilde{\mu}$ yield more differential topological information than the individual perturbations μ_κ ? If not, express the cohomotopy invariant associated with $\tilde{\mu}$ in terms of the invariant associated with μ_κ and topological invariants of X .

These questions are interesting as soon as $b_1 \geq b_+ - 1$ (even for $b_+ > 1!$) and they are also interesting *for the classical invariant*, because for non-constant perturbations κ one gets new Seiberg-Witten type moduli spaces. The universal wall-crossing formula for the *full Seiberg-Witten invariant*¹ [OT] should be a formal consequence of a universal cohomotopy invariant jump formula. These questions will be completely answered in this article.

Another motivation for proposing a new formalism was the need to have *well defined* invariants, with clear *functorial properties* with respect to diffeomorphisms of 4-manifolds. The point here is that the definition of the stable cohomotopy group used in [BF] depends on the choice of a presentation $(E, B \times \mathbb{C}^n)$ of $\text{ind}(\mathcal{D}) \in K(B)$ (see [BF] p. 8-9). Since in general such a presentation has homotopically non-trivial automorphisms, the obtained cohomotopy groups cannot be regarded as invariants of the K-theory element $\text{ind}(\mathcal{D}) \in K(B)$. This makes it difficult to control the functorial properties of the Bauer-Furuta stable cohomotopy groups with respect to homeomorphisms or diffeomorphisms of 4-manifolds, and to understand in which sense the constructed class is well defined. Using Segal cocycles instead of finite rank presentations ([BF] p. 7-8) does not remove the problem, because of monodromy phenomena in the space of Segal cocycles². A similar difficulty concerns the concept “Thom spectrum of a virtual bundle”, used by Bauer-Furuta (see [BF] p. 8) and other authors in order to give a geometric interpretation of the Bauer-Furuta classes. One can indeed associate a Thom spectrum to a *fixed presentation* $(E, B \times \mathbb{C}^n)$ of a K-theory element $x \in K(B)$, but unfortunately *not to x itself*. For 4-manifolds with $b_1 = 0$, the Bauer-Furuta class gives a well defined invariant, which can easily be identified with the image of our invariant under a boundary morphism of cohomotopy groups. The two invariants are equivalent when $b_1 = 0$, $b_+ > 1$.

Our new point of view has the following advantages:

- (1) The new cohomotopy Seiberg-Witten invariants are finer than the full classical Seiberg-Witten invariants *in all cases*, including the case of manifolds with $b_1 \geq b_+ - 1$ and including the invariants associated with non-constant perturbations $\kappa : B \rightarrow i\mathbb{H}_+^2 \setminus \{0\}$. In the case $b_1 \geq b_+ - 1$ we prove a universal cohomotopy invariant jump formula; the universal wall-crossing formula for the classical invariant is a formal consequence of this result.

¹The full Seiberg-Witten invariant is an element in $\Lambda^*(H^1(X; \mathbb{Z}))$ [OT]. The numerical Seiberg-Witten invariant (the original invariant introduced by Witten) is the degree 0 term of the full invariant.

²Contrary to what is often stated in the literature, the space of possible Dirac operators associated with a fixed equivalence class of $Spin^c$ -structures is not contractible (see section 3.4). So even if one considers only $Spin^c$ -Dirac operators, one does not get a contractible space of Segal cocycles.

- (2) Our invariant belongs to a cohomotopy group which is intrinsically associated with the base 4-manifold (and a set of topological data), and has clear functorial properties with respect to homeomorphisms of 4-manifolds.

1.2. Summary of results. In the first section we construct a graded cohomotopy group associated with a K-theory element $x \in K(B)$. To every representative $(E, F) \in x$ we associate the graded group ${}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+)$, where ${}_{S^1}\alpha_B^*(\cdot, \cdot)$ stands for the S^1 -equivariant stable cohomotopy group functor on the category of pointed S^1 -spaces over B ; it is obtained by stabilizing with spaces of the form $(\eta \oplus \xi_0)_B^+$, where η is a complex, and ξ_0 a real bundle. Note that we do not use all characters of S^1 in the stabilizing process; for this reason we do not use the standard notation ${}_{S^1}\omega_B^*$ found in the literature [CJ]. We define $\alpha^*(x)$ to be the inductive limit of the graded groups ${}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+)$ with respect to the category $\mathcal{T}(x)$ of representatives (E, F) of x . Since $\mathcal{T}(x)$ is not a small filtering category (see [AM], and section 5.1 below), this limit cannot be obtained using the classical construction. It will be constructed in two steps: First we stabilize the graded group ${}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+)$ with respect to standard representative enlargements $(E, F) \mapsto (E \oplus U, F \oplus U)$, and we obtain a new graded group $\hat{\alpha}^*(E, F)$, which still depends on the fixed pair (E, F) . The groups $\hat{\alpha}^*(E, F)$, $\hat{\alpha}^*(E', F')$ defined by two representatives (E, F) , (E', F') of x are *non-canonically* isomorphic. The group $\alpha^*(x)$ will be the quotient of $\hat{\alpha}^*(E, F)$ by the equivalence relation generated by the inductive limit of the automorphism groups $\text{Aut}(E \oplus U) \times \text{Aut}(F \oplus U)$. We give an explicit description of $\alpha^*(x)$ as a quotient of the group $\hat{\alpha}^*(E, F)$ associated with any representative (E, F) by the action of the image of the J -homomorphism ${}_{S^1}J : K^{-1}(B) \rightarrow {}_{S^1}\alpha^0(B)^\times$ in the group of units ${}_{S^1}\alpha^0(B)^\times$ of the ground ring ${}_{S^1}\alpha^0(B) := {}_{S^1}\alpha_B^0(B_{+B}, B_{+B})$. In other words, we are able to control the effect of bundle automorphisms in our inductive limit and we obtain a graded group which is intrinsically associated with the K-theory element x . We believe that this construction is of independent interest from the point of view of homotopy theory.

A way to understand the role of the graded group $\alpha^*(x)$ is the following: Because of the presence of homotopically non-trivial bundle automorphisms, one cannot define the projectivization $\mathbb{P}(x)$ of a K-theory element $x \in K(B)$ (neither in the category of topological spaces nor in the category of spectra). *The graded group $\alpha^*(x)$ plays the role of what should be the cohomotopy group of a formal projectivization of the K-theory element x .*

In the second section we first introduce a distinguished class of non-linear maps μ between Hilbert bundles over a compact base B . The \mathbb{C} -linear part of the linearization of such a map μ at the zero section is a linear Fredholm operator, so it defines a K-theory element $x \in K(B)$. The goal of the section is the construction of an invariant $\{\mu\} \in \alpha^*(x)$. This invariant is constructed using *finite dimensional approximations* of the map μ . In order to get these approximations we make use of the retractions $\rho_A : \mathcal{A}^+ \setminus S(\mathcal{A}^\perp) \rightarrow \mathcal{A}^+$ associated with finite dimensional subspaces A of a Hilbert space \mathcal{A} , as in [BF]. This method to construct finite dimensional approximations applies to a very large class of non-linear maps, whereas Furuta's original method based on L^2 -orthogonal projections on direct sums of eigenspaces (see [Fu2]) is limited to maps whose linearizations are elliptic differential operators.

The main difference between our definition and the construction of the Bauer-Furuta classes given in [BF], is that

- (1) our construction uses only spaces fibered over the base B . In particular we avoid using Thom spaces,
- (2) we treat the real and the complex summands in our finite dimensional approximations separately.

Therefore, from this point of view, our construction is *closer to the original ideas of Furuta* [Fu2]. Having the finite dimensional approximation, a representative of the invariant is an element in a group of the form ${}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+)$ obtained by a simple geometric construction, which we call *the cylinder construction*³. The obtained class carries more information than the one defined in [BF]. In 3.4 we show that the Seiberg-Witten map associated with a Spin^c -structure τ on a Riemannian 4-manifold M with $b_+(M) > 0$ yields a non-linear Fredholm map sw_κ (depending on a twisting map $\kappa : B = H^1(X; \mathbb{R})/H^1(X; \mathbb{Z}) \rightarrow i\mathbb{H}_+^2 \setminus \{0\}$) which belongs to our distinguished class of maps. Hence the general theory applies and yields a cohomotopy Seiberg-Witten invariant $\{sw_\kappa\} \in \alpha^{b_+(M)-1}(\text{ind}(\not{D}))$, which only depends of the homotopy class of κ . Our construction of the bundle map sw_κ is different from (and more “economical” than) the one given in [BF]. In this section we also explain why the space of Dirac operators associated with a fixed equivalence class of Spin^c -structures is not contractible, so there is no way to distinguish a contractible class of Segal cocycles defining the K -theory element $\text{ind}(\not{D})$. This makes clear why the cohomotopy groups defined in [BF] (which depend essentially on the choice of a Segal cocycle) cannot be regarded as intrinsic (functorial) invariants of the base manifold.

In the third section we prove several fundamental properties of the invariant $\{\mu\} \in \alpha^*(x)$ in our general, abstract framework:

- (1) We study the image of our invariant under the Hurewicz morphism, and we prove that the Poincaré dual of this image can be identified with the virtual fundamental class of the vanishing locus. In other words, the full homology invariant associated with the virtual fundamental class of the “moduli space” (i.e. the S^1 -quotient of the vanishing locus of μ) can be identified with the Hurewicz image of the cohomotopy invariant. Moreover, the Hurewicz morphism is an isomorphism when the “expected dimension” vanishes.
- (2) We prove a formal universal cohomotopy invariant jump formula for our refined cohomotopy invariants.
- (3) We prove a general product formula for the invariant $\{\mu_1 \times \mu_2\}$ associated with a product of maps; in this formula we allow one of the factors to have zeros on the S^1 -fixed point locus. When both factors are nowhere vanishing on their fixed point loci, we prove a vanishing result for the Hurewicz image of the invariant.

Specialized to the Seiberg-Witten map, these properties automatically yield important results for the new cohomotopy Seiberg-Witten invariants. The first result shows that the cohomotopy Seiberg-Witten invariant is refinement of the classical *full* Seiberg-Witten invariants in all cases. Combined with the second property, this also yields a universal invariant jump formula for the full classical Seiberg-Witten

³After completing the first version of this article we found out about the results [C] on Leray-Schauder index theory. Under the assumption that the group-action is free on a neighborhood of the zero-set of the vector field, one can define a refinement of the usual Poincaré-Hopf vector field index, which is probably related to our refinement of the Bauer-Furuta class.

invariant in the case $b_1(X) \geq b_+(X) - 1$. The third result gives a formula for the cohomotopy invariant of a connected sum of two 4-manifolds, even in the case when one term of the sum has $b_+ = 0$. The vanishing result in (2) reproves the classical vanishing theorem for the Seiberg-Witten invariant of a direct sum in the case where both summands X_i have $b_+(X_i) > 0$. These applications in Seiberg-Witten theory are not detailed in this article.

Acknowledgment: The first author would like to thank Tammo tom Dieck for useful and interesting discussions at the beginning of this project. We also thank Michael Crabb for pointing us to his paper [C] and for suggesting a relation between our invariant and his refinement of the Poincaré-Hopf vector field index.

2. COHOMOTOPY GROUPS ASSOCIATED WITH ELEMENTS IN $K(B)$

2.1. Definition of ${}_{S^1}\alpha_B^*(X, Y)$. Let B be a compact topological space endowed with the trivial S^1 -action. Let \mathcal{C}_B be the category defined in the following way: the objects of \mathcal{C}_B are vector bundles over B of the form

$$\xi = \eta \oplus \xi_0 ,$$

where η is a complex vector bundle endowed with the standard S^1 -action and ξ_0 is a real vector bundle endowed with the trivial S^1 -action; for two objects $\xi = \eta \oplus \xi_0$, $\xi' = \eta' \oplus \xi'_0$ a morphism $\nu : \xi \rightarrow \xi'$ is a pair (i, ζ) consisting of an S^1 -equivariant bundle embedding $i = \iota \oplus i_0 : \xi \rightarrow \xi'$ and a complement $\zeta = \nu \oplus \zeta_0$ of $i(\xi)$ in ξ' . Composition of morphisms is defined in a natural way. A morphism $u = (i, \zeta) : \xi \rightarrow \xi'$ defines a push-forward morphism $A(u) : A(\xi) \rightarrow A(\xi')$, where $A(\xi) := A(\eta) \times A(\xi_0)$ is the automorphism group of ξ . We obtain in this way a functor $A : \mathcal{C}_B \rightarrow \mathcal{G}r$. In the terminology of section 5.1, the pair (\mathcal{C}_B, A) is a category with automorphism push-forward.

Let $X \rightarrow B, Y \rightarrow B$ be two pointed S^1 -spaces over B . The assignment

$$\xi \mapsto {}_{S^1}\pi_B^0(X \wedge_B \xi_B^+, Y \wedge_B \xi_B^+)$$

(where ${}_{S^1}\pi_B^0(X, Y)$ stands for the set of homotopy classes of S^1 -equivariant base point preserving maps over B) is functorial with respect to morphisms in \mathcal{C}_B : for a morphism $u = (i, \zeta) : \xi \rightarrow \xi'$, the push-forward class $u_*([f])$ is defined using $i \circ f \circ i^{-1}$ on $i(\xi)$ and id_ζ on its complement ζ . Therefore this assignment defines a functor ${}_{S^1}\pi_B^0(X \wedge_B \cdot, Y \wedge_B \cdot) : \mathcal{C}_B \rightarrow \mathcal{S}ets$. It is not clear at all that an inductive limit of this functor exists, because $\mathcal{O}b(\mathcal{C}_B)$ is neither a filtering nor a small category (see section 5.1).

Proposition 2.1. *Let $\xi = \eta \oplus \xi_0 \in \mathcal{O}b(\mathcal{C}_B)$, $\mathbf{a} = (\alpha, a_0) \in A(\xi)$, and $u = (i, \zeta)$ the standard morphism $\eta \oplus \xi_0 = \xi \rightarrow \tilde{\xi} := (\eta \oplus \eta) \oplus (\xi_0 \oplus \xi_0)$ defined by $(y, x) \mapsto ((y, 0), (x, 0))$. For every $[f] \in {}_{S^1}\pi_B^0(X \wedge_B \xi_B^+, Y \wedge_B \xi_B^+)$ one has*

$$u_*(\mathbf{a}_*([f])) = u_*([f]) .$$

Proof: Identifying $\tilde{\xi}$ with $\xi \oplus \xi$ one can write $u_*(\mathbf{a}_*[f]) = [g]$ where g is the composition

$$(\text{id}_X \wedge_B [\mathbf{a} \oplus \text{id}_\xi]_B^+) \circ (f \wedge_B \text{id}_{\xi_B^+}) \circ (\text{id}_X \wedge_B [\mathbf{a}^{-1} \oplus \text{id}_\xi]_B^+) : X \wedge_B [\xi \oplus \xi]_B^+ \rightarrow Y \wedge_B [\xi \oplus \xi]_B^+ .$$

Let R_t be the automorphism of $\xi \oplus \xi$ defined by the matrix

$$r_t := \begin{pmatrix} \cos(t\frac{\pi}{2}) & -\sin(t\frac{\pi}{2}) \\ \sin(t\frac{\pi}{2}) & \cos(t\frac{\pi}{2}) \end{pmatrix} .$$

For an automorphism \mathfrak{b} of ξ note that $r_t \circ (\mathfrak{b} \oplus \text{id}_\xi) \circ r_t^{-1}$ defines a homotopy between $\mathfrak{b} \oplus \text{id}_\xi$ and $\text{id}_\xi \oplus \mathfrak{b}$. This shows that g is homotopic to the map

$$g' := (\text{id}_X \wedge_B [\text{id}_\xi \oplus \mathfrak{a}]_B^+) \circ (f \wedge_B \text{id}_{\xi_B^+}) \circ (\text{id}_X \wedge_B [\text{id}_\xi \oplus \mathfrak{a}^{-1}]_B^+) = f \wedge_B \text{id}_{\xi_B^+}$$

which is a representative of the class $u_*([f])$. \blacksquare

We define the stable cohomotopy group ${}_{S^1}\alpha_B^0(X, Y)$ by

$${}_{S^1}\alpha_B^0(X, Y) := \varinjlim_{(n, m) \in \mathbb{N}^2} {}_{S^1}\pi_B^0(X \wedge_B [\underline{\mathbb{C}}^n \oplus \underline{\mathbb{R}}^m]_B^+, Y \wedge_B (\underline{\mathbb{C}}^n \oplus \underline{\mathbb{R}}^m)_B^+).$$

In this formula and in the rest of the paper we use the notation \underline{V} for the trivial bundle $B \times V$ over the base B . This inductive limit has a natural Abelian group structure (see [CJ] p. 168 for the non-equivariant case).

Proposition 2.2. *The functor ${}_{S^1}\pi_B^0(X \wedge_B \cdot, Y \wedge_B \cdot) : \mathcal{C}_B \rightarrow \mathcal{S}ets$ admits an inductive limit, which can be identified with ${}_{S^1}\alpha_B^0(X, Y)$.*

Proof: Let \mathcal{N}^2 be the small category associated with the ordered set $(\mathbb{N} \times \mathbb{N}, \leq)$ and consider the functor $\Theta : \mathcal{N}^2 \rightarrow \mathcal{C}_B$ which assigns to a pair (n, m) the trivial bundle $\underline{\mathbb{C}}^n \oplus \underline{\mathbb{R}}^m$ over B , and to an inequality $(n, m) \leq (n', m')$ the standard morphism between the corresponding trivial bundles. Using the terminology of section 5.1, \mathcal{N} is a small filtering category, and Θ is a cofinal functor from \mathcal{N} to the category (\mathcal{C}_B, A) , which is a category with automorphism push-forward. By definition ${}_{S^1}\alpha_B^0(X, Y)$ is just the limit of the composed functor ${}_{S^1}\pi_B^0(X \wedge_B \cdot, Y \wedge_B \cdot) \circ \Theta$. On the other hand, Proposition 2.1 shows that the functor ${}_{S^1}\pi_B^0(X \wedge_B \cdot, Y \wedge_B \cdot)$ satisfies the “trivial stable actions” axioms TSA, Θ SA. The result follows therefore from Proposition 5.11 in section 5.1. \blacksquare

Note that Proposition 2.2 implicitly yields a canonical map

$$c_\xi : {}_{S^1}\pi_B^0(X \wedge_B \xi_B^+, Y \wedge_B \xi_B^+) \rightarrow {}_{S^1}\alpha_B^0(X, Y)$$

for every $\xi \in \mathcal{O}(\mathcal{C}_B)$, such that the system $(c_\xi)_{\xi \in \mathcal{O}(\mathcal{C}_B)}$ satisfies the universal property of the inductive limit.

As in the non-equivariant case we put

$${}_{S^1}\alpha_B^p(X, Y) := {}_{S^1}\alpha_B^0(X \wedge_B (\underline{\mathbb{R}}^N)_B^+, Y \wedge_B (\underline{\mathbb{R}}^{N+p})_B^+) \quad (N, N+p \geq 0).$$

Each ${}_{S^1}\alpha_B^p(X, Y)$ is a bimodule over the ring

$${}_{S^1}\alpha^0(B) := {}_{S^1}\alpha^0(B_+, S^0) = {}_{S^1}\alpha_B^0(B_{+B}, B_{+B}),$$

and ${}_{S^1}\alpha_B^*(X, Y) := \bigoplus_{p \in \mathbb{Z}} {}_{S^1}\alpha_B^p(X, Y)$ is a graded bimodule over the graded ring ${}_{S^1}\alpha^*(B) = \bigoplus_{S^1}\alpha^p(B)$, where

$${}_{S^1}\alpha^p(B) := {}_{S^1}\alpha^p(B_+, S^0) = {}_{S^1}\alpha^0(B_+, S^p).$$

Right and left multiplication with elements in ${}_{S^1}\alpha^0(B)$ coincide (see [CJ] p. 172).

Remark 2.3. *In the special case when Y is of the form $Y = \zeta_B^+$ with $\zeta \in \mathcal{C}_B$, one has a canonical identification*

$${}_{S^1}\alpha_B^0(X, \zeta_B^+) = {}_{S^1}\alpha^0\left(X \wedge_B [\zeta']_B^+ / \infty, V^+\right),$$

where $\zeta \oplus \zeta' = \underline{V}$, and V has the form $\mathbb{C}^k \oplus \mathbb{R}^l$. In the terminology of [BF] the latter group is a stable cohomotopy group formed with respect to the universum generated by the S^1 -representations \mathbb{C} and \mathbb{R} .

2.2. The computation of ${}_{S^1}\alpha^k(B_+, V^+)$. Let $S^1 \rightarrow O(V)$ be an orthogonal representation of S^1 . Our next goal is the computation of the group ${}_{S^1}\alpha^k(B_+, V^+)$ for $k \geq 0$. In particular, we obtain explicit descriptions of the positive summands ${}_{S^1}\alpha^k(B) = {}_{S^1}\alpha^k(B_+, [\mathbb{R}^k]^+)$ of the graded ring ${}_{S^1}\alpha^*(B)$.

Replacing V by $V \oplus \mathbb{R}^k$, we can reduce the problem to the case $k = 0$. One has

$${}_{S^1}\alpha^0(B_+, V^+) = \varinjlim_{(n,m) \in \mathbb{N}^2} [B_+ \wedge [\mathbb{C}^n \oplus \mathbb{R}^m]^+, V^+ \wedge [\mathbb{C}^n \oplus \mathbb{R}^m]^+]_0^{S^1},$$

where $[\cdot, \cdot]_0^{S^1}$ stands for the set of homotopy classes of S^1 -equivariant maps between two pointed S^1 -spaces.

According to Hauschild's splitting theorem (Satz 3.4 in [H]) there is a natural identification

$$[B_+ \wedge [\mathbb{C}^n \oplus \mathbb{R}^m]^+, [V \oplus \mathbb{C}^n \oplus \mathbb{R}^m]^+]_0^{S^1} = \quad (1)$$

$$[B_+ \wedge [\mathbb{R}^m]^+, [V^{S^1}]^+ \wedge [\mathbb{R}^m]^+]_0 \times \left[B_+ \wedge \left[\frac{[\mathbb{C}^n \oplus \mathbb{R}^m]^+}{[\mathbb{R}^m]^+} \right], V^+ \wedge [\mathbb{C}^n \oplus \mathbb{R}^m]^+ \right]_0^{S^1}$$

where the projection on the first factor is given by restriction to the fixed point set. There exists a homeomorphism of S^1 -spaces

$$\frac{[\mathbb{C}^n \oplus \mathbb{R}^m]^+}{[\mathbb{R}^m]^+} \approx S(\mathbb{C}^n)_+ \wedge S^{m+1}.$$

Indeed, one has

$$\begin{aligned} \frac{[\mathbb{C}^n \oplus \mathbb{R}^m]^+}{[\mathbb{R}^m]^+} &\approx S(\mathbb{C}^n \oplus \mathbb{R}^{m+1}) / S(\mathbb{R}^{m+1}) \approx \\ &\approx S(\mathbb{C}^n) \times D(\mathbb{R}^{m+1}) \cup D(\mathbb{C}^n) \times S(\mathbb{R}^{m+1}) / D(\mathbb{C}^n) \times S(\mathbb{R}^{m+1}) \approx S(\mathbb{C}^n)_+ \wedge S^{m+1}. \end{aligned}$$

Using the natural identification

$$B_+ \wedge [S(\mathbb{C}^n)_+ \wedge S^{m+1}] \approx S(\mathbb{C}^n)_+ \wedge [B_+ \wedge S^{m+1}] \approx S(\mathbb{C}^n) \times [B_+ \wedge S^{m+1}] / S(\mathbb{C}^n) \times \{*\}$$

and denoting by \tilde{V}_n the associated bundle $S(\mathbb{C}^n) \times_{S^1} V$ over $\mathbb{P}(\mathbb{C}^n)$ we find

$$\begin{aligned} \left[B_+ \wedge \left[\frac{[\mathbb{C}^n \oplus \mathbb{R}^m]^+}{[\mathbb{R}^m]^+} \right], V^+ \wedge [\mathbb{C}^n \oplus \mathbb{R}^m]^+ \right]_0^{S^1} &\cong \\ &\cong \left[S(\mathbb{C}^n) \times [B_+ \wedge S^{m+1}] / S(\mathbb{C}^n) \times \{*\}, V^+ \wedge [\mathbb{C}^n \oplus \mathbb{R}^m]^+ \right]_0^{S^1} \cong \\ &\cong {}_{S^1}\pi_{S(\mathbb{C}^n)}^0 \left(S(\mathbb{C}^n) \times [B_+ \wedge S^{m+1}], S(\mathbb{C}^n) \times [V \oplus \mathbb{C}^n \oplus \mathbb{R}^m]^+ \right) \cong \\ &\cong \pi_{\mathbb{P}(\mathbb{C}^n)}^0 \left(\mathbb{P}(\mathbb{C}^n) \times [B_+ \wedge S^{m+1}], [\tilde{V}_n \oplus \mathcal{O}_{\mathbb{P}(\mathbb{C}^n)}(1)^{\oplus n} \oplus \mathbb{R}^m]_{\mathbb{P}(\mathbb{C}^n)}^+ \right) \cong \\ &\cong \pi_{\mathbb{P}(\mathbb{C}^n)}^0 \left([\mathbb{P}(\mathbb{C}^n) \times [B_+ \wedge S^1]] \wedge_{\mathbb{P}(\mathbb{C}^n)} \underline{\mathcal{S}}^m, [\tilde{V}_n \oplus \mathcal{O}_{\mathbb{P}(\mathbb{C}^n)}(1)^{\oplus n}]_{\mathbb{P}(\mathbb{C}^n)}^+ \wedge_{\mathbb{P}(\mathbb{C}^n)} \underline{\mathcal{S}}^m \right). \end{aligned}$$

The limit over m of this set is

$$\omega_{\mathbb{P}(\mathbb{C}^n)}^0 \left(\mathbb{P}(\mathbb{C}^n) \times [B_+ \wedge S^1], [\tilde{V}_n \oplus \mathcal{O}_{\mathbb{P}(\mathbb{C}^n)}(1)^{\oplus n}]_{\mathbb{P}(\mathbb{C}^n)}^+ \right).$$

Now note that

$$\tilde{V}_n \oplus \underline{\mathbb{C}} \oplus T_{\mathbb{P}(\mathbb{C}^n)} \cong \tilde{V}_n \oplus \mathcal{O}_{\mathbb{P}(\mathbb{C}^n)}(1)^{\oplus n} .$$

Therefore, applying the duality isomorphism given in Proposition 12.41 [CJ] to the map $\pi : \mathbb{P}(\mathbb{C}^n) \rightarrow \{*\}$, one gets

$$\begin{aligned} \omega_{\mathbb{P}(\mathbb{C}^n)}^0 \left(\mathbb{P}(\mathbb{C}^n) \times [B_+ \wedge S^1], [\tilde{V}_n \oplus \mathcal{O}_{\mathbb{P}(\mathbb{C}^n)}(1)^{\oplus n}]_{\mathbb{P}(\mathbb{C}^n)}^+ \right) &\cong \omega^0(B_+ \wedge S^1, \pi_*([\tilde{V}_n \oplus \underline{\mathbb{C}}]_{\mathbb{P}(\mathbb{C}^n)}^+)) \\ &\cong \omega^0(B_+ \wedge S^1, T(\tilde{V}_n \oplus \underline{\mathbb{C}})) \cong \omega^0(B_+ \wedge S^1, T(\tilde{V}_n) \wedge S^2) \cong \omega^0(B_+, T(\tilde{V}_n) \wedge S^1) , \end{aligned}$$

where $T(\cdot)$ stands for the Thom space functor. This shows that

$$\varinjlim_{(n,m) \in \mathbb{N}^2} \left[B_+ \wedge \left[[\mathbb{C}^n \oplus \mathbb{R}^m]^+ / [\mathbb{R}^m]^+ \right], V^+ \wedge [\mathbb{C}^n \oplus \mathbb{R}^m]^+ \right]_0^{S^1} \cong \omega^0(B_+, T(ES^1 \times_{S^1} V) \wedge S^1)$$

where $ES^1 \times_{S^1} V$ is the vector bundle associated with the universal S^1 -bundle $ES^1 \rightarrow BS^1 = \mathbb{P}^\infty$ and the fiber V . Using formula (1) we obtain the following

Proposition 2.4. *One has a natural group isomorphism*

$${}_{S^1}\alpha^0(B_+, V^+) \cong \omega^0(B_+, [V^{S^1}]^+) \times \omega^0(B_+, T(ES^1 \times_{S^1} V) \wedge S^1) . \quad (2)$$

where the projection on the first factor is given by restriction to the fixed point set. In particular

$${}_{S^1}\alpha^k(B) \cong \omega^k(B) \times \omega^k(B_+, \mathbb{P}_+^\infty \wedge S^1) .$$

Remark 2.5. *The second summand in the decomposition*

$${}_{S^1}\alpha^0(B) \cong \omega^0(B) \times \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1)$$

is called “the free summand” in [CK]. The projection ${}_{S^1}\alpha^0(B) \rightarrow \omega^0(B)$ is given by restriction to the fixed point set, hence it is a ring homomorphism. Therefore the free summand $\omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1)$ is an ideal of ${}_{S^1}\alpha^0(B)$, and one has a natural ring isomorphism

$$\omega^0(B) \simeq {}_{S^1}\alpha^0(B) / \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1) .$$

Corollary 2.6. *Suppose that B is a finite CW complex. Restriction to the fixed point set defines an isomorphism*

$$\varinjlim_{N \in \mathbb{N}} {}_{S^1}\alpha^k(B_+, [\mathbb{C}^N]^+) \xrightarrow{\cong} \omega^k(B) .$$

Proof: Indeed, taking $V = \mathbb{C}^N \oplus \mathbb{R}^k$, the second summand in (2) is:

$$\omega^0(B_+, T(ES^1 \times_{S^1} [\mathbb{C}^N \oplus \mathbb{R}^{k+1}])) = \varinjlim_{l \in \mathbb{N}} \pi^0(B_+ \wedge [\mathbb{R}^l]^+, T(ES^1 \times_{S^1} [\mathbb{C}^N \oplus \mathbb{R}^{k+1+l}]))$$

Recall that the Thom space of a real vector bundle of rank r over a CW complex X admits a CW decomposition consisting of a single 0-dimensional cell and cells of dimension $\geq r$. Therefore, for N sufficiently large any map $B_+ \wedge [\mathbb{R}^l]^+ \rightarrow T(ES^1 \times_{S^1} [\mathbb{C}^N \oplus \mathbb{R}^{k+1+l}])$ is homotopically trivial. \blacksquare

2.3. The groups $\alpha^*(x)$ associated with an element $x \in K(B)$. Fix an element $x \in K(B)$. We define a category $\mathcal{T}(x)$ in the following way: the objects of $\mathcal{T}(x)$ are the presentations of x . For two such presentations (E, F) , (E', F') , a morphism $\tau : (E, F) \rightarrow (E', F')$ is a system $\tau = (i, j, E_1, F_1, k)$ consisting of bundle monomorphisms $j : E \hookrightarrow E'$, $i : F \hookrightarrow F'$, complements E_1 and F_1 of $i(E)$ and $j(F)$ in E' and F' respectively, and an isomorphism $k : E_1 \rightarrow F_1$.

With every $(E, F) \in x$ we associate the graded group ${}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+)$. We claim that a morphism $\tau : (E, F) \rightarrow (E', F')$ induces a morphism

$$\tau_* : {}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+) \longrightarrow {}_{S^1}\alpha_B^*(S(E')_{+B}, [F']_B^+) .$$

Note first that, for Euclidean or Hermitian vector spaces V, W , one has a contraction

$$S(V \oplus W) \rightarrow S(V)_+ \wedge W^+$$

induced by the map

$$\begin{aligned} c : S(V \oplus W) &= [S(V) \times D(W)] \cup_{S(V) \times S(W)} [D(V) \times S(W)] \longrightarrow \\ &\longrightarrow S(V) \times D(W) / S(V) \times S(W) \simeq S(V) \times W^+ / S(V) \times \infty_W = S(V)_+ \wedge W^+ . \end{aligned}$$

It is useful to have explicit analytic formulae for the contraction map c . One can define W^+ in two equivalent ways: as the one-point compactification of W , and as the quotient $D(W)/S(W)$. Accordingly, the contraction maps $c, c' : S(V \oplus W) \rightarrow S(V)_+ \wedge W^+$ are given by the formulae:

$$c(v, w) = \begin{cases} \left(\frac{1}{\|v\|} v, \frac{1}{\sqrt{1-\|w\|^2}} w \right) & v \neq 0 \\ * & v = 0 \end{cases} , \quad c'(v, w) = \begin{cases} \left(\frac{1}{\|v\|} v, w \right) & v \neq 0 \\ * & v = 0 . \end{cases} \quad (3)$$

To save on notations we will still write c instead of c' when the second definition of W^+ is used.

Therefore, in the presence of a morphism $\tau = (i, j, E_1, F_1, k) : (E, F) \rightarrow (E', F')$ one gets a map

$$S(E')_{+B} = S(i(E) \oplus E_1)_{+B} \xrightarrow{c} S(i(E))_{+B} \wedge_B (E_1)_B^+ ,$$

which is well defined up to homotopy (the section ${}_{+B}$ on the left is mapped fiberwise to the distinguished section on the right). We obtain morphisms

$$\begin{aligned} & {}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+) \xrightarrow{(i,j)\simeq} {}_{S^1}\alpha_B^*(S(i(E))_{+B}, j(F)_B^+) = \\ &= {}_{S^1}\alpha_B^*(S(i(E))_{+B} \wedge_B (F_1)_B^+, j(F)_B^+ \wedge_B (F_1)_B^+) = {}_{S^1}\alpha_B^*(S(i(E))_{+B} \wedge_B (F_1)_B^+, (F')_B^+) \\ & \xrightarrow{k} {}_{S^1}\alpha_B^*(S(i(E))_{+B} \wedge_B (E_1)_B^+, (F')_B^+) \xrightarrow{c^*} {}_{S^1}\alpha_B^*(S(E')_{+B}, (F')_B^+) . \end{aligned}$$

The composition of these maps will be denoted by τ_* . One checks that τ_* is a morphism of ${}_{S^1}\alpha^*(B)$ -modules and that, for any two composable morphisms τ, τ' , one has

$$(\tau' \circ \tau)_* = \tau'_* \circ \tau_* .$$

In other words, the assignment $(E, F) \mapsto {}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+)$ is functorial, so it defines a functor $\mathbf{a}_x^* : \mathcal{T}(x) \rightarrow \mathcal{A}b^*$, where $\mathcal{A}b^*$ is the category of graded Abelian groups.

Example: Suppose that the stable class $\varphi \in {}_{S^1}\alpha_B^0(S(E)_{+B}, F_B^+)$ is represented by an S^1 -equivariant map $f : S(E) \rightarrow F_B^+$ over B (or, equivalently, by an S^1 -equivariant map $S(E)_{+B} \rightarrow F_B^+$ of pointed spaces over B). Let U be a complex

vector bundle over B and let τ be the obvious morphism $(E, F) \rightarrow (E \oplus U, F \oplus U)$. Then f defines a map

$$[S(E) \times_B U_B^+] / S(E) \times_B \infty_U \longrightarrow F_B^+ \times_B U_B^+ / F_B^+ \times_B \infty_U$$

which, composed from the right with the contraction

$$S(E \oplus U) \rightarrow [S(E) \times_B U_B^+] / S(E) \times_B \infty_U$$

and from on left with the contraction

$$F_B^+ \times_B U_B^+ / F_B^+ \times_B \infty_U \rightarrow F_B^+ \times_B U_B^+ / [F_B^+ \times_B \infty_U \cup \infty_F \times_B U_B^+] = (F \oplus U)_B^+$$

gives an S^1 -equivariant map $S(E \oplus U) \rightarrow (F \oplus U)_B^+$ over B . This map represents $\tau_*(\varphi) \in {}_{S^1}\alpha_B^0(S(E \oplus U)_{+B}, (F \oplus U)_B^+)$.

Let $a \in \text{Aut}(E)$ be a unitary gauge transformation of the bundle E . Composing with the induced automorphisms $S(a)$ of the sphere bundles $S(E)_{+B}$ defines a morphism

$${}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+) \xrightarrow{S(a)^*} {}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+) .$$

On the other hand, a defines an element $[a_B^+] \in {}_{S^1}\pi_B^0(E_B^+, E_B^+)$, whose stable class $\{a_B^+\}$ is a unit in the ground ring ${}_{S^1}\alpha^0(B)$ and defines multiplication automorphisms

$${}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+) \xrightarrow{m(a)} {}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+) .$$

Clearly these automorphisms depend only on the homotopy class of a .

Proposition 2.7. *Let $\varphi \in {}_{S^1}\alpha^*(S(E)_{+B}, F_B^+)$ and $a \in \text{Aut}(E)$. Let τ be the obvious morphism $\tau : (E, F) \rightarrow (E \oplus E, F \oplus E)$. In ${}_{S^1}\alpha^*(S(E \oplus E)_{+B}, [F \oplus E]_B^+)$ is holds*

$$\tau_*(S(a)^*(\varphi)) = \tau_*(m(a)(\varphi)) .$$

Proof: For simplicity we prove the statement only in degree 0. We may assume that a is a unitary automorphism with respect to a Hermitian structure on E . Suppose that φ is represented by

$$[f] \in {}_{S^1}\pi_B^0(S(E)_{+B} \wedge_B \xi_B^+, F_B^+ \wedge_B \xi_B^+) .$$

We will prove that the natural representatives

$$p, q \in {}_{S^1}\text{Map}_B(S(E \oplus E)_{+B} \wedge_B \xi_B^+ \wedge_B E_B^+, (F \oplus E)_B^+ \wedge_B \xi_B^+ \wedge_B E_B^+)$$

of $\tau_*(S(a)^*([f]))$, $\tau_*(m(a)([f]))$ are homotopic, so they define the same element in

$${}_{S^1}\pi_B^0(S(E \oplus E)_{+B} \wedge_B \xi_B^+ \wedge_B E_B^+, (F \oplus E)_B^+ \wedge_B \xi_B^+ \wedge_B E_B^+) .$$

We suppose for simplicity that ξ is trivial, to save on notations. Consider the contraction map $c : S(E \oplus E)_{+B} \rightarrow S(E)_{+B} \wedge_B E_B^+$ defined by the first formula in (3), and introduce the maps

$$\Psi, \chi : S(E)_{+B} \wedge_B E_B^+ \wedge_B E_B^+ \longrightarrow F_B^+ \wedge_B E_B^+ \wedge_B E_B^+$$

defined by

$$\Psi := [f \circ S(a)] \wedge_B \text{id}_{E_B^+} \wedge_B \text{id}_{E_B^+} , \chi := f \wedge_B \text{id}_{E_B^+} \wedge_B a_B^+ .$$

Using our definitions it is easy to see that $p = \Psi \circ (c \wedge_B \text{id}_{E_B^+})$, $q = \chi \circ (c \wedge_B \text{id}_{E_B^+})$. Use the same method as in the proof of Proposition 2.1 (conjugation with the rotations of $E \oplus E$ defined by the matrices r_t) to construct a homotopy

$$\chi = f \wedge_B (\text{id}_E \oplus a)_B^+ \simeq f \wedge_B (a \oplus \text{id}_E)_B^+ = f \wedge_B a_B^+ \wedge_B \text{id}_{E_B^+} := \chi' .$$

It suffices to construct a homotopy between $\Psi \circ (c \wedge_B \text{id}_{E_B^+})$, and $\chi' \circ (c \wedge_B \text{id}_{E_B^+})$, and for this it suffices to construct a homotopy between the maps $\Psi_0 \circ c$ and $\chi'_0 \circ c$, where

$$\begin{aligned} \Psi_0 &:= [f \circ S(a)] \wedge_B \text{id}_{E_B^+} = (f \wedge_B \text{id}_{E_B^+}) \circ (S(a) \wedge_B \text{id}_{E_B^+}) , \\ \chi'_0 &:= f \wedge_B a_B^+ = (f \wedge_B \text{id}_{E_B^+}) \circ (\text{id}_{E_B^+} \wedge_B a_B^+) . \end{aligned}$$

Note that $(S(a) \wedge_B \text{id}_{E_B^+}) \circ c = c \circ S(a \oplus \text{id}_E)$, and $(\text{id}_{S(E)} \wedge_B a_B^+) \circ c = c \circ S(\text{id}_E \oplus a)$. In these formulae we use the fact that a is a unitary. On the other hand, using again conjugation with the rotations defined by the matrices r_t , we see that $S(a \oplus \text{id}) \simeq S(\text{id}_E \oplus a)$. Therefore

$$\begin{aligned} \Psi_0 \circ c &= (f \wedge_B \text{id}_{E_B^+}) \circ c \circ S(a \oplus \text{id}) \simeq (f \wedge_B \text{id}_{E_B^+}) \circ c \circ S(\text{id}_E \oplus a) = \\ &= (f \wedge_B \text{id}_{E_B^+}) \circ (\text{id}_{S(E)} \wedge_B a_B^+) \circ c = \chi'_0 , \end{aligned}$$

which completes the proof. \blacksquare

A similar statement holds for the action of an automorphism $b \in \text{Aut}(F)$. Denote by $[b_B^+]_*$ the automorphism of ${}_{S^1}\alpha^*(S(E)_{+B}, F_B^+)$ defined by composition with b_B^+ .

Proposition 2.8. *The automorphisms $[b_B^+]_*$, $m(b)$ coincide on ${}_{S^1}\alpha^*(S(E)_{+B}, F_B^+)$.*

The proof uses similar arguments as the proof of Proposition 2.7 but is substantially easier. \blacksquare

An automorphism $c \in \text{Aut}(U)$ defines a automorphism $\sigma(c)$ of the graded group $\alpha^*(S(E \oplus U)_{+B}, [F \oplus U]_B^+)$ defined by $f \mapsto [\text{id}_F \oplus c]_B^+ \circ f \circ S(\text{id}_E \oplus c)^{-1}$.

Corollary 2.9. *Let $\tau : (E \oplus U, F \oplus U) \rightarrow (E \oplus U \oplus E \oplus U, F \oplus U \oplus E \oplus U)$ be the natural morphism. Then for any $\varphi \in \alpha^*(S(E \oplus U)_{+B}, [F \oplus U]_B^+)$ one has*

$$\tau_*(\sigma(c)(\varphi)) = \tau_*(\varphi) .$$

Proof: Indeed, one has

$$\tau_* \circ \{[\text{id}_F \oplus c]_B^+\}_* = \tau_* \circ m(c) , \quad \tau_* \circ \{S(\text{id}_E \oplus c)^{-1}\}_* = \tau_* \circ (m(c))^{-1} .$$

On the other hand the morphism τ_* is ${}_{S^1}\alpha^0(B)$ -linear. \blacksquare

Consider now the category \mathcal{U}_B of all finite rank complex vector bundles over B . A morphism $\nu : U \rightarrow U'$ in the category \mathcal{U}_B is a pair (i, U_1) consisting of a bundle embedding $i : U \rightarrow U'$ and a complement U_1 of $i(U)$ in U' . This category can be regarded in an obvious way as a category with automorphism push-forward (see section 5.1). The assignment $U \mapsto {}_{S^1}\alpha_B^*(S(E \oplus U)_{+B}, (F \oplus U)_B^+)$ is functorial with respect to morphisms in \mathcal{U}_B , so it defines a functor $\mathfrak{a}_{E,F}^* : \mathcal{U}_B \rightarrow \mathcal{A}b^*$. Since \mathcal{U}_B is not a small category, it is not clear whether this functor has an inductive limit (see sections 2.1, 5.1). We put

$$\hat{\alpha}^*(E, F) := \varinjlim_{n \in \mathbb{N}} {}_{S^1}\alpha_B^*(S(E \oplus \mathbb{C}^n)_{+B}, (F \oplus \mathbb{C}^n)_B^+) . \quad (4)$$

Proposition 2.10. *The functor $\mathfrak{a}_{E,F}^*$ admits an inductive limit which can be identified with $\hat{\alpha}^*(E, F)$.*

Proof: Let \mathcal{N} be the category associated with the ordered set (\mathbb{N}, \leq) and let $\Theta : \mathcal{N} \rightarrow \mathcal{U}_B$ be the cofinal functor $n \mapsto \underline{\mathbb{C}}^n$ (see section 5.1). By Corollary 2.3, the functor $\mathfrak{a}_{E,F}^*$ satisfies the trivial stable action axiom Θ SA. The result follows now from Proposition 5.11 in section 5.1. \blacksquare

In particular one has canonical morphisms $c_U : {}_{S^1}\alpha_B^*(S(E \oplus U)_{+B}, (F \oplus U)_B^+) \rightarrow \hat{\alpha}^*(E, F)$ for any complex bundle U , and the system $(c_U)_U$ is $\mathfrak{a}_{E,F}^*$ -compatible and satisfies the universal property of the inductive limit. Note that $\hat{\alpha}^*(E, F)$ has a natural structure of a graded ${}_{S^1}\alpha^*(B)$ bimodule. By Propositions 2.7 and 2.8 we get:

Remark 2.11. *The action of the gauge groups $\text{Aut}(E \oplus U)$, $\text{Aut}(F \oplus U)$ on $\hat{\alpha}^*(E, F)$ is induced by the canonical ${}_{S^1}\alpha^0(B)^\times$ -action defined by its module structure via the morphisms $\text{Aut}(E \oplus U) \rightarrow {}_{S^1}\alpha^0(B)^\times$, $\text{Aut}(F \oplus U) \rightarrow {}_{S^1}\alpha^0(B)^\times$ defined by $a \mapsto a_B^+$.*

A morphism $\tau = (i, j, E_1, F_1, k) : (E, F) \rightarrow (E', F')$ between two presentations (E, F) , (E', F') of x induces a sequence of morphisms $(E \oplus \underline{\mathbb{C}}^n, F \oplus \underline{\mathbb{C}}^n) \rightarrow (E' \oplus \underline{\mathbb{C}}^n, F' \oplus \underline{\mathbb{C}}^n)$, so it induces a morphism $\hat{\tau}_* : \hat{\alpha}^*(E, F) \xrightarrow{\cong} \hat{\alpha}^*(E', F')$. It is easy to see that $\hat{\tau}_*$ is an isomorphism: it suffices to note that there exists an isomorphism $\theta : (E', F') \rightarrow (E \oplus U, F \oplus U)$ (with $U := E_1$) such that $\theta \circ \tau$ is the standard morphism $(E, F) \rightarrow (E \oplus U, F \oplus U)$, and to apply Proposition 2.10. Therefore we obtain a functor $\hat{\mathfrak{a}}_x^* : \mathcal{T}(x) \rightarrow \mathcal{A}b^*$ whose associated morphisms $\hat{\mathfrak{a}}_x^*(\tau) = \hat{\tau}_*$ are all isomorphisms. According to Proposition 5.8 an inductive limit of this functor exists and can be identified with a quotient of $\hat{\alpha}^*(E, F)$, for any fixed presentation (E, F) of x . Therefore we can make

Definition 2.12. *Define*

$$\alpha^*(x) := \varinjlim_{(E,F) \in x} \hat{\alpha}^*(E, F) .$$

Remark 2.13. *This inductive limit is also an inductive limit of the functor \mathfrak{a}_x^* introduced at the beginning of this section. The existence of the inductive limit of this functor is a non-trivial statement.*

We introduce now the notations

$$\mathbb{A}(E) := \varinjlim_{N \in \mathbb{N}} \text{Aut}(E \oplus \mathbb{C}^N), \quad \mathbb{A}(F) := \varinjlim_{N \in \mathbb{N}} \text{Aut}(F \oplus \mathbb{C}^N) .$$

The two groups $\mathbb{A}(E)$, $\mathbb{A}(F)$ act on the graded group $\hat{\alpha}^*(E, F)$ via the group morphisms $l : \mathbb{A}(E) \rightarrow {}_{S^1}\alpha^0(B)^\times$, $r : \mathbb{A}(F) \rightarrow {}_{S^1}\alpha^0(B)^\times$ (see Remark 2.11), so the two actions commute. Let $\mathbb{Z}[\mathbb{A}(E)]$, $\mathbb{Z}[\mathbb{A}(F)]$ be the group rings of $\mathbb{A}(E)$, $\mathbb{A}(F)$, $I[\mathbb{A}(E)]$, $I[\mathbb{A}(F)]$ the augmentation ideals, and $\lambda : \mathbb{Z}[\mathbb{A}(E)] \rightarrow {}_{S^1}\alpha^0(B)$, $\rho : \mathbb{Z}[\mathbb{A}(F)] \rightarrow {}_{S^1}\alpha^0(B)$ the ring morphisms associated with the group morphisms l, r . Using Proposition 5.8 we get

Remark 2.14. *For every presentation $(E, F) \in x$ there is a canonical isomorphism*

$$\alpha(x) \xrightarrow{\cong} \hat{\alpha}^*(E, F) / \lambda(I[\mathbb{A}(E)])\hat{\alpha}^*(E, F) + \rho(I[\mathbb{A}(F)])\hat{\alpha}^*(E, F) .$$

In the next section we will see that $\mathbb{A}(E)$, $\mathbb{A}(F)$ are both isomorphic to $K^{-1}(B)$ and we will identify the images $\lambda(I[\mathbb{A}(E)])$, $\rho(I[\mathbb{A}(F)])$ of the two ideals in ${}_{S^1}\alpha^0(B)$ with the image of the ideal $I[K^{-1}(B)]$ under the ring morphism $\mathbb{Z}[K^{-1}(B)] \rightarrow {}_{S^1}\alpha^0(B)$ induced by the J -map $K^{-1}(B) \rightarrow {}_{S^1}\alpha^0(B)^\times$.

2.4. The S^1 -equivariant J -map and the description of $\alpha^*(x)$. Let $\pi : E \rightarrow B$ be a Hermitian vector bundle over a compact basis, and let $a, b \in \text{Aut}(E)$ be two unitary automorphisms. We define a map

$$\Delta_E(a, b) : S(E)_{+B} \wedge_B \underline{S}^1 \longrightarrow E_B^+$$

in the following way: We use the models

$$S(E)_{+B} \wedge_B \underline{S}^1 \cong S(E) \times [0, 1] / S(E) \times \{0, 1\}, \quad E_B^+ \cong D(E) /_B S(E)$$

for the two spaces, and define

$$\Delta_E(a, b)([e, t]) := \begin{cases} [(1-2t)a(e)] & \text{for } 0 \leq t \leq \frac{1}{2} \\ [(2t-1)b(e)] & \text{for } \frac{1}{2} \leq t \leq 1. \end{cases}$$

Consider the contraction map

$$\mathbf{c}_E : E_B^+ \longrightarrow S(E)_{+B} \wedge_B \underline{S}^1$$

induced by $e \mapsto [(\frac{1}{\|e\|}e, \|e\|)]$. One has

$$\{\Delta_E(a, b)\} = \{b_B^+\} - \{a_B^+\}. \quad (5)$$

Definition 2.15. *The J -homomorphism associated with a Hermitian bundle E is the morphism $J_E : \pi_0(\text{Aut}(E)) \rightarrow {}_{S^1}\alpha_B^0(B)^\times$ defined by $J_E([a]) := \{a_B^+\}$.*

We introduce the map

$$\Theta_E : \pi_0(\text{Aut}(E)) \longrightarrow {}_{S^1}\alpha_B^{-1}(S(E)_{+B}, E_B^+), \quad \Theta_E([a]) := \{\Delta_E(\text{id}_E, a)\}.$$

Let $\partial_E : {}_{S^1}\alpha_B^{-1}(S(E)_{+B}, E_B^+) \rightarrow {}_{S^1}\alpha_B^0(E_B^+, E_B^+)$ be the connecting morphism in the long exact cohomotopy sequence:

$$\cdots \rightarrow {}_{S^1}\alpha_B^{-1}(S(E)_{+B}, E_B^+) \xrightarrow{\partial_E} {}_{S^1}\alpha_B^0(E_B^+, E_B^+) \rightarrow {}_{S^1}\alpha_B^0(B_{+B}, E_B^+) \rightarrow \cdots \quad (6)$$

associated with E_B^+ and the cofiber sequence

$$S(E)_{+B} \longrightarrow D(E)_{+B} \longrightarrow E_B^+.$$

Since ∂_E acts by composition with the contraction \mathbf{c}_E , we see that the diagram

$$\begin{array}{ccc} \pi_0(\text{Aut}(E)) & \xrightarrow{\Theta_E} & {}_{S^1}\alpha_B^{-1}(S(E)_{+B}, E_B^+) \\ J_E \downarrow & & \downarrow \partial_E \\ {}_{S^1}\alpha^0(B)^\times & \xrightarrow{\cdot - 1} & {}_{S^1}\alpha^0(B) = {}_{S^1}\alpha_B^0(E_B^+, E_B^+) \end{array} \quad (7)$$

is commutative.

Remark 2.16. *Let $\omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1) \subset {}_{S^1}\alpha^0(B)$ be the free summand of the ring ${}_{S^1}\alpha^0(B)$ (see Proposition 2.4). For any $[a] \in \pi_0(\text{Aut}(E))$ it holds*

$$J_E([a]) - 1 \in \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1).$$

Indeed, $\omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1)$ is the kernel of the morphism $\rho : {}_{S^1}\alpha^0(B) \rightarrow \omega^0(B)$ given by restriction to the fixed point set. Therefore

$$\rho(J_E([a])) = \rho(\{a_B^+\}) = \{(a_B^+)^{S^1}\} = \{\text{id}_{B_{+B}}\}.$$

■

Proposition 2.17. *One has*

(1)

$$\varinjlim_N \pi_0(\text{Aut}(E \oplus \underline{\mathbb{C}}^N)) = K^{-1}(B)$$

(2) *The system of morphisms $(\partial_{E \oplus \underline{\mathbb{C}}^N})_{N \in \mathbb{N}}$ defines an isomorphisms*

$$\partial : \varinjlim_N S^1 \alpha_B^{-1}(S(E \oplus \underline{\mathbb{C}}^N)_{+B}, [E \oplus \underline{\mathbb{C}}^N]_B^+) \longrightarrow \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1) .$$

Proof: Let Φ be a complex bundle on B . For any automorphism $a \in \text{Aut}(\Phi)$ we construct a bundle Φ_a over $B \times S^1$ in the following way: we consider the bundle $\Phi \times [0, 1]$ over $B \times [0, 1]$ and we identify $\Phi \times \{0\}$ with $\Phi \times \{1\}$ via a . This bundle comes with an obvious identification $\Phi_a|_{B \times \{0\}} \simeq p_B^*(\Phi)|_{B \times \{0\}}$, so the difference $[\Phi_a] - [p_B^*(\Phi)]$ defines an element $k_\Phi(a) \in K(B \times S^1, B \times \{0\})$. It is easy to see that the obtained map $k_\Phi : \text{Aut}(\Phi) \rightarrow K(B \times S^1, B \times \{0\}) = K^{-1}(B)$ is a group morphism. Taking the limit over N of the system of morphisms $k_{E \oplus \underline{\mathbb{C}}^N}$ we obtain a morphism

$$\kappa_E : \varinjlim_N \pi_0(\text{Aut}(E \oplus \underline{\mathbb{C}}^N)) \rightarrow K^{-1}(B) .$$

Let E' be a complement of E and fix an isomorphism $E' \oplus E \cong \underline{\mathbb{C}}^n$. The assignment $a \mapsto \text{id}_{E'} \oplus a$ defines an *injective* morphism

$$i_{E'} : \varinjlim_N \pi_0(\text{Aut}(E \oplus \underline{\mathbb{C}}^N)) \rightarrow \varinjlim_N \pi_0(\text{Aut}(\underline{\mathbb{C}}^{n+N})) .$$

Similarly, we obtain an obvious *injective* morphism

$$j_E : \varinjlim_N \pi_0(\text{Aut}(\underline{\mathbb{C}}^N)) \rightarrow \varinjlim_N \pi_0(\text{Aut}(E \oplus \underline{\mathbb{C}}^N)) .$$

Hence we have morphisms

$$\varinjlim_N \pi_0(\text{Aut}(\underline{\mathbb{C}}^N)) \xrightarrow{j_E} \varinjlim_N \pi_0(\text{Aut}(E \oplus \underline{\mathbb{C}}^N)) \xrightarrow{i_{E'}} \varinjlim_N \pi_0(\text{Aut}(\underline{\mathbb{C}}^{n+N})) \xrightarrow{\kappa_{\underline{\mathbb{C}}^n}} K^{-1}(B) .$$

The composition $i_{E'} \circ j_E$ is clearly an isomorphism. Moreover, it is well-known that $\kappa_{\underline{\mathbb{C}}^n}$ is an isomorphism, for every $n \in \mathbb{N}$. Since $i_{E'}$ is injective, we see that $\kappa_E = \kappa_{\underline{\mathbb{C}}^n} \circ i_{E'}$ is injective. On the other hand, $\kappa_{\underline{\mathbb{C}}^n} \circ i_{E'} \circ j_E = \kappa_E \circ j_E$ is an isomorphism, so κ_E is also surjective.

For the second isomorphism, we take the direct limit over N in the cohomotopy exact sequence (6) associated with $E \oplus \underline{\mathbb{C}}^N$. We have

$$\varinjlim_N S^1 \alpha_B^k([E \oplus \underline{\mathbb{C}}^N]_B^+, [E \oplus \underline{\mathbb{C}}^N]_B^+) = S^1 \alpha^k(B) .$$

On the other hand, the system of morphisms defined by restriction to the fixed point set (see section 2.2) defines a morphism

$$r_E^k : \varinjlim_N S^1 \alpha_B^k(B_{+B}, [E \oplus \underline{\mathbb{C}}^N]_B^+) \rightarrow \omega^k(B_+, S^0) = \omega^k(B) .$$

Using again a complement E' of E as above, we obtain morphisms

$$\begin{aligned} \varinjlim_{N \in \mathbb{N}} S^1 \alpha^k(B_+, [\mathbb{C}^N]^+) &= \varinjlim_{N \in \mathbb{N}} S^1 \alpha_B^k(B_{+B}, B \times [\mathbb{C}^N]^+) \rightarrow \\ &\rightarrow \varinjlim_N S^1 \alpha_B^k(B_{+B}, [E \oplus \underline{\mathbb{C}}^N]_B^+) \rightarrow \varinjlim_N S^1 \alpha_B^k(B_{+B}, [\underline{\mathbb{C}}^{n+N}]_B^+) \xrightarrow{r_{\underline{\mathbb{C}}^n}^k} \omega^k(B) . \end{aligned}$$

The morphism $\varinjlim_{N \in \mathbb{N}} S^1 \alpha_B^k(B_{+B}, B \times [\mathbb{C}^N]^+) \rightarrow \varinjlim_N S^1 \alpha_B^k(B_{+B}, [\underline{\mathbb{C}}^{n+N}]_B^+)$ is an isomorphism, and $\varinjlim_N S^1 \alpha_B^k(B_{+B}, [E \oplus \underline{\mathbb{C}}^N]_B^+) \rightarrow \varinjlim_N S^1 \alpha_B^k(B_{+B}, [\underline{\mathbb{C}}^{n+N}]_B^+)$ is injective.

Moreover, by Corollary 2.6, the map $r_{\underline{\mathbb{C}}^n}^k$ is an isomorphism. Now the same arguments as above show that r_E^k is an isomorphism. The limit of (6) becomes

$${}_{S^1}\alpha^{-1}(B) \xrightarrow{\rho^{-1}} \omega^{-1}(B) \rightarrow \varinjlim_N {}_{S^1}\alpha_B^{-1}(S(E \oplus \underline{\mathbb{C}}^N)_{+B}, [E \oplus \underline{\mathbb{C}}^N]_B^+) \xrightarrow{\partial} {}_{S^1}\alpha^0(B) \xrightarrow{\rho} \omega^0(B)$$

But the map

$$\rho^{-1} : {}_{S^1}\alpha^{-1}(B) = {}_{S^1}\alpha^0(B_+ \wedge S^1) \rightarrow \omega^0(B_+ \wedge S^1) = \omega^{-1}(B)$$

is also induced by restriction to the fixed point set, so it is surjective by Remark 2.5 applied to the basis $B_+ \wedge S^1$. Therefore ∂ induces an isomorphism

$$\varinjlim_N {}_{S^1}\alpha_B^{-1}(S(E \oplus \underline{\mathbb{C}}^N)_{+B}, [E \oplus \underline{\mathbb{C}}^N]_B^+) \xrightarrow{\cong} \ker(\rho) = \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1).$$

■

Taking the inductive limit with respect to N of the diagram (7) written for $E \oplus \underline{\mathbb{C}}^N$, we obtain the commutative diagram

$$\begin{array}{ccc} K^{-1}(B) & \xrightarrow{\Theta} & \varinjlim_N {}_{S^1}\alpha_B^{-1}(S(E \oplus \underline{\mathbb{C}}^N)_{+B}, [E \oplus \underline{\mathbb{C}}^N]_B^+) = \hat{\alpha}^{-1}(E, E) \\ \downarrow J & & \simeq \downarrow \partial \\ {}_{S^1}\alpha^0(B)^\times & \xrightarrow{\cdot - 1} & \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1) \xrightarrow{\iota} {}_{S^1}\alpha^0(B). \end{array} \quad (8)$$

Remark 2.18. *The map $\iota \circ \partial \circ \Theta : K^{-1}(B) \rightarrow {}_{S^1}\alpha^0(B)$ satisfies the identity*

$$[\iota \circ \partial \circ \Theta](a + b) = [\iota \circ \partial \circ \Theta](a)[\iota \circ \partial \circ \Theta](b) + [\iota \circ \partial \circ \Theta](a) + [\iota \circ \partial \circ \Theta](b).$$

It is the “free J -map” in the terminology of Crabb-Knapp ([CK], p. 88, p.93).

Corollary 2.19. *The map $J : K^{-1}(B) \rightarrow {}_{S^1}\alpha^0(B)^\times$ is injective.*

Proof: It suffices to note that $\partial \circ \Theta$ is injective by Corollary 2.5 in [CK]. ■

The group morphism J extends to a ring morphism $\tilde{J} : \mathbb{Z}[K^{-1}(B)] \rightarrow {}_{S^1}\alpha^0(B)$.

Question: *Does the subgroup*

$$\tilde{J}(I[K^{-1}(B)]) = \langle \{J(u) - 1 \mid u \in K^{-1}(B)\} \rangle = \langle \text{im}(\partial \circ \Theta) \rangle \subset \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1)$$

coincide with the free summand $\omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1)$?

We come back to the description of $\alpha^*(x)$: Using Remarks 2.11 and 2.14 one gets the following descriptions of $\alpha^*(x)$.

Proposition 2.20. *For every presentation $(E, F) \in x$ there exist canonical isomorphisms*

$$\alpha^*(x) \cong \hat{\alpha}^*(E, F) / \tilde{J}(I[K^{-1}(B)]) \hat{\alpha}^*(E, F).$$

Since $\tilde{J}(I[K^{-1}(B)])$ is contained in $\omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1)$, which is an ideal of ${}_{S^1}\alpha^0(B)$, we get epimorphisms

$$\alpha^*(x) \longrightarrow \hat{\alpha}^*(E, F) / \omega^0(B_+, \mathbb{P}_+^\infty \wedge S^1) \cdot \hat{\alpha}^*(E, F).$$

2.5. Stabilization. In this section we will show that the morphism

$$\tau_* : {}_{S^1}\alpha^k(S(E)_{+B}, F_B^+) \rightarrow {}_{S^1}\alpha^k(S(E')_{+B}, [F']_B^+) \quad (9)$$

associated with a morphism $\tau : (E, F) \rightarrow (E', F')$ in the category $\mathcal{T}(x)$ is an isomorphism as soon as the rank f of F is sufficiently large. In other words, for fixed k , the groups $\alpha^k(x)$ can be computed using only presentations (E, F) with a priori bounded ranks.

Proposition 2.21. *Suppose that B is a finite CW complex. The stabilization morphism (9) is an isomorphism for $2f \geq \dim(B) - k$.*

Proof: A morphism τ defines a bundle U and isomorphisms $E' \cong E \oplus U$, $F' \cong F \oplus U$. The long exact sequence associated with the cofiber sequence over B

$$S(U)_{+B} \longrightarrow SE'_{+B} \xrightarrow{c} S(E)_{+B} \wedge_B U_B^+,$$

and the target space $[F']_B^+$ contains the segment

$$\rightarrow {}_{S^1}\alpha_B^{k-1}(S(U)_{+B}, [F']_B^+) \xrightarrow{\partial} {}_{S^1}\alpha_B^k(S(E)_{+B} \wedge_B U_B^+, [F']_B^+) \xrightarrow{c^*} {}_{S^1}\alpha_B^k(SE'_{+B}, [F']_B^+).$$

The morphism τ_* is defined by c^* via the identification ${}_{S^1}\alpha_B^k(S(E)_{+B}, F_B^+) = {}_{S^1}\alpha_B^k(S(E)_{+B} \wedge_B U_B^+, [F']_B^+)$, so it is an isomorphism as soon as

$${}_{S^1}\alpha_B^{k-1}(S(U)_{+B}, [F']_B^+) = {}_{S^1}\alpha_B^k(S(U)_{+B}, [F']_B^+) = 0.$$

Suppose for simplicity $k \geq 0$. A class $u \in {}_{S^1}\alpha_B^k(S(U)_{+B}, [F']_B^+)$ is represented by an S^1 -equivariant pointed map over B

$$\varphi : S(U)_{+B} \wedge_B \xi_B^+ = S(U) \times_B \xi^+ /_B S(U) \times_B \infty_\xi \longrightarrow [F' \oplus \mathbb{R}^k \oplus \xi]_B^+,$$

where $\xi = \eta \oplus \xi_0$ is the sum of a complex and a real vector bundle. We may suppose that ξ_0 is an oriented bundle, so that all our bundles become oriented bundles. We will prove that any such map is homotopic to the map φ_∞ which maps the left hand space fiberwise onto the section $\infty_{F' \oplus \mathbb{R}^k \oplus \xi}$. Denote by $q : \mathbb{P}(U) \rightarrow B$ the bundle projection and put

$$\tilde{F}' := q^*(F')(1), \quad \tilde{\xi} := q^*(\eta)(1) \oplus q^*(\xi_0).$$

A map φ as above induces a pointed bundle map $\tilde{\varphi} : \tilde{\xi}_{\mathbb{P}(U)}^+ \longrightarrow [\tilde{F}' \oplus \mathbb{R}^k \oplus \tilde{\xi}]_{\mathbb{P}(U)}^+$ over $\mathbb{P}(U)$, and the assignment $\varphi \mapsto \tilde{\varphi}$ is a bijection. But by Corollary 5.15 in section 5.2, any such pointed bundle map is homotopic to the fiberwise constant bundle map as soon as $\dim_{\mathbb{R}}(\mathbb{P}(U)) + \text{rk}(\tilde{\xi}) < \text{rk}_{\mathbb{R}}(\tilde{F}') + k + \text{rk}(\tilde{\xi})$. This condition is equivalent to $2f > \dim(B) - k - 2$. Similarly, we will have ${}_{S^1}\alpha_B^{k-1}(S(U)_{+B}, [F']_B^+) = 0$ as soon as $2f > \dim(B) - k - 1$. \blacksquare

2.6. The cohomotopy Euler class of an element in $K(B)$. Let $x \in K(B)$ and consider a presentation $(E, F) \in x$. The map $o_{(E,F)} : S(E)_{+B} \rightarrow F_B^+$ which sends the section $+_B$ of $S(E)_{+B}$ to the infinity section of F_B^+ and maps any point $e_b \in S(E_b)$ to 0_b is an S^1 -equivariant map of pointed spaces over B , hence it defines an element $\{o_{(E,F)}\} \in {}_{S^1}\alpha_B^0(S(E)_{+B}, F_B^+)$.

One has a canonical isomorphism (see [CJ] Proposition 12.40)

$${}_{S^1}\alpha_B^0(S(E)_{+B}, F_B^+) \cong {}_{S^1}\alpha_{S(E)}^0(S(E)_{+S(E)}, \pi^*(F)_{S(E)}^+),$$

where $\pi : S(E) \rightarrow B$ is the obvious projection. Under this isomorphism the class $\{o_{(E,F)}\}$ maps to the equivariant Euler class of the bundle $\pi^*(F)$ over $S(E)$. This

class is the pull-back of the equivariant Euler class $\gamma(F) \in S^1 \alpha^0(B_{+B}, F_B^+)$ of the bundle F under the projection $S(E)_{+S(E)} \rightarrow B_{+B}$.

For any morphism $\tau = (i, j, E_1, F_1, k) : (E, F) \rightarrow (E', F')$ in the category $\mathcal{T}(x)$ one has $\tau_*(\{o_{(E,F)}\}) = \{o_{(E',F')}\}$. Therefore the assignment $(E, F) \mapsto -\{o_{(E,F)}\}$ defines a *tautological element* $\gamma(x) \in \alpha^*(x)$. This element will be called the equivariant cohomotopy Euler class of x .

3. COHOMOTOPY INVARIANTS ASSOCIATED WITH CERTAIN NON-LINEAR MAPS BETWEEN HILBERT BUNDLES

3.1. The cylinder construction. Let (E, F) be a pair of Hermitian vector bundles over a compact basis B . Let V, W be Euclidean vector spaces, and let $\mu : E \times V \rightarrow [F \times W]_B^+$ be an S^1 -equivariant map over B . We suppose that μ is fiberwise differentiable and its fiberwise differential is continuous on $E \times V$. The equivariance property implies that

$$\mu(0^E \times V) \subset [0^F \times W]_B^+. \quad (10)$$

We assume that μ has the following properties:

P1: (properness) There exist positive constants c, C such that $\|\mu(e, v)\| > c$ for all pairs $(e, v) \in E \times V$ with $\|(e, v)\| \geq C$.

P2: (restriction to the S^1 -fixed point set)

(1) There exists a direct sum decomposition $W = H \oplus W_0$ such that

$$\mu(0_y^E, v) = h(y) + l(v), \quad \forall y \in B, \quad \forall v \in V,$$

where $l : V \xrightarrow{\cong} W_0 \subset W$ is a linear isomorphism, which does not depend on y , and $h : B \rightarrow H$ is a continuous map.

(2) There exists $\varepsilon_0 > 0$ such that

$$\|h(y)\| = \|\text{p}_H(\mu(0_y^E, v))\| \geq \varepsilon_0 \quad \forall (y, v) \in B \times V. \quad (11)$$

We fix an orientation \mathcal{O} of H , and set $b := \dim(H)$. Choose numbers $R \geq C$ and $\varepsilon \leq \min(c, \varepsilon_0)$. The restriction μ_R of μ to $D_R(E) \times D_R(V)$ satisfies

$$\|\mu(e, v)\| \geq \varepsilon \quad \forall (e, v) \in \partial[D_R(E) \times D_R(V)] \cup [0^E \times D_R(V)].$$

Therefore, μ_R defines an S^1 -equivariant morphism of pairs over B

$$\begin{aligned} \mu_{R,\varepsilon} : (D_R(E) \times D_R(V), \partial[D_R(E) \times D_R(V)] \cup [0^E \times D_R(V)]) &\longrightarrow \\ &\longrightarrow \left([F \times W]_B^+, [F \times W]_B^+ \setminus \overset{\circ}{D}_\varepsilon(F \times W) \right). \end{aligned}$$

The first space $D_R(E) \times D_R(V)$ of the pair on which $\mu_{R,\varepsilon}$ is defined can be regarded as a ‘‘cylinder bundle’’ over B , whose base is the complex disk bundle $D(E)$; the second space of this pair is the union of the boundary of this cylinder bundle with the core $0^E \times D_R(V)$. Using polar coordinates in $D_R(E)$ we obtain a map $S(E) \times [0, R] \rightarrow D_R(E)$, hence a map

$$\rho : S(E) \times [0, R] \times D_R(V) = S(E) \times D_R(\mathbb{R} \oplus V) \rightarrow D_R(E) \times D_R(V),$$

which maps

$$[S(E) \times \{0, R\} \times D_R(V)] \cup [S(E) \times [0, R] \times S_R(V)] = S(E) \times S_R(\mathbb{R} \oplus V)$$

onto the the second component of the pair on which $\mu_{R,\varepsilon}$ is defined. Here we used suitable models $D(\mathbb{R} \oplus V)$, $S(\mathbb{R} \oplus V)$ for the disc and the sphere in $\mathbb{R} \oplus V$. Therefore, composing $\mu_{R,\varepsilon}$ with ρ we get an S^1 -equivariant map of pairs over B

$$(S(E) \times [0, R] \times D_R(V), S(E) \times (\{0, R\} \times D_R(V) \cup [0, R] \times S_R(V))) = \\ (S(E) \times D_R(\mathbb{R} \oplus V), S(E) \times S_R(\mathbb{R} \oplus V)) \rightarrow ([F \times W]_B^+, [F \times W]_B^+ \setminus \mathring{D}_\varepsilon(F \times W))$$

which we denote by the same symbol $\mu_{R,\varepsilon}$. Collapsing fiberwise over B the second terms of the two pairs, and composing with the natural isomorphism

$$[F \times W]_B^+ /_B [F \times W]_B^+ \setminus \mathring{D}_\varepsilon(F \times W) \simeq [F \times W]_B^+,$$

one gets an S^1 -equivariant map of pointed spaces over B

$$\mu_{R,\varepsilon} : S(E) \times [\mathbb{R} \oplus V]_B^+ /_B S(E) \times \{\infty\} = S(E)_{+B} \wedge_B [B \times (\mathbb{R} \oplus V)]_B^+ \longrightarrow [F \times W]_B^+.$$

Using the isomorphism $l : V \xrightarrow{\cong} W_0$ and an orientation preserving isomorphism $\mathbb{R}^b \simeq H$, we obtain an element

$$\{\mu\} \in {}_{S^1} \alpha_B^{b-1}(S(E)_{+B}, F_B^+),$$

which is obviously independent of the choice of the pair (R, ε) . This element will be called the *cohomotopy invariant* of μ .

3.2. General properties of the invariant $\{\mu\}$.

3.2.1. *A vanishing property.* Let $\mu : E \times V \rightarrow F \times W$ a map satisfying **P1**, **P2**.

Proposition 3.1. *If $\mu|_{D_C(E) \times D_C(V)}$ is nowhere vanishing, then $\{\mu\} = 0$.*

Proof: We take $\varepsilon \leq \inf\{\|\mu(e, v)\| \mid \|e\| \leq C, \|v\| \leq C\}$, and we note that the $\{[F \times W]_B^+\} /_B \{[F \times W]_B^+ \setminus \mathring{D}_\varepsilon(F \times W)\}$ -valued pointed map induced by $\mu_{R,\varepsilon}$ is fiberwise constant. ■

3.2.2. *Homotopy invariance.* Let $\mu', \mu'' : E \times V \rightarrow [F \times W]_B^+$ two maps satisfying properties **P1**, **P2** with constants C', c', ε'_0 , and $C'', c'', \varepsilon''_0$. We suppose that the property **P2** of the two maps holds for the same decomposition $W = H \oplus W_0$ of W and for the same isomorphism $l : V \rightarrow W_0$. We introduce the notations

$$\tilde{B} := B \times [0, 1], \quad \tilde{E} := E \times [0, 1] = p_B^*(E), \quad \tilde{F} := F \times [0, 1] = p_B^*(F).$$

Proposition 3.2. *Suppose there exists $C \geq \max(C', C'')$ and a continuous S^1 -equivariant map $\tilde{\mu} : D_C(\tilde{E}) \times D_C(V) \rightarrow [\tilde{F} \times W]_{\tilde{B}}^+$ over \tilde{B} whose restriction to*

$$\partial [D_C(\tilde{E}) \times D_C(V)] \cup [0^{\tilde{E}} \times D_R(V)]$$

is nowhere vanishing. Then $\{\mu'\} = \{\mu''\}$ in ${}_{S^1} \alpha_B^{b-1}(S(E)_{+B}, F_B^+)$.

Proof: The stable classes $\{\mu'\}$, $\{\mu''\}$ can be computed using the the cylinder $D_C(\tilde{E}) \times D_C(V)$ and taking

$$\varepsilon \leq \min \left(\varepsilon'_0, \varepsilon''_0, c', c'', \inf \left\{ \|\tilde{\mu}(y)\| \mid y \in \partial [D_C(\tilde{E}) \times D_C(V)] \cup [0^{\tilde{E}} \times D_R(V)] \right\} \right)$$

Applying the cylinder construction with parameters C, ε to the map $\tilde{\mu}$ we obtain a homotopy between the corresponding representatives of the classes $\{\mu'\}$, $\{\mu''\}$. ■

3.2.3. *A product formula.* Let V_i, W_i be Euclidean spaces, E_i, F_i Hermitian bundles over a compact base B ($i = 1, 2$) and $\mu_i : E_i \times V_i \rightarrow [F_i \times W_i]_B^+$ S^1 -equivariant maps over B satisfying the properties **P1**, **P2** (1) of section 3.1 with constants C, c . Let $W_i = H_i \oplus W_{0,i}$ be the corresponding direct sum decompositions, and $l_i : V_i \xrightarrow{\cong} W_{0,i}$, $h_i : B \rightarrow H_i$ the maps given by **P2** (1). Fix orientations on the H_i , and put

$V := V_1 \oplus V_2$, $W := W_1 \oplus W_2$, $H := H_1 \oplus H_2$, $W_0 := W_{0,1} \oplus W_{0,2}$, $l := l_1 \oplus l_2$, and consider the bundles $E := E_1 \oplus E_2$, $F := F_1 \oplus F_2$. We have a product map

$$\mu : E \times V = [E_1 \times V_1] \oplus [E_2 \times V_2] \longrightarrow [F \times W]_B^+ = [F_1 \times W_1]_B^+ \wedge_B [F_2 \times W_2]_B^+$$

over B . This map satisfies properties **P1**, **P2** (1) with the map

$$h = (h_1, h_2) : B \rightarrow H .$$

Note that μ will also satisfy **P2** (2) as soon as one of the two maps μ_1, μ_2 has this property. Suppose that μ_1 also satisfies property **P2** (2) with constant ε_0 and denote by

$$\{\mu_1\} \in {}_{S^1}\alpha_B^{b_1-1}(S(E_1)_{+B}, [F_1]_B^+)$$

the corresponding stable class. The map μ_2 defines a map $[E_2 \oplus V_2]_B^+ \longrightarrow [F_2 \oplus W_2]_B^+$ hence a class $\{\mu_2^+\} \in {}_{S^1}\alpha_B^{b_2}([E_2]_B^+, [F_2]_B^+)$. One can then form the product

$$\{\mu_1\} \wedge_B \{\mu_2^+\} \in {}_{S^1}\alpha_B^{b-1}(S(E_1)_{+B} \wedge_B [E_2]_B^+, F_B^+) .$$

Consider now the contraction map ${}_1c : S(E_1 \oplus E_2)_{+B} \rightarrow S(E_1)_{+B} \wedge_B [E_2]_B^+$ introduced in section 2.3 (see formula (3)). Using the identifications

$$[E_2]_B^+ = D_R(E_2) /_B S_R(E_2) = E_2 /_B E_2 \setminus \dot{D}_R(E_2) ,$$

we can use as model for the contraction ${}_1c$ any map of the form ${}_1c_R^{\mathfrak{R}}$ given by

$${}_1c_R^{\mathfrak{R}}(e_1, e_2) := \left[\frac{1}{\|e_1\|} e_1, \mathfrak{R}e_2 \right] , \quad (\mathfrak{R} \geq R) .$$

Proposition 3.3. *Under the above assumptions it holds $\{\mu\} = {}_1c^* (\{\mu_1\} \wedge_B \{\mu_2^+\})$.*

Proof: The class $\{\mu\}$ is represented by the map of pairs

$$\begin{aligned} \mu_R : (S(E) \times [0, R] \times D_R(V), S(E) \times ([0, R] \times S_R(V) \cup \{0, R\} \times D_R(V))) \longrightarrow \\ \longrightarrow ([F \times W]_B^+, [F \times W]_B^+ \setminus D_\varepsilon(F \times W)) \end{aligned}$$

which is defined by

$$\mu_R(e_1, e_2, \rho, v_1, v_2) = [\mu_1(\rho e_1, v_1), \mu_2(\rho e_2, v_2)] .$$

The class ${}_1c^* (\{\mu_1\} \wedge_B \{\mu_2^+\})$ is represented by the map $\nu_R^{\mathfrak{R}}$ between the same pairs defined by

$$\nu_R^{\mathfrak{R}}(e_1, e_2, \rho, v_1, v_2) = \left[\mu_1\left(\rho \frac{1}{\|e_1\|} e_1, v_1\right), \mu_2(\mathfrak{R}e_2, v_2) \right] .$$

Composing $\mu_R, \nu_R^{\mathfrak{R}}$ with the projection

$$p : [F \times W]_B^+ \longrightarrow [F \times W]_B^+ /_B [F \times W]_B^+ \setminus D_\varepsilon(F \times W)$$

we obtain two maps

$$m_0, m_1 : S(E) \times [0, R] \times D_R(V) \longrightarrow [F \times W]_B^+ / [F \times W]_B^+ \setminus D_\varepsilon(F \times W) \simeq [F \times W]_B^+$$

which map $S(E) \times ([0, R] \times S_R(V) \cup \{0, R\} \times D_R(V))$ onto the infinity section in the right hand bundle. The natural homotopy between these maps is the map

$$m : [0, 1] \times S(E) \times [0, R] \times D_R(V) \longrightarrow [F \times W]_B^+ / [F \times W]_B^+ \setminus D_\varepsilon(F \times W)$$

given by

$$m(t, e_1, e_2, \rho, v_1, v_2) = \left[\mu_1 \left(\rho \left[1 - t + t \frac{1}{\|e_1\|} \right] e_1, v_1 \right), \mu_2 \left([(1-t)\rho + t\mathfrak{R}]e_2, v_2 \right) \right]$$

Claim: For any $R \geq \sqrt{2}C$ and sufficiently large $\mathfrak{R} \geq R$ it holds

- (1) the map m is well defined and continuous at the points $(t, e_1, e_2, \rho, v_1, v_2)$ with $e_1 = 0$.
- (2) the map m maps $[0, 1] \times S(E) \times ([0, R] \times S_R(V) \cup \{0, R\} \times D_R(V))$ to the infinity section in the right hand bundle.

In fact we show that for $e_2 \in [E_2]_y$, one has

$$\lim_{u \rightarrow (t, 0_{b_1}^{E_1}, e_2, \rho, v_1, v_2)} m(u) = \infty_y,$$

so m maps the locus $e_1 = 0$ to the infinity section. Let $\eta_R > 0$ be sufficiently small, such that $\|\mu_1(e_1, v_1)\| > \varepsilon_0$ for every $(e_1, v_1) \in D_{\eta_R}(E_1) \times D_R(V_1)$. One has

$$\lim_{e_1 \rightarrow 0} \left\| \rho \left[1 - t + t \frac{1}{\|e_1\|} \right] e_1 \right\| = \rho t.$$

When $\rho t < \eta_R$, the first component of $m(t, e_1, e_2, \rho, v_1, v_2)$ will already have a norm larger than ε_0 . When $\rho t \geq \eta_R$, we obtain (using $\|e_1\|^2 + \|e_2\|^2 = 1$):

$$\lim_{e_1 \rightarrow 0} \|[(1-t)\rho + t\mathfrak{R}] e_2\| = (1-t)\rho + t\mathfrak{R} \geq \eta_R \left(\frac{1}{t} - 1 \right) + t\mathfrak{R} \geq 2\sqrt{\eta_R \mathfrak{R}} - \eta_R,$$

which will be larger than R when \mathfrak{R} is sufficiently large. The second part of the claim is obvious for the spaces $[0, 1] \times S(E) \times [0, R] \times S_R(V)$, $[0, 1] \times S(E) \times \{0\} \times D_R(V)$. For $\rho = R$ we obtain

$$\left\| \rho \left[1 - t + t \frac{1}{\|e_1\|} \right] e_1 \right\|^2 + \|[(1-t)\rho + t\mathfrak{R}] e_2\|^2 \geq R^2 (\|e_1\|^2 + \|e_2\|^2) = R^2 \geq 2C^2,$$

so at least one of the two norms is $\geq C$.

Using the claim, it follows that m descend to an homotopy between two representatives of the classes $\{\mu\}$ and ${}_1c^* (\{\mu_1\} \wedge_B \{\mu_2^+\})$. \blacksquare

An interesting case is the one when also μ_2 satisfies property **P2** (2). In this case the cylinder construction applies to μ_2 and one can write

$$\{\mu_2^+\} = \partial_2(\{\mu_2\}),$$

where $\{\mu_2\} \in {}_{S^1}\alpha_B^{b_2-1}(S(E_2)_{+B}, [F_2]_B^+)$ is the invariant associated with μ_2 , and ∂_2 is the connecting morphism in the long exact sequence associated with the cofiber sequence

$$S(E_2)_{+B} \longrightarrow D(E_2)_{+B} \longrightarrow [E_2]_B^+.$$

Let ${}_2c : S(E_1 \oplus E_2)_{+B} \rightarrow [E_1]_B^+ \wedge_B S(E_2)_{+B}$ be the standard contraction. In this case, our multiplication formula becomes

Corollary 3.4. *Suppose that both maps μ_1, μ_2 satisfy properties **P1**, **P2**. Then*

$$\{\mu\} = {}_1c^* (\{\mu_1\} \wedge_B \partial_2(\{\mu_2\})) = {}_2c^* (\partial_1(\{\mu_1\}) \wedge_B \{\mu_2\}) .$$

Another corollary is obtained when μ_2 is defined by a pair of linear isomorphisms $E_2 \rightarrow F_2, V_2 \rightarrow W_2$. The corresponding formula will play an important role in the proof of the coherence Lemma 3.10 comparing the invariants associated to two finite dimensional approximations of an admissible bundle map between Hilbert bundles.

Proposition 3.5. *Let $\mu : E \times V \rightarrow F \times W$ be a map satisfying the properties **P1**, **P2** with constants C, c, ε_0 and maps $l : V \rightarrow W_0, h : B \rightarrow H$. Let $a : E' \rightarrow F'$ be an isomorphism of complex vector bundles over B , and let $b : V' \rightarrow W'$ be an isomorphism of real vector spaces. Put $\tilde{E} := E \oplus E', \tilde{F} := F \oplus F', \tilde{V} := V \oplus V', \tilde{W} := W \oplus W'$, and define*

$$\tilde{\mu}(e, e', v, v') = \iota[\mu(e, v) \wedge_B (a(e'), b(v'))] ,$$

where ι is the obvious identification

$$\iota : [F \times W]_B^+ \wedge_B (F' \times W')_B^+ \rightarrow [(F \oplus F') \times (W \oplus W')]_B^+ .$$

Then

- (1) $\tilde{\mu}$ satisfies **P1** with constants C, γ (for sufficiently small $0 < \gamma < c$), and **P2** with constant ε_0 and maps $\tilde{l} := l \oplus b, \tilde{h} := h$.
- (2) $\{\tilde{\mu}\} = \tau_*(\{\mu\})$, where τ denotes the obvious morphism $(E, F) \rightarrow (\tilde{E}, \tilde{F})$.

The second statement follows directly from Proposition 3.3. The first statement (which is specific to case when the second factor is a linear isomorphism) is proved as follows: Since the closed set $\mu^{-1}(D_c(F \times W))$ is contained in the open disk $\mathring{D}_C(E \times V)$, there exists $r > 0$ such that $\|\mu(e, v)\| > c$ as soon as $\|(e, v)\| \geq C - r$. For a point (e, e', v, v') with $\|(e, e', v, v')\| \geq C$ one has either $\|(e, v)\| \geq C - r$, or $\|(e', v')\| \geq r$. In the first case one obtains $\|\mu(e, v)\| > c$, whereas in the second we get $\|(a(e'), b(v'))\| \geq c'r$ for a constant c' . ■

3.3. A class of non-linear maps between Hilbert bundles. Suppose now that \mathcal{V}, \mathcal{W} are real Hilbert spaces, and that \mathcal{E}, \mathcal{F} are complex Hilbert bundles over the compact basis B , and let $\mu : \mathcal{E} \times \mathcal{V} \rightarrow \mathcal{F} \times \mathcal{W}$ be a continuous S^1 -equivariant map over B which is fiberwise \mathcal{C}^∞ , and whose fiberwise derivatives are continuous on $\mathcal{E} \times \mathcal{V}$. We assume that the fiberwise differentials

$$d_y := d_{0_y} \mu_y = \mathcal{E}_y \times \mathcal{V} \longrightarrow \mathcal{F}_y \times \mathcal{W} , \quad y \in B$$

at the origins of the fibers $\mathcal{E}_y \times \mathcal{V}$ are Fredholm. The linear operator d_y has the form $d_y = (\delta_y, l_y)$, where $\delta_y : \mathcal{E}_y \rightarrow \mathcal{F}_y$ and $l_y : \mathcal{V} \rightarrow \mathcal{W}$ are defined by the derivatives of the restrictions $\mu|_{\mathcal{E}_y \times \{0^\mathcal{V}\}}, \mu|_{\{0_y^\mathcal{E}\} \times \mathcal{V}}$. Note that the continuous family $\delta := (\delta_y)_{y \in B}$ of complex Fredholm operators defines an element $\text{ind}(\delta) \in K(B)$. Let $d : \mathcal{E} \times \mathcal{V} \rightarrow \mathcal{F} \times \mathcal{W}$ the fiberwise linear map defined by the family of Fredholm operators $(d_y)_{y \in B}$. We suppose that μ also has the properties

P1: (properness) There exist positive constants c, C such that $\|\mu(e, v)\| > c$ for all pairs $(e, v) \in \mathcal{E} \times \mathcal{V}$ with $\|(e, v)\| \geq C$.

P2: (behavior near the S^1 -fixed point set)

- (1) \mathcal{W} splits orthogonally as $\mathcal{W} = H \oplus \mathcal{W}_0$, where H is a finite dimensional subspace, and for every $y \in B$ one has

$$\mu(0_y^E, v) = h(y) + l(v) \quad \forall y \in B, \quad \forall v \in \mathcal{V},$$

where $l : \mathcal{V} \xrightarrow{\cong} \mathcal{W}_0 \subset \mathcal{W}$ is a linear isometry.

In particular the operator l_y coincides with l , so is independent of y .

- (2) There exists $\varepsilon_0 > 0$ such that for every $y \in B$ one has

$$\|h(y)\| = \|\mathbb{D}_H(\mu(0_y^E, v))\| \geq \varepsilon_0.$$

P3: (linear+compactness) The difference $k := \mu - d$ is globally compact, in the sense that for every $R > 0$ the image $k(D_R(\mathcal{E} \times \mathcal{V}))$ of the disk bundle $D_R(\mathcal{E} \times \mathcal{V})$ is relatively compact in the total space $\mathcal{F} \times \mathcal{W}$.

Note that one has the identity

$$k(0_y^E, v) = h(y) \in H, \quad \forall y \in B. \quad (12)$$

In the next section we will see that the left hand of the Seiberg-Witten equations on a 4-manifold M defines a map satisfying properties $\mathcal{P}_1 - \mathcal{P}_3$. A different construction of such a map can be found in [BF].

3.4. The Seiberg-Witten map in dimension 4. Let M be closed oriented 4-manifold, and let L be a Hermitian line bundle on M . We fix the following data:

- (1) A closed complement \mathcal{S} of the closed subspace $iB_{\text{DR}}^1(M) = d(iA^0(M))$ of $iA^1(M)$.
- (2) A closed complement \mathcal{V} of the finite dimensional space

$$i\mathbb{H}^1 := \mathcal{S} \cap \ker(d : iA^1(M) \rightarrow iA^2(M)) \simeq iH^1(M, \mathbb{R})$$

in \mathcal{S}

- (3) A complement $i\mathbb{H}^2$ of $d(iA^1(M))$ in $\ker(d : iA^2(M) \rightarrow iA^3(M))$. This complement will come with an isomorphism $i\mathbb{H}^2 \simeq iH^2(M, \mathbb{R})$.
- (4) An affine subspace \mathcal{A} of the space of connections $\mathcal{A}(L)$ modeled after \mathcal{S} .

Therefore, \mathcal{A} is a slice to the orbits of the right action of the gauge group \mathcal{G} on the space of connections:

$$a \cdot g := a + 2g^{-1}dg$$

The quotient $\bar{\mathcal{A}} := \mathcal{A}/\mathcal{V}$ is an affine space modeled after $iH^1(M, \mathbb{R})$. Consider the finite dimensional Lie group

$$G := \{u \in \mathcal{C}^\infty(M, S^1) \mid u^{-1}du \in \mathcal{S}\}.$$

One has an obvious short exact sequence

$$\{1\} \longrightarrow S^1 \longrightarrow G \xrightarrow{\lambda} 2\pi iH^1(M; \mathbb{Z}) \longrightarrow \{1\},$$

where λ is defined by $u \mapsto [u^{-1}du]_{\text{DR}}$. The choice of a point $x_0 \in M$ defines a left splitting $\text{ev}_{x_0} : G \rightarrow S^1$ whose kernel is isomorphic to $2\pi iH^1(M; \mathbb{Z})$ and which will be denoted by G_{x_0} . In the affine space \mathcal{A} we have a natural $i\mathbb{H}^1$ -invariant (hence G_{x_0} -invariant) subset \mathcal{A}_0 defined by

$$\mathcal{A}_0 := \{a \in \mathcal{A} \mid F_a \in i\mathbb{H}^2\}.$$

The curvature F_{a_0} of a connection $a_0 \in \mathcal{A}_0$ is independent of a_0 , because it coincides with the representative in $i\mathbb{H}^2$ of the de Rham class $-2\pi i c_1^{\text{DR}}(L)$; this 2-form will be denoted by F_0 . Note that \mathcal{A}_0 is a G_{x_0} -invariant complete system of representatives

for the quotient $\bar{\mathcal{A}} = \mathcal{A}/\mathcal{V}$. The space \mathcal{A}/G_{x_0} can be regarded as an affine bundle over the torus

$$\text{Pic}(L) := \bar{\mathcal{A}}/G_{x_0} ,$$

which is naturally a $iH^1(X; \mathbb{R})/4\pi iH^1(X; \mathbb{Z})$ -torsor. The fibers of the affine bundle

$$\pi : \mathcal{A}/G_{x_0} \longrightarrow \text{Pic}(L)$$

are affine \mathcal{V} -spaces. Since the quotient \mathcal{A}_0/G_{x_0} is a section of this affine bundle, we can regard it as a \mathcal{V} -vector bundle over $\text{Pic}(L)$ with \mathcal{A}_0/G_{x_0} as zero section. This vector bundle is actually trivial: indeed, the map $(a_0, v) \mapsto a_0 + v \in \mathcal{A}$ is G_{x_0} -equivariant, and it descends to a trivialization $\text{Pic}(L) \times \mathcal{V} \rightarrow \mathcal{A}/G_{x_0}$.

Remark 3.6. *Choosing a Riemannian metric g on M gives canonical choices for the three objects $S, T, i\mathbb{H}^2$ above, namely*

$$\mathcal{S} = \ker(d^* : iA^1(M) \longrightarrow iA^0(M)) , \quad \mathcal{V} := d^*(iA^2(M)) , \quad i\mathbb{H}^2 = i\mathbb{H}_g^2 ,$$

where the subscript g on the right denotes the respective g -harmonic space. With these choices, \mathcal{A}_0 is just the set of g -Yang-Mills connections in the slice \mathcal{A} .

Let g be a Riemannian metric on M , $\mathfrak{c} \in \text{Spin}^c(M)$ an equivalence class of Spin^c -structures, and let $\tau : Q \rightarrow P_g$ be a Spin^c -structure on M representing the class \mathfrak{c} . Denote by Σ^\pm , $\Sigma := \Sigma^+ \oplus \Sigma^-$ the spinor bundles of τ , $L = \det(\Sigma^\pm)$ the determinant line bundle, and $\gamma : \Lambda^1 \rightarrow \text{End}_0(\Sigma)$ the Clifford map [OT]. Note that the gauge group $\text{Aut}(Q)$ of Q acts on the space of Spin^c -structures $\tau : Q \rightarrow P_g$ representing \mathfrak{c} (or, equivalently, on the space of Clifford maps $\gamma : \Lambda^1 \rightarrow \text{End}_0(\Sigma)$ which are compatible with \mathfrak{c}). Therefore, the space of Spin^c -Dirac operators which are compatible with the pair (g, \mathfrak{c}) has a very complicated topology. Note that, for the construction of a Dirac operator one needs a concrete Spin^c -structure τ (or, equivalently, a concrete Clifford map γ), not only an equivalence class \mathfrak{c} .

The gauge group \mathcal{G} and its subgroup G_{x_0} act from the left on the vector spaces of sections $A^0(\Sigma^\pm)$ by the formula

$$(g, \Psi) \mapsto g^{-1}\Psi .$$

Since G_{x_0} acts freely on the affine quotient space $\bar{\mathcal{A}}$ we get two flat vector bundles $\bar{\mathcal{A}} \times_{G_{x_0}} A^0(\Sigma^\pm)$ over $\text{Pic}(L)$ with standard fibers $A^0(\Sigma^\pm)$. In order to use our general formalism we make the following definitions:

$$B := \text{Pic}(L) , \quad \mathcal{E} := \bar{\mathcal{A}} \times_{G_{x_0}} A^0(\Sigma^+) , \quad \mathcal{F} := \bar{\mathcal{A}} \times_{G_{x_0}} A^0(\Sigma^-) , \quad \mathcal{W} := iA_+^2(M) .$$

Let $\kappa : B \rightarrow i\mathbb{H}_g^+$ be a smooth map. The κ -twisted Seiberg-Witten map is the map from $A^0(\Sigma^+) \times \mathcal{A}$ to $A^0(\Sigma^-) \times iA_+^2$ given by

$$(\Psi, a) \mapsto (\not{D}_a \Psi, (F_a - F_0 + \kappa(\pi(a)))^+ - \gamma^{-1}((\Psi \bar{\Psi})_0)) .$$

Via the identification $B \times \mathcal{V} = \mathcal{A}/G_{x_0}$ this map descends to an S^1 -equivariant map

$$sw_\kappa : \mathcal{E} \times \mathcal{V} \longrightarrow \mathcal{F} \times \mathcal{W} .$$

The restriction of sw_κ to the fiber over $y = [a_0] \in B$ is given by the formula

$$sw_\kappa(\Psi, v) = \left(\not{D}_{a_0} \Psi + \frac{1}{2} \gamma(v) \Psi , d^+ v + \kappa(y) - \gamma^{-1}((\Psi \bar{\Psi})_0) \right) .$$

The linearization of this map at the zero section in the bundle $\mathcal{E} \times \mathcal{V}$ over B is a fiberwise linear bundle map given by

$$d(\Psi, v) = (\not{D}_{a_0} \Psi, d^+ v) .$$

Hence sw_β decomposes as

$$sw_\kappa = d + c_\kappa ,$$

where c_κ is the sum of a quadratic map c and the fiberwise constant map defined by κ . Denote by w_τ the expected dimension of the Seiberg-Witten moduli space corresponding to τ :

$$w_\tau := \frac{1}{4}(c_1(L)^2 - 3\sigma(M) - 2e(M))$$

We define Sobolev L_k^2 -completions of the spaces \mathcal{V} , \mathcal{W} in the usual way. The construction of Sobolev norms on the bundles \mathcal{E} , \mathcal{F} is more delicate, because these bundles are quotients with respect to group \mathcal{G}_0 , which does not operate by L_k^2 -isometries⁴. For a point $y = [a_0] \in B$ (with $a_0 \in \mathcal{A}_0$) one identifies the fiber \mathcal{E}_y , \mathcal{F}_y with $\{a_0\} \times A^0(\Sigma^\pm)$ and uses the covariant derivatives associated with ∇_{a_0} to define the L_k^2 -norm on \mathcal{E}_y . A gauge transformation $g \in G_0$ defines an isometry $\{a_0\} \times A^0(\Sigma^\pm) \rightarrow \{a_0 \cdot g\} \times A^0(\Sigma^\pm)$, so in this way one obtains a well defined Sobolev norm on the fiber \mathcal{E}_y .

Lemma 3.7. *With respect to suitable Sobolev completions, the following holds:*

- (1) sw_κ is smooth.
- (2) The fiberwise linear map d is fiberwise Fredholm of index $w_\tau - b_1 + 1$, and c_β is a compact map.
- (3) There exists positive constants c, C such that

$$\|(\Psi, v)\| \geq C \Rightarrow \|sw_\kappa(\Psi, v)\| > c .$$

- (4) The map $c_\kappa = sw_\kappa - d$ is compact.

Therefore the Seiberg-Witten map sw_κ satisfies always the properties $\mathcal{P}1$, $\mathcal{P}2$ (1) and $\mathcal{P}3$ in section 3.3. It also satisfies $\mathcal{P}2$ (2) for all maps $\kappa : B \rightarrow i\mathbb{H}_g^+ \setminus \{0\}$.

The first and the third statements in the lemma are easy to see. The crucial properness assertion (2) is stated in [Fu1], [Fu2]. A proof of the analogue statement for another version of the Seiberg-Witten map can be found in [BF]. A detailed proof for our version, and an analogue properness property in a different gauge theoretic context can be found in [B]. Similar methods can be also used to treat the 3-dimensional Casson-Seiberg-Witten theory.

3.5. Finite dimensional approximation. We will need the following simple geometric construction. Let \mathcal{A} be a (real or complex) Hilbert space, and $A \subset \mathcal{A}$ a finite dimensional subspace. Following [BF] we introduce, for every $\varepsilon > 0$ the retraction

$$\rho_{\varepsilon, A} : \mathcal{A}^+ \setminus S_\varepsilon(A^\perp) \rightarrow \mathcal{A}^+$$

in the following way. For every $a \in \mathcal{A} \setminus \{0\}$ put

$$s_{\varepsilon, a} := \frac{\|a\|^2 - \varepsilon^2}{2\|a\|^2} , \quad c_{\varepsilon, a} = s_{\varepsilon, a}a , \quad r_{\varepsilon, a} := \frac{\|a\|^2 + \varepsilon^2}{2\|a\|} .$$

Let $S_{\varepsilon, a} \subset \mathbb{R}a + A^\perp$ be the hypersphere of $\mathbb{R}a + A^\perp$ defined by the equation

$$\|b - c_{\varepsilon, a}\|^2 + \|a'\|^2 = r_{\varepsilon, a}^2 .$$

The hypersphere $S_{\varepsilon, a}$ has the properties

$$a \in S_{\varepsilon, a} , \quad S_\varepsilon(A^\perp) \subset S_{\varepsilon, a} .$$

⁴We are grateful to Markus Bader for pointing out this subtlety to us.

Consider also the spherical calotte:

$$C_{\varepsilon,a} := \{ta + a' \in S_{\varepsilon,a} \mid t > 0\} \subset S_{\varepsilon,a} .$$

Denote by $C_{\varepsilon,\infty} \subset [A^\perp]^+$ the exterior of the sphere $S_\varepsilon(A^\perp) \subset A^\perp$ (including ∞), and by $C_{\varepsilon,0}$ its interior. Now note that

$$\mathcal{F}_{\varepsilon,A} := \{C_{\varepsilon,a} \mid a \in A^+\}$$

is a foliation of $\mathcal{A}^+ \setminus S_\varepsilon(A^\perp)$ with closed leaves; the leaves are all diffeomorphic to the standard disk of A^\perp . The retraction $\rho_{\varepsilon,A}$ assigns the point $a \in A^+$ to any point of the leaf $C_{\varepsilon,a} \subset \mathcal{A}^+$. Note that for any $z \in \mathcal{A}$ one has the implication

$$(z \in \mathcal{A}^+ \setminus S_\varepsilon(A^\perp), \|z\| \geq \varepsilon) \Rightarrow \|\rho_{\varepsilon,A}(z)\| \geq \|z\| \quad (13)$$

(equality is obtained when $\|z\| = \varepsilon$ or $z \in A$). A second important property of the retraction $\rho_{\varepsilon,A}$ is

$$z \in \mathcal{A} \setminus A^\perp \Rightarrow (\rho_{\varepsilon,A}(z) = \lambda_{\varepsilon,z} p_A(z) \text{ with } \lambda_{\varepsilon,z} \geq 1) . \quad (14)$$

Any \mathbb{R} -linear isometry u of \mathcal{A} which leaves the subspace A invariant will also leave invariant the foliation $\mathcal{F}_{\varepsilon,A}$. Therefore

Remark 3.8. $\rho_{\varepsilon,A}$ is equivariant with respect to any \mathbb{R} -linear isometry of \mathcal{A} which leaves the subspace A invariant.

These retractions play a fundamental role in the following construction of finite dimensional approximations. This construction is a refinement of the one developed in [BF]. The main difference is that we have to work over a base B , and that we treat the real and complex summands separately.

Consider again an S^1 -equivariant map $\mu : \mathcal{E} \times \mathcal{V} \rightarrow \mathcal{F} \times \mathcal{W}$ over B satisfying the properties $\mathcal{P}1$, $\mathcal{P}2$, $\mathcal{P}3$ of section 3.3. Recall from section 3.3 that we denoted by d the linearization of μ at the 0-section and by δ and l the complex and the real components of d . We may suppose that the \mathbb{R} -linear operator l induces an isometry $\mathcal{V} \rightarrow \mathcal{W}_0$. A finite rank subbundle $F \subset \mathcal{F}$ will be called *admissible* if it is mapped surjectively onto the linear space defined by the family of cokernels $(\text{coker}(\delta_y))_{y \in B}$. A finite dimensional subspace $W \subset \mathcal{W}$ will be called admissible if it contains H . A pair (F, W) will be called admissible if F and W are both admissible; in this case, for every $y \in B$ the product $F_y \times W$ is mapped surjectively onto $\text{coker}(d_y)$.

For every admissible pair $\pi = (F, W)$ the preimage $d^{-1}(F \times W)$ is a finite rank subbundle of $\mathcal{E} \times \mathcal{V}$ which splits as

$$d^{-1}(F \times W) = \delta^{-1}(F) \times l^{-1}(W) .$$

We denote by W_0 the orthogonal complement of H in W , and put $V := l^{-1}(W) = l^{-1}(W_0)$, $E := \delta^{-1}(F) \subset \mathcal{E}$. The pair (E, F) represents $\text{ind}(\delta) \in K(B)$. We get *topological* orthogonal direct sum decompositions

$$\mathcal{F} = F \oplus F^\perp , \quad \mathcal{E} = E \oplus E^\perp , \quad \mathcal{W} = W \oplus W^\perp = H \oplus W_0 \oplus W^\perp , \quad \mathcal{V} = V \oplus V^\perp .$$

The product $F \times W$ is a finite dimensional Hilbert subbundle of $\mathcal{F} \times \mathcal{W}$ whose orthogonal complement is $F^\perp \times W^\perp$. The retraction

$$\rho_{\varepsilon, F \times W} : [\mathcal{F} \times \mathcal{W}]_B^+ \setminus S_\varepsilon(F^\perp \times W^\perp) \longrightarrow [F \times W]_B^+$$

is defined fiberwise. We will see that, for sufficiently small $\varepsilon > 0$ and sufficiently large admissible pairs $\pi = (F, W)$, the image of the restriction $\mu|_{E \times V}$ does not

intersect $S_\varepsilon(F^\perp \times W^\perp)$. Therefore we can define a map

$$\mu_{\varepsilon,\pi} := \{\rho_{\varepsilon,F \times W} \circ \mu\}|_{E \times V} : E \times V \longrightarrow [F \times W]_B^+,$$

which belongs to the class studied in section 3.1. Such a map will be called a *finite dimensional approximation* of μ . The result we need is very much similar to the first part of Lemma 2.3 in [BF]. We know that the preimage $\mu^{-1}(D_c(\mathcal{F} \times \mathcal{W}))$ is contained in the disk bundle $D_C(\mathcal{E} \times \mathcal{V}) \subset D_C(\mathcal{E}) \times D_C(\mathcal{V})$. The image $k(D_C(\mathcal{E}) \times D_C(\mathcal{V}))$ is relatively compact in the total space $\mathcal{F} \times \mathcal{W}$, because k is compact by property $\mathcal{P}3$. Now fix $\eta > 0$ and let M_η be a finite subset of $\mathcal{F} \times \mathcal{W}$ such that $k(D_C(\mathcal{E}) \times D_C(\mathcal{V}))$ is contained in the union of the balls of radius η with centers in M_η .

A pair $\pi := (F, W)$ will be called η -admissible if it is admissible and $F \times W$ contains the finite set M_η . The set of η -admissible pairs is non-empty and cofinal in the set of pairs of finite dimensional subspaces (F, W) .

Lemma 3.9. (*Finite dimensional approximations*) *Let $0 < \eta < \frac{\varepsilon}{4}$. Then*

- (1) *For any η -admissible pair $\pi = (F, W)$ one has*

$$\text{im}(\mu|_{E \times V}) \cap S_c(F^\perp \times W^\perp) = \emptyset,$$

so the finite dimensional approximation

$$\mu_{c,\pi} := \{(\rho_{c,F \times W}) \circ \mu\}|_{E \times V} : E \times V \longrightarrow (F \times W)_B^+$$

is defined.

- (2) *The restriction $\mu_{c,\pi}|_{D_C(F) \times D_C(V)}$ takes values in $F \times W$.*
(3) *For any η -admissible pair $\pi = (F, W)$ the finite dimensional approximation $\mu_{c,\pi}$ satisfies the conditions **P1**, **P2** (see section 3.1) with the same constants C, c, ε_0 , isometry $l : \mathcal{V} \rightarrow \mathcal{W}_0 \subset \mathcal{W}$ and the same map $h : B \rightarrow H$ as μ .*

Proof: 1. If the intersection $\text{im}(\mu|_{E \times V}) \cap S_c(F^\perp \oplus W^\perp)$ was not empty, there would exist a point $(e, v) \in E \times V$ such that $\mu(e, v) \in S_c(F^\perp \times W^\perp)$. Since $S_c(F^\perp \times W^\perp) \subset D_c(\mathcal{F} \times \mathcal{W})$, it follows $(e, v) \in D_C(\mathcal{E}) \times D_C(\mathcal{V})$. Therefore

$$\mu(e, v) = d(e, v) + k(e, v) \in F \times W_0 + k(D_C(\mathcal{E}) \times D_C(\mathcal{V})).$$

But any element in the second set $k(D_C(\mathcal{E}) \times D_C(\mathcal{V}))$ is η -close to an element in $M_\eta \subset F \times W$, so $\mu(e, v)$ is η -close to $F \times W$. Since $\eta < \frac{\varepsilon}{4}$, this contradicts $\mu(e, v) \in S_c(F^\perp \oplus W^\perp)$.

2. The same argument shows that $\mu(D_C(E) \times D_C(V))$ does not intersect the complement of $D_c(F^\perp \oplus W^\perp)$ in $F^\perp \oplus W^\perp$.

3. We have to check that, for an η -admissible pair $\pi = (F, W)$, the finite dimensional approximation $\mu_{c,\pi}$ has the two properties **P1**, **P2** in section 3.1. For a point $(e, v) \in E \times V$ with $\|(e, v)\| \geq C$ it holds $\|\mu(e, v)\| > c$ so, by (13), we have

$$\|\rho_{c,F \times W}(\mu(e, v))\| \geq \|\mu(e, v)\| > c. \quad (15)$$

On the other hand, for any $y \in B, v \in V$ one has $\mu(0_y^E, v) = h(y) + l(v) \in \{0_y^F\} \times W$, hence

$$\mu_{c,\pi}(0_y^E, v) = \rho_{c,F \times W}(\mu(0_y^E, v)) = \mu(0_y^E, v) = h(y) + l(v). \quad \blacksquare$$

3.6. Compatibility properties.

Lemma 3.10. (*Coherence Lemma*) *Let $0 < \eta < \frac{c}{4}$, let $\pi = (F, W)$, $\tilde{\pi} = (\tilde{F}, \tilde{W})$ be two η -admissible pairs with $\pi \subset \tilde{\pi}$, and let F', W' be the orthogonal complements of F, W in \tilde{F}, \tilde{W} respectively. The map*

$$\mu_{c,\pi,\tilde{\pi}} := \iota \circ \left\{ [\mu_{c,\pi} \circ (\mathfrak{p}_E, \mathfrak{p}_V)] \wedge_B [(\mathfrak{p}_{F'}, \mathfrak{p}_{W'}) \circ (\delta, l)]_B^+ \right\} : \tilde{E} \times \tilde{V} \rightarrow \tilde{F} \times \tilde{W}$$

*satisfies properties **P1**, **P2** with constants C, γ (for a sufficiently small γ with $0 < \gamma < c$), ε_0 , and one has $\{\mu_{c,\pi}\} = \{\mu_{c,\tilde{\pi}}\}$.*

Proof: The first statement follows from Proposition 3.5. We use the same method as in the proof of Lemma 2.3 in [BF] to construct a homotopy between the restriction of the two maps to the product $D_C(\tilde{E}) \times D_C(V)$ and we will apply the homotopy invariance property of our invariant (see Proposition 3.2). The main difference compared to [BF] is that we have to control the restriction to the S^1 -fixed point set, but we do not need an extension of the homotopy to the whole $\tilde{E} \times \tilde{V}$. For completeness we include detailed arguments adapted to our situation.

Proof: Denote by E', V' the orthogonal complements of E, V in \tilde{E}, \tilde{V} . We define the map

$$H : [0, 4] \times [D_C(\tilde{E}) \times D_C(\tilde{V})] \longrightarrow [\mathcal{F} \times \mathcal{W}] \setminus \left[\tilde{F}^\perp \times \tilde{W}^\perp \setminus \mathring{D}_c(\tilde{F}^\perp \times \tilde{W}^\perp) \right] \quad (16)$$

by the formula ⁵

$$H_t = \begin{cases} d + [(1-t) \text{id}_{\mathcal{F} \times \mathcal{W}} + t \mathfrak{p}_{F \times W}] \circ k & \text{for } 0 \leq t \leq 1, \\ d + \mathfrak{p}_{F \times W} \circ k \circ [(2-t) \text{id}_{\tilde{E} \times \tilde{V}} + (t-1) \mathfrak{p}_{E \times V}] & \text{for } 1 \leq t \leq 2, \\ \mathfrak{p}_{F \times W} \circ k \circ \mathfrak{p}_{E \times V} + [d - (t-2) \mathfrak{p}_{F \times W} \circ d \circ \mathfrak{p}_{E' \times V'}] & \text{for } 2 \leq t \leq 3, \\ \mathfrak{p}_{F' \times W'} \circ d + [(4-t) \mathfrak{p}_{F \times W} + (t-3) \rho_{c, F \times W}] \circ \mu \circ \mathfrak{p}_{E \times V} & \text{for } 3 \leq t \leq 4. \end{cases}$$

Claim: *H is a well defined, continuous, S^1 -equivariant map over B.*

This follows from:

a) For a point $(t, \tilde{e}, \tilde{v}) \in [0, 4] \times D_C(\tilde{E}) \times D_C(\tilde{V})$, the term $\rho_{c, F \times W}(\mu(\mathfrak{p}_{E \times V}(\tilde{e}, \tilde{v})))$ is finite, so the convex combination in the fourth branch is defined and finite.

Indeed, recall that the retraction $\rho_{c, F \times W}$ is finite on the complement of the leaf $[F^\perp \times W^\perp] \setminus D_c(F^\perp \times W^\perp)$. Therefore it suffices to note that $k(D_C(\mathcal{E}) \times D_C(\mathcal{V}))$ is η -close to $F \times W$ and $d(E \times V) \subset F \times W$, so the point $\mu(\mathfrak{p}_{E \times V}(\tilde{e}, \tilde{v}))$ is η -close to $F \times W$ for $(\tilde{e}, \tilde{v}) \in D_C(\tilde{E}) \times D_C(\tilde{V})$. Therefore

$$\mu(\mathfrak{p}_{E \times V}(\tilde{e}, \tilde{v})) \notin [F^\perp \times W^\perp] \setminus \mathring{D}_c(F^\perp \times W^\perp).$$

b) The formulae given for the four components of H agree on the intersections of their domains.

c) H takes values in $[\mathcal{F} \times \mathcal{W}] \setminus \left[\tilde{F}^\perp \times \tilde{W}^\perp \setminus \mathring{D}_c(\tilde{F}^\perp \times \tilde{W}^\perp) \right]$.

Indeed, for $(t, \tilde{e}, \tilde{v}) \in [0, 4] \times D_C(\tilde{E}) \times D_C(\tilde{V})$ we see as in the proof of a) that the right hand term of H_t must be η -close to $\tilde{F} \times \tilde{W}$, so $H([0, 4] \times D_C(\tilde{E}) \times D_C(\tilde{V}))$ avoids $[F^\perp \times W^\perp] \setminus \mathring{D}_c(F^\perp \times W^\perp)$.

⁵The third branch of the homotopy was omitted in [BF].

The map H has the following properties:

- (1) H_0 coincides with the restriction $\mu|_{D_C(\tilde{E}) \times D_C(\tilde{V})}$.
- (2) H_4 coincides with the map $\mu_{c,\pi,\tilde{\pi}}$ composed with the inclusion $\tilde{F} \times \tilde{V} \hookrightarrow [\mathcal{F} \times \mathcal{W}]_B^+ \setminus S_c(\tilde{F}^\perp \times \tilde{W}^\perp)$.
- (3) One has

$$H_t(0_{\tilde{E}}, \tilde{v}) = h(y) + l(\tilde{v}), \quad \forall t \in [0, 4] \quad \forall y \in B \quad \forall \tilde{v} \in D_C(\tilde{V}). \quad (17)$$

Formula (17) follows from (12) and the fact that l is an isometry, so it commutes with orthogonal projections.

- (4) $H([0, 4] \times \partial(D_C(\tilde{E}) \times D_C(\tilde{V}))) \cap [\tilde{F}^\perp \times \tilde{W}^\perp] = \emptyset$.

Indeed, for $(\tilde{e}, \tilde{v}) \in \partial(D_C(\tilde{E}) \times D_C(\tilde{V}))$ we get $\|H_0(\tilde{e}, \tilde{v})\| = \|\mu(\tilde{e}, \tilde{v})\| \geq c$, whereas $\|\mu(\tilde{e}, \tilde{v})\|$ is η -close to $F \times W \subset \tilde{F} \times \tilde{W}$. Moreover, for $t \in [0, 1]$ it holds $\|H_t(\tilde{e}, \tilde{v}) - H_0(\tilde{e}, \tilde{v})\| = t\|(\mathfrak{p}_{F^\perp \times W^\perp} \circ k)(\tilde{e}, \tilde{v})\| \leq \eta$. For $t \geq 2$ we have

$$\mathfrak{p}_{F' \times W'} \circ h_t = \mathfrak{p}_{F' \times W'} \circ d,$$

so $H_t(\tilde{e}, \tilde{v})$ can belong to $\tilde{F}^\perp \times \tilde{W}^\perp$ only when $\mathfrak{p}_{F' \times W'} \circ d(\tilde{e}, \tilde{v}) = 0$, i.e. when $(\tilde{e}, \tilde{v}) \in E \times V$. For such a pair we find

$$H_t(\tilde{e}, \tilde{v}) = (d + \mathfrak{p}_{F \times W} \circ k)(\tilde{e}, \tilde{v}) = \mu(\tilde{e}, \tilde{v}) - (\mathfrak{p}_{F^\perp \times W^\perp} \circ k)(\tilde{e}, \tilde{v}) \quad \forall t \in [1, 3],$$

$$H_t(\tilde{e}, \tilde{v}) \in [\mathfrak{p}_{F \times W}(\mu(\tilde{e}, \tilde{v})), \rho_{c,F}(\mu(\tilde{e}, \tilde{v}))] \quad \forall t \in [3, 4],$$

so $H_t(\tilde{e}, \tilde{v})$ is a non-vanishing vector of $F \times W$ (more precisely a positive multiple of $\mathfrak{p}_{F \times W}(\mu(\tilde{e}, \tilde{v})) = \mu(\tilde{e}, \tilde{v}) - (\mathfrak{p}_{F^\perp \times W^\perp} \circ k)(\tilde{e}, \tilde{v})$) for any $t \in [1, 4]$.

These properties have the following important consequence:

Remark: *The composition $\rho_{c,\tilde{\pi}} \circ H$ is nowhere vanishing on the space*

$$[0, 4] \times \left\{ \partial \left[D_C(\tilde{E}) \times D_C(\tilde{V}) \right] \cup \left[0^{\tilde{E}} \times \tilde{V} \right] \right\}.$$

This follows from the fact that the vanishing locus of the retraction $\rho_{c,\tilde{\pi}}$ is the leaf $\tilde{D}_c(\tilde{F}^\perp \times \tilde{W}^\perp) \subset \tilde{F}^\perp \times \tilde{W}^\perp$. On the other hand we have

$$\rho_{c,\tilde{\pi}} \circ H_0 = \mu_{c,\tilde{\pi}}|_{D_C(\tilde{E}) \times D_C(\tilde{V})}, \quad \rho_{c,\tilde{\pi}} \circ H_4 = \mu_{c,\pi,\tilde{\pi}}|_{D_C(\tilde{E}) \times D_C(\tilde{V})}$$

It suffices now to apply Proposition 3.2. ■

Using Proposition 3.5 and Lemma 3.10 we obtain

Corollary 3.11. *Let $\mu : \mathcal{E} \times \mathcal{V} \rightarrow \mathcal{F} \times \mathcal{W}$ be an S^1 -equivariant map over a compact base B satisfying $\mathcal{P}1$, $\mathcal{P}2$, $\mathcal{P}3$, and let $0 < \eta < \frac{c}{4}$. Fix an orientation \mathfrak{o} of the finite dimensional summand H of \mathcal{W} . The elements*

$$\{\mu_{c,\pi}\} \in S^1 \alpha_B^{b-1}(S(E)_{+B}, F_B^+)$$

associated with η -admissible pairs $\pi = (F, W)$ define a unique class

$$\{\mu\} \in \alpha^{b-1}(\text{ind}(\delta))$$

which depends only on the map μ and the orientation \mathfrak{o} .

In particular, using finite dimensional approximations associated with constants $C' \geq C$ and $0 < c' \leq c$ (and parameter $0 < \eta < \frac{c'}{4}$), one obtains the same class.

Proposition 3.12. *Suppose that the restriction $\mu|_{D_C(\mathcal{E}) \times D_C(\mathcal{V})}$ is nowhere vanishing. Then $\{\mu\} = 0$.*

Proof: Since $\mu|_{D_C(\mathcal{E}) \times D_C(V)}$ is nowhere vanishing, it is easy to see that there exists $\gamma > 0$ such that $\|\mu(e, v)\| > \gamma$ for every $(e, v) \in D_C(\mathcal{E}) \times D_C(V)$. Indeed, if not there would exist a sequence $(e_n, v_n) \in D_C(\mathcal{E}) \times D_C(V)$ such that $\|\mu(e_n, v_n)\| \rightarrow 0$. Let $K \subset \mathcal{F} \times \mathcal{W}$ be a compact subspace which contains $k(D_C(\mathcal{E}) \times D_C(V))$. Since $d = (\delta, l)$ is a continuous family of Fredholm operators, it follows that $d^{-1}(K) \cap [D_C(\mathcal{E}) \times D_C(V)]$ is compact. Therefore $(e_n, v_n)_n$ admits a subsequence which converges in this intersection. The limit will be a vanishing point of μ , which contradicts the assumption.

Use now the constant $c' := \min(\gamma, c)$ (instead of c) in the construction of the finite dimensional approximations of μ . The obtained maps $\mu_{c', \pi}$ are nowhere vanishing on $D_C(E) \times D_C(V)$, and our assertion follows from the vanishing property Proposition 3.1 proved in the finite dimensional case. \blacksquare

4. FUNDAMENTAL PROPERTIES OF THE COHOMOTOPY INVARIANTS

4.1. The Hurewicz image of the cohomotopy invariant.

4.1.1. *The relative Hurewicz morphism.* Let B be a compact space, and let E, F be Hermitian bundles of ranks e, f over B . Let k be an integer and $u \in S^1 \alpha_B^k(S(E)_{+B}, F_B^+)$ a stable class. Suppose for simplicity $k \geq 0$. Consider a representative

$$\varphi : S(E)_{+B} \wedge_B \xi_B^+ \rightarrow F_B^+ \wedge_B [\mathbb{R}^k]_B^+ \wedge \xi_B^+$$

of this stable class, where $\xi = \eta \oplus \xi_0$ is the direct sum of a complex vector bundle η and a real vector bundle ξ_0 . We may suppose that the real summand ξ_0 of ξ is orientable. We choose an orientation of ξ_0 ; in this way all our bundles become oriented bundles. The space $S(E)_{+B} \wedge_B \xi_B^+$ can be identified with the fiberwise quotient $\{S(E) \times_B \xi_B^+ \} /_B \{S(E) \times_B \infty_\xi\}$. Composing φ with the canonical projection one obtains a map of pairs over B

$$\tilde{\varphi} : (S(E) \times_B \xi_B^+, S(E) \times_B \infty_\xi) \rightarrow ([F \oplus \mathbb{R}^k \oplus \xi]_B^+, \infty_{F \oplus \mathbb{R}^k \oplus \xi}) .$$

Consider now the projection $\pi : \mathbb{P}(E) \rightarrow B$ and the following bundles over $\mathbb{P}(E)$:

$$\tilde{F} := \pi^*(F)(1) , \quad \tilde{\xi} := \pi^*(\eta)(1) \oplus \pi^*(\xi_0) .$$

The map $\tilde{\varphi}$ descends to a morphism of pointed sphere bundles over $\mathbb{P}(E)$

$$\bar{\varphi} : \tilde{\xi}_{\mathbb{P}(E)}^+ \longrightarrow [\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi}]_{\mathbb{P}(E)}^+ .$$

Denote by s the real rank of ξ . Let

$$t_{\tilde{\xi}} \in H^s(\tilde{\xi}_{\mathbb{P}(E)}^+, \infty_{\tilde{\xi}}; \mathbb{Z}) , \quad t_{\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi}} \in H^{2f+k+s}([\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi}]_{\mathbb{P}(E)}^+, \infty_{\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi}}; \mathbb{Z})$$

be the Thom classes of the oriented bundles $\tilde{\xi}, \tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi}$. The formula

$$\bar{\varphi}^*(t_{\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi}}) = P_{\mathbb{P}(E)}^*(h_{\bar{\varphi}}) \cup t_{\tilde{\xi}}$$

defines a cohomology class $h_{\bar{\varphi}} \in H^{2f+k}(\mathbb{P}(E); \mathbb{Z})$ which is independent of the chosen orientation of ξ_0 and of the representative φ of the stable class u . For $k \leq 0$ one has a similar construction, but uses a $[\mathbb{R}^{-k}]_B^+$ factor on the left side.

The assignment $u = [\varphi] \mapsto h_{\bar{\varphi}}$ defines a morphism

$$h : S^1 \alpha_B^k(S(E)_{+B}, F_B^+) \rightarrow H^{2f+k}(\mathbb{P}(E); \mathbb{Z}) ,$$

which we call the *relative Hurewicz morphism* over B .

Denote by $q : \tilde{\xi} \rightarrow \mathbb{P}(E)$ the bundle projection, and by $\hat{\varphi}$ the section in the pull-back $[q^*(\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi})]_{\tilde{\xi}}^{\perp}$ over $\tilde{\xi}$ defined by $\bar{\varphi}$. Since the vanishing locus $Z(\hat{\varphi})$ of this section is compact, one can define its *localized Euler class* class $[\hat{\varphi}] \in H_{d+2e-2-2f-k}(\tilde{\xi}; \mathbb{Z})$, which coincides with the fundamental class $[Z(\hat{\varphi})]$ of the compact oriented submanifold $[Z(\hat{\varphi})]$ when $\hat{\varphi}$ is smooth and transversal to the zero section [Br].

Remark 4.1. *(The geometric interpretation of the Hurewicz morphism) Suppose that B is an oriented n -dimensional compact manifold. Then*

$$PD_{\mathbb{P}(E)}(h(u)) = [\iota_*]^{-1}([\hat{\varphi}]) ,$$

where

$$\iota_* : H_{n+2e-2-2f-k}(\mathbb{P}(E); \mathbb{Z}) \rightarrow H_{n+2e-2-2f-k}(\tilde{\xi}; \mathbb{Z}) .$$

is the isomorphism induced by the zero section of $\tilde{\xi}$. If $\hat{\varphi}$ is smooth and transversal to the zero section, then

$$PD_{\mathbb{P}(E)}(h(u)) = [\iota_*]^{-1}([Z(\hat{\varphi})]) .$$

Proof: The localized Euler class $[\hat{\varphi}] \in H_{n+2e-2-2f-k}(\tilde{\xi}; \mathbb{Z})$ is defined as the cap product $\hat{\varphi}^*(t_{q^*(\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi})}) \cap [\tilde{\xi}]$, where $[\tilde{\xi}]$ stands for the fundamental class of $\tilde{\xi}$ in cohomology with compact supports [Br]. We get

$$\begin{aligned} [\hat{\varphi}] &:= \hat{\varphi}^*(t_{q^*(\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi})}) \cap [\tilde{\xi}] = \bar{\varphi}^*(t_{\tilde{F} \oplus \mathbb{R}^k \oplus \tilde{\xi}}) \cap [\tilde{\xi}] = [p_{\mathbb{P}(E)}^*(h(u)) \cup t_{\tilde{\xi}}] \cap [\tilde{\xi}] = \\ &= p_{\mathbb{P}(E)}^*(h(u)) \cap \iota_*([\mathbb{P}(E)]) = \iota_*(h(u) \cap [\mathbb{P}(E)]) = \iota_*(PD_{\mathbb{P}(E)}(h(u))) . \end{aligned}$$

■

Let $\nu = (i, E_1) : E \rightarrow E'$ be a morphism in the category \mathcal{U}_B of complex vector bundles over B (see section 2.3). Such a morphism induces an isomorphism $E' \cong E \oplus E_1$. The complement $\mathbb{P}(E') \setminus \mathbb{P}(E_1)$ can be identified with the total space of the complex vector bundle $\pi^*(E_1)(1) \rightarrow \mathbb{P}(E)$. Multiplication with the Thom class $t_{\pi^*(E_1)(1)}$ defines a morphism

$$\begin{aligned} H^*(\mathbb{P}(E); \mathbb{Z}) &\longrightarrow H^{*+2e_1}(\pi^*(E_1)(1)_{\mathbb{P}(E)}^+, \infty_{\pi^*(E_1)(1)}; \mathbb{Z}) \cong \\ &\cong H^{*+2e_1}(\mathbb{P}(E'), \mathbb{P}(E_1); \mathbb{Z}) \longrightarrow H^{*+2e_1}(\mathbb{P}(E'); \mathbb{Z}) , \end{aligned}$$

which will be denoted by a_ν .

Now fix an element $x \in K(B)$. A morphism $\tau = (i, j; E_1, F_1, l) : (E, F) \rightarrow (E', F')$ in the category $\mathcal{T}(x)$ defines morphisms

$$\begin{aligned} a_{(i, E_1)} &: H^{2f+k}(\mathbb{P}(E); \mathbb{Z}) \rightarrow H^{2f'+k}(\mathbb{P}(E'); \mathbb{Z}) , \\ \mathbb{P}(i)_* &: H_k(\mathbb{P}(E); \mathbb{Z}) \rightarrow H_k(\mathbb{P}(E'); \mathbb{Z}) . \end{aligned}$$

For an integer $k \in \mathbb{Z}$ we define

$$H^k(x; \mathbb{Z}) := \varinjlim_{(E, F) \in x} H^{2f+k}(\mathbb{P}(E); \mathbb{Z}) , \quad H_k(x; \mathbb{Z}) := \varinjlim_{(E, F) \in x} H_k(\mathbb{P}(E); \mathbb{Z}) .$$

Using the same methods as in sections 2.1, 2.3 (stabilizing first with respect to trivial bundle enlargements) we see that these inductive limits exist in \mathcal{Ab} .

Remark 4.2. (1) *One has $H_*(x; \mathbb{Z}) = H_*(B; \mathbb{Z}) \otimes \mathbb{Z}[t]$.*

(2) For a compact n -dimensional CW complex B there exist isomorphisms

$$H^k(x; \mathbb{Z}) \simeq \bigoplus_{\substack{s-k \in 2\mathbb{Z} \\ \max(0, k-2\iota(x)+2) \leq s \leq n}} H^s(B; \mathbb{Z}) ,$$

where $\iota(x) \in \mathbb{Z}$ is the index of x . In particular, putting $n(x) := 2\iota(x) - 2 + n$, one has $H^{n(x)}(x; \mathbb{Z}) = H^n(B; \mathbb{Z})$.

The integer $n(x) := 2\iota(x) - 2 + n$ will be called *the dimension of the formal projectivization of x* .

Remark 4.3. Suppose that B is a compact connected oriented manifold of dimension n . The system of Poincaré duality isomorphisms $PD_{\mathbb{P}(E)}$ defines isomorphisms

$$PD_x : H^k(x; \mathbb{Z}) \xrightarrow{\simeq} H_{n(x)-k}(x; \mathbb{Z}) .$$

Remark 4.4. The system of Hurewicz morphisms

$$h : {}_{S^1} \alpha_B^k(S(E)_{+B}, F_B^+) \rightarrow H^{2f+k}(\mathbb{P}(E); \mathbb{Z})$$

defines a morphisms of graded groups $h_x : \alpha^*(x) \rightarrow H^*(x; \mathbb{Z})$. If B is a compact connected oriented manifold, one also gets a morphism $PD_x \circ \chi_x : \alpha^*(x) \rightarrow H_*(x; \mathbb{Z})$, which we call *the homological Hurewicz morphism*.

The result below has the following important consequence: for a moduli problem with vanishing “expected dimension”, the cohomotopy invariant yields the same information as the classical (co)homological invariant. Recall that our cohomotopy invariant $\{\mu\}$ associated with a map satisfying properties $\mathcal{P}1 - \mathcal{P}3$ belongs to $\alpha^{b-1}(x)$, where $x := \text{ind}(\delta)$, $b := \dim(H)$ (see section 3.3). The *expected dimension* $w(\mu) := 2\iota(x) + \dim(B) - b - 1$ of the moduli problem associated with μ vanishes if and only if $b - 1 = n(x)$.

Proposition 4.5. Suppose that B is a finite CW complex of dimension n . Then the Hurewicz morphism

$$h_x^{n(x)} : \alpha^{n(x)}(x) \longrightarrow H^{n(x)}(x; \mathbb{Z}) = H^n(B; \mathbb{Z}) .$$

is an isomorphism.

Proof: Suppose $n(x) \geq 0$ for simplicity. Fix a stabilizing bundle ξ . Using the same method and the same notations as in section 4.1.1 we see that the set

$${}_{S^1} \pi^0(S(E)_{+B} \wedge_B \xi_B^+, F_B^+ \wedge_B [\mathbb{R}^{n(x)}]_B^+ \wedge \xi_B^+)$$

can be identified with the set of pointed bundle maps

$$\bar{\varphi} : \tilde{\xi}_{\mathbb{P}(E)}^+ \longrightarrow [\tilde{F} \oplus \mathbb{R}^{n(x)} \oplus \tilde{\xi}]_{\mathbb{P}(E)}^+$$

over $\mathbb{P}(E)$. The latter set can be identified with $H^{\dim_{\mathbb{R}}(\mathbb{P}(E))}(\mathbb{P}(E); \mathbb{Z}) = H^n(B; \mathbb{Z})$ by Proposition 5.15 via the map $\bar{\varphi} \mapsto h_{\bar{\varphi}}$. The obtained bijections

$${}_{S^1} \pi^0(S(E)_{+B} \wedge_B \xi_B^+, F_B^+ \wedge_B [\mathbb{R}^{n(x)}]_B^+ \wedge \xi_B^+) \simeq H^n(B; \mathbb{Z})$$

are compatible with morphisms $\xi \rightarrow \xi'$ in the category \mathcal{C}_B and with morphisms $(E, F) \rightarrow (E', F')$ in the category $\mathcal{T}(x)$. Therefore we get a bijection $\alpha^{n(x)}(x) \rightarrow H^n(B; \mathbb{Z})$, which coincides with the Hurewicz map by the definition. \blacksquare

4.1.2. *A comparison theorem.* The main result of this section states: the virtual fundamental class of the moduli space of solutions associated with a map μ satisfying properties $\mathcal{P}1$, $\mathcal{P}2$, $\mathcal{P}3$ can be identified with the image of the cohomotopy invariant under the homological Hurewicz map. Applied to Seiberg-Witten theory, this implies that the *full* Seiberg-Witten type invariant coincides with the Hurewicz image of the cohomotopy Seiberg-Witten invariant.

We begin with the finite dimensional case. Let B be a compact oriented manifold, $p : E \rightarrow B$, $q : F \rightarrow B$ Hermitian bundles over B , let V , W be Euclidean spaces, and let $\mu : E \times V \rightarrow [F \times W]_B^+$ be an S^1 -equivariant map over B satisfying properties **P1**, **P2** of section 3.1. The invariant $\{\mu\} \in S^1 \alpha_B^{b-1}(S(E)_{+B}, F_B^+)$ is defined by a map of pairs

$$(S(E) \times D_R(\mathbb{R} \oplus V), S(E) \times S_R(\mathbb{R} \oplus V)) \rightarrow ([F \times W]_B^+, [F \times W]_B^+ \setminus \mathring{D}_\varepsilon(F \times W))$$

induced by the restriction $\mu_{R,\varepsilon} : D_R(E) \times D_R(V) \rightarrow (F \times W)_B^+$ of μ to a sufficiently large cylinder $D_R(E) \times D_R(V)$. The vanishing locus of μ (regarded as section in the bundle $(p^*(F) \times V) \times W \rightarrow E \times V$) is an S^1 -invariant compact space contained in the open subspace $\mathring{D}_R(E) \times \mathring{D}_R(V) \setminus [0^E \times_B D_R(V)]$ of the cylinder. Its S^1 -quotient can be identified with the vanishing locus of the section $\mathring{\mu}_{R,\varepsilon}$ induced by $\mu_{R,\varepsilon}$ on the S^1 -quotient $\mathbb{P}(E) \times \mathring{D}_R(\mathbb{R} \oplus V)$ of $S(E) \times \mathring{D}_R(\mathbb{R} \oplus V)$. Using Remark 4.1 one obtains

Corollary 4.6. *Suppose that B is a compact oriented manifold. Via the isomorphism $H_*(\mathbb{P}(E) \times \mathring{D}_R(\mathbb{R} \oplus V); \mathbb{Z}) \simeq H_*(\mathbb{P}(E); \mathbb{Z})$ the Poincaré dual $PD_{\mathbb{P}(E)}(h(\{\mu\}))$ coincides with the virtual fundamental class associated with the section $\mathring{\mu}_{R,\varepsilon}$. If this section is smooth and transversal to the zero section, then $PD_{\mathbb{P}(E)}(h(\{\mu\}))$ can be identified with the fundamental class of the vanishing locus $Z(\mathring{\mu}_{R,\varepsilon}) \subset \mathbb{P}(E) \times \mathring{D}_R(\mathbb{R} \oplus V)$.*

Note that μ is nowhere vanishing outside the cylinder $D_R(E) \times D_R(V)$, so the vanishing loci of μ and $\mu_{R,\varepsilon}$ can be identified. The vanishing locus $M := Z(\mathring{\mu}_{R,\varepsilon}) \cong Z(\mu)/S^1$ will be called the “moduli space” associated with the map μ .

Let $p : \mathcal{E} \rightarrow B$, $q : \mathcal{F} \rightarrow B$ be complex Hilbert bundles over B , let \mathcal{V} , \mathcal{W} be real Hilbert spaces, and let $\mu : \mathcal{E} \times \mathcal{V} \rightarrow \mathcal{F} \times \mathcal{W}$ be an S^1 -equivariant map over B satisfying properties $\mathcal{P}1$, $\mathcal{P}2$, $\mathcal{P}3$ in section 3.3. Denote by $\pi : \mathbb{P}(\mathcal{E}) \rightarrow B$ the natural projection. The map $\mu_{R,\varepsilon}$ descends to a smooth section $\mathring{\mu}_{R,\varepsilon}$ in the bundle

$$\pi^*(\mathcal{F})(1) \times \mathring{D}_R(\mathbb{R} \oplus \mathcal{V}) \times \mathcal{W} \rightarrow \mathbb{P}(\mathcal{E}) \times \mathring{D}_R(\mathbb{R} \oplus \mathcal{V}) ,$$

and again one can identify the moduli space $\mathcal{M} := Z(\mu)/S^1$ of μ with the vanishing locus $Z(\mathring{\mu}_{R,\varepsilon})$ of this section. Using the same argument as in the proof of Proposition 3.12, we see that the moduli space \mathcal{M} is compact. Suppose now that

\mathcal{P}_4 : B is a compact, smooth, connected, oriented manifold, μ is smooth and the fiberwise differential of $k := \mu - d$ at any point is a compact operator.

This condition is always satisfied in practical gauge theoretical situations; indeed, the map k is usually given by the composition of a smooth map $\mathcal{E} \times \mathcal{V} \rightarrow \mathcal{F}_1 \times \mathcal{W}_1$ with a map $\mathcal{F}_1 \times \mathcal{W}_1 \rightarrow \mathcal{F} \times \mathcal{W}$ over B defined by a smooth family of compact operators. The condition \mathcal{P}_4 implies that $\mathring{\mu}_{R,\varepsilon}$ is a smooth Fredholm section on the Banach manifold $\mathbb{P}(\mathcal{E}) \times \mathring{D}_R(\mathbb{R} \oplus \mathcal{V})$. In order to give sense to the virtual

fundamental class of the moduli space \mathcal{M} we have to trivialize the determinant line bundle $\det(\text{index}(D\check{\mu}_{R,\varepsilon}))$ over \mathcal{M} . Equivalently, it suffices to trivialize the line bundle $\det(\text{index}(D\mu))$ over $Z(\mu)$. In these formulae the symbol D stands for the family of intrinsic derivatives of a section at its zero locus, and μ is regarded as a section of the bundle $[p^*(\mathcal{F}) \times \mathcal{V}] \times \mathcal{W} \rightarrow \mathcal{E} \times \mathcal{V}$. For a point $(e, v) \in Z(\mu)$ with $p(e) = y$ one has a natural identification

$$\det(\text{index}(D_{(e,v)}\mu)) = \Lambda^n(T_y(B)) \otimes \det(\text{index}(d_{(e,v)}\mu|_{\mathcal{E}_y \times \mathcal{V}})) ,$$

where $n := \dim(B)$ and $\mu|_{\mathcal{E}_y \times \mathcal{V}} : \mathcal{E}_y \times \mathcal{V} \rightarrow \mathcal{F}_y \times \mathcal{W}$ is the restriction of μ to the fiber over y . By the condition $\mathcal{P}4$, the differential of this restriction is congruent with the operator $d_y = (\delta_y, l)$ modulo a compact operator. Therefore (since the family $\delta = (\delta_y)_{y \in B}$ has a canonical complex orientation, and B is oriented) one obtains a trivialization of $\det(\text{index}(D\mu))$ for every orientation \mathfrak{o} of $\text{coker}(l) = H$. This is precisely the orientation parameter involved in the definition of the cohomotopy invariant $\{\mu\}$. Fix such an orientation \mathfrak{o} . Using the results in [Br], we obtain a virtual fundamental class in Čech homology $[\mathcal{M}]^{\text{vir}} \in \check{H}_w(\mathcal{M}; \mathbb{Z})$, where

$$w = w(\mu) := n + 2\iota(\text{ind}(\delta)) - b - 1 = n(\text{ind}(\delta)) - (b - 1)$$

is the expected dimension of our moduli problem (the index of the section $\check{\mu}_{R,\varepsilon}$).

Put $x := \text{ind}(\delta)$, and note that the group

$$H_w(x; \mathbb{Z}) = \bigoplus_{0 \leq 2i \leq w} H_{w-2i}(B; \mathbb{Z}) \otimes t^i .$$

can be identified with $H_w(\mathbb{P}(\mathcal{E}) \times \mathring{D}_R(\mathbb{R} \oplus V); \mathbb{Z}) = H_w(\mathbb{P}(\mathcal{E}); \mathbb{Z})$.

Definition 4.7. *The full homological invariant of μ is the image $\{\mu\}_{\mathbb{H}}$ of the class $[\mathcal{M}]^{\text{vir}}$ in the group $H_w(\text{ind}(\delta); \mathbb{Z})$.*

Theorem 4.8. *Suppose that conditions $\mathcal{P}1 - \mathcal{P}4$ hold. Then*

$$\{\mu\}_{\mathbb{H}} = PD_x \circ h_x(\{\mu\}) .$$

Proof: As in section 3.5 choose a finite dimensional approximation $\mu_{c,\pi}$ of π , associated with an η -admissible pair (F, W) . Define $\mu_{c,\pi,\infty} : D_C(\mathcal{E}) \times D_C(\mathcal{V}) \rightarrow \mathcal{F} \times \mathcal{W}$ by

$$\mu_{c,\pi,\infty}(e, v) = \mu_{c,\pi}(p_E(e), p_V(v)) + p_{F^\perp \times W^\perp} \circ d \circ p_{E^\perp \times V^\perp} .$$

This map takes finite values by Lemma 3.9. We claim that there exists a smooth homotopy

$$\mathcal{H} : [0, 4] \times D_C(\mathcal{E}) \times D_C(\mathcal{V}) \rightarrow \mathcal{F} \times \mathcal{W}$$

between $\mu|_{D_C(\mathcal{E}) \times D_C(\mathcal{V})}$ and $\mu_{c,\pi,\infty}$ in the space of S^1 -equivariant Fredholm maps over B , such that for $0 \leq t \leq 4$ the map \mathcal{H}_t has no zeroes in $\partial[D_C(\mathcal{E}) \times D_C(\mathcal{V})] \cup 0^\mathcal{E} \times D_C(\mathcal{V})$. To obtain such a homotopy it suffices to replace $\check{E}, \check{V}, \check{F}, \check{W}$ in the definition of the homotopy H used in the proof of Lemma 3.10 by $\mathcal{E}, \mathcal{V}, \mathcal{F}, \mathcal{W}$, and to compose the resulting map from the right with a smooth homeomorphism $\theta : [0, 4] \rightarrow [0, 4]$ having the properties

$$\theta(i) = i , \quad \theta^{(k)}(i) = 0 \text{ for } i \in \{0, 1, 2, 3, 4\} , \quad k \geq 1$$

(to assure differentiability). Using the homotopy invariance of the virtual class [Br], we can identify $\{\mu\}_{\mathbb{H}}$ with the image of the virtual class $[\mu_{c,\pi,\infty}]^{\text{vir}}$ in $H_w(\mathbb{P}(\mathcal{E}); \mathbb{Z})$.

On the other hand, by the ‘‘associativity property’’ of the virtual class (see Proposition 14 (4) in [Br]) and Corollary 4.6, the latter is just the image of $PD_{\mathbb{P}(E)}(h(\{\mu_{c,\pi}\}))$ via the embedding $\mathbb{P}(E) \rightarrow \mathbb{P}(\mathcal{E})$. But $PD_{\mathbb{P}(E)}(h(\{\mu_{c,\pi}\}))$ is a representative of $PD_x \circ h_x(\{\mu\})$. ■

4.2. Cohomotopy invariant jump formulae.

4.2.1. *General results.* Let

$$M \longrightarrow N \longrightarrow P$$

be a cofiber sequence of pointed S^1 -spaces over a compact basis B . For every pointed S^1 -space Y over B there is an associated long exact sequence of cohomotopy groups

$$\cdots \rightarrow {}_{S^1}\alpha_B^k(P, Y) \rightarrow {}_{S^1}\alpha_B^k(N, Y) \rightarrow {}_{S^1}\alpha_B^k(M, Y) \xrightarrow{\partial} {}_{S^1}\alpha_B^{k+1}(P, Y) \rightarrow \cdots \quad (18)$$

The connecting morphism

$$\partial : {}_{S^1}\alpha_B^k(M, Y) = {}_{S^1}\alpha^{k+1}(M \wedge_B S^1, Y) \longrightarrow {}_{S^1}\alpha^{k+1}(P, Y)$$

is given by composition with the contraction map $\mathbf{c} : P \rightarrow M \wedge_B \underline{S}^1$ induced by a fixed homotopy equivalence between P and the mapping cone of the map $M \rightarrow N$. For the cofiber sequence

$$S(\xi)_{+B} \longrightarrow D(\xi)_{+B} \longrightarrow \xi_B^+$$

associated with a vector bundle ξ over a compact basis B , the morphism ∂ can be described in the following way. The obvious isomorphisms

$$S(\xi)_{+B} \wedge_B \underline{S}^1 \cong S(\xi) \times [0, 1] / S(\xi) \times \{0, 1\}, \quad \xi_B^+ \cong S(\xi) \times [0, 1] / \sim$$

(where \sim is the equivalence relation generated by $(v, 0) \sim (v', 0)$, $(v, 1) \sim (v', 1)$) allow us to use $S(\xi) \times [0, 1] / S(\xi) \times \{0, 1\}$, $S(\xi) \times [0, 1] / \sim$ as models for $S(\xi)_{+B} \wedge_B \underline{S}^1$ and ξ_B^+ . Using these models, the morphism ∂ is given by composition with the contraction map

$$\mathbf{c}_\xi : S(\xi) \times [0, 1] / \sim \longrightarrow S(\xi) \times [0, 1] / S(\xi) \times \{0, 1\} \quad (19)$$

induced by the identity of $S(\xi) \times [0, 1]$.

Consider now an oriented b -dimensional real vector space H and the cofiber sequence over B associated with the trivial bundle $\underline{H} = B \times H$ over B :

$$S(\underline{H})_{+B} \longrightarrow D(\underline{H})_{+B} \longrightarrow \underline{H}_B^+.$$

Let E be a Hermitian vector bundle over B . Taking smash product with $S(E)_{+B}$ over B yields the following cofiber sequence over B

$$S(E)_{+B} \wedge_B S(\underline{H})_{+B} \rightarrow S(E)_{+B} \rightarrow S(E)_{+B} \wedge_B \underline{H}_B^+$$

Since $S(E)_{+B} \wedge_B S(\underline{H})_{+B} = [S(E) \times S(H)]_{+B}$, the associated long exact cohomotopy sequence is

$$\begin{aligned} \cdots \rightarrow {}_{S^1}\alpha_B^{-1}(S(E)_{+B} \wedge_B \underline{H}_B^+, [F \oplus \underline{H}]_B^+) &\rightarrow {}_{S^1}\alpha_B^{-1}(S(E)_{+B}, [F \oplus \underline{H}]_B^+) \rightarrow \\ &\rightarrow {}_{S^1}\alpha_B^{-1}([S(E) \times S(H)]_{+B}, [F \oplus \underline{H}]_B^+) \xrightarrow{\partial} \\ &\rightarrow {}_{S^1}\alpha_B^0(S(E)_{+B}, [F]_B^+) \rightarrow {}_{S^1}\alpha_B^0(S(E)_{+B}, [F \oplus \underline{H}]_B^+) \rightarrow \cdots \quad (20) \end{aligned}$$

Note that one has canonical base change isomorphisms

$$s^1 \alpha_B^k([S(E) \times S(H)]_{+B}, [F \oplus \underline{H}]_B^+) \simeq s^1 \alpha_{\tilde{B}}^k(S(\tilde{E})_{+\tilde{B}}, [\tilde{F} \oplus \underline{H}]_{\tilde{B}}^+) . \quad (21)$$

associated with the projection

$$p : \tilde{B} = B \times S(H) \rightarrow B$$

(see [CJ] Proposition 5.37, Proposition 12.40 for the non-equivariant case).

A map $\kappa : B \rightarrow S(H)$ defines a section $j_\kappa^E : S(E)_{+B} \rightarrow [S(E) \times S(H)]_{+B}$ over B of the projection $[S(E) \times S(H)]_{+B} \rightarrow S(E)_{+B}$, so it defines a splitting of the exact sequence (4.2.1).

Lemma 4.9. *Let $m \in s^1 \alpha_B^{-1}([S(E) \times S(H)]_{+B}, [F \oplus \underline{H}]_B^+)$, and let $\kappa_0, \kappa_1 : B \rightarrow S(H)$ be two maps. One has the identity*

$$(j_{\kappa_1}^E)^*(m) - (j_{\kappa_0}^E)^*(m) = d(\kappa_0, \kappa_1) \cdot \partial(m) ,$$

where $d(\kappa_0, \kappa_1) \in s^1 \alpha_B^{-1}(B_{+B}, \underline{H}_B^+) = s^1 \alpha_B^{b-1}(B_{+B}, B_{+B})$ is the difference class of the maps κ_0, κ_1 regarded as sections in the sphere bundle $S(\underline{H})$.

Proof: The difference class $d(\kappa_0, \kappa_1)$ is defined by the map

$$\Delta : B_{+B} \wedge_B \underline{S}^1 = B \times [0, 1] /_{B} B \times \{0, 1\} \longrightarrow D_\epsilon(\underline{H}) / S_\epsilon(\underline{H}) = \underline{H}_B^+$$

induced by

$$(b, t) \mapsto \begin{cases} [(1-2t)\kappa_0(b)] & \text{for } 0 \leq t \leq \frac{1}{2} \\ [(2t-1)\kappa_1(b)] & \text{for } \frac{1}{2} \leq t \leq 1 . \end{cases}$$

The connecting morphism ∂_H in the long exact sequence

$$s^1 \alpha_B^{-1}(B_{+B}, \underline{H}_B^+) \xrightarrow{\partial_H} s^1 \alpha_B^0(B_{+B}, S(\underline{H})_{+B}) \rightarrow s^1 \alpha_B^0(B_{+B}, B_{+B}) \rightarrow s^1 \alpha_B^0(B_{+B}, \underline{H}_B^+)$$

is defined via the identifications

$$\begin{aligned} s^1 \alpha_B^{-1}(B_{+B}, \underline{H}_B^+) &= s^1 \alpha_B^0(B_{+B} \wedge_B \underline{S}^1, \underline{H}_B^+) \\ s^1 \alpha_B^0(B_{+B}, S(\underline{H})_{+B}) &= s^1 \alpha_B^0(B_{+B} \wedge_B \underline{S}^1, S(\underline{H})_{+B} \wedge_B \underline{S}^1) , \end{aligned}$$

by left composition with the contraction $\mathbf{c}_H : \underline{H}_B^+ \rightarrow S(\underline{H})_{+B} \wedge_B \underline{S}^1$. The image of $d(\kappa_0, \kappa_1)$ under ∂_H is just the difference $\{\kappa_1\} - \{\kappa_0\} \in s^1 \alpha_B^0(B_{+B}, S(\underline{H})_{+B})$.

One has obviously

$$(j_{\kappa_1}^E)^*(m) - (j_{\kappa_0}^E)^*(m) = m \circ (\{\kappa_1\} - \{\kappa_0\}) = m \circ \partial_H(d(\kappa_0, \kappa_1)) .$$

We know that $\partial_H(d(\kappa_0, \kappa_1))$ is represented by $\mathbf{c}_H \circ \Delta$ and the connecting operator ∂ in the exact sequence (4.2.1) acts by right composition with the same contraction \mathbf{c}_H . Therefore

$$\begin{aligned} (j_{\kappa_1}^E)^*(m) - (j_{\kappa_0}^E)^*(m) &= m \circ (\mathbf{c}_H \circ \Delta) = \partial(m) \circ d(\kappa_0, \kappa_1) = \partial(m) \circ (d(\kappa_0, \kappa_1) \cdot \{\text{id}_{B_{+B}}\}) \\ &= (d(\kappa_0, \kappa_1) \cdot \partial(m)) \circ \{\text{id}_{B_{+B}}\} = d(\kappa_0, \kappa_1) \cdot \partial(m) . \end{aligned}$$

Here we have used the fact that the composition multiplication \circ is $s^1 \alpha^*(B)$ -bilinear. \blacksquare

This lemma has an important analogue for the groups $\alpha^*(x)$ associated with a K-theory element x . For a compact space P we put

$$\alpha^*(P; x) = \varinjlim_{(E, F) \in x} s^1 \alpha_B^*(S(E)_{+B} \wedge_B \underline{P}_{+B}, F_B^+) .$$

where the inductive limit is taken with respect to the category $\mathcal{T}(x)$. Using the methods used in section 2.3 for the definition of the groups $\alpha^*(x)$, and the results in section 5.1, we see that this inductive limit exists; it can be constructed by taking first the limit of ${}_{S^1}\alpha_B^*(S(E \oplus \mathbb{C}^n)_{+B} \wedge_B \underline{P}_{+B}, [F \oplus \mathbb{C}^n]_B^+)$ over n , and factorizing the result by the action of $\tilde{J}(I[K^{-1}(B)]) \subset {}_{S^1}\alpha^0(B)$. The graded group $\alpha^*(P; x)$ comes with an obvious homomorphism $\alpha^*(P; x) \rightarrow \alpha^*(p_B^*(x))$, where $p_B : B \times P \rightarrow B$ is the projection on the first summand.

Taking the inductive limit of the connection morphisms $\partial = \partial_{E,F}$ in (4.2.1) with respect to the category $\mathcal{T}(x)$, one gets a morphism

$$\partial_x := \varinjlim_{(E,F) \in x} \partial_{E,F} : \alpha^{b-1}(S(H); x) \longrightarrow \alpha^0(x) . \quad (22)$$

which is intrinsically associated with x .

Let $\kappa : B \rightarrow S(H)$ be a fixed map. The system of morphisms

$$(j_\kappa^E)^* : {}_{S^1}\alpha_B^*([S(E) \times S(H)]_{+B}, F_B^+) \rightarrow {}_{S^1}\alpha_B^*(S(E)_{+B}, F_B^+)$$

induces a morphism $j_\kappa^* : \alpha^*(S(H); x) \rightarrow \alpha^*(x)$.

Corollary 4.10. *Let $m \in \alpha^{b-1}(S(H); x)$, and let $\kappa_0, \kappa_1 : B \rightarrow S(H)$ be two maps. One has the identity*

$$(j_{\kappa_1})^*(m) - (j_{\kappa_0})^*(m) = d(\kappa_0, \kappa_1) \cdot \partial_x(m) .$$

4.2.2. The universal perturbation and the invariant jump formulae. Let E, F be Hermitian vector bundles over a compact basis B , let V, W be Euclidean vector spaces, and let $\mu : E \times V \rightarrow [F \times W]_B^+$ be an S^1 -equivariant map over B satisfying the properties **P1** and **P2** (1) with $h = 0$. In other words,

$$\mu(0_y^E, v) = l(v) , \quad \forall y \in B \quad \forall v \in V ,$$

where $l : V \xrightarrow{\cong} W_0 \subset W$ is a linear embedding. The cylinder construction cannot be applied to such a map, because μ has vanishing points on the core $0^E \times D^R(V)$ of any cylinder $D_R(E) \times D_R(V)$. We orient the orthogonal complement H of W_0 in W , and we denote by b its dimension. Let $\epsilon > 0$. For every map $\kappa : B \rightarrow S_\epsilon(H)$ we define the perturbation

$$\mu_\kappa : E \times V \rightarrow [F \times W]_B^+$$

by putting $\mu_\kappa(e, v) := T_{\kappa(y)}(\mu(e, v))$ for $e \in E_y$. Here $T_{\kappa(y)}$ denotes the automorphism of $[F \times (H \oplus W_0)]_B^+$ which extends the translation

$$(f, w) \mapsto (f, w + \kappa(y)) .$$

Remark 4.11. *If $\epsilon > 0$ is sufficiently small, the map μ_κ satisfies the properties **P1**, **P2** of section 3.1, so the cylinder construction applies and yields a stable class $\{\mu_\kappa\} \in {}_{S^1}\alpha_B^{b-1}(S(E)_{+B}, F_B^+)$.*

Proof: Suppose that μ satisfies the property **P1** with constants C, c . Choose $\epsilon < \frac{c}{2}$. The map μ_κ satisfies **P1** with constants $C, c' := \frac{c}{2}$, and **P2** with constant $\epsilon_0 = \epsilon$. ■

Another way to construct a map satisfying properties **P1**, **P2** is to let κ vary in the sphere $S_\epsilon(H)$ and consider the *universal perturbation*

$$\tilde{\mu} : \tilde{E} \times V \longrightarrow \tilde{F} \times W$$

over the basis $\tilde{B} := B \times S_\varepsilon(H)$ (where $\tilde{E} := p_B^*(E)$, $\tilde{F} := p_B^*(F)$) which acts as μ_κ over $B \times \{\kappa\}$. This map also satisfies properties **P1**, **P2** with the same constants as any μ_κ , so that the cylinder construction applies and yields a class $\{\tilde{\mu}\} \in S^1 \alpha_{\tilde{B}}^{b-1}(S(\tilde{E})_{\tilde{B}}, \tilde{F}_{\tilde{B}}^+)$. Our next goal is to understand this class $\{\tilde{\mu}\}$. The essential point is to identify the image of $\{\tilde{\mu}\} \in S^1 \alpha_{\tilde{B}}^{b-1}(S(\tilde{E})_{+\tilde{B}}, \tilde{F}_{\tilde{B}}^+)$ under the connecting morphism ∂ .

Recall from section 2.6 that $\{o_{(E,F)}\} \in S^1 \alpha_B^0(S(E)_{+B}, F_B^+)$ is the class of the obvious pointed map $S(E)_{+B} \rightarrow F_B^+$ over B which maps $+_B$ to the infinity section, and $S(E)$ to the trivial section.

Proposition 4.12. *(The ∂ -image of the invariant of the universal perturbation)*
Via the identification

$$S^1 \alpha_B^0(S(E)_{+B} \wedge_B \underline{H}_B^+, [F \oplus \underline{H}]_B^+) = S^1 \alpha_B^0(S(E)_{+B}, F_B^+)$$

one has

$$\partial(\{\tilde{\mu}\}) = -\{o_{(E,F)}\} .$$

Proof: As in section 3.1 fix $R > C$ and $\varepsilon < \min(\varepsilon_0, c') = \min(\varepsilon, \frac{c'}{2})$. Let $\tau_0 < R$ be sufficiently small such that $\mu(e, v)$ remains finite for every $(e, v) \in D_{\tau_0}(E) \times D_R(V)$.

Step 1. We replace $\tilde{\mu}|_{D_R(\tilde{E}) \times D_R(V)}$ by a map $\tilde{\mu}_\tau$ which represents the same class $\{\mu\}$ and coincides with the κ -independent map μ outside the smaller cylinder $D_\tau(E) \times D_R(V)$.

Define $\tilde{\mu}_\tau : D_R(\tilde{E}) \times D_R(V) \rightarrow [\tilde{F} \times W]_{\tilde{B}}^+$ by the formula

$$\tilde{\mu}_\tau(e, \kappa, v) := \begin{cases} (1 - \frac{1}{\tau}\|e\|)(\kappa + l(v)) + \frac{1}{\tau}\|e\|\mu(e, v) & \text{for } 0 \leq \|e\| \leq \tau \\ \mu(e, v) & \text{for } \|e\| \geq \tau . \end{cases}$$

The maps $\tilde{\mu}_\tau$ and $\tilde{\mu}$ coincide on the core $0^{\tilde{E}} \times D_R(V)$ of the cylinder $D_R(\tilde{E}) \times D_R(V)$ and they differ by the translation T_κ outside $D_\tau(\tilde{E}) \times D_R(V)$. We define a homotopy between $\tilde{\mu}_\tau$ and $\tilde{\mu}|_{D_R(\tilde{E}) \times D_R(V)}$ by putting

$$\tilde{\mu}_\tau^t(e, \kappa, v) := \begin{cases} (1-t)\tilde{\mu}_\tau(e, \kappa, v) + t\tilde{\mu}(e, \kappa, v) & \text{for } \|e\| \leq \tau \\ T_{t\kappa} \circ \mu(e, v) & \text{for } \|e\| \geq \tau . \end{cases}$$

Claim: If τ is sufficiently small, then $\|\tilde{\mu}_\tau^t\| \geq c'$ on $\partial [D_R(\tilde{E}) \times D_R(V)]$ for every $t \in [0, 1]$.

The claim is not obvious only for points $(e, v) \in D_\tau(\tilde{E}) \times S_R(V)$. One has the identity

$$\begin{aligned} \tilde{\mu}_\tau^t(e, \kappa, v) &= (1-t) \left\{ \left(1 - \frac{1}{\tau}\|e\|\right) \kappa + l(v) + \frac{1}{\tau}\|e\| [\mu(e, v) - l(v)] \right\} + t\mu(e, v) + t\kappa = \\ &= l(v) + \left(1 - \frac{1-t}{\tau}\|e\|\right) \kappa + \left[t + \frac{(1-t)}{\tau}\|e\| \right] [\mu(e, v) - l(v)] . \end{aligned}$$

The first two terms belong to orthogonal complements, so for $e \in D_\tau(E)$ one has

$$\|\tilde{\mu}_\tau^t(e, \kappa, v)\| \geq \|l(v)\| - \|\mu(e, v) - l(v)\| .$$

Since $\mu(0_y^E, v) = l(v)$, and μ is fiberwise differentiable with globally continuous derivatives on $E \times V$, it holds

$$\lim_{\tau \rightarrow 0} \left\{ \sup \{ (\mu(e, v) - l(v)) \mid 0 \leq \|e\| \leq \tau, \|v\| \leq R \} \right\} = 0 .$$

On the other hand, for $\|v\| = R$ one has $\|l(v)\| = \|\mu(0_y^E, v)\| > c$. This proves the claim.

Using the Claim and $\|\tilde{\mu}_\tau^t(e, \kappa, v)\| = \|\kappa\| = \epsilon > 0$ we see that $(\tilde{\mu}_\tau^t)_{t \in [0,1]}$ defines a homotopy between $\tilde{\mu}_\tau$ and $\tilde{\mu}|_{D_R(\tilde{E}) \times D_R(V)}$ in the space of maps for which the cylinder construction applies. Therefore

$$\{\tilde{\mu}\} = \{\tilde{\mu}_\tau\} \in {}_{S^1} \alpha_B^{b-1}([S(E) \times S(H)]_{+B}, F_B^+) \text{ for all sufficiently small } \tau > 0. \quad (23)$$

Step 2. We compute the class $-\partial(\{\tilde{\mu}_\tau\})$.

Regard $\{\tilde{\mu}_\tau\}$ as an element in the group

$${}_{S^1} \alpha_B^{-1}([S(E) \times S(H)]_{+B}, [F \oplus \underline{H}]_B^+) = {}_{S^1} \alpha_B^0([S(E)_{+B} \wedge_B \underline{S}(H)_{+B} \wedge_B \underline{S}^1, [F \oplus \underline{H}]_B^+) .$$

As explained at the beginning of this section the morphism ∂ is given by composition with the contraction map

$$\mathbf{c}_H : H^+ = S_\epsilon(H) \times [0, R] / \sim_H \rightarrow S(H)_+ \wedge S^1 = S_\epsilon(H) \times [0, R] / S(H) \times \{0, R\}$$

induced by the identity of $S_\epsilon(H) \times [0, R]$. The morphism $-\partial$ is defined by composition with \mathbf{c}' , where \mathbf{c}' is induced by the map $(\kappa, \rho) \rightarrow (\kappa, R - \rho)$.

The class $\{\tilde{\mu}_\tau\}$ is represented by the map

$$\tilde{m}_\tau : S(E) \times S_\epsilon(H) \times [0, R] \times D_R(V) \longrightarrow [F \times W]_B^+ / [F \times W]_B^+ \setminus D_\epsilon(F \times W)$$

given by

$$\tilde{m}_\tau(e, \kappa, \rho, v) = [\tilde{\mu}_\tau(\rho e, \kappa, v)] .$$

As we have seen in section 3.1, this map induces a map

$$S(E)_{+B} \wedge_B \underline{S}(H)_{+B} \wedge_B \underline{S}^1 \wedge_B \underline{V}_B^+ \longrightarrow [F \times W]_B^+ / [F \times W]_B^+ \setminus D_\epsilon(F \times W)$$

because it has the following properties

- (1) $\tilde{m}_\tau(e, \kappa, 0, v)$ and $m_\tau(e, \kappa, R, v)$ belong always to the infinity section of the right hand space,
- (2) $\tilde{m}_\tau(e, \kappa, \rho, v)$ belongs to the infinity section of the right hand space when $\|v\| = R$.

The class $-\partial(\{\tilde{\mu}_\tau\})$ is defined by the map

$$\tilde{m}'_\tau : S(E) \times S_\epsilon(H) \times [0, R] \times D_R(V) \rightarrow [F \times W]_B^+ / [F \times W]_B^+ \setminus D_\epsilon(F \times W)$$

given by

$$\tilde{m}'_\tau(e, \kappa, \rho, v) = \tilde{m}_\tau(e, \kappa, R - \rho, v) .$$

This map descends to a map

$$S(E)_{+B} \wedge_B \underline{H}_B^+ \wedge_B \underline{V}_B^+ \rightarrow [F \times W]_B^+ / [F \times W]_B^+ \setminus D_\epsilon(F \times W)$$

because it has the following properties:

- (1) $\tilde{m}'_\tau(e, \kappa, 0, v)$ and $\tilde{m}'_\tau(e, \kappa, R, v)$ are independent of κ .
- (2) $\tilde{m}'_\tau(e, \kappa, R, v)$ belongs always to the infinity section of the right hand space.
- (3) $\tilde{m}'_\tau(e, \kappa, \rho, v)$ belongs to the infinity section of the right hand space when $\|v\| = R$.

These three conditions characterize the maps of pointed spaces over B defined on $S(E) \times S_\epsilon(H) \times [0, R] \times D_R(V)$ which descend to $S(E)_{+B} \wedge_B \underline{H}_B^+ \wedge_B \underline{V}_B^+$.

Step 2 (a). We deform the map \tilde{m}'_τ in the space of maps satisfying the three properties above, by composing it with a 1-parameter family of contractions in the ρ -direction.

For $t \in [0, 1]$ define the map

$$[\tilde{m}'_\tau]^t : S(E) \times S_\epsilon(H) \times [0, R] \times D_R(V) \rightarrow [F \times W]_B^+ /_B [F \times W]_B^+ \setminus D_\epsilon(F \times W)$$

by

$$[\tilde{m}'_\tau]^t(e, \kappa, \rho, v) = \tilde{m}_\tau \left(e, \kappa, (1-t + t\frac{\tau}{R})(R-\rho), v \right) .$$

The family $([\tilde{m}'_\tau]^t)_{t \in [0,1]}$ defines a homotopy in the space of maps satisfying properties (1), (2), (3) above. The main point in checking (1) is the fact that the map \tilde{m}_τ is constant with respect to κ for $\rho \in [\tau, R]$. Therefore it holds

$$-\partial(\{\tilde{\mu}_\tau\}) = \{[\tilde{m}'_\tau]^0\} = \{[\tilde{m}'_\tau]^1\} .$$

Putting $\tilde{m}''_\tau := [\tilde{m}'_\tau]^1$, one has

$$\begin{aligned} \tilde{m}''_\tau(e, \kappa, \rho, v) &= \tilde{m}_\tau(e, \kappa, \frac{\tau}{R}(R-\rho), v) = \left[1 - \frac{R-\rho}{R} \right] (\kappa + l(v)) + \frac{R-\rho}{R} \mu(\tau \frac{R-\rho}{R} e, v) \\ &= \frac{\rho}{R} \kappa + l(v) + \frac{R-\rho}{R} \left(\mu(\tau \frac{R-\rho}{R} e, v) - l(v) \right) . \end{aligned}$$

Step 2 (b). We remark that the family of maps \tilde{m}''_τ has a uniform limit as $\tau \rightarrow 0$ and we compute this limit explicitly .

Using arguments as in the proof of the claim above, we see that

$$\lim_{\tau \rightarrow 0} \frac{R-\rho}{R} \left(\mu(\tau \frac{R-\rho}{R} e, v) - l(v) \right) = 0$$

uniformly. Therefore $\tilde{m}'' := \lim_{\tau \rightarrow 0} \tilde{m}''_\tau$ operates by $\tilde{m}''(e, \kappa, \rho, v) = \frac{\rho}{R} \kappa + l(v)$. It is now easy to see that the map

$$S(E)_{+B} \wedge_B \underline{H}_B^+ \wedge_B \underline{V}_B^+ \rightarrow [F \times W]_B^+ /_B [F \times W]_B^+ \setminus D_\epsilon(F \times W) = F_B^+ \wedge_B \underline{H}_B^+ \wedge_B [\underline{W}_0]_B^+$$

induced by \tilde{m}'' is homotopic to the smash product over B of the obvious map $S(E)_{+B} \rightarrow F_B^+$ (which represents $o(E, F)$) with $l_B^+ : \underline{V}_B^+ \rightarrow [\underline{W}_0]_B^+$, and $\text{id} : \underline{H}_B^+ \rightarrow \underline{H}_B^+$. \blacksquare

For a map $\kappa : B \rightarrow S_\epsilon(H)$ one has

$$\{\mu_\kappa\} = (j_\kappa^E)^*(\{\tilde{\mu}\}) . \quad (24)$$

This formula shows that the individual invariant $\{\mu_\kappa\}$ associated with a map $\kappa : B \rightarrow S_\epsilon(H)$ is determined by the invariant associated with the universal perturbation $\tilde{\mu}$ and the homotopy class of κ . Using Corollary 4.10 we obtain

Corollary 4.13. *(Cohomotopy invariant jump formula) One has*

$$\{\mu_{\kappa_0}\} - \{\mu_{\kappa_1}\} = o_{(E,F)} \cdot d(\kappa_0, \kappa_1) ,$$

where $d(\kappa_0, \kappa_1) \in {}_{S^1}\alpha_B^{-1}(B_{+B}, \underline{H}_B^+)$ is the difference class of the maps κ_0, κ_1 regarded as sections in the sphere bundle $S_\epsilon(\underline{H})$.

Suppose now that $b = 1$. In this case $S_\epsilon(H)$ has two elements κ_0, κ_1 , and the difference class $d(\kappa_0, \kappa_1)$ is just the unit element of ${}_{S^1}\alpha_B^0(B_{+B}, B_{+B})$. Therefore, in this case, our result gives

Corollary 4.14. *(Cohomotopy wall crossing) Suppose $b = 1$. Then the two classes $\{\mu_{\kappa_0}\}, \{\mu_{\kappa_1}\}$ associated with the two perturbations $\mu_{\kappa_0}, \mu_{\kappa_1}$ of μ are related by the formula*

$$\{\mu_{\kappa_0}\} - \{\mu_{\kappa_1}\} = \{o(E, F)\} .$$

We can now extend our results to the infinite dimensional case. Let B be an oriented compact manifold, \mathcal{E}, \mathcal{F} complex Hilbert bundles over B , \mathcal{V}, \mathcal{W} real Hilbert spaces, and $\mu : \mathcal{E} \times \mathcal{V} \rightarrow \mathcal{F} \times \mathcal{W}$ an S^1 -equivariant, fiberwise differentiable map over B satisfying properties $\mathcal{P}1, \mathcal{P}3$ and $\mathcal{P}2$ (1) with $h = 0$. Then we have an orthogonal decomposition $\mathcal{W} = H \oplus \mathcal{W}_0$, and $\mu(0_y^{\mathcal{E}}, v) = l(v)$ for every $v \in \mathcal{V}$, where $l : \mathcal{V} \rightarrow \mathcal{W}_0$ is a linear isometry. We fix an orientation of the finite dimensional summand H . Defining in the same way as in the finite dimensional framework the universal perturbation $\tilde{\mu}$, one gets a stable class

$$\{\tilde{\mu}\} \in \alpha^*(S_\epsilon(H); x) ,$$

where $x \in K(B)$ is the index of the complex part of the fiberwise linearization of μ at the zero section. Recall that the Euler class $\gamma(x) \in \alpha^0(x)$ is defined by the system of stable classes $-\{o_{(E,F)}\} \in {}_{S^1}\alpha_B^0(S(E)_{+B}, F_B^+)$ defined by the obvious maps $S(E)_{+B} \rightarrow F_B^+$ (see section 2.6). Using the results obtained above and taking inductive limit over $\mathcal{T}(x)$, we obtain

Corollary 4.15. (1) *The image of $\{\tilde{\mu}\}$ under the morphism*

$$\partial_x : \alpha^{b-1}(S_\epsilon(H); x) \rightarrow \alpha^0(x)$$

is given by

$$\partial_x(\{\tilde{\mu}\}) = \gamma(x) .$$

(2) *Let $\kappa_0, \kappa_1 : B \rightarrow S(H)$ two maps. Then*

$$\{\mu_{\kappa_1}\} - \{\mu_{\kappa_0}\} = d(\kappa_0, \kappa_1) \cdot \gamma(x) .$$

(3) *Suppose $b = 1$ and write $S_\epsilon(H) = \{\kappa_0, \kappa_1\}$. Then*

$$\{\mu_{\kappa_1}\} - \{\mu_{\kappa_0}\} = \gamma(x) .$$

4.3. A product formula and a vanishing theorem. In this section we give the infinite dimensional analogue of the product formula proven in section 3.2.3.

Let $\mathcal{V}_i, \mathcal{W}_i$ be real Hilbert spaces, $\mathcal{E}_i, \mathcal{F}_i$ complex Hilbert bundles over a compact base B ($i = 1, 2$), and let $\mu_i : E_i \times V_i \rightarrow [F_i \times W_i]_B^+$ be S^1 -equivariant maps over B , satisfying the properties $\mathcal{P}1, \mathcal{P}2$ (1) and $\mathcal{P}3$ of section 3.3 with constants C, c . Let $\mathcal{W}_i = H_i \oplus \mathcal{W}_{0,i}$ be the corresponding orthogonal sum decompositions, $l_i : \mathcal{V}_i \xrightarrow{\cong} \mathcal{W}_{0,i}$ isometries, $x_i \in K(B)$ the K-theory elements defined by the corresponding families δ_i of Fredholm operators, and $h_i : B \rightarrow H_i$ the maps given by $\mathcal{P}2$ (1). We introduce the notations:

$\mathcal{V} := \mathcal{V}_1 \oplus \mathcal{V}_2, \mathcal{W} := \mathcal{W}_1 \oplus \mathcal{W}_2, H := H_1 \oplus H_2, \mathcal{W}_0 := \mathcal{W}_{0,1} \oplus \mathcal{W}_{0,2}, l := l_1 \oplus l_2$, and consider the Hilbert bundles $\mathcal{E} := \mathcal{E}_1 \oplus \mathcal{E}_2, \mathcal{F} := \mathcal{F}_1 \oplus \mathcal{F}_2$. The product map

$$\mu : \mathcal{E} \times \mathcal{V} = [\mathcal{E}_1 \times \mathcal{V}_1] \oplus [\mathcal{E}_2 \times \mathcal{V}_2] \longrightarrow [\mathcal{F} \times \mathcal{W}]_B^+ = [\mathcal{F}_1 \times \mathcal{W}_1]_B^+ \wedge_B [\mathcal{F}_2 \times \mathcal{W}_2]_B^+$$

also satisfies properties $\mathcal{P}1, \mathcal{P}2$ (1) (with associated map $h = (h_1, h_2) : B \rightarrow H$) and $\mathcal{P}3$; it satisfies $\mathcal{P}2$ (2) as soon as one of the two maps μ_1, μ_2 does.

Suppose that μ_1 satisfies property $\mathcal{P}2$ (2). In this case the construction of section 3.3 applies and yields an invariant

$$\{\mu_1\} \in \alpha^{b_1-1}(x_1).$$

The finite dimensional approximations of the map μ_2 define classes

$$\{(\mu_2)_{c,\pi_2}^+\} \in S^1 \alpha_B^{b_2}([E_2]_B^+, [F_2]_B^+).$$

It can be shown that a compatibility result similar to Proposition 3.10 holds, so that one obtains an invariant

$$\{\mu_2^+\} \in \alpha^{b_2}(x_2^+) := \varinjlim_{(E_2, F_2) \in x_2} S^1 \alpha_B^{b_2}([E_2]_B^+, [F_2]_B^+).$$

Here the inductive limit on the right is taken over the category $\mathcal{T}(x_2)$ and is constructed using the same methods as in the definition of the groups $\alpha^*(x)$ (see section 2.3). The direct limit of the obvious products

$$S^1 \alpha_B^{b_1-1}(S(E_1)_{+B}, [F_1]_B^+) \times S^1 \alpha_B^{b_2}([E_2]_B^+, [F_2]_B^+) \rightarrow$$

$S^1 \alpha_B^{b_1+b_2-1}(S(E_1)_{+B} \wedge_B [E_2]_{B_2}^+, [F_1 \oplus F_2]_B^+) \xrightarrow{1c^*} S^1 \alpha_B^{b_1+b_2-1}(S(E_1 \oplus E_2)_{+B}, [F_1 \oplus F_2]_B^+)$ gives a well defined product

$$\cdot : S^1 \alpha^{b_1-1}(x_1) \times \alpha^{b_2}(x_2^+) \rightarrow S^1 \alpha^{b_1+b_2-1}(x_1 + x_2).$$

Using finite dimensional approximations of μ of the form

$$\mu_{c,\pi_1 \times \pi_2} = (\mu_1)_{c,\pi_1} \times (\mu_2)_{c,\pi_2}$$

and applying Proposition 3.3 we obtain

Remark 4.16. *Under the assumptions and with the notations above, the invariant of the product map $\mu = \mu_1 \times \mu_2$ is given by the formula*

$$\{\mu_1 \times \mu_2\} = \{\mu_1\} \cdot \{\mu_2^+\}.$$

Note that in this formula the map μ_2 is allowed to have S^1 -invariant zeroes. In the case when both maps μ_i satisfy $\mathcal{P}2$ (2) (so they are nowhere zero on their S^1 -fixed point loci) one has the following important vanishing result for the Hurewicz image of the invariant associated with a product map:

Proposition 4.17. *Put $x := x_1 + x_2 \in K(B)$ and let $h_x : \alpha^*(x) \rightarrow H^*(x; \mathbb{Z})$ be the Hurewicz morphism associated with x . Suppose that both maps μ_i satisfy properties $\mathcal{P}1$, $\mathcal{P}2$ (1), $\mathcal{P}2$ (2) and $\mathcal{P}3$, and that B is a finite CW complex. Then*

$$h_x(\{\mu_1 \times \mu_2\}) = 0 .$$

Proof: Let $m_i := (\mu_i)_{c, \pi_i}$ be finite dimensional approximations of μ_i and put $m := m_1 \times m_2$. Applying the cylinder construction to this maps we get a representative

$$m_R : S(E_1 \oplus E_2)_{+B} \wedge_B [\mathbb{R} \oplus \underline{V}_1 \oplus \underline{V}_2]_B^+ \rightarrow [F_1 \oplus F_2 \oplus \underline{W}_1 \oplus \underline{W}_2]_B^+$$

of the class $\{\mu_1 \times \mu_2\}$. Put $E := E_1 \oplus E_2$, $F := F_1 \oplus F_2$, $V := V_1 \oplus V_2$, $W := W_1 \oplus W_2$, and $b = b_1 + b_2$. Let

$$\bar{m}_R : [\mathbb{R} \oplus \underline{V}]_{\mathbb{P}(E)}^+ \rightarrow [\tilde{F} \oplus \underline{W}]_{\mathbb{P}(E)}^+$$

be the associated sphere bundle map, constructed as in section 4.1.1. We denote by

$$p : [\mathbb{R} \oplus \underline{V}]_{\mathbb{P}(E)}^+ \rightarrow \mathbb{P}(E) , \quad q : [\tilde{F} \oplus \underline{W}]_{\mathbb{P}(E)}^+ \rightarrow \mathbb{P}(E)$$

the two bundle projections, and by $h := h_{\bar{m}_R} \in H^{2f+b_1+b_2-1}(\mathbb{P}(E); \mathbb{Z})$ the corresponding Hurewicz class, which is defined by the equality

$$(\bar{m}_R)^*(t_{\tilde{F} \oplus \underline{W}}) = p^*(h) \cup t_{\mathbb{R} \oplus \underline{V}} \quad (25)$$

in $H^*([\mathbb{R} \oplus \underline{V}]_{\mathbb{P}(E)}^+, \infty_{\mathbb{R} \oplus \underline{V}}; \mathbb{Z})$. Since both maps μ_i satisfy property **P2**, it follows that, for a sufficiently small neighborhood \mathcal{P} of $\mathbb{P}(E_1) \cup \mathbb{P}(E_2)$ in $\mathbb{P}(E)$, the map \bar{m}_R maps $p^{-1}(\mathcal{P})$ to the infinity section of the right hand bundle. We can suppose that \mathcal{P} is a standard compact neighborhood of this union, i.e. it has the form

$$\mathcal{P} = \mathbb{P}(E) \setminus \left\{ [e_1, e_2] \in \mathbb{P}(E) \mid e_i \neq 0, \ln \frac{\|e_1\|}{\|e_2\|} \in (-s, s) \right\}$$

for sufficiently large $s > 0$. The pull-back class $(\bar{m}_R)^*(t_{\tilde{F} \oplus \underline{W}})$ can be regarded as an element in $H^*([\mathbb{R} \oplus \underline{V}]_{\mathbb{P}(E)}^+, \infty_{\mathbb{R} \oplus \underline{V}} \cup p^{-1}(\mathcal{P}); \mathbb{Z})$, which can be identified with $H^{*-(\dim(V)+1)}(\mathbb{P}(E), \mathcal{P}; \mathbb{Z})$ via the relative Thom isomorphism over the pair $(\mathbb{P}(E), \mathcal{P})$. Therefore, the equality

$$(\bar{m}_R)^*(t_{\tilde{F} \oplus \underline{W}}) = p^*(h') \cup t_{\mathbb{R} \oplus \underline{V}} \quad (26)$$

in $H^*([\mathbb{R} \oplus \underline{V}]_{\mathbb{P}(E)}^+, \infty_{\mathbb{R} \oplus \underline{V}} \cup p^{-1}(\mathcal{P}); \mathbb{Z})$ defines a class $h' \in H^*(\mathbb{P}(E), \mathcal{P}; \mathbb{Z})$, and h is just the image of h' via the morphism $C^* : H^*(\mathbb{P}(E), \mathcal{P}; \mathbb{Z}) \rightarrow H^*(\mathbb{P}(E); \mathbb{Z})$ associated with the map $C : (\mathbb{P}(E), \emptyset) \rightarrow (\mathbb{P}(E), \mathcal{P})$. Put now

$$\mathbb{P}_0 := \mathbb{P}(E) \setminus (\mathbb{P}(E_1) \cup \mathbb{P}(E_2)) , \quad \mathcal{P}_0 := \mathcal{P} \setminus (\mathbb{P}(E_1) \cup \mathbb{P}(E_2)) ,$$

and denote by h'_0 the image of h' via the morphism $I^* : H^*(\mathbb{P}(E), \mathcal{P}; \mathbb{Z}) \rightarrow H^*(\mathbb{P}_0, \mathcal{P}_0; \mathbb{Z})$ defined by the map $I : (\mathbb{P}_0, \mathcal{P}_0) \rightarrow (\mathbb{P}(E), \mathcal{P})$. The main point in the proof of our proposition is that the restriction

$$\bar{m}_R|_{\mathbb{P}_0} : p^{-1}(\mathbb{P}_0) \rightarrow q^{-1}(\mathbb{P}_0) .$$

is equivariant with respect to the free S^1 -action $(\zeta, [e_1, e_2]) \mapsto [\zeta e_1, e_2]$ on \mathbb{P}_0 and the obvious lift of this action in the bundle $\tilde{F}|_{\mathbb{P}_0}$. This is just because μ is the product of two S^1 -equivariant maps μ_i . Therefore, $\bar{m}_R|_{\mathbb{P}_0}$ descends to a bundle map

$$[\bar{m}_R]_0 : p^{-1}(\mathbb{P}_0)/S^1 \longrightarrow q^{-1}(\mathbb{P}_0)/S^1$$

over $\mathbb{Q}_0 := \mathbb{P}_0/S^1$. The two sphere bundles above coincide with the fibrewise compactifications $[\underline{\mathbb{R}} \oplus \underline{\mathbb{V}}]_{\mathbb{Q}_0}^+$, $[\tilde{F}_0 \oplus \underline{W}]_{\mathbb{Q}_0}^+$, where \tilde{F}_0 is the S^1 -quotient of \tilde{F} , regarded as a bundle over \mathbb{Q}_0 . We denote by p_0, q_0 the corresponding bundle projections on \mathbb{Q}_0 . Put $\mathcal{Q} := \mathcal{P}/S^1$, $\mathcal{Q}_0 := \mathcal{Q} \cap \mathbb{Q}_0$. Using the relative Thom isomorphism over the pair $(\mathbb{Q}_0, \mathcal{Q}_0)$, it follows that the equality

$$[\tilde{n}_R]_0^*(t_{\tilde{F}_0 \oplus \underline{W}}) = p_0^*(k_0) \cup t_{\underline{\mathbb{R}} \oplus \underline{\mathbb{V}}}$$

defines a class $k_0 \in H^*(\mathbb{Q}_0, \mathcal{Q}_0; \mathbb{Z})$. Taking the pull-back of this equality via the projection $\Pi_0 : (\mathbb{P}_0, \mathcal{P}_0) \rightarrow (\mathbb{Q}_0, \mathcal{Q}_0)$, (and comparing the obtained formula with a similar equality satisfied by h'_0), we see that $\Pi_0^*(k_0) = h'_0$. Therefore

$$h = C^* \circ I^{*-1} \circ \Pi_0^*(k_0) = C^* \circ \Pi^* \circ [J^*]^{-1}(k_0) , \quad (27)$$

where

$$\Pi : (\mathbb{P}(E), \mathcal{P}) \rightarrow \left(\mathbb{P}(E)/_{S^1}, \mathcal{Q} \right) , \quad J : (\mathbb{Q}_0, \mathcal{Q}_0) \rightarrow \left(\mathbb{P}(E)/_{S^1}, \mathcal{Q} \right)$$

denote the obvious maps. In this formula we used the identity $J \circ \Pi_0 = \Pi \circ I$, and that the maps I, J induce isomorphisms in cohomology, by the excision theorem. The result follows now directly from Lemma 4.18 below. \blacksquare

Lemma 4.18. *The morphism*

$$U^* : H^* \left(\mathbb{P}(E)/_{S^1}, \mathcal{Q}; \mathbb{Z} \right) \longrightarrow H^*(\mathbb{P}(E); \mathbb{Z})$$

induced by the map $U := \Pi \circ C : (\mathbb{P}(E), \emptyset) \rightarrow \left(\mathbb{P}(E)/_{S^1}, \mathcal{Q} \right)$, vanishes.

Proof: By the excision and homotopy invariance theorem one has

$$H^* \left(\mathbb{P}(E)/_{S^1}, \mathcal{Q}; \mathbb{Z} \right) = H^* \left(\mathbb{P}(E)/_{S^1} \setminus \mathring{\mathcal{Q}}, \mathcal{Q} \setminus \mathring{\mathcal{Q}}; \mathbb{Z} \right) ,$$

where $\mathring{\mathcal{Q}}$ is the interior of \mathcal{Q} . One has a natural homeomorphism

$$\mathbb{P}(E)/_{S^1} \setminus \mathring{\mathcal{Q}} \cong [\mathbb{P}(E_1) \times_B \mathbb{P}(E_2)] \times [-s, s] , \quad [e_1, e_2] \mapsto \left([e_1], [e_2], \ln \frac{\|e_1\|}{\|e_2\|} \right) ,$$

and this homeomorphism identifies $\mathcal{Q} \setminus \mathring{\mathcal{Q}}$ with $[\mathbb{P}(E_1) \times \mathbb{P}(E_2)] \times \{-s, s\}$. Multiplication with the Thom class of the trivial bundle

$$\mathbb{P}(E_1) \times_B \mathbb{P}(E_2) \times (-s, s) \rightarrow \mathbb{P}(E_1) \times_B \mathbb{P}(E_2)$$

defines an isomorphism

$$H^i(\mathbb{P}(E_1) \times_B \mathbb{P}(E_2); \mathbb{Z}) \xrightarrow{\cong} H^{i+1} \left(\mathbb{P}(E)/_{S^1} \setminus \mathring{\mathcal{Q}}, \mathcal{Q} \setminus \mathring{\mathcal{Q}}; \mathbb{Z} \right) = H^{i+1} \left(\mathbb{P}(E)/_{S^1}, \mathcal{Q}; \mathbb{Z} \right) .$$

Step 1. When B is a point, the statement of the Lemma is obvious because in this case both spaces $\mathbb{P}(E_1) \times_B \mathbb{P}(E_2)$ and $\mathbb{P}(E)$ have trivial cohomology in odd dimensions.

Step 2. For a general basis, note that U induces a morphism of the Leray spectral sequences associated with the projections

$$\mathbb{P}(E) \longrightarrow B , \quad \left(\mathbb{P}(E)/_{S^1}, \mathcal{Q} \right) \longrightarrow B .$$

But the Leray spectral sequence for the relative cohomology of the pair $(\mathbb{P}(E)/_{S^1}, \mathcal{Q})$ can be identified with the spectral sequence for the cohomology with compact supports of $\mathbb{P}(E)/_{S^1} \setminus \mathcal{Q}$. It suffices to note that the induced spectral sequence morphism vanishes at the $E_1^{p,q}$ -level, by *Step 1*. ■

5. APPENDIX

5.1. Inductive limits of functors. We recall the following important

Definition 5.1. ([AM] p. 148) *A filtering category is category \mathcal{C} with the properties*

- F1. *For every pair (O, O') of objects, there exists an object O'' and morphisms $O \rightarrow O'', O' \rightarrow O''$.*
- F2. *For every two morphisms $u, v : O \rightarrow O'$ there exists an object O'' and a morphism $w : O' \rightarrow O''$ such that $w \circ u = w \circ v$.*

For *small* filtering categories one has the following basic fact:

Proposition 5.2. ([AM], p. 149-150) *Let \mathcal{A} be one of the categories $\mathcal{S}ets$, $\mathcal{A}b$ or $\mathcal{G}r$, and let \mathcal{C} be a filtering small category. Then any functor $F : \mathcal{C} \rightarrow \mathcal{A}$ has an inductive limit, which can be constructed in the classical way: one factorizes the disjoint union $\coprod_{O \in \text{Ob}(\mathcal{C})} F(O)$ by the equivalence relation*

$$(O, x) \sim (O', x') \text{ if } \exists u : O \rightarrow O'', u' : O' \rightarrow O'' \text{ with } F(u)(x) = F(u')(x'). \quad (28)$$

When $\mathcal{A} = \mathcal{A}b$ or $\mathcal{G}r$, one endows the obtained set of equivalence classes with the operation induced by the group operations on the summands $F(O)$ of the disjoint union.

We will say that \mathcal{C} is *weakly filtering* if it satisfies F1 and the following weak form of the axiom F2.

- $\tilde{F}2$. For every two morphisms $u, v : O \rightarrow O'$ there exists an object O'' and morphisms $w, z : O' \rightarrow O''$ such that $w \circ u = z \circ v$.

Lemma 5.3. *Suppose that \mathcal{C} is weakly filtering and small. Then the relation \sim defined in (28) is still an equivalence relation, and the conclusion of Proposition 5.2 holds for $\mathcal{A} = \mathcal{S}ets$.*

Proof: It suffices to check that \sim is transitive. Let $x \in F(O)$, $x' \in F(O')$, $x'' \in F(O'')$ with $x \sim x'$, $x' \sim x''$. Therefore there exists morphisms $u : O \rightarrow \hat{O}$, $u' : O' \rightarrow \hat{O}$, $v' : O' \rightarrow \tilde{O}$, $v'' : O'' \rightarrow \tilde{O}$ such that $F(u)(x) = F(u')(x')$ and $F(v')(x') = F(v'')(x'')$. By F1 there exists morphisms $\hat{w} : \hat{O} \rightarrow O_0$, $\tilde{w} : \tilde{O} \rightarrow O_0$. We apply $\tilde{F}2$ to the morphisms $\hat{w}u'$, $\tilde{w}v' : O' \rightarrow O_0$. We obtain morphisms \hat{z} , $\tilde{z} : O_0 \rightarrow O_1$ such that $\hat{z}\hat{w}u' = \tilde{z}\tilde{w}v'$. Therefore

$$\begin{aligned} F(\hat{z}\hat{w}u)(x) &= F(\hat{z}\hat{w})(F(u)(x)) = F(\hat{z}\hat{w})(F(u')(x')) = F(\hat{z}\hat{w}u')(x') = \\ &= F(\tilde{z}\tilde{w}v')(x') = F(\tilde{z}\tilde{w})(F(v')(x')) = F(\tilde{z}\tilde{w})(F(v'')(x'')) = F(\tilde{z}\tilde{w}v'')(x''), \end{aligned}$$

hence $x \sim x''$. ■

For $\mathcal{A} = \mathcal{A}b$ or \mathcal{G} one cannot endow the quotient of the disjoint union by this equivalence relation with a coherent group structure using only the weakly filtering condition.

Unfortunately, we will need inductive limits of functors defined on index categories which are not small. In this case the disjoint union considered in Remark

5.2 might not be a set. However, there exists a simple situation when the existence of an inductive limit is guaranteed:

Lemma 5.4. *Let \mathcal{C} be a weakly filtering category, $Q \in \text{Ob}(\mathcal{C})$ a fixed object and $F : \mathcal{C} \rightarrow \mathcal{A}$ a functor such that $F(u)$ is surjective for every morphism $u : Q \rightarrow O$.*

- (1) *Suppose $\mathcal{A} = \text{Sets}$.*
 - (a) *The relation on $F(Q)$ defined by*

$$y \approx y' \text{ if } \exists u, v : Q \rightarrow O \text{ such that } F(u)(y) = F(v)(y') \quad (29)$$
is an equivalence relation. Put $L := F(Q)/\approx$.
 - (b) *For any $O \in \text{Ob}(\mathcal{C})$ there exists a unique map $f_O : F(O) \rightarrow L$ defined by $f_O(x) = [y]$ for any pair $(x, y) \in F(O) \times F(Q)$ for which there exist morphisms $u : O \rightarrow \hat{O}$, $v : Q \rightarrow \hat{O}$ with $F(u)(x) = F(v)(y)$. The system $(f_O)_{O \in \text{Ob}(\mathcal{C})}$ is F -compatible (i.e. it holds $f_{O'} \circ F(w) = f_O$ for any morphism $w : O \rightarrow O'$).*
 - (c) *The system $(f_O)_{O \in \text{Ob}(\mathcal{C})}$ satisfies the universal property of the inductive limit, so the inductive limit of F exists and can be identified with L .*
- (2) *Suppose $\mathcal{A} = \text{Ab}$ or Gr .*
 - (a) *Let H be a smallest normal subgroup of $F(Q)$ which contains the elements $x'x^{-1}$ with $x \approx x'$. Put $L := F(Q)/H$.*
 - (b) *The system of morphism $(f_O : F(O) \rightarrow L)_{O \in \text{Ob}(\mathcal{C})}$ defined in a similar way as in (1) is F -compatible and satisfies the universal property of the inductive limit. Therefore the inductive limit of F exists and can be identified with L .*

Proof: (1) (a) is clear. For (b) we have to prove that the map f_O is well defined. Let $y \in F(Q)$, $y' \in F(Q)$, $u : O \rightarrow \hat{O}$, $v : Q \rightarrow \hat{O}$, $u' : O \rightarrow \hat{O}'$, and $v' : Q \rightarrow \hat{O}'$ such that $F(u)(x) = F(v)(y)$ and $F(u')(x) = F(v')(y)$. We can find an object \tilde{O} and morphisms $w : \hat{O} \rightarrow \tilde{O}$, $w' : \hat{O}' \rightarrow \tilde{O}$. Since \mathcal{C} is weakly filtering, the exist morphisms $z : \tilde{O} \rightarrow O_0$, $z' : \tilde{O} \rightarrow O_0$ such that $z w u = z' w' u'$. This implies

$$F(z w v)(y) = F(z w)(F(u)(x)) = F(z' w')(F(u')(x)) = F(z w' v')(y'),$$

so $y \approx y'$.

The F -compatibility of the system $(f_O)_{O \in \text{Ob}(\mathcal{C})}$ and the fact that this system satisfies the universal property of the inductive limit are easily verified.

(2) Follows easily from (1). ■

Definition 5.5. ([AM] p. 149) *Let \mathcal{N} , \mathcal{C} be categories. A functor $\Theta : \mathcal{N} \rightarrow \mathcal{C}$ is called*

- (1) *cofinal, if*
 - C1. *For any $O \in \text{Ob}(\mathcal{C})$ there exists $n \in \text{Ob}(\mathcal{N})$ and $u : O \rightarrow \Theta(n)$.*
 - C2. *For every $n \in \text{Ob}(\mathcal{N})$, $O \in \text{Ob}(\mathcal{C})$, and $u : \Theta(n) \rightarrow O$, there exists $m \in \text{Ob}(\mathcal{N})$, $\nu : n \rightarrow m$ and $v : O \rightarrow \Theta(m)$ such that $vu = \Theta(\nu)$.*
- (2) *cofinal in the sense of Artin-Mazur ([AM] p. 149), if*
 - C1. *holds,*
 - C2. *For every $O \in \text{Ob}(\mathcal{C})$, $n \in \text{Ob}(\mathcal{N})$ and $u, v : O \rightarrow \Theta(n)$, there exists a morphism $\mu : n \rightarrow m$ in \mathcal{N} such that $\Theta(\mu)u = \Theta(\mu)v$.*

Lemma 5.6. (1) *If \mathcal{N} is filtering and Θ is cofinal in the sense of Artin-Mazur, then Θ is cofinal and \mathcal{C} is filtering.*

- (2) If \mathcal{C} is filtering and Θ is cofinal, then Θ is cofinal in the sense of Artin-Mazur.
- (3) Suppose $\Theta : \mathcal{N} \rightarrow \mathcal{C}$ is cofinal, and \mathcal{N}, \mathcal{C} are both small and filtering. For any functor $F : \mathcal{C} \rightarrow \mathcal{A}$ (with $\mathcal{A} = \mathcal{S}ets, \mathcal{A}b$ or $\mathcal{G}r$) the canonical morphism

$$\varinjlim_{n \in \mathcal{O}b(\mathcal{N})} F(\Theta(n)) \rightarrow \varinjlim_{O \in \mathcal{O}b(\mathcal{C})} F(O)$$

is an isomorphism.

Proof: 1. Let $u : \Theta(n) \rightarrow O$ be a morphism. Using C1, we can find a morphism $w : O \rightarrow \Theta(m)$; since \mathcal{N} is filtering, we can find morphisms $\eta : n \rightarrow k, \kappa : m \rightarrow k$. Therefore, we get two morphisms $\Theta(\eta), \Theta(\kappa)wu : \Theta(n) \rightarrow \Theta(k)$. By $\tilde{\mathcal{C}}2$, there exists $\mu : k \rightarrow l$ such that $\Theta(\mu)\Theta(\eta) = \Theta(\mu)\Theta(\kappa)wu$. This shows $[\Theta(\mu\kappa)w]u = \Theta(\mu\eta)$, so C2 holds with $v = \Theta(\mu\kappa)w$ and $\nu = \mu\eta$. The fact that \mathcal{C} is filtering is stated in [AM] p. 149.

2. Let $u, v : O \rightarrow \Theta(n)$ be two morphisms. Since \mathcal{C} is filtering, there exists $w : \Theta(n) \rightarrow O'$ with $wu = vw$. By C2, we can find $m \in \mathcal{O}b(\mathcal{N}), \nu : n \rightarrow m$ and $v' : O' \rightarrow \Theta(n)$, such that $v'w = \Theta(\nu)$. We will have $\Theta(\nu)u = v'wu = v'vw = \Theta(\nu)v$, which proves $\tilde{\mathcal{C}}2$.

3. See Proposition 1.8 in [AM] p. 150. ■

Example 1. Let B be a compact space and let \mathcal{U}_B be the category of complex vector bundles over B . A morphism $U \rightarrow U'$ is a pair $u = (i, U_1)$ consisting of a bundle embedding $i : U \rightarrow U'$ and a complement U_1 of $i(U)$ in U' (see section 2.3). The category \mathcal{U}_B satisfies F1 but not F2, so it is *not* filtering. Let \mathcal{N} be category associated with the ordered set (\mathbb{N}, \leq) . Then the functor $\Theta : \mathcal{N} \rightarrow \mathcal{U}_B$ which associates to n the trivial bundle $\underline{\mathbb{C}}^n$ and to an inequality $n \leq m$ the standard morphism $\underline{\mathbb{C}}^n \rightarrow \underline{\mathbb{C}}^m$ is cofinal. This follows from the fact that any vector bundle on B possesses a complement. Note however that Θ is not cofinal in the sense of Artin-Mazur.

Example 2. For a category \mathcal{C} and an object $Q \in \mathcal{O}b(\mathcal{C})$ we will denote by \mathcal{C}_Q the category whose objects are morphism $u : Q \rightarrow O$ and whose morphisms are

$$\text{Hom}(Q \xrightarrow{u} O, Q \xrightarrow{v} O') := \{w : O \rightarrow O' \mid w \circ u = v\}.$$

A morphism $u : Q \rightarrow Q'$ induces in an obvious way a pull-back functor $u^* : \mathcal{C}_{Q'} \rightarrow \mathcal{C}_Q$. If \mathcal{C} is filtering then \mathcal{C}_Q is filtering and the target functor functor $T : \mathcal{C}_Q \rightarrow \mathcal{C}$ is both cofinal and cofinal in the sense of Artin-Mazur.

Definition 5.7. A category with automorphism push-forward is a pair (\mathcal{U}, A) , where \mathcal{U} is a category and $A : \mathcal{U} \rightarrow \mathcal{G}r$ a functor, such that

F1. holds in \mathcal{U} .

S1. $A(O) = \text{Aut}(O)$ for every $O \in \mathcal{O}b(\mathcal{C})$.

S2. For any $u : O \rightarrow O'$ and $a \in \text{Aut}(O)$ one has $A(u)(a) \circ u = u \circ a$

S3. For every two morphisms $u, v : O \rightarrow O'$ in \mathcal{U} there exists an object O'' , a morphism $w : O' \rightarrow O''$ and an automorphism $a \in A(O'')$ such that $a \circ w \circ u = w \circ v$.

Note that when (\mathcal{U}, A) is a category with automorphism push-forward, then \mathcal{U} is weakly filtering (use S3).

Example 3. Defining the automorphism push-forward functors in the obvious way, the categories $\mathcal{U}_B, \mathcal{C}_B, \mathcal{T}(x)$ introduced in this article become categories with automorphism push-forward.

Let (\mathcal{U}, A) be a category with automorphism push-forward, $Q \in \mathcal{Ob}(\mathcal{U})$ a fixed object, and $F : \mathcal{U} \rightarrow \mathcal{Ab}$ a functor such that $F(u)$ is an isomorphism for any morphism $u : Q \rightarrow O$. We know by Lemma 5.4 that the inductive limit of F exists and is a quotient of $F(Q)$. We need an explicit description of this quotient. For every object $u : Q \rightarrow O$ in the category \mathcal{U}_Q the group $A(T(u))$ acts on $F(Q)$ via the isomorphism $F(u) : F(Q) \rightarrow F(T(u))$. A morphism $w : T(u) \rightarrow T(v)$ can be regarded as an element in $\text{Hom}_{\mathcal{U}_Q}(u, v)$ and defines a group morphism $A(w) : A(T(u)) \rightarrow A(T(v))$ which intertwines the actions of these groups on $G(Q)$.

Proposition 5.8. *Let (\mathcal{U}, A) be a category with automorphism push-forward, $Q \in \mathcal{Ob}(\mathcal{U})$ a fixed object, and $F : \mathcal{U} \rightarrow \mathcal{Ab}$ a functor such that $F(u)$ is an isomorphism for any $u : Q \rightarrow O$. Let \mathcal{N} be a small filtering category and $\Theta : \mathcal{N} \rightarrow \mathcal{U}_Q$ a functor satisfying the cofinality axiom C1. Put*

$$\mathbb{A} := \varinjlim_{n \in \mathcal{Ob}(\mathcal{N})} A(T(\Theta(n))) .$$

Then \mathbb{A} acts on $F(Q)$ in a natural way, the inductive limit $\varinjlim_{O \in \mathcal{Ob}(\mathcal{U})} F(O)$ exists and can be identified with the quotient $F(Q)/I[\mathbb{A}]F(Q)$.

Proof: By Lemma 5.4 the inductive limit of G exists and can be identified with a quotient $F(Q)/H$. Here H is the group generated by the elements of the form $x - x'$ where $x, x' \in F(Q)$ are such that there exists $u, u' : Q \rightarrow O$ with $F(u)(x) = F(u')(x')$. We claim that the set of such pairs (x, x') coincides with the set of pairs of the form $(\mathbf{a}x', x')$ with $x' \in F(Q)$, $\mathbf{a} \in \mathbb{A}$.

Indeed, if $F(u)(x) = F(u')(x')$, choose $v : O \rightarrow \hat{O}$ and $a \in A(\hat{O})$ such that $vu' = avu$. The morphism vu can be regarded as an object in the category \mathcal{U}_Q . Since Θ satisfies the axiom C1, there exists $n \in \mathcal{Ob}(\mathcal{N})$ and a morphism $vu \rightarrow \Theta(n)$ in \mathcal{U}_Q , i.e. a morphism $w : \hat{O} \rightarrow T(\Theta(n))$ such that $wvu = \Theta(n)$. We obtain

$$\begin{aligned} F(\Theta(n))(x) &= F(wvu)(x) = F(wvu')(x') = F(wavu)(x') = F(A(w)(a)wvu)(x') = \\ &= A(w)(a)(F(wvu)(x')) = A(w)(a)(F(\Theta(n))(x')) , \end{aligned}$$

which shows that $x = \mathbf{a}x'$, where \mathbf{a} is the class of $A(w)(a) \in A(T(\Theta(n)))$ in \mathbb{A} . Conversely let $\mathbf{a} = [a] \in \mathbb{A}$ be represented by $a \in A(T(\Theta(n)))$ and suppose that $x = \mathbf{a}x'$. This means $F(\Theta(n))(x) = a(F(\Theta(n))(x'))$ so, putting $u := \Theta(n)$, $u' := a\Theta(n)$ one has $F(u)(x) = F(u')(x')$. \blacksquare

Let (\mathcal{U}, A) be a category with automorphism push-forward, and let $G : \mathcal{C} \rightarrow \mathcal{A}$ be a functor, where \mathcal{A} is one of the categories *Sets*, *Gr* or *Ab*.

Definition 5.9. *We say that the stabilized automorphisms act trivially on G if*

TSA. For every $O \in \mathcal{Ob}(\mathcal{C})$, $x \in G(O)$ and $a \in A(O)$ there exists a morphism $u : O \rightarrow O'$ such that $G(u)(G(a)(x)) = G(u)(x)$.

In the presence of functor $\Theta : \mathcal{N} \rightarrow \mathcal{U}$, we say that the Θ -stabilized automorphisms act trivially on G if

OSA. For every $n \in \mathcal{Ob}(\mathcal{N})$, $x \in G(\Theta(n))$ and $a \in A(\Theta(n))$ there exists a morphism $\nu : n \rightarrow m$ such that $G(\Theta(\nu))(G(a)(x)) = G(\Theta(\nu))(x)$.

Remark 5.10. *If Θ is cofinal and G satisfies ΘSA , then it also satisfies TSA . If \mathcal{C} is filtering, then any functor $G : \mathcal{C} \rightarrow \mathcal{A}$ satisfies TSA . If, moreover, Θ is cofinal, then G also satisfies ΘSA .*

Let (\mathcal{U}, A) be a category with automorphism push-forward, and let $G : \mathcal{U} \rightarrow \mathcal{A}$ be a functor. Let \mathcal{N} be a *small filtering* category and $\Theta : \mathcal{N} \rightarrow \mathcal{U}$ a cofinal functor such that ΘSA holds. Consider the classical inductive limit $L_\Theta := \varinjlim_{n \in \mathcal{Ob}(\mathcal{N})} G(\Theta(n))$.

For every $O \in \mathcal{Ob}(\mathcal{U})$ we define a morphism $f_O : G(O) \rightarrow L_\Theta$ by $f_O(x) := [G(v)(x)]$ where $v : O \rightarrow \Theta(n)$ is a morphism (whose existence is guaranteed by C1).

Proposition 5.11. *Under the assumptions and with the notations above it holds*

- (1) *For any $O \in \mathcal{Ob}(\mathcal{U})$ the map f_O is well defined. The system of maps $(f_O)_{O \in \mathcal{Ob}(\mathcal{U})}$ is G -compatible i.e. for any $u : O \rightarrow O'$ one has $f_{O'} \circ G(u) = f_O$. When $\mathcal{A} = \mathcal{Ab}$ or \mathcal{Gr} , the map f_O is a group morphism.*
- (2) *The system $(f_O)_{O \in \mathcal{Ob}(\mathcal{U})}$ satisfies the universal property of the inductive limit, therefore the functor G admits an inductive limit in \mathcal{A} which can be identified with L_Θ .*

We agree to write $u(x), v(x) \dots$, instead of $G(u)(x), G(v)(x) \dots$, to save on notations.

Proof: 1. Let $v : O \rightarrow \Theta(n), v' : O \rightarrow \Theta(n')$ be two morphisms. Since \mathcal{N} is filtering, there exists morphisms $\nu : n \rightarrow m, \nu' : n' \rightarrow m$. Applying axiom S3 to the morphisms $\Theta(\nu)v, \Theta(\nu')v'$, we get a morphism $w : \Theta(m) \rightarrow \hat{O}$ and an automorphism $a \in A(\hat{O})$ such that $w\Theta(\nu')v' = aw\Theta(\nu)v$. Now we apply the axiom C2 to w and we get morphisms $u : \hat{O} \rightarrow \Theta(k), \mu : m \rightarrow k$ such that $uw = \Theta(\mu)$. We have

$$\Theta(\mu\nu')v' = uw\Theta(\nu')v' = uaw\Theta(\nu)v = A(u)(a)uw\Theta(\nu)v = A(u)(a)\Theta(\mu\nu)v .$$

Using the axiom ΘSA we obtain a morphism $\eta : k \rightarrow l$ such that

$$\Theta(\eta) [A(u)(a)\Theta(\mu\nu)v(x)] = \Theta(\eta) [\Theta(\mu\nu)v(x)] .$$

Therefore $\Theta(\eta\mu\nu')(v'(x)) = \Theta(\eta\mu\nu)(v(x))$, which shows that $v(x)$ and $v'(x')$ define the same element in L_Θ . The second and the third claim are obvious.

2. Let $\Lambda \in \mathcal{Ob}(\mathcal{A})$ and $(g_O)_{O \in \mathcal{Ob}(\mathcal{U})}, g_O : G(O) \rightarrow \Lambda$ a system of G -compatible morphisms. Using the system $(g_{\Theta(n)})_{n \in \mathcal{Ob}(\mathcal{N})}$ (which is $G \circ \Theta$ -compatible) we get a unique morphism $g : L_\Theta \rightarrow \Lambda$ such that $g \circ c_n = g_{\Theta(n)}$ for every $n \in \mathcal{Ob}(\mathcal{N})$, where $c_n : G(\Theta(n)) \rightarrow L_\Theta$ is the canonical morphism. It remains to prove that $g \circ f_O = g_O$ for every $O \in \mathcal{Ob}(\mathcal{U})$. Let $x \in G(O)$ and choose $v : G(O) \rightarrow \Theta(n)$. One has

$$g \circ f_O(x) = g(c_n(v(x))) = g_{\Theta(n)}(v(x)) = g_O(x) .$$

■

5.2. Bundle maps between pointed sphere bundles. Let X be a CW complex and $Y \subset X$ a subcomplex. For two sections s', s'' in an oriented r -sphere bundle over X which coincide over Y , we denote by $o_Y(s', s'') \in H^r(X, Y; \mathbb{Z})$ the *primary obstruction* to the existence of a homotopy between s' and s'' in the space of sections which coincide with $s'|_Y = s''|_Y$ on Y [S].

Let $\pi_\zeta : \zeta \rightarrow \mathcal{B}$ be an oriented real bundle of rank r over a CW complex \mathcal{B} . Denote by $\pi_\zeta^+ : \zeta_B^+ =: \hat{\mathcal{B}} \rightarrow \mathcal{B}$ the bundle projection of the associated sphere bundle, and consider the pull-back bundle $\hat{\zeta} := [\pi_\zeta^+]^*(\zeta)$ on $\hat{\mathcal{B}}$. The sphere bundle $\hat{\zeta}_B^+ = [\pi_\zeta^+]^*(\zeta_B^+)$ comes with a tautological section θ_ζ and an ‘‘infinite’’ section s_ζ^∞ . These sections coincide on the subspace $\infty_\zeta \subset \hat{\mathcal{B}}$. We endow the space $\hat{\mathcal{B}}$ with a CW structure in the following way: First, on the subspace ∞_ζ we copy the CW structure from the base \mathcal{B} via s_ζ^∞ . Second, for every k -cell $e \subset \mathcal{B}$ we put $\hat{e} := \pi_\zeta^{-1}(e)$. The attaching map corresponding to \hat{e} is defined in the following way: let $u : D^k \rightarrow \bar{e} \subset \mathcal{B}$ the attaching map of e . The pullback bundle $u^*(\zeta)$ is trivial, so it can be identified with $D^k \times \mathbb{R}^r = D^k \times \mathring{D}^r$. The induced map $D^k \times \mathring{D}^r \rightarrow \pi_\zeta^{-1}(\bar{e}) \subset \zeta$ can be extended to map $\hat{u} : D^k \times D^r \rightarrow [\pi_\zeta^+]^{-1}(\bar{e}) \subset \zeta^+$ in an obvious way. Let t_ζ be the Thom class of the bundle ζ . We claim

Lemma 5.12. *With respect to such a cellular structure on $\hat{\mathcal{B}}$ one has $o_{\infty_\zeta}(s_\zeta^\infty, \theta_\zeta) = t_\zeta$ in $H^r(\hat{\mathcal{B}}, \infty_\zeta; \mathbb{Z})$.*

Proof: Let $P : \mathbb{E} \rightarrow \mathbb{B} := BSO(r)$ be the universal vector bundle with structure group $SO(r)$ and a fixed CW structure on the classifying space \mathbb{B} . Since $H^r(\mathbb{B}, \infty_{\mathbb{E}}; \mathbb{Z}) \simeq H^0(\mathbb{B}; \mathbb{Z}) \simeq \mathbb{Z}$, there exists an integer N such that $o_{\infty_{\mathbb{E}}}(s_{\mathbb{E}}^\infty, \theta_{\mathbb{E}}) = Nt_{\mathbb{E}}$. Let $f : \mathcal{B} \rightarrow \mathbb{B}$ a cellular map which induces the bundle ζ . This map is covered by a bundle map $\hat{f} : \hat{\mathcal{B}} \rightarrow \hat{\mathbb{B}}$, which is obviously cellular and maps the subcomplex ∞_ζ of $\hat{\mathcal{B}}$ into the subcomplex $\infty_{\mathbb{E}}$ of $\hat{\mathbb{B}}$. Using the functorial properties of the relative obstruction class and of the Thom class, we obtain $o_{\infty_\zeta}(s_\zeta^\infty, \theta_\zeta) = Nt_\zeta$. The integer N can be computed using any bundle ζ , so we will choose the bundle $\mathbb{R}^r \rightarrow \{*\}$. The tautological section is just the identity of $[\mathbb{R}^r]^+$. It’s easy to see that both classes can be identified with the generator of $H^r([\mathbb{R}^r]^+, \infty; \mathbb{Z})$. ■

Corollary 5.13. *Let ζ be an oriented r -bundle over a CW complex \mathcal{B} , and let s be a section in ζ_B^+ which coincides with s_ζ^∞ on a subcomplex $\mathcal{A} \subset \mathcal{B}$. Then $o_{\mathcal{A}}(s_\zeta^\infty, s) = s^*(t_\zeta)$ in $H^r(\mathcal{B}, \mathcal{A}; \mathbb{Z})$.*

Proof: Note that, with respect to the cellular decomposition of $\hat{\mathcal{B}}$ considered above, the section $s : \mathcal{B} \rightarrow \hat{\mathcal{B}}$ is a cellular map and maps to subcomplex \mathcal{A} into the subcomplex ∞_ζ . It suffices to apply the functorial property of the relative obstruction classes with respect to cellular maps. ■

Corollary 5.14. *Let ζ be an oriented r -bundle over a finite CW complex \mathcal{B} of dimension $n \leq r$, and let $\mathcal{A} \subset \mathcal{B}$ be a subcomplex. Then the map $o_{\mathcal{A}} : s \mapsto s^*(t_\zeta)$ defines a bijection between the set $\Gamma_{\mathcal{A}}(\zeta_B^+)$ of homotopy classes of sections in ζ_B^+ which coincide with s_ζ^∞ on \mathcal{A} , and $H^r(\mathcal{B}, \mathcal{A}; \mathbb{Z})$.*

Proof: Injectivity: Since $\dim(\mathcal{B}) \leq r$, for a section $s \in \Gamma_{\mathcal{A}}(\zeta_B^+)$ the only obstruction to the existence of a homotopy between s_ζ^∞ and s is the primary obstruction $o_{\mathcal{A}}(s_\zeta^\infty, s)$. To prove surjectivity, consider, for any r -cell $e \subset \mathcal{B} \setminus \mathcal{A}$, a section s_e which coincides with s_ζ^∞ on $\bar{e} \setminus e$ and has a single vanishing point, which is non-degenerate. The pull-back $s_e^*(t_\zeta)$ is a generator of $H^r(\mathcal{B}, \mathcal{B} \setminus e; \mathbb{Z}) \cong \mathbb{Z}$. ■

Corollary 5.15. *Let ζ_0, ζ_1 be two oriented bundles of ranks r_0, r_1 over an n -dimensional complex \mathcal{B} .*

- (1) If $n+r_0 < r_1$, any pointed bundle map $f : [\zeta_0]_{\mathcal{B}}^+ \rightarrow [\zeta_1]_{\mathcal{B}}^+$ over \mathcal{B} is homotopic (in the space of pointed bundle maps over \mathcal{B}) to the fiberwise constant map f^∞ which maps $[\zeta_0]^+$ into ∞_{ζ_1} .
- (2) If $n+r_0 = r_1$, then a pointed bundle map $f : [\zeta_0]_{\mathcal{B}}^+ \rightarrow [\zeta_1]_{\mathcal{B}}^+$ over \mathcal{B} is homotopic to f^∞ if and only if the class $h_f \in H^n(\mathcal{B}; \mathbb{Z})$, defined by the condition $f^*(t_{\zeta_1}) = [\pi_{\zeta_0}^+]^*(h_f) \cup t_{\zeta_0}$, vanishes. Moreover, the assignment $f \mapsto h_f$ defines a bijection between the set of homotopy classes of pointed bundle maps $[\zeta_0]_{\mathcal{B}}^+ \rightarrow [\zeta_1]_{\mathcal{B}}^+$ and $H^n(\mathcal{B}; \mathbb{Z})$.

Proof: It suffices to apply Corollary 5.14 to the pull-back bundle $\tilde{\zeta}_1 := [\pi_{\zeta_0}^+]^*(\zeta_1)$ over $\tilde{\mathcal{B}} := [\zeta_0]_{\mathcal{B}}^+$ and to identify the space of pointed bundle maps $[\zeta_0]_{\mathcal{B}}^+ \rightarrow [\zeta_1]_{\mathcal{B}}^+$ with the space of those sections in $[\tilde{\zeta}_1]_{\tilde{\mathcal{B}}}^+$ which coincide with $s_{\tilde{\zeta}_1}^\infty$ on $\infty_{\zeta_0} \subset \tilde{\mathcal{B}}$. Then use the Thom isomorphism $\cdot \cup t_{\zeta_0} : H^n(\mathcal{B}; \mathbb{Z}) \rightarrow H^{r_1}(\tilde{\mathcal{B}}, \infty_{\zeta_0}; \mathbb{Z})$. ■

REFERENCES

- [AM] Artin, M.; Mazur, B.: *Etale homotopy*, Lecture Notes in Math. 100, Springer Verlag, 1969.
- [B] Bader, M.: Cohomotopy invariants in gauge theoretic Gromov-Witten theory, PhD Thesis, Zürich, 2007, in preparation.
- [B1] Bauer, S.: *A stable cohomotopy refinement of Seiberg-Witten invariants: II*, Invent. math 155, 21-40, 2004.
- [B2] Bauer, S.: *Refined Seiberg-Witten invariants*, Different faces of Geometry, 1-46, Int. Math. Ser. (N. Y.), Kluwer/Plenum, New York, 2004.
- [B3] Bauer, S.: *On connected sums of four dimensional manifolds*, Preprintreihe 2000, Univ. Bielefeld, <http://www.mathematik.uni-bielefeld.de/sfb343/preprints/pr00001.ps.gz>.
- [BF] Bauer, S.; Furuta, M.: *A stable cohomotopy refinement of Seiberg-Witten invariants: I*, Invent. math 155, 1-19, 2004.
- [Br] Brussee, R.: *The canonical class and the C^∞ -properties of Kahler surfaces*, New York J. Math. 2, 103-146, 1996.
- [C] Crabb, M.: *The fibrewise Leray-Schauder index*, J. fixed point theory appl. 1, 3-30, 2007.
- [CJ] Crabb, M.; James, I.: *Fiberwise homotopy theory*, Springer Verlag, 1998.
- [CK] Crabb, M.; Knapp, K.: *On the codegree of negative multiplies of the Hopf bundle*, Proc. of the Royal Soc. of Edinburgh, 107A, 87-107, 1987.
- [CS] Crabb, M.; Sutherland.: *The space of sections of a sphere bundle I*, Proc. Edinburgh Math. Soc. 29, 383-403, 1986.
- [tD] tom Dieck, T.: *Transformation Groups*, De Gruyter, 1987.
- [Fu1] Furuta, M.: *Monopole equations and the 11/8 conjecture*, Math. Res. Lett. 8, 279-291, 2001.
- [Fu2] Furuta, M.: *Stable homotopy version of Seiberg-Witten invariant*, preprint, MPI Bonn, 1997, <http://www.mpim-bonn.mpg.de>.
- [H] Hauschild, H.: *Zerspaltung äquivarianter Homotopiemengen*, Math. Ann. 230, 279-292, 1977.
- [OT] Okonek, Ch.; Teleman, A. : *Seiberg-Witten invariants for manifolds with $b_+ = 1$ and the universal wall crossing formula*, Internat. J. Math. 7, no. 6, 811-832, 1996.
- [S] Steenrod, N.: *The topology of Fibre Bundles*, Princeton University Press, 1951.

Christian Okonek:

Institut für Mathematik, Universität Zürich, Winterthurerstrasse 150, CH-8057 Zürich,
e-mail: okonek@math.unizh.ch

Andrei Teleman:

CMI, Université de Provence, 39 Rue F. Joliot-Curie, F-13453 Marseille Cedex 13, e-mail:
teleman@cmi.univ-mrs.fr