



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2007

---

## **Aperiodic colorings and tilings of Coxeter groups**

Dranishnikov, A ; Schroeder, Viktor

**Abstract:** We construct a limit aperiodic coloring of hyperbolic groups. Also we construct limit strongly aperiodic strictly balanced tilings of the Davis complex for all Coxeter groups.

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-21531>

Journal Article

Accepted Version

Originally published at:

Dranishnikov, A; Schroeder, Viktor (2007). Aperiodic colorings and tilings of Coxeter groups. *Groups, Geometry, and Dynamics*, 1(3):311-328.

# APERIODIC COLORINGS AND TILINGS OF COXETER GROUPS

ALEXANDER DRANISHNIKOV AND VIKTOR SCHROEDER

ABSTRACT. We construct a limit aperiodic coloring of hyperbolic groups. Also we construct limit strongly aperiodic strictly balanced tilings of the Davis complex for all Coxeter groups.

## §0 INTRODUCTION

In [BDS] we constructed a quasi-isometric embedding of hyperbolic groups into a finite product of binary trees. First we implemented such construction for hyperbolic Coxeter groups [DS]. As a byproduct we obtained aperiodic tilings with finitely many tiles of the Davis complex for these groups. Our tilings are limit strongly aperiodic and the set of tiles can be taken to be aperiodic. As result we obtain limit strongly aperiodic tilings of the hyperbolic spaces  $\mathbf{H}^n$  which admits cocompact reflection groups. Thus 2-dimensional examples come from regular  $p$ -gons,  $p \geq 5$  in the hyperbolic plane. In dimension 3 there exists a right-angled regular hyperbolic dodecahedron ([A]). In dimension 4 there exists a right-angled hyperbolic 120-cell ([C],[D2],[PV]). The examples exist up to dimension 8 [VS]. Of course the dimension of the hyperbolic spaces is limited by Vinberg's theorem ( $\leq 29$ ) [V]. Existence of aperiodic sets of tiles for these cases also follows from results of Block and Weinberger [BW]. A new part of our results is the limit strong aperiodicity of tilings. Also the Block-Weinberger aperiodic tilings are unbalanced. In this paper we construct strongly balanced tilings which are limit strongly aperiodic. A strongly aperiodic tiling of  $\mathbf{H}^2$  was recently constructed by Goodman-Strauss [GS] (his tiling has even finite strongly aperiodic set of tiles).

First we obtain our tilings of the Davis complex as a tiling by color with geometrically the same tile (the chamber). Then we take a geometric resolution of the tiling by color. If

---

1991 *Mathematics Subject Classification.* Primary 20H15.

*Key words and phrases.* aperiodic coloring, aperiodic tiling, Coxeter group, Davis complex.

The authors were partially supported by NSF grants DMS-0604494 and SNF grant 200020-103594

a discrete group  $G$  acts by isometries properly cocompactly on a metric space  $X$ , there is an universal way to construct an aperiodic tiling of  $X$  by means of an aperiodic coloring of  $G$ . We consider an orbit  $Gx$  and consider the Voronoi cells  $V_{gx}$  for  $g \in G$  where

$$V_y = \{x \in X \mid d(y, x) \leq d(y, Gx)\}.$$

Clearly, all cells are isometric to each other and an aperiodic coloring of  $G$  defines an aperiodic tiling of  $X$  by color. In the case of Coxeter groups one can consider colorings of the walls instead of groups. This allows us to construct strictly balanced aperiodic tiling.

This paper is arranged as follows. First we consider coloring of discrete groups (§1). Then we extend this to a coloring of spaces, in particular trees, on which the group acts (§2). Then we apply this to the case of trees of walls in a Coxeter group (§3). Finally we construct a strictly balanced limit aperiodic tilings (§5). In (§4) we give a brief account of topology on the space of tilings.

It is a pleasure to thank Victor Bangert and Mark Sapir for useful discussion about the Morse-Thue sequence. Also we would like to thank the Max-Planck Institut für Mathematik in Bonn for its hospitality.

## §1 APERIODIC COLORING OF GROUPS

DEFINITION. A coloring of a set  $X$  by the set of colors  $F$  is a map  $\phi : X \rightarrow F$ .

We consider the product topology on the set of all colorings  $F^X$  of  $X$  where  $F$  is taken with the discrete topology.

A group  $G$  acts (from the left) on the space of functions  $F^G$  by the formula  $(g\phi)(x) = \phi(g^{-1}x)$ .

DEFINITION. A coloring  $\phi : \Gamma \rightarrow F$  of a discrete group is called *aperiodic* if  $\phi \neq g\phi$  for all  $g \in \Gamma \setminus e$ . It means that the orbit  $\Gamma\phi$  of  $\phi$  under the left action of  $\Gamma$  on  $F^\Gamma$  is full i.e. every the isotropy group of  $\phi$  is trivial.

If  $\phi = g\phi$  for some  $g \in \Gamma$  we call  $\phi$  *g-periodic*. Clearly, the  $g$ -periodicity is equivalent to the  $g^{-1}$ -periodicity.

Note that every group admits an aperiodic coloring by two elements  $\delta_e : \Gamma \rightarrow \{0, 1\}$  with  $\delta_e(e) = 1$  and  $\delta_e(g) = 0$  for  $g \neq e$ . This coloring is not interesting since it fails to satisfy the following condition.

DEFINITION. A coloring  $\phi : \Gamma \rightarrow F$  of a discrete group is called *limit aperiodic* if the action of  $\Gamma$  on the closure  $\overline{\Gamma\phi}$  of the orbit  $\Gamma\phi$  is free, i.e., every coloring  $\psi \in \overline{\Gamma\phi}$  is aperiodic.

We consider the question whether every finitely generated group admits a limit aperiodic coloring by finitely many colors.

REMARK. There is a weaker version of this question. We say that a coloring  $\phi \in F^\Gamma$  is *weakly aperiodic* if the orbit  $\Gamma\phi$  is infinite. A coloring  $\phi$  is called *limit weakly aperiodic* if every coloring in the closure  $\psi \in \overline{\Gamma\phi}$  is weakly aperiodic. We note that the weakly aperiodic version of this question has an affirmative answer. Namely V. Uspenskii proved [Us] that for every discrete group  $\Gamma$  the topological dynamical system  $(F^\Gamma, \Gamma)$  has a compact infinite  $\Gamma$ -invariant set  $X \subset F^\Gamma$  with a minimal action on it. We recall that an action is *minimal* if every orbit is dense. Thus, no orbit in  $X$  can be finite and hence every element  $\phi \in X$  is limit weakly aperiodic.

We don't deal with weakly aperiodic colorings in this paper. We just note that weakly aperiodic colorings correspond to aperiodic tilings and aperiodic coloring correspond to strongly aperiodic tilings (see §4).

**Proposition 1.** *Let  $H \subset G$  be a finite index subgroup. Then the group  $G$  admits a limit aperiodic coloring by finitely many colors if and only if  $H$  does.*

*Proof.* Let  $\phi : G \rightarrow F$  be a limit aperiodic coloring of  $G$ . Let  $n = [G : H]$  and let  $G = \coprod_{i=1}^n Hy_i$  be the decomposition of  $G$  into the right  $H$ -cosets. Let  $\phi y$  denote the result of the right  $y$ -action, that is  $(\phi y)(x) = \phi(xy)$ .

We define  $\psi : H \rightarrow F^n$  by the formula  $\psi(x) = ((\phi y_1)(x), \dots, (\phi y_n)(x))$ . Assume that  $\lim_k (h_k \psi) = \psi'$  for a sequence  $h_k \in H$  and  $\psi'(ax) = \psi'(x)$  for some  $a \in H$  and for all  $x \in H$ . Taking a subsequence we may assume that the sequence  $h_k \phi$  is convergent. Since  $\lim_k (h_k \phi)$  is not  $a$ -periodic, there is  $z \in G$  such that  $(h_k \phi)(az) \neq (h_k \phi)(z)$  for infinitely many  $k$ . Let  $z \in Hy_i$ . Then for all sufficiently large  $k$  we have the equality  $(h_k \psi)(ax) = (h_k \psi)(x)$  for  $x = zy_i^{-1} \in H$ . Hence  $(h_k \phi y_i)(ax) = (h_k \phi y_i)(x)$ . Therefore we have a contradiction:

$$(h_k \phi)(az) = \phi(h_k^{-1} a z y_i^{-1} y_i) = (h_k \phi y_i)(ax) = (h_k \phi y_i)(x) = \phi(h_k^{-1} x y_i) = (h_k \phi)(z).$$

In the other direction we may assume that  $H$  is normal. If it is not, we take a smaller normal subgroup of finite index  $H'$ . By the above  $H'$  admits a limit aperiodic coloring. Let  $\psi : H \rightarrow F$  be a limit aperiodic coloring of  $H$  and let  $n = [G : H]$ . We define a coloring  $\phi : G \rightarrow F \times \{1, \dots, n\}$  by the formula  $\phi(x) = (\psi(y_i^{-1}x), i)$  for  $x \in Hy_i$  where  $G = \coprod_{i=1}^n Hy_i$  is the decomposition of  $G$  into the left  $H$ -cosets (= the right  $H$ -cosets). Assume that  $\phi' = \lim_k g_k \phi$  is  $a$ -periodic for some  $a$ . Since all  $H$ -cosets are colored by different colors,  $a$  must be in  $H$ . We may assume that all  $g_k \in y_j H$  for some fixed  $j$ . Let  $a' = y_j^{-1} a y_j$ . Since  $\psi$  is limit aperiodic, there is  $z \in H$  such that  $(y_j^{-1} g_k \psi)(z) \neq (y_j^{-1} g_k \psi)(a'z)$ . We take  $x = y_j z$ . Then for infinitely many  $k$ ,  $g_k \phi(x) = g_k \phi(ax)$ . Note that  $g_k \phi(x) = \phi(g_k^{-1} y_j z) = \psi(g_k^{-1} y_j z)$  by the definition of  $\psi$

and the choice  $y_1 = e$ . Thus,  $g_k\phi(x) = (y_j^{-1}g_k\psi)(z)$ . On the other hand  $g_k\phi(x) = g_k\phi(ax) = \phi(g_k^{-1}ay_jz) = \phi(g_k^{-1}y_ja'z) = (y_j^{-1}g_k\psi)(a'z)$ . Contradiction.  $\square$

EXAMPLE. The following coloring of  $\mathbf{Z}$  is not limit aperiodic:

$$\phi(n) = \begin{cases} 1 & \text{if } n = \pm k^2, k \in \mathbf{N} \\ 0 & \text{otherwise.} \end{cases}$$

We consider the sequence of colorings  $\psi_m(x) = \phi(x + m^2)$ ,  $\psi_m \in \mathbf{Z}\phi$ , and show that has the constant 0-coloring is the limit of  $\psi_m$ . We need to show that  $\lim_{m \rightarrow \infty} \psi_m(x) = 0$  for every  $x \in \mathbf{N}$ . Since the equation  $x + m^2 = \pm k^2$  has only finitely many integral solutions  $(m, k)$ , the result follows.

Since every weakly aperiodic coloring of  $\mathbf{Z}$  is aperiodic, the existence of a limit aperiodic coloring of  $\mathbf{Z}$  follows from the existence of non-periodic minimal set for the shift action of  $\mathbf{Z}$  on  $\{0, 1\}^{\mathbf{Z}}$  (see the above REMARK).

An explicit example of a limit aperiodic coloring of  $\mathbf{Z}$  can be given by means of the Morse-Thue sequence  $m : \mathbf{N} \rightarrow \{0, 1\}$ .

DEFINITION [Mor],[Th]. Morse-Thue sequence  $m(i)$ . Consider a substitution rule:  $0 \rightarrow 01$  and  $1 \rightarrow 10$ . Then start from 0 to perform this substitutions

$$0 \rightarrow 01 \rightarrow 0110 \rightarrow 01101001 \rightarrow \dots$$

By taking the limit we obtain a sequence of 0 and 1 called the *Morse-Thue sequence*.

**Theorem 1** [HM],[Th]. *The Morse-Thue sequence contains no string of type  $WWW$  where  $W$  is any word in 0 and 1.*

We consider the coloring  $\phi : \mathbf{Z} \rightarrow \{0, 1\}$  defined as  $\phi(x) = m(|x|)$ .

**Proposition 2.** *The coloring  $\phi$  of  $\mathbf{Z}$  is limit aperiodic.*

*Proof.* Assume the contrary: there is a sequence  $\{n_k\}$  tending to infinity and  $a \in \mathbf{N}$  such that  $\psi(x + a) = \psi(x)$  where  $\psi(x) = \lim_{k \rightarrow \infty} \phi(x + n_k)$  for all  $x \in \mathbf{Z}$ . We may assume that all  $n_k > 0$ . Then there is  $k_0$  such that for all  $x \in [1, 3a]$  and all  $k > k_0$ , we have  $\psi(x) = \phi(x + n_k)$ . Let  $w = \psi(1), \psi(2), \dots, \psi(a)$ . Note that  $\psi(1), \psi(2), \dots, \psi(3a) = www$ . On the other hand  $\psi(x) = \phi(x + n_k) = m(x + n_k)$  for  $x \in [1, 3a]$ . Thus we have a "cube"  $www$  in the Morse-Thue sequence. Contradiction  $\square$

QUESTION: Actually this coloring of  $\mathbf{Z}$  has the following stronger property: given  $n \in \mathbf{Z} \setminus 0$  and  $m \in \mathbf{Z}$  there exists a  $q \in \mathbf{Z}$  with  $|q - m| \leq 3|n|$  such that  $\phi(q) \neq \phi(q + n)$ .

One can ask, if every finitely generated group has a finite coloring with a similar property. To be precise, let  $G$  be a finitely generated group, and  $d$  be the word metric with respect to a finite generating set.

Does there exist a finite coloring  $\phi : G \rightarrow F$  and a constant  $\lambda > 0$ , such that for every element  $g \in G \setminus e$  and every  $h \in G$  there exists  $b \in B_{\lambda d_g(h)}(h)$  with  $\phi(gb) \neq \phi(b)$ ? Here  $d_g(h) = d(gh, h)$  is the displacement of  $g$  at  $h$  and  $B_r(h)$  the distance ball of radius  $r$  with center  $h$ .

Such a coloring  $\phi$  is not  $g$  invariant and one can see this aperiodicity already by considering the coloring only on a distance ball  $B_r(h)$ , where  $h$  is an arbitrary point and the radius is a fixed constant times the displacement  $d_g(h)$ . A coloring with this property can be considered as a natural generalization of the Morse-Thue coloring to the group  $G$ . Such a coloring is in some sense "as aperiodic as possible" and in particular limit aperiodic.

## §2 APERIODIC COLORING OF HYPERBOLIC GROUPS

It is very plausible that every finitely generated group has limit aperiodic colorings by finitely many colors. On the other hand a random coloring is not limit aperiodic. In this section we construct such colorings for torsion free finitely generated hyperbolic groups.

In the sequel  $G$  is a finitely generated group, and  $\|\cdot\|$  the norm with respect to a finite set of generators.

**Lemma 1.** *Let  $a \in G$  be an element of infinite order in a hyperbolic group. Then there is  $n = n(a)$  such that for every  $g \in G$  there is  $k \leq n$  with  $\|ga^k\| \neq \|g\|$ .*

*Proof.* It is known that the sequences  $\{a^k\}$  and  $\{a^{-k}\}$  define two different points on the Gromov boundary  $\partial_\infty G$  of  $G$ . Let  $\xi : \mathbf{R} \rightarrow K$  be a geodesic in the Cayley graph  $K$  joining these two points. Note that the action of  $a$  on  $K$  leaves these points at infinity invariant. Let  $d = d(e, im\xi)$ , then  $d(a^k, im\xi) = d(e, a^{-k}im\xi) \leq d + \delta$  for every  $k$  where  $G$  is  $\delta$ -hyperbolic. The last inequality follows from the fact that a degenerated triangle in  $K$  defined by the geodesics  $im\xi$  and  $a^{-k}(im\xi)$  is  $\delta$ -thin. Take  $n$  such that  $\|a^n\| > 2\|a\| + 10d + 14\delta$ . Assume that there is  $g$  such that  $\|g\| = \|ga\| = \|ga^2\| = \dots = \|ga^n\|$ . Consider the geodesic  $g(im(\xi))$ . Let  $w \in g(im(\xi))$  and  $w' \in g(im(\xi))$  be the closest points in  $g(im(\xi))$  to  $g$  and  $ga^n$  respectively.

Since the triangle  $\langle e, w, w' \rangle$  is  $\delta$ -thin, the geodesic segment  $[w, w']$  contains a point  $z$  such that  $d(z, y) < \delta$  and  $d(z, y') < \delta$  where  $y \in [e, w]$  and  $y' \in [e, w']$ . Thus,

$$(1) \quad 2\|z\| - \|y\| - \|y'\| \leq 2\delta.$$

Denote by  $z_k$ ,  $k = 0, \dots, n$ , a point in  $g(\text{im}(\xi))$  such that  $d(z_k, ga^k) \leq d + \delta$ . There is  $i$  such that  $z \in [z_i, z_{i+1}] \subset g(\text{im}(\xi))$ . Then  $d(z, ga^i) \leq 3d + 3\delta + \|a\|$ . Thus,

$$(2) \quad \|g\| = \|ga^i\| \leq \|z\| + 3d + 3\delta + \|a\|.$$

In view of (1) and (2) and the facts  $\|w\| - \|g\|, \|w'\| - \|g\| \leq d + \delta$  we obtain a contradiction:  $\|a^n\| = d(g, ga^n) \leq d(w, w') + 2d + 2\delta = d(w, z) + d(z, w') + 2d + 2\delta \leq d(w, y) + d(y', w') + 2d + 4\delta = \|w\| - \|y\| + \|w'\| - \|y'\| + 2d + 4\delta \leq 2\|g\| - \|y\| - \|y'\| + 4d + 6\delta \leq 2\|z\| - \|y\| - \|y'\| + 2\|a\| + 10d + 12\delta \leq 2\|a\| + 10d + 14\delta$ .  $\square$

**EXAMPLE.** The group  $\mathbf{Z}^2$  does not have the above property with respect to the generators  $(0, \pm 1)$  and  $(\pm 1, 0)$ . Take  $a = (1, -1)$  and  $g_n = (n, n)$ . Then  $\|(n, n) - k(1, -1)\| = 2n$  for all  $k \leq n$ .

A geodesic segment in a finitely generated group is the corresponding sequence of vertices in a geodesic segment in the Cayley graph. A geodesic segment  $[x_1, \dots, x_k]$  is called *radial* if  $\|x_1\| < \|x_2\| < \dots < \|x_k\|$ .

To construct the limit aperiodic coloring we consider a square free sequence  $\nu : \mathbf{N} \rightarrow \{0, 1, 2\}$ , i.e. a sequence which does not contain any subsequence of the form  $WW$ , where  $W$  is a word in  $0, 1, 2$ . It is possible to construct a square free sequence in the following way: take the Thue-Morse sequence  $0110100110010110\dots$  and look at the sequence of words of length 2 that appear:  $01, 11, 10, 01, 10, 00, 01, 11, 10 \dots$ . Replace  $01$  by  $0$ ,  $10$  by  $1$ ,  $00$  by  $2$  and  $11$  by  $2$  to get the following:  $021012021\dots$ . Then this sequence is square-free [HM].

**Theorem 2.** *Every finitely generated torsion free hyperbolic group admits a limit aperiodic coloring by 9 colors.*

*Proof.* The set of colors will be the set of pairs  $(m, n)$  where  $m, n \in \{0, 1, 2\}$ . Let  $G$  be a group taken with the word metric with respect a finite generating set  $S$ . We define  $\phi(g) = (\nu(\|g\|), \|g\| \bmod 3)$  for every  $g \in G$ .

*Claim :* *Let  $[x_1, x_2, \dots, x_k]$  be a radial geodesic segment. Let  $g \in G$  be such that  $\phi(x_i) = \phi(gx_i)$  for all  $i \in \{1, \dots, k\}$ . Assume in addition that  $d(gx_{i_0}, x_{i_0}) < \frac{k}{2}$  for some  $i_0, 1 \leq i_0 \leq k$ . Then  $\|gx_i\| = \|x_i\|$  for all  $i$ .*

We first show that  $[gx_1, \dots, gx_k]$  is also a radial geodesic segment. Since multiplication from the left is an isometry,  $[gx_1, \dots, gx_k]$  is a geodesic segment and in particular  $-1 \leq \|gx_{i+1}\| - \|gx_i\| \leq 1$ . Since  $g$  preserves the coloring we have  $\|gx_{i+1}\| \equiv \|gx_i\| + 1 \pmod 3$ . This two relations imply  $\|gx_{i+1}\| = \|gx_i\| + 1$ , hence  $[gx_1, \dots, gx_k]$  is a radial geodesic segment and thus  $\|gx_i\| = \|x_i\| + q$  for some fixed integer  $q$ . By our assumption  $|q| < \frac{k}{2}$ . Assume  $0 < q$ . Since  $g$  preserves the colors we obtain the equality

$$(\nu(\|x_1\|), \dots, \nu(\|x_q\|)) = (\nu(\|x_{q+1}\|), \dots, \nu(\|x_{2q}\|)),$$

a contradiction to the square freeness of  $\nu$ . In a similar way we obtain a contradiction if  $q < 0$ . This implies  $q = 0$  and hence the claim.

Now assume that there is a sequence  $g_l \in G$  with  $\|g_l\| \rightarrow \infty$  such that the limit  $\lim_{l \rightarrow \infty} \phi(g_l x) = \psi(x)$  exists for every  $x \in G$ .

Assume that there is  $b \in G \setminus \{e\}$  such that  $\psi(x) = \psi(bx)$  for all  $x$ .

Let  $n$  be taken from Lemma 1 for  $a = b$ . We may assume that there is  $l_0$  such that for  $l > l_0$ ,  $\psi(y) = \phi(g_l y)$  for all  $y \in B_{4n\|b\|}(e)$ . Consider a radial geodesic segment  $x_1, \dots, x_k$  of the length  $k - 1$  with  $k = 3n\|b\|$  and with endpoint  $x_k = g_l$ . Such a segment clearly exists for all  $l$  large enough.

Let  $s$  be the smallest natural number such that  $\|g_l b^s\| \neq \|g_l\|$ , thus  $s \leq n$ . Both segments  $g_l^{-1}([x_1, \dots, x_k])$  and  $g_l^{-1}(g_l b^s g_l^{-1}[x_1, \dots, x_k])$  lie in  $B_{4n\|b\|}(e)$ . Thus  $\phi(x) = \phi(g_l g_l^{-1}(x)) = \psi(g_l^{-1}(x)) = \psi(b^s g_l^{-1}(x)) = \phi(g_l b^s g_l^{-1}x)$  for all  $x \in [x_1, \dots, x_k]$ . Furthermore we compute  $d(g_l b^s g_l^{-1}x_k, x_k) = d(g_l b^s, g_l) = \|b^s\| \leq n\|b\| < \frac{k}{2}$ . We apply the claim to  $g = g_l b^s g_l^{-1}$  to obtain the contradiction:  $\|g_l b^s\| = \|g_l b^s g_l^{-1}x_k\| = \|x_k\| = \|g_l\|$ .  $\square$

REMARK. It is still an open problem whether every hyperbolic group contains a torsion free subgroup of finite index.

### §3 APERIODIC COLORING OF $G$ -SPACES

We note that in the definitions from the beginning of §1 one can replace a group  $G$  by a space  $X$  with a  $G$ -action. Thus,  $G$  acts on the space of colorings  $F^X$  by the same formula  $(g\phi)(x) = \phi(g^{-1}x)$ . Let  $K$  be the kernel of the action. We can speak about  $G$ -aperiodic colorings of  $X$  as ones with  $\phi \neq g\phi$  for all  $g \in G \setminus K$ . Similarly one can define *limit  $G$ -aperiodic* colorings of  $X$  as those whose orbit  $G\phi$  has only  $G$ -aperiodic coloring in its closure  $\overline{G\phi}$  with respect to the product topology  $F^X$ .

The following is an analog of Proposition 1.

**Proposition 3.** *Let  $H \subset G$  be a finite index subgroup and let  $G$  act on  $X$ . Then  $X$  admits a limit  $G$ -aperiodic coloring by finitely many colors if and only if it admits a limit  $H$ -aperiodic coloring by finitely many colors.*

It is possible to extend Theorem 2 to the case of an isometric action on a hyperbolic space  $X$ . We consider only the case when  $X$  is a simplicial tree. Thus every edge has length equal to 1.

If  $x_0$  is a root of  $X$ , we denote by  $\|x\| = d(x, x_0)$  for  $x \in X$ . We prove the following analog of Lemma 1.

**Lemma 2.** *Let  $G$  act on a simplicial rooted tree  $X$  and let  $a \in G$  operate without fixed points. Then for every  $g \in G$  there is  $k \leq 2$  with  $\|ga^k x_0\| \neq \|g x_0\|$ .*



*Proof.* Assume that  $\|ga^2x_0\| = \|gax_0\| = \|gx_0\|$ . Since  $d(gx_0, gax_0) = d(hgx_0, hgax_0) = d(gax_0, ga^2x_0)$  for  $h = gag^{-1}$ , the points  $z = gx_0$ ,  $h(z) = gax_0$ , and  $h^2(z) = ga^2x_0$  have common predecessor  $y$  in the tree which is the common midpoint of the geodesic segments  $[z, h(z)]$  and  $[h(z), h^2(z)]$ . Then  $hy = y$  and hence  $g^{-1}y$  is a fixed point for  $a$ :  $a(g^{-1}y) = g^{-1}y$ . This contradicts to the assumption.

□

Let  $x_0 \in X$  be a base point in a tree  $X$ . We consider the coloring of the set of vertices of  $X$  defined like in the proof of Theorem 2:  $\phi(x) = (\nu(\|x\|), \|x\| \bmod 3)$ .

**Proposition 4.** *Suppose that a group  $G$  acts by isometries on a simplicial tree  $X$  with the above coloring  $\phi$  and let  $\psi \in \overline{G\phi}$ . Then  $b\psi \neq \psi$  for every  $b \in G$  with unbounded orbit  $\{b^k x_0 \mid k \in \mathbb{N}\}$ . Moreover,  $b\psi \neq \psi$  on the orbit  $Gx_0$ .*

*Proof.* First we note that a similar as in the proof of Theorem 2 claim take place:

Let  $[z_1, \dots, z_k]$  be a radial geodesic segment in  $X$ . Let  $g \in G$  such that  $\phi(gz_i) = \phi(z_i)$  for all  $i$  and  $d(gz_{i_0}, z_{i_0}) < \frac{k}{2}$  for some  $i_0$ ,  $1 \leq i_0 \leq k$ . Then  $\|gz_i\| = \|z_i\|$ .

The proof is exactly the same as in Theorem 2.

Assume that there is a sequence  $g_l \in G$  with  $\|g_l\| \rightarrow \infty$  with the limit  $g_l^{-1}\phi$  equal to a  $b$ -periodic coloring  $\psi$  such that  $\{b^k x_0\}$  is infinite. That is the limit  $\lim_{k \rightarrow \infty} \phi(g_k x) = \psi(x)$  exists for every  $x \in X$ .

Since the orbit  $b^m x_0$  is infinite we can apply Lemma 2 for  $a = b$ . Let  $[y_1, \dots, y_{8k}]$  be a radial segment with  $y_1 = x_0$  where  $k = 6\|b(x_0)\|$ . By the definition of the pointwise limit we may assume that there is  $l_0$  such that for  $l > l_0$ ,  $\psi(y) = \phi(g_l y)$  for all  $y \in [y_1 \dots y_{8k}] \cup b[y_1 \dots y_{8k}] \cup b^2[y_1 \dots y_{8k}]$ . Then the image  $g_l[y_1, \dots, y_{8k}]$  contains either a radial segment  $[z_1, \dots, z_k]$  of the length  $k - 1$  with  $z_k = g_l(x_0)$  or it contains a radial segment  $[z_1, \dots, z_{6k}]$  with  $d(z_1, g_l(x_0)) < k$ .

Let  $s$  be the smallest natural number such that  $\|g_l b^s x_0\| \neq \|g_l x_0\|$ . Thus,  $s \leq 2$ .

Then  $\phi(x) = \phi(g_l g_l^{-1}(x)) = \psi(g_l^{-1}(x)) = \psi(b^s g_l^{-1}(x)) = \phi(g_l b^s g_l^{-1}x)$  for all  $x \in [z_1, \dots, z_k]$ . Furthermore, in the first case we compute  $d(g_l b^s g_l^{-1} z_k, z_k) = d(b^s x_0, x_0) = \|b^s x_0\| \leq 2\|b x_0\| < \frac{k}{2}$ . We apply the claim to  $g = g_l b^s g_l^{-1}$  with  $i_0 = k$  to obtain the contradiction:  $\|g_l b^s x_0\| = \|g_l b^s g_l^{-1} z_k\| = \|z_k\| = \|g_l x_0\|$ .

In the second case  $d(z_1, g_l b^s g_l^{-1} z_1) \leq$

$$d(z_1, g_l(x_0)) + d(g_l(x_0), g_l b^s(x_0)) + d(g_l b^s(x_0), g_l b^s g_l^{-1} z_1) \leq 2k + 2\|b(x_0)\| \leq 3k.$$

We apply the claim with  $g = g_l b^s g_l^{-1}$ ,  $i_0 = 1$ , and  $6k$  instead of  $k$ . Let  $i$  be such that  $g_l y_i = z_1$ . Then  $\|z_i\| = \|g_l x_0\|$ . Since  $\phi(\|g_l b^s y_{i-1}\|) = \phi(\|g_l y_{i-1}\|) \neq \phi(\|g_l y_i\| - 1)$ , we obtain that  $\|g_l b^s y_{i-1}\| \neq \|g_l y_i\| - 1 = \|z_1\| - 1$ . Hence  $\|g_l b^s y_{i-1}\| = \|z_1\| + 1$  and

$\|g_l b^s x_0\| = \|z_1\| + i - 1$ . Then we obtain a contradiction:  $\|g_l b^s x_0\| = \|z_1\| + i - 1 = \|g_l b^s g_l^{-1} z_1\| + i - 1 = \|g_l b^s g_l^{-1} z_i\| = \|z_i\| = \|g_l x_0\|$ .  $\square$

As a consequence we obtain the following:

**Corollary 1.** *Suppose that a group  $G$  acts on a rooted simplicial tree  $X$  such that  $Gx_0$  is a full orbit. Then there is a limit  $G$ -aperiodic coloring of  $X$  by 9 colors. Moreover, the restriction of this coloring to the orbit  $Gx_0$  is also limit  $G$ -aperiodic.*

**Theorem 3.** *Suppose that a group  $G$  acts by isometries on simplicial trees  $X_1, \dots, X_n$  in such a way that the induced action on the product  $\prod X_i$  is free. Then  $G$  admits a limit aperiodic coloring by  $9n$  colors.*

*Proof.* First we note that the fixed point theorem for CAT(0) spaces implies that  $G$  must be torsion free.

Let  $K_i$  be the kernel of the representation  $G \rightarrow \text{Aut}(X_i)$  and let  $G_i = G/K_i$ . We fix a base point  $x_0^i$  in each tree and consider colorings  $\phi^i : X_i \rightarrow F_i$ ,  $|F_i| = 9$  from Proposition 4. This defines a map  $\phi : \prod X_i \rightarrow \prod F_i$ . Let  $\phi' : G \rightarrow \prod F_i$  be the restriction to the orbit:  $\phi'(g) = \phi(gx_0)$  where  $x_0 = (x_0^1, \dots, x_0^n)$ .

Assume that  $\psi = \lim g_k^{-1} \phi'$  be  $a$ -periodic:  $a\psi(x) = \psi(x)$ ,  $a \in G \setminus \{e\}$ . Then  $\psi = (\psi_1, \dots, \psi_n)$  where  $\psi_i = \lim g_k^{-1} \phi'_i$  and  $\phi'_i = \phi_i|_{Gx_0^i}$ . Since  $a$  operates without fixed points on  $X$ , there exists  $i$  such that  $a$  has no fixed points on  $X_i$ . Then by Proposition 4  $a\psi \neq \psi$  which contradicts with the assumption.  $\square$

We recall the definition of Coxeter groups A *Coxeter matrix*  $M = (m_{ij})$  is a symmetric square matrix with 1 on the diagonal and all other entries are from  $\mathbf{N}_+ = \{0\} \cup \mathbf{N}$ . A *Coxeter group*  $\Gamma$  with a generating set  $S$  is a group with a presentation

$$\langle S \mid (uv)^{m_{ij}} = 1, m_{ij} \in M \rangle$$

where  $M$  is a Coxeter matrix. Here we use the convention  $a^0 = 1$ . Traditionally the literature on Coxeter groups uses  $\infty$  instead of 0.

A Coxeter group is called *right-angled* if all  $m_{ij}$  are 0s or 2s.

**Theorem 4.** *Every Coxeter group  $\Gamma$  admits a limit aperiodic coloring by finitely many colors.*

*Proof.* Since every Coxeter group contains a finite index torsion free subgroup, in view of Proposition 1 it suffices to prove it for a finite index subgroup. We apply Januszkiewicz's construction of equivariant isometric embedding of a torsion free finite index normal subgroup  $\Gamma' \subset \Gamma$  into the finite product of trees [DJ] and then apply Theorem 3.  $\square$

Let  $K$  be the Cayley graph of a Coxeter group  $(\Gamma, S)$ . Then every generator  $c \in S$  every conjugate  $w = gcg^{-1}$ ,  $g \in \Gamma$ , acts on  $K$  by reflection. Let  $M_w$  be the set of fixed point of  $w$ . We call it the *wall (or mirror)* of the reflection  $w$ . Clearly, that  $uM_w$  is a wall for all  $u$  and  $w$ . Therefore  $\Gamma$  acts on the set of all walls  $\mathcal{W}$ . According to [DJ] all walls can be partitioned in finitely many classes  $\mathcal{W} = \mathcal{W}_1 \cup \dots \cup \mathcal{W}_m$  such that the walls from each class  $\mathcal{W}_i$  form a vertex set of a simplicial tree  $T_i$ . Moreover, all sets  $\mathcal{W}_i$  are invariant under a normal finite index subgroup  $\Gamma'$  which acts by isometries on each  $T_i$  in such a way that the  $\Gamma'$ -action on the product  $\prod_{i=1}^m T_i$  is free.

**Theorem 5.** *The set  $\mathcal{W}$  of all walls in a Coxeter group admits a limit  $\Gamma$ -aperiodic coloring by finitely many colors.*

*Proof.* On every tree  $T_i$  we consider a coloring from Proposition 6 with 9 each time different colors. Thus we use  $9m$  colors. This defines a coloring  $\phi = \cup \phi_i$  of  $\mathcal{W}$ . Let  $\psi$  be a limit coloring of  $\mathcal{W}$ . Then, clearly,  $\psi = \cup \psi_i$  where each  $\psi_i$  is a limit coloring for  $\phi_i$ . Let  $b \in \Gamma'$ . Since the action of  $\Gamma'$  on the product  $\prod_{i=1}^m T_i$  is free and  $b$  is of infinite order, we obtain that  $b$  has an infinite orbit on some  $T_i$ . By Proposition 6  $b\psi_i \neq \psi_i$  and hence  $b\psi \neq \psi$ . Proposition 5 completes the proof.  $\square$

#### §4 SPACE OF TILINGS

We recall a definition of tiling of a metric space from [BW]. Let  $X$  be a metric space. A set of tiles  $(\mathcal{T}, \mathcal{F})$  is a finite collection of  $n$ -dimensional complexes  $t \in \mathcal{T}$  and a collection of subcomplexes  $f \in \mathcal{F}$  of dimension  $< n$ , together with an opposition function  $o : \mathcal{F} \rightarrow \mathcal{F}$ ,  $o^2 = id$ . A space  $X$  is tiled by the set  $(\mathcal{T}, \mathcal{F})$  if

- (1)  $X = \cup_{\lambda} t_{\lambda}$  where each  $t_{\lambda}$  is isometric to one of the tiles in  $\mathcal{F}$ ;
- (2)  $t_{\lambda} \setminus \cup_{f \in t_{\lambda}} f = Int(t_{\lambda})$  in  $X$  for every  $\lambda$ ;
- (3) If  $Int(t_{\lambda} \cup t_{\lambda'}) \neq Int(t_{\lambda}) \cup Int(t_{\lambda'})$  then  $t_{\lambda}$  and  $t_{\lambda'}$  intersect along  $f \in t_{\lambda}$  and  $o(f) \in t_{\lambda'}$ ;
- (4) There are no free faces of  $t_{\lambda}$ .

Strictly speaking a tiling of  $X$  is a collection  $\alpha = \{\phi_{\lambda}\}$  of isometries  $\phi_{\lambda} : t_{\lambda} \rightarrow t$ ,  $t \in \mathcal{T}$ , satisfying the above axioms. For every tiling  $\alpha$  there is a minimal (or reduced) set of tiles  $(\mathcal{T}_{\alpha}, \mathcal{F}_{\alpha}) \subset (\mathcal{T}, \mathcal{F})$ .

Let  $X$  be a metric space with a base point  $x_0$ . Assume that  $diam t \leq 1$  for all  $t \in \mathcal{T}$ . Let  $\alpha$  be a tiling  $X = \cup_{\lambda} t_{\lambda}$  of  $X$ . We denote by  $\alpha_n = \{t_{\lambda} \mid t_{\lambda} \subset B_n(x_0)\}$  where  $B_r(x)$  stands for the closed ball of radius  $r$  centered at  $x$ . For a metric space  $Y$  we denote by  $expY$  the space of compact subsets of  $Y$  taken with the Hausdorff metric. Also for  $m \in \mathbf{N}$  denote by  $exp_m Y$  the  $m$ th hyperpower of  $Y$ , i.e., subspace of  $expY$  that consists of subsets of cardinality  $\leq m$ . Note that  $\alpha_n$  defines a point in  $exp(expB_n(x_0))$ . Actually, there is

$k = k(n)$  such that  $\alpha_n$  lies in  $\exp_k(\exp B_n(x_0))$ . Clearly, the sequence  $\alpha_n$  completely defines the tiling  $\alpha$ .

The space of tilings was defined by many authors (see for example [BBG],[S],[SW]). Here we give an alternative definition. Let  $tl(X, \mathcal{T})$  denote the set of all  $\mathcal{T}$ -tilings of  $X$ . We introduce the topology on  $tl(X, \mathcal{T})$  as a subspace topology

$$tl(X, \mathcal{T}) \subset \prod_{n=1}^{\infty} \exp(\exp B_n(x_0)).$$

Let  $\mathcal{F}$  be a finite family of compact subsets in a metric space  $Y$ . We denote by  $\exp^{\mathcal{F}} Y$  the subspace of  $\exp Y$  whose points are isometric copies of elements of  $\mathcal{F}$ .

Note that  $tl(X, \mathcal{T}) \subset \prod_{n=1}^{\infty} \exp_{k(n)} \exp^{\mathcal{T}}(B_n)$ .

The following Proposition is well-known [S].

**Proposition 7.** *The space  $tl(X, \mathcal{T})$  is compact.*

*Proof.* Since  $\exp^{\mathcal{T}}(B_n)$  is compact, it suffices to show that  $tl(X, \mathcal{T})$  is a closed subset in  $\prod_{n=1}^{\infty} \exp_{k(n)} \exp^{\mathcal{T}}(B_n)$ . For that it suffices to show that the set  $\{\alpha_n \mid \alpha \in tl(X, \mathcal{T})\}$  is closed in  $\exp_{k(n)} \exp^{\mathcal{T}}(B_n)$  for every  $n$ .  $\square$

Let  $G \subset Iso(X)$  be a subgroup of the group of isometries of  $X$ . Clearly  $G$  acts on  $tl(X, \mathcal{T})$ . We say that a tiling  $\alpha \in tl(X, \mathcal{T})$  is *strongly  $G$ -aperiodic* if  $g\alpha \neq \alpha$  for all  $g \in G \setminus \{e\}$ . A tiling  $\alpha$  is called *aperiodic* if the group  $Iso(\alpha) \subset Iso(X)$  of isometries of  $\alpha$  does not act cocompactly on  $X$ . A tiling  $\alpha$  is *limit strongly  $G$ -aperiodic* if every tiling  $\beta \in \overline{G\alpha}$  is strongly aperiodic. If  $G = Iso(X)$  we use the terms *strongly aperiodic* and *limit strongly aperiodic*.

## §5 APERIODIC TILING OF DAVIS COMPLEX

Here we recall the definition of the Davis complex [D1]. Let  $\Gamma$  be a Coxeter group with generating set  $S$ . The nerve  $N = N(\Gamma, S)$  is the simplicial complex defined in the following way: the vertices of  $N$  are elements of  $S$ . Different vertices  $s_1, \dots, s_k$  span a simplex  $\sigma$  if and only if the set  $s_1, \dots, s_k$  generates a finite subgroup  $\Gamma_\sigma$  of  $\Gamma$ . By  $N'$  we denote the barycentric subdivision of  $N$ . The cone  $C = Cone N'$  over  $N'$  is called a *chamber* for  $\Gamma$ . The Davis complex  $X = X(\Gamma, S)$  is the image of a simplicial map  $q : \Gamma \times C \rightarrow X$  defined by the following equivalence relation on the vertices:  $a \times v_\sigma \sim b \times v_\sigma$  provided  $a^{-1}b \in \Gamma_\sigma$  where  $\sigma$  is a simplex in  $N$  and  $v_\sigma$  is the barycenter of  $\sigma$ . We identify  $C$  with the image  $q(e \times C)$ . The group  $\Gamma$  acts simplicially on  $X$  with the orbit space equivalent to the chamber. Thus, the Davis complex is obtained by gluing the chambers  $\gamma C$ ,  $\gamma \in \Gamma$  along their boundaries. Note that  $X$  admits an equivariant cell structure with

the vertices  $X^{(0)}$  equal the cone points of the chambers and with the 1-skeleton  $X^{(1)}$  isomorphic to the Cayley graph of  $\Gamma$ . A conjugate  $r = wsw^{-1}$  of every generator  $s \in S$  is a reflection. The fixed point set  $M_r$  of a reflection  $r$  is called the wall of  $r$ . Note that walls defined in §3 are obtained from the walls in Davis' complex by the restriction to the Cayley graph.

**Proposition 8.** *Every finite coloring  $\phi : \mathcal{W} \rightarrow F$  of the set of walls of the Davis complex  $X$  defines a tiling  $\bar{\phi}$  of  $X$  with  $o(f) = f$ .*

*Proof.* The set of tiles  $\mathcal{T}$  of  $\bar{\phi}$  is the set of chambers with all possible colorings of their faces. The set of faces  $\mathcal{F}$  is the set of all possible colored faces of the chambers. Set  $o(f) = f$ . Then all conditions hold.  $\square$

We call the tiling  $\bar{\phi}$  as a *tiling by coloring*  $\phi$ .

Let  $(\mathcal{T}, \mathcal{F})$  be a set of tiles. A function  $w : \mathcal{F} \rightarrow \mathbf{Z}$  is called a *weight function* if  $w(o(f)) = -w(f)$  for every  $f \in \mathcal{F}$ . We recall a definition from [BW].

**Definition.** *A finite set of tiles  $(\mathcal{T}, \mathcal{F})$  is unbalanced if there is a weight function  $w$  such that  $\sum_{f \in t} w(f) > 0$  for all  $t \in \mathcal{T}$ .*

*It is called semibalanced if  $\sum_{f \in t} w(f) \geq 0$  for all  $t \in \mathcal{T}$ .*

We call a set of tiles *strictly balanced* if for every nontrivial weight function  $w$  there are tiles  $t_+$  and  $t_-$  such that  $\sum_{f \in t_+} w(f) > 0$  and  $\sum_{f \in t_-} w(f) < 0$ .

A tiling is called *strictly balanced (unbalanced)* if its minimal set of tiles is strictly balanced (unbalanced).

We now associate to every wall in the Davis complex an orientation. A wall divides the Davis complex into two components. Roughly speaking the orientation says which of the components is left and which is right. Let  $\bar{\phi}$  be tiling of the Davis complex  $X$  by coloring of the walls  $\phi$  with the set of tiles  $(\mathcal{T}, \mathcal{F})$ . The orientation of the walls define a new tiling  $\phi'$  of  $X$  with the set of tiles  $(\mathcal{T}', \mathcal{F}')$  where  $\mathcal{F}' = \mathcal{F}_+ \cup \mathcal{F}_-$  where  $\mathcal{F}_+$  and  $\mathcal{F}_-$  are copies of  $\mathcal{F}$ . The face  $f \in t_\lambda$  has sign  $+$ , if  $Int(t_\lambda)$  is left of the wall and sign  $-$ , if  $Int(t_\lambda)$  is right of the wall. The opposition function  $o : \mathcal{F}' \rightarrow \mathcal{F}'$  maps  $f_+$  to  $f_-$ . We call such tiling a *geometric resolution* of a tiling by coloring. This new tiling is not any more a tiling by coloring. A geometric meaning of this resolution is that we deform all faces of a given color and a given sign in the same direction by the same pattern. For the faces of the same color but of opposite sign we take an opposite deformation.

The following is obvious.

**Lemma 3.** *Assume that a coloring  $\phi : \mathcal{W} \rightarrow F$  is limit aperiodic. Then the tiling by color  $\bar{\phi}$  as well as it's any geometric resolution is limit strongly aperiodic.*

Note that in the Davis complex every wall has a canonical orientation, by deciding that the base chamber  $C$  is in the left component. Thus we can indicate the chosen orientation itself by a sign. A wall gets the sign  $+$ , if the orientation of the wall is the canonical one and  $-$  otherwise.

In [BW] unbalanced tilings of some nonamenable spaces are constructed. In particular all hyperbolic Coxeter groups admit such tilings. We can derive this fact using geometric resolutions.

**Proposition 9.** *Every coloring of the walls for a hyperbolic Coxeter group admits an unbalanced geometric resolution.*

*Proof.* We assign  $+$  to every wall. The hyperbolicity implies that for every chamber  $C'$  the numbers of faces of  $C'$  whose walls separate  $C'$  from the base chamber  $C$  is strictly less than the number of faces whose walls do not separate  $C'$  and  $C$ . Then for every chamber  $C'$  the faces whose walls do not separate  $C'$  from  $C$  obtain sign  $+$ , all other  $-$ . We define a weight function by sending a positive face to  $+1$  and a negative face to  $-1$ .  $\square$

We note that every unbalanced tiling is aperiodic. This fact can be derived formally from proposition 4.1 [BW]. Since the proof there has some omissions we present a proof below.

**Proposition 10.** *Let  $(\mathcal{T}, \mathcal{F})$  be the set of tiles of a geometric realization of a tiling by coloring of the Davis complex  $X$  of a Coxeter group  $\Gamma$ . Suppose that the set of tiles  $(\mathcal{T}, \mathcal{F})$  is unbalanced. Then any  $(\mathcal{T}, \mathcal{F})$ -tiling  $\alpha$  is aperiodic.*

*Proof.* Let  $G$  be a group of isometries of  $\alpha$ . Then  $G \subset \Gamma$ . Hence  $G$  is a matrix group. By Selberg Lemma it contains a torsion free subgroup  $G'$  of finite index. Then the orbit space  $X/G'$  is compact and admits a  $(\mathcal{T}, \mathcal{F})$ -tiling (Note that by taking  $X/G$  as in [BW] we cannot always obtain a tiling because of free faces). Then we obtain a contradiction:

$$0 < \sum_{t \in X/G'} \sum_{f \in t} w(f) = \sum_{f \in X/G'} (w(f) + w(o(f))) = 0.$$

$\square$

**Theorem 6.** *For every Coxeter group  $\Gamma$  for every coloring  $\phi : \mathcal{W} \rightarrow F$  with the property that walls of the same color do not intersect, there is a strictly balanced geometric resolution. Additionally, every limit tiling of this resolution is strictly balanced.*

*Proof.* First we construct a strictly balanced geometric resolution of  $\phi$ . Consider the set of walls  $\mathcal{W}_c = \phi^{-1}(c)$  of the same color  $c \in F$ . Since the walls of the same color do not

intersect, they are ordered by level from the base chamber. (The level  $lev$  is defined by induction. If one removes the walls  $\mathcal{W}_c$  from  $X$ , the space is divided into components. Walls from  $\mathcal{W}_c$  that bound the component of the base chamber are of level one. Then drop the walls of level one and repeat the procedure to get new walls of level one and call them of level two and so on). We give the walls  $\mathcal{W}_c$  signs in an alternate fashion by the level  $(-1)^{lev(M)}$ :  $- + - + - + - + - \dots$

We show that this geometric resolution is strictly balanced. Let  $w : F_+ \cup F_- \rightarrow \mathbf{Z}$  be a nontrivial weight function with  $w(f_+) = -w(f_-)$ . We show that there are chambers  $C_+$  and  $C_-$  such that

$$\sum_{f \in C_+} w(f) > 0 \quad \text{and} \quad \sum_{f \in C_-} w(f) < 0.$$

Because of the symmetry it suffices to show the first. Since  $w$  is nontrivial, there exists a face  $f^0$  which is the common face of two adjacent tiles  $t_\lambda$  and  $t_{\lambda'}$  such that  $w(f^0) \neq 0$ . Let  $M_0$  denote the wall that contains  $f^0$ . Now there are four cases corresponding to the parity of the sign of  $w(f_+^0)$  and the sign of  $M_0$ . We discuss only one, and to be fair not the easiest of the cases:  $w(f_+^0) > 0$  and the orientation of  $M_0$  is negative.

We assume that  $f^0$  in  $t_\lambda$  has sign  $-$  and in  $t_{\lambda'}$  has sign  $+$ . We take a number  $k$  larger than the number of walls separating  $t_{\lambda'}$  and the base chamber  $C$ . Let  $c \in F$  be a color. We call  $c$  *even* if  $w(c_+) > 0$ , *odd* if  $w(c_+) < 0$ , and *neutral* if  $w(c_+) = 0$ . We define

$$\begin{aligned} \mathcal{W}_{ev}^{2k} &= \{M \in \mathcal{W} \mid \phi(M) \text{ is even and } lev(M) = 2k\} \\ \mathcal{W}_{odd}^{2k+1} &= \{M \in \mathcal{W} \mid \phi(M) \text{ is even and } lev(M) = 2k + 1\} \\ \mathcal{W}_0^{k+1} &= \{M \in \mathcal{W} \mid \phi(M) \text{ is neutral and } lev(M) = k + 1\}. \end{aligned}$$

Claim 1: *The set of walls  $\mathcal{W}_{ev}^{2k} \cup \mathcal{W}_{odd}^{2k+1} \cup \mathcal{W}_0^{k+1} \cup \{M_0\}$  bounds a bounded set  $D$  containing the chamber  $t_{\lambda'}$ .*

Clearly, it bounds a convex set in the Hadamard space  $X$ . If it is unbounded, then there is a geodesic ray from  $t_{\lambda'}$  to the visual boundary which does not intersect any of our mirrors. Since we have only finitely many colors, there is a color  $c$  such that this ray intersects infinitely many walls of this color. By the choice of  $k$  the first of this crossed walls has level  $\leq k + 1$ . To get to infinity the ray must cross walls of color  $c$  with all levels  $\geq k + 1$ . Thus, one of the intersected walls is contained in our set.

Claim 2: If  $f$  occurs as a face of a tile  $t_\mu \subset D$  such that  $f \subset \partial D$ , then  $w(f) > 0$ .

We consider cases.

- (i) If  $M_f = M_0$ , then  $t_\mu$  lies on the same side of  $M_0$  as  $t_{\lambda'}$ . Then  $w(f) = w(f_+) > 0$ .
- (ii) If  $f$  is a neutral face, then  $w(f) = 0$  anyway.
- (iii) If  $f$  is of even color. Then  $f$  is contained in a wall  $M$  from  $\mathcal{W}_{ev}^{2k}$  then  $t_\mu$  lies on the same side of the wall  $M_f$  as the base chamber. Since  $M$  has orientation  $+$ . Hence  $f$  as a face of  $t_\mu$  gets the sign  $+$ . Hence  $w(f) = w(f_+) > 0$ .

(iv) A similar argument applies for  $f$ , if  $f$  is of an odd color.

According to the Claim 1 we have  $D = \cup_{i=1}^k C_i$  where  $C_1, \dots, C_k$  is a finite collection of chambers. Then

$$\sum_{i=1}^k \sum_{f' \in C_i} w(f') = \sum_{f' \in \partial D} w(f') \geq 0$$

by Claim 2. Since  $f_+^0$  is in the last set of faces, we see that the expression is indeed  $> 0$ . Therefore,  $\sum_{f' \in C_i} w(f') > 0$  for some  $i$ .

This finishes the proof of the first step. Thus we have constructed a strictly balanced geometric resolution of  $\phi$ .

Actually the proof of the first step shows more: If we chose for any given color  $c$  an orientations of walls  $\mathcal{W}_c$  in the alternate way  $+ - + - + - \dots$  or  $- + - + - + \dots$  (and maybe for different colors in a different way), then the resulting geometric resolution is strictly balanced. Let us call such choice of orientations as *allowed*. The levels of walls depend on the base chamber. If we define levels with respect to a different chamber, all parities of the levels will be either preserved or changed to opposite. as a consequence we obtain the following: if the orientation of tiling by coloring  $\phi$  is allowed, then also the orientation of the tiling by coloring  $g\phi$  is allowed for every  $g \in \Gamma$ . Thus also all limit tilings of tiling constructed in the step 1 are strictly balanced.  $\square$

**Corollary 2.** *For every Coxeter group  $\Gamma$  there is a strictly balanced strictly aperiodic tiling of the Davis complex such that every limit tiling is strictly balanced and strictly aperiodic.*

*Proof.* We apply Theorem 6 to a coloring from Theorem 5.  $\square$

Note that in the proof of Corollary 2 we used that  $\Gamma$  is the isometry group of the Davis complex.

In 2-dimensional jigsaw tiling puzzles a geometric resolution is usually realized by adding rounded tabs out on the sides of the pieces with corresponding blank cut into intervening sides to receive the tabs of adjacent pieces. This procedure destroys the convexity of the pieces. We show that in the case of the hyperbolic plane  $\mathbf{H}^2$  we can modify this construction to obtain aperiodic and strictly balanced tiling with convex tiles. Compare also the papers [MM],[Moz].

**Theorem 7.** (1) *For every  $n \geq 3$  there is a strictly balanced limit strongly aperiodic tiling of  $\mathbf{H}^2$  by convex  $2n$ -gons.*

(2) *For every  $n \geq 3$  there is a finite set of tiles  $(\mathcal{T}, \mathcal{F})$  that consists of convex  $2n$ -gons with limit strongly aperiodic tiling of  $\mathbf{H}^2$  such that every  $(\mathcal{T}, \mathcal{F})$ -tiling of  $\mathbf{H}^2$  is aperiodic.*



*Proof.* (1) Identify  $\mathbf{H}^2$  with the Davis complex for the right-angled Coxeter group  $\Gamma$  generated by reflections at a regular right-angled  $2n$ -gon. Coloring the sides of the  $2n$ -gon in two colors  $a$  and  $b$  in an alternating fashion induces a coloring of the walls  $\psi : \mathcal{W} \rightarrow \{a, b\}$  such that the walls of the same color do not intersect. The walls of the same color  $c$  define a tree  $T_c$  with an action of  $\Gamma$  on it such that the induced  $\Gamma$ -action on the product  $T_a \times T_b$  is free (see [BDS] or [DJ]). By Theorem 3 we can refine  $\psi$  to a limit aperiodic coloring  $\psi : \mathcal{W} \rightarrow \{a_i, b_i\}_{i=1, \dots, 9}$ . We apply Theorem 6 to obtain limit strictly balanced geometric resolution  $\phi'$ . By Lemma 3 it will be limit strongly  $\Gamma$ -aperiodic tiling. It's easy to see that the tiling is limit strongly aperiodic with respect to entire isometry group of  $\mathbf{H}^2$ .

Now we define a modification of the tiling  $\phi'$ . Consider a vertex of a translate of the  $2n$ -gon. It is the intersection point of an  $a_i$ -wall with a  $b_j$ -wall. Denote it by  $O_{ij}$ . The orientations on these walls define local coordinate system. We move the vertex  $O_{ij}$  by small amount using these coordinates. We chose a small different numbers  $d_{ij}$ ,  $i, j \in \{1, \dots, 9\}$  and move  $O_{ij}$  to the distance  $d_{ij}$  in the direction of the diagonal of the positive quadrant. After this deformation we obtain a finite number of new convex tiles, which (for generic deformations) only allow tilings of  $\mathbf{H}^2$  compatible with the matching rule defined by  $\phi'$ . New tiling has all desired properties.

(2) We take the above coloring  $\phi$  of the walls and take a geometric resolution from Proposition 9. Apply Proposition 10 to complete the proof.  $\square$

An interesting question is under what conditions the set of tiles  $(\mathcal{T}, \mathcal{F})$  of a geometric resolution  $\phi'$  of a tiling by coloring of the Davis complex  $X$  is (strongly) aperiodic. Clearly, it is strongly aperiodic whenever its orbit  $\Gamma\phi'$  in  $tl(X, \mathcal{T})$  is dense.

## REFERENCES

- [A] E. Andreev, *On the intersection of faces of acute-angled polyhedra*, Math. Notes **8** (1971), 761-764.
- [BW] J. Block, S. Weinberger, *Aperiodic tilings, positive scalar curvature and amenability of spaces*, J. Amer. Math. Soc. **5 no. 4** (1992), 907-921.
- [BBG] J. Bellissard, R. Benedetti, and J.-M. Gambaudo, *Spaces of tilings, finite telescopic approximations and gap-labelling*, Comm. Math. Phys. **261 no 1.** (2006), 1-41.
- [BDS] S. Buyalo, A. Dranishnikov, and V. Schroeder, *Embedding of hyperbolic groups into products of binary trees*, Invent. Math. (2007).
- [C] H. Coxeter, *Regular honeycombs in hyperbolic space*, Twelve Geometric Essays (1968), Southern Univ. Press, Carbondale, II, 200-214.
- [D1] M. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Annals of math.(2) **117, no 2** (1983), 293-324.
- [D2] M. Davis, *A hyperbolic 4-manifold*, Proc. Amer. Math. Soc. **93** (1985), 325-328.

- [DJ] A. Dranishnikov and T. Januszkiewicz, *Every Coxeter group acts amenably on a compact space*. *Proceedings of the 1999 Topology and Dynamics Conference (Salt Lake City, UT)*, Topology Proceedings **24** (1999), 135-141 (arXiv:math.GR/0402398).
- [DS] A. Dranishnikov and V. Schroeder, *Embedding of hyperbolic Coxeter groups into products of binary trees and aperiodic tilings*, Preprint MPIM2005-50 (2005).
- [GS] C. Goodman-Strauss, *A strongly aperiodic set of tiles in the hyperbolic plane*, Invent. Math. **159** (2005), 119-132.
- [HM] G.A. Hedlund, M. Morse, *Unending chess, symbolic dynamics and a problem in semigroup*, Duke Math. J. **11** (1944), 1-7.
- [MM] G.A. Margulis, S. Mozes, *Aperiodic tilings of the hyperbolic plane by convex polygons*, Israel J. Math. **107** (1998), 319-325.
- [Mor] M. Morse, *Recurrent geodesics on surface of negative curvature*, Trans. Amer. Math. Soc. **22** (1921), 84-100.
- [Moz] S. Mozes, *Aperiodic tilings*, Invent. Math. **128 no. 3** (1997), 603-611.
- [SW] L. Sadun and R.F. Williams, *Tiling spaces are Cantor set fiber bundles*, Ergod. Theory Dyn. Syst. **23** (2003), 307-316.
- [PV] L. Potyagailo, E. Vinberg, *On right-angled reflection groups in hyperbolic spaces*, Comment. Math. Helv. **80 no. 1** (2005), 63-73.
- [S] L. Sadun, *Tiling spaces are inverse limits*, J. Math. Phys. **44 no. 11** (2003), 5410-5411.
- [Th] A. Thue, *Über unendliche Zeichenreihen*, Norske vid Selsk Skr. Mat. Nat. Kl. **7** (1906), 1-22.
- [Us] V.V. Uspenskii, *private communication*.
- [V] E.B. Vinberg, *Absence of crystallographic groups of reflections in Lobachevsky spaces of large dimension.*, Trudy Moskov. Mat. Obshch **47** (1984), 68-102.
- [VS] E.B. Vinberg and O.V. Shvartsman, *Discrete groups of motions of spaces of constant curvature. Geometry, II*, Encyclopaedia Math. Sci **29** (1993).

UNIVERSITY OF FLORIDA, DEPARTMENT OF MATHEMATICS, P.O. BOX 118105, 358 LITTLE HALL, GAINESVILLE, FL 32611-8105, USA; UNIVERSITY OF ZURICH, DEPARTMENT OF MATHEMATICS, WINTERTHURERSTRASSE 190, CH-8057 ZURICH, SWITZERLAND

*E-mail address:* dranish@math.ufl.edu; vschroed@math.unizh.ch