**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2022

Coping under stress: Prefrontal control predicts stress burden during the COVID-19 crisis

Monninger, Maximilian ; Pollok, Tania M ; Aggensteiner, Pascal-M ; Kaiser, Anna ; Reinhard, Iris ; Hermann, Andrea ; Meyer-Lindenberg, Andreas ; Brandeis, Daniel ; Banaschewski, Tobias ; Holz, Nathalie E

Abstract: The coronavirus (COVID-19) pandemic has confronted millions of people around the world with an unprecedented stressor, affecting physical and mental health. Accumulating evidence suggests that emotional and cognitive self-regulation is particularly needed to effectively cope with stress. Therefore, we investigated the predictive value of affective and inhibitory prefrontal control for stress burden during the COVID-19 crisis. Physical and mental health burden were assessed using an online survey, which was administered to 104 participants of an ongoing at-risk birth cohort during the first wave in April 2020. Two follow-ups were carried out during the pandemic, one capturing the relaxation during summer and the other the beginning of the second wave of the crisis. Prefrontal activity during emotion regulation and inhibitory control were assessed prior to the COVID-19 crisis. Increased inferior frontal gyrus activity during emotion regulation predicted lower stress burden at the beginning of the first and the second wave of the crisis. In contrast, inferior and middle frontal gyrus activity during inhibitory control predicted effective coping only during the summer, when infection rates decreased but stress burden remained unchanged. These findings remained significant when controlling for sociodemographic and clinical confounders such as stressful life events prior to the crisis or current psychopathology. We demonstrate that differential stress-buffering effects are predicted by the neural underpinnings of emotion regulation and cognitive regulation at different stages during the pandemic. These findings may inform future prevention strategies to foster stress coping in unforeseen situations.

DOI: <https://doi.org/10.1016/j.euroneuro.2021.11.007>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-215488>

Journal Article

Published Version

The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Monninger, Maximilian; Pollok, Tania M; Aggensteiner, Pascal-M; Kaiser, Anna; Reinhard, Iris; Hermann, Andrea; Meyer-Lindenberg, Andreas; Brandeis, Daniel; Banaschewski, Tobias; Holz, Nathalie

E (2022). Coping under stress: Prefrontal control predicts stress burden during the COVID-19 crisis. *European Neuropsychopharmacology*, 56:13-23.
DOI: <https://doi.org/10.1016/j.euroneuro.2021.11.007>

Coping under stress: Prefrontal control predicts stress burden during the COVID-19 crisis

Maximilian Monninger^a, Tania M. Pollok^a,
 Pascal-M. Aggensteiner^a, Anna Kaiser^a, Iris Reinhard^b,
 Andrea Hermann^{c,d,e}, Andreas Meyer-Lindenberg^f,
 Daniel Brandeis^{a,g,h,i}, Tobias Banaschewski^a, Nathalie E. Holz^{a,*}

^aDepartment of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, J5, Mannheim 68159, Germany

^bDepartment of Biostatistics, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, J5, Mannheim 68159, Germany

^cDepartment of Psychotherapy and Systems Neuroscience, Justus Liebig University, Giessen, Germany

^dBender Institute of Neuroimaging, Justus Liebig University Giessen, Germany

^eCenter for Mind, Brain and Behavior, Phillips University Marburg and Justus Liebig University Giessen, Germany

^fDepartment of Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim / Heidelberg University, J5, Mannheim 68159, Germany

^gDepartment of Child and Adolescent Psychiatry and Psychotherapy, Psychiatric Hospital, University of Zurich, Neumünsterallee 9, Zurich 8032, Switzerland

^hCenter for Integrative Human Physiology, University of Zurich, Winterthurerstr. 190, Zurich 8057, Switzerland

ⁱNeuroscience Center Zurich, University of Zurich and ETH Zurich, Winterthurerstr. 190, Zurich 8057, Switzerland

Received 17 August 2021; received in revised form 11 November 2021; accepted 15 November 2021

Available online xxx

KEYWORDS

Covid-19;
 Stress;
 fMRI;

Abstract

The coronavirus (COVID-19) pandemic has confronted millions of people around the world with an unprecedented stressor, affecting physical and mental health. Accumulating evidence suggests that emotional and cognitive self-regulation is particularly needed to effectively cope

* Corresponding author at: Department of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, J 5, 68159 Mannheim, Germany.

E-mail address: nathalie.holz@zi-mannheim.de (N.E. Holz).

Longitudinal study;
Affective control;
Cognitive control

with stress. Therefore, we investigated the predictive value of affective and inhibitory prefrontal control for stress burden during the COVID-19 crisis. Physical and mental health burden were assessed using an online survey, which was administered to 104 participants of an ongoing at-risk birth cohort during the first wave in April 2020. Two follow-ups were carried out during the pandemic, one capturing the relaxation during summer and the other the beginning of the second wave of the crisis. Prefrontal activity during emotion regulation and inhibitory control were assessed prior to the COVID-19 crisis. Increased inferior frontal gyrus activity during emotion regulation predicted lower stress burden at the beginning of the first and the second wave of the crisis. In contrast, inferior and middle frontal gyrus activity during inhibitory control predicted effective coping only during the summer, when infection rates decreased but stress burden remained unchanged. These findings remained significant when controlling for sociodemographic and clinical confounders such as stressful life events prior to the crisis or current psychopathology. We demonstrate that differential stress-buffering effects are predicted by the neural underpinnings of emotion regulation and cognitive regulation at different stages during the pandemic. These findings may inform future prevention strategies to foster stress coping in unforeseen situations.

© 2021 The Author(s). Published by Elsevier B.V.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1. Introduction

The coronavirus (COVID-19) pandemic is related to physical health impairments and dramatic changes to everyday life for hundreds of millions of people around the world. With the social contact restrictions beginning in March 2020 in Germany, nearly all schools were closed overnight, working environments changed radically, and social life was restricted probably as never before. Early studies investigating the impact of these initial “first-level threats” (imminent threat) on mental health burden reported elevated levels of perceived stress, anxiety, and depressive symptoms during the first wave of the COVID-19 pandemic (Benke et al., 2020a, b). While sustained socioeconomic “second-level threats” continued throughout the time when infection rates lowered during the summer, renewed first-level health threats emerged at the beginning of the second wave of the pandemic in late 2020.

Appropriate coping in response to these time-dependent challenges in the face of an ongoing crisis like the COVID-19 pandemic requires the balanced use of emotional regulation and cognitive control strategies (Feder et al., 2019). Indeed, so far, several studies investigated the effect of emotion regulation strategies and cognitive control on well-being and mental health during the pandemic. Findings indicate that adaptive emotion regulation strategies predicted increased well-being, whereas maladaptive emotion regulation strategies and deficits in cognitive control predicted higher levels of anxiety, greater depressive symptoms, reduced quality of life, risky health behavior, and diminished well-being during the pandemic (Appelhans et al., 2021; Breau et al., 2021; Brehl et al., 2021; Low et al., 2020; Panayiotou et al., 2021; Weissman et al., 2021; Yang et al., 2020). While affective control, such as reappraisal, involves the adaptive modulation of emotions in response to unpleasant stimuli (Gross, 2015; Holley et al., 2017), cognitive control includes the inhibition of inappropriate or ineffective behavior (Aron, 2007). Both are part of the executive function system (Holley et al.,

2017) and reflected mainly in the prefrontal cortex (PFC; Davidson et al., 2000; Ochsner et al., 2002). Affective control strategies are associated with an increased activity primarily in the inferior frontal gyrus (IFG), the middle frontal gyrus (MFG), and the ventro-, medial- and dorsolateral PFC (Gross, 2015; Kanske et al., 2011; Ochsner et al., 2002). Cognitive inhibitory control is linked to activity in the fronto-basal-ganglia circuit including the MFG and the IFG (Aron, 2007, 2003, 2004; Verbruggen and Logan, 2008), indicating shared neuronal underpinnings within the IFG and MFG.

The present longitudinal study investigated the predictive value of prefrontal activity during an emotion regulation and an inhibitory control task assessed prior to the COVID-19 pandemic for stress burden during the COVID-19 crisis in 104 participants of an ongoing at-risk cohort study following participants since birth. Approximately four weeks after the initial lockdown in Germany in March 2020, participants rated their current stress burden in an online survey (baseline during the first wave). Six to eight weeks later (first follow-up) as well as six months later (at the beginning of the second wave of the pandemic; second follow-up), participants completed the same questionnaire. We expected that higher affective and cognitive control predict lower stress burden during the COVID-19 pandemic. Given the ongoing uncertainties about the course and duration of the pandemic and its unprecedented nature, we had no specific a-priori hypotheses about how neural activity predict stress burden at different stages of the pandemic. However, based on previous findings (Brehl et al., 2021; Weissman et al., 2021; Yang et al., 2020), we speculate that the ability to control emotions in order to overcome perceived stress burden would be highly necessary at the beginning of the first wave of the COVID-19 crisis, when participants were confronted with direct threat due to high infection and death numbers (first-level threat). By contrast and based on previous findings assessed during the course of the pandemic (Appelhans et al., 2021), we anticipated that cognitive control would become increasingly important

in order to cope with the ongoing uncertainties during the pandemic (second-level threat).

negative affect. The participants were carefully instructed to either view the depicted scenario from a more positive or at least a less negative point of view (e.g. a person in jail might be a famous

2. Experimental procedures

2.1. Sample

The present investigation was conducted within the Mannheim Study of Children at Risk, an ongoing longitudinal study of the long-term outcomes of early psychosocial and biological risk factors following participants since birth (Laucht et al., 2000). The initial sample consisted of 384 children born between 1986 and 1988 in the Rhine-Neckar region of Germany. Infants were recruited from two obstetric and six children's hospitals and were included according to a two-factorial design intended to enrich and control the risk status of the sample (distribution in the current sample: 37 (35.6%) participants without psychosocial risk, 35 (33.7%) with low psychosocial risk, and 32 (30.8%) with high psychosocial risk at birth).

Up to the time of the lockdown on March 23rd 2020, 165 participants completed an ongoing assessment wave, including a comprehensive questionnaire package on physical and mental health, a structured clinical interview (SCID-5-CV German version; (Beesdo-Baum et al., 2019)), and a magnetic resonance imaging (MRI) session, including an emotion regulation and stop-signal task (see below). In April 2020 (during the first wave), two months later (first follow up during summer), and six months later (second follow-up in December), participants were invited to take part in an online COVID-19 questionnaire, adapted and modified from the Coronavirus Health and Impact Survey (CRISIS, (Nikolaidis et al., 2020)). 133 (80.61%) participants completed the COVID-19 baseline assessment, of whom 128 (96.24%) also responded to the first follow-up questionnaire and 116 (87.21%) to the second follow-up questionnaire. Functional MRI data were available for 104 participants for the baseline and first follow-up questionnaire, and for 95 participants for the second follow-up questionnaire. The study was approved by the Ethics Committee of the University Heidelberg, Germany, written informed consent from all participants was obtained, and participants were financially compensated.

2.2. Stress burden during COVID-19

To assess the impact and threat of COVID-19 on physical and mental health, we used four items ('The impact of COVID-19 on my physical health is?', 'COVID-19 is a potential threat to my physical health') and mental health ('The impact of COVID-19 on my mental health is?', 'COVID-19 is a potential threat to my mental health') rated on a 10-point Likert scale (0: completely disagree; 10: completely agree). We calculated a sum score for each assessment (baseline, first follow-up, second follow-up).

2.3. Emotion regulation

2.3.1. Emotion regulation task

Prior to the COVID-19 pandemic, participants performed an adapted and modified version of an emotion regulation task (Hermann et al., 2017) during functional MRI. In brief, participants were asked to either watch aversive ('Look_{negative}') or neutral ('Look_{neutral}') pictures from the International Affective Picture System (IAPS; Lang et al., 1999) or to reappraise negative pictures ('Reappraisal'). In the reappraisal condition, participants were asked to use the strategy of reappraisal to decrease the intensity of their

Table 1 Results of the ROI analyses for the emotion regulation task and the stop-signal task.

		MNI Coordinates					
	ROI	k	P _{FWE-corr}	T	X	Y	Z
Emotion Regulation Task							
Left	IFG	509	<.001	10.46	−36	26	−1
Right	IFG	338	<.001	9.91	51	26	−1
Left	MFG	406	<.001	10.54	−42	2	53
Right	MFG	174	<.001	8.03	51	11	44
Stop-Signal Task							
Left	IFG	50	.001	4.76	−48	41	8
Right	IFG	34	<.001	5.23	51	38	14
Left	MFG	200	<.001	7.57	−27	29	50
Right	MFG	200	<.001	7.48	30	20	50

Note: FWE-corr = family-wise error corrected; IFG = Inferior frontal gyrus; k = Cluster size; MFG = Middle frontal gyrus; MNI = Montreal Neurological Institute.

actor) or to rationalize the presented picture (e.g. due to enormous advances in modern medicine, a very premature baby may have an entirely normal life). During the neutral conditions, participants were instructed to simply watch the depicted scenarios without actively changing their emotional state evoked by the pictures.

The task consisted of a randomized block design, in which every block started with a jittered 3 s presentation of the instruction form (i.e. 'Look' or 'Reappraise'). Subsequently, participants viewed 20 s blocks of negative or neutral pictures (each picture was presented for 5 s without an interstimulus interval) according to the presented condition, and were asked to rate the intensity of currently perceived negative feelings on a 7-point Likert scale (1 = no negative feelings at all; 7 = extremely negative feelings) via a button press (max 4 s). After that, a white fixation cross on black background was presented during the inter-trial-interval up to a total block duration of 30 s. The total task comprised four blocks per condition (12 blocks in total) and lasted for 6 min 37 s. The blocks were arranged in four runs with a randomized presentation of all conditions within each run, leading to a maximum of two presentations of the same condition in succession.

2.4. Inhibitory control

2.4.1. Inhibitory control task

Participants completed the stop-signal task (SST) to assess cognitive inhibitory control during functional MRI (Rubia et al., 2003). Each trial began with a fixation cross (500 ms), which was followed by an arrow pointing to the left or right (go-signal). In the go-trials (75% of a total number of 160 trials), participants were instructed to react to the arrow as quickly and accurately as possible by pressing either the left or the right button according to the previously shown arrow. Stop-trials (25% of a total number of 160 trials) were designed as trials in which a go-signal was followed by an arrow pointing upwards (stop-signal). During the stop trials, participants were instructed to inhibit their response (yielding successful vs. unsuccessful inhibition). The delay between a go-signal and a stop-signal (stop-signal delay; SSD) started at 250 ms and was dynamically changed according to the participant's performance using an adaptive algorithm. The SSD latency increased by 50 ms whenever the participant correctly inhibited the previous response (max. 900 ms), making it more difficult to stop. In contrast, the SSD latency decreased by 50 ms whenever the previous stop-trial was incorrectly answered (min. 50 ms). Using this procedure, an approximately equal number of successful and unsuccessful stop-trials can

be achieved (inhibitory control in the present sample was on average 60.79% (SD = 12.68)). Moreover, we excluded participants with a negative stop-signal reaction time (SSRT). Prior to scanning, all participants were given clear instructions. Total scanning time was 6 min 37 s.

2.5. Functional mri data acquisition and preprocessing

Functional MRI data collection consisted of a localizer scan followed by a blood oxygen level-dependent (BOLD)-sensitive T2*-weighted echo-planar imaging (EPI) sequence and structural T1-weighted sequence using a 3T-scanner (PrismaFit; Siemens) with a standard 32-channel head coil. For functional imaging, a total of 186 vol with 36 slices covering the whole brain (matrix 64×64, resolution 3.0 × 3.0 × 3.0 mm with 1 mm gap, repetition time = 2100 ms, echo time = 35 ms, flip angle = 90°) were acquired for each task. The slices were inclined 20° from the anterior/posterior commissure level. The first 11 vol of the emotion regulation task and the first six volumes of the SST were discarded to allow longitudinal magnetization to reach equilibrium.

2.6. Functional MRI data analyses and statistical analyses

Statistical parametric mapping version 12 (SPM12) implemented in MATLAB R2017b was used to analyze functional MRI data. Preprocessing included slice time correction of the volumes to the first slice, realignment to correct for movement artifacts, co-registration of functional and anatomical data, spatial normalization to standard Montreal Neurological Institute (MNI) space, and spatial smoothing with a Gaussian filter of 8 mm full-width at half maximum. Motion parameters were examined to ensure head movement did not exceed 3 mm.

For the emotion regulation task, individual first-level contrasts based on onsets and durations of each condition were convolved with the canonical hemodynamic response function (HRF) in order to model the BOLD time course by using general linear models. Six motion parameters were included as regressors of no interest. For the group-level analysis, individual contrast images of the Reappraisal > Look_{neutral} condition were entered into a random-effects analysis to assess emotion regulation.

Similarly, for the SST, individual first-level contrasts for the onsets of the four event types (correct go-trials, incorrect go-trials,

successful inhibition, and unsuccessful inhibition) were modeled using delta functions convolved with the canonical HRF. Again, six motion estimation parameters were included as regressors of no interest. Second-level random effects analyses were modeled for the contrast of interest, i.e. successful inhibition was determined by contrasting successful stop-trials with unsuccessful inhibition and correct go-trials [successful inhibition > (unsuccessful inhibition - correct go-trials)]. Whole-brain differences in activation for the contrasts of interest for both tasks were estimated using t-tests at a family-wise error (FWE) corrected p-value of 0.05.

The IFG and the MFG were defined as regions of interest (ROIs) based on previous studies reporting a strong involvement in emotion regulation and inhibitory control, and their potential relevance in coping under stress (Berretz et al., 2021; Gavazzi et al., 2020; Golde et al., 2020; Rosenbaum et al., 2018). To extract predefined ROIs, anatomical masks implemented in the Wake Forest University (WFU) PickAtlas v2.4 (Maldjian et al., 2003) were used, where a $p < .05$ FWE correction (minimum of 10 adjacent voxels) was applied. Mean contrast values of each participant were extracted from the FWE-corrected significant clusters in the contrasts of interest (Table 1) and exported to SPSS Statistics 25. The predictive power of IFG and MFG activity during affective and inhibitory control for the impact of COVID-19 on mental and physical health was investigated using linear regression models controlling for all covariates (see below). Moreover, a conservative Bonferroni correction was applied for the hemispheres, two regions of interest (MFG and IFG), and for affective and cognitive control (α of 0.05 divided by 8 tests, resulting in $\alpha_{adj} = 0.0063$). Additional sensitivity analyses were performed to control for the robustness of significant effects.

2.7. Covariates

Several important covariates were added to the main analyses, including gender, psychosocial adversity and obstetric risk at birth, current diagnoses of a mental disorder (assessed prior to the pandemic), as well as current life events prior to and during the COVID-19 pandemic.

2.7.1. Psychosocial adversity

Psychosocial adversity was assessed using a standardized parent interview according to an enriched family adversity index (Rutter and Quinton, 1977) at the participants' age of 3 months. The interview comprised 11 items covering characteristics of the family environment, the parents, and the parents' partnership (e.g., presence of parental psychiatric disorders, overcrowding in the home, ongoing parental conflicts, or unwanted pregnancy) during a period of one year prior to the assessment. A sum score of psychosocial adversities was calculated by adding up the presence of all items.

2.7.2. Obstetric risk at birth

Obstetric adversity was assessed using a standardized parent interview conducted at the participants' age of 3 months. A sum score of obstetric risk factors was computed by adding up the presence of nine adverse conditions during pregnancy, delivery, and in the early postnatal phase, such as preterm birth or low birth weight (Laucht et al., 2000).

2.7.3. Life events

Life events were recorded using a modified version of the Munich Events List (MEL) (Maier-Diewald, 1983) within the online questionnaire at all three time points. The MEL covers several areas of acute and chronic, positive and negative stressors, including marriage, delivery of a child, but also negative health outcomes, illness of a relative or job loss. We adjusted the items depending on the time point, i.e., presence of life events between the regular assessment wave and prior to the COVID-19 crisis for the COVID-19 baseline

assessment, and life events between baseline assessment and first follow-up as well as between the first and second follow-up for both COVID-19 follow-up assessments, respectively.

2.8. Sensitivity analyses

Further sensitivity analyses were calculated to control for the robustness of the predictive value of the neural activity, including parenthood, household income, current work status, workplace changes, and whether the participants were critical workers for the COVID-19 response as additional covariates. To control for a confounding impact of a COVID-19 infection, participants were asked in a separate question whether they have been infected by the virus ($n = 1$).

3. Results

3.1. Descriptive data and changes of COVID-19 impact

Descriptive data are depicted in Table 2. There were significant differences for the reported physical and mental impact of the COVID-19 pandemic between the baseline assessment and the follow-up assessments ($F = 11.812$, $P < .001$). Post-hoc analyses revealed no significant differences between the baseline assessment and the first follow-up ($P = .894$), but significant differences emerged between the baseline and the second follow-up ($T = -4.306$, $P < .001$), and between the first and second follow-up ($T = -3.912$, $P < 0.001$), suggesting a continuous burden on mental and physical health since the beginning of the COVID-19 outbreak, with a further increased stress burden at the beginning of the second wave of the pandemic.

3.2. COVID-19-related stress burden at the baseline assessment

Higher right IFG activity during the emotion regulation task was associated with a lower stress burden at the baseline assessment ($\beta = -.281$, $T = -3.023$, $P = .003$, Fig. 1a), after controlling for gender and psychosocial and obstetric risk factors at birth, current mental disorder prior to the COVID-19 pandemic, and current life events. In contrast, left IFG and bilateral MFG activity during emotion regulation were not related to stress burden at the beginning of the COVID-19 crisis ($P > 0.121$). In addition, inhibition-related right MFG activity was negatively associated with the COVID-19 burden at the baseline assessment ($\beta = -.215$, $T = -2.220$, $P = .029$), although this was not significant after correction for multiple testing. No significant relationships emerged with regard to left MFG and IFG activity during inhibition.

3.3. COVID-19-related stress burden at the first follow-up assessment

During the first follow-up assessment in the summer, neither IFG nor MFG during emotion regulation was related to the COVID-19 impact (all P s > 0.61), suggesting that affective control did not predict stress burden at this point

Fig. 1 A. Association of right IFG activity and COVID-19 impairments at the baseline assessment. Higher affective control was related to decreased stress burden during the first wave of the COVID-19 pandemic. B. and C. Association of right IFG activity and right MFG activity and COVID-19 impairments at the first follow-up assessment during the summer. Higher cognitive activity was related to decreased stress burden caused by the COVID-19 pandemic. D. Association of right IFG activity and COVID-19 impairments at the second follow-up assessment. Higher affective control was related to decreased stress burden at the beginning of the second wave of the pandemic.

of time. In contrast, lower right IFG and right MFG activity during inhibitory control was associated with a higher COVID-19 stress burden at the first follow-up assessment, even when controlling for the above-mentioned covariates (right IFG: $\beta = -.348$, $T = -3.638$, $P < .001$; right MFG: $\beta = -.375$, $T = -3.884$, $P < .001$, Fig. 1b&c). Moreover, a negative association for left MFG activity emerged, which did not survive correction for multiple comparisons after including the above-mentioned covariates ($\beta = -.250$, $T = -2.582$, $P = .011$). Inhibition-related left IFG activity was not related to stress burden at the first follow-up ($P = .093$).

3.4. COVID-19-related stress burden at the second follow-up assessment

Higher right IFG activity during the emotion regulation task was associated with a lower stress burden at the second follow-up assessment ($\beta = -.327$, $T = -3.246$, $P = .002$, Fig. 1d), after controlling for the above-mentioned covariates. In contrast, left IFG and MFG activity during emotion regulation were not related to stress burden during the second wave of the COVID-19 pandemic (all P s > 0.086). Likewise, inhibition-related activity was not related to COVID-19 burden at the second follow-up (all P s > 0.105).

3.5. Sensitivity analyses

Separate subsequent sensitivity analyses were calculated including additional control variables such as parenthood, income, current work status, critical worker status, and workplace changes due to the COVID-19 pandemic. To control for a confounding impact of a COVID-19 infection, we performed an additional sensitivity analysis after excluding one participant, who reported a COVID-19 infection. None of the sensitivity analyses changed the results (Tables 3 and 4).

4. Discussion

To the best of our knowledge, this is the first study to report a differential predictive value of emotion- and inhibition-related prefrontal control for stress burden at the beginning of and during the COVID-19 crisis, respectively (Fig. 2).

Specifically, we found a significant negative relationship between right IFG activity during emotion regulation assessed prior to the COVID-19 pandemic and stress burden approximately four weeks after the lockdown in Germany. In light of unforeseen first-level threat entailing immediate danger to health and emotional challenges due to the dramatic, unprecedented social contact restrictions at the beginning of the crisis with high levels of emotional arousal, fear and the imminent feeling of loneliness (Bauerle et al., 2020; Brooks et al., 2020; Dubey et al.,

2020; Odriozola-Gonzalez et al., 2020), affective coping was particularly needed (Groarke et al., 2020; Restubog et al., 2020; Weissman et al., 2021). Our results support this assumption, indicating that participants who are characterized by higher affective control were able to cope better with these initial threats and uncertainties of an unprecedented stressor like the COVID-19 pandemic. At the first follow-up assessment, which occurred during the summer, the first-level threat decreased due to loosened restrictions and lower infection rates. Also, emotional habituation may occur if participants are confronted with the same stressor for a longer amount of time (Grissom and Bhatnagar, 2009). These reasons both may contribute to a decreased necessity for affective coping. Notably, towards the second wave of the pandemic, when infection rates increased dramatically even above those during the first wave and a second lockdown was imposed, affective coping was again required to face this immediate, even more intense, threat.

In contrast, we found an increased need for cognitive control strategies when facing second-level threat, i.e. socioeconomic uncertainties. Specifically, higher cognitive inhibition-related activity in the right MFG and IFG predicted lower COVID-19 distress only at the first follow-up (when restrictions had eased), but was unrelated to stress burden at the beginning of the first wave (baseline) and the second wave (second follow-up). While there was an immediate threat at the beginning of both the first and the second wave, the progression of the pandemic is characterized by ongoing socioeconomic challenges, uncertainties about one's own future financial and work capabilities, reduced working hours, and the length of the crisis (Cutler and Summers, 2020; Fegert et al., 2020). These sustained and secondary challenges (Pedrosa et al., 2020) require increased cognitive coping and flexibility over time, thus explaining the negative association of the MFG and IFG activity with stress burden at the first follow-up.

Given the unique prospective design of our study, we were able to demonstrate the strong predictive value of neural activity for stress coping during the pandemic, irrespective of the presence of several important factors which have previously been reported to affect stress coping. Specifically, the findings were robust against control for the presence of other current stressful life events (Undheim and Sund, 2017), socioeconomic disadvantages (McEwen and Gianaros, 2010), the occurrence of early life psychosocial adversities (Quinlan et al., 2017; Sheffler et al., 2019), or the presence of a mental disorder before the crisis (Joormann and Gotlib, 2010), thus highlighting the superior role of neural self-regulation for coping under stress.

Coping under stress has become a growing research field of interest, and is specifically important for promoting resilience (Feder et al., 2019; Feder et al., 2009). Broadly defined, resilience means staying healthy despite the exposure to stressful events, chronic adversities or traumatic experiences (Fletcher and Sarkar, 2013; Vinkers et al., 2020). Important personal factors promoting resilience encompass an individual's adaptive abilities, such as appropriate emotion regulation, inhibitory control, or cognitive flexibility, but also include an individual's social integration, a person's genetic make-up, or the individual neurobiological constitution (Campbell-Sills et al., 2006; Feder et al., 2019). While a variety of studies investigated the beneficial impact of adequate coping during stress and resilience either after a traumatic event or in response to an artificially induced acute stressor (Caston and Mauss, 2011; Cavanagh et al., 2014; Gloria and Steinhardt, 2016; Min et al., 2013; Roos et al., 2017; Shapero et al., 2019), predictive assertions on future coping with a naturally occurring stressor are lacking. Our results provide first hints that bridge this gap and provide important implications not only for affective and cognitive coping with potential further waves of increasing COVID-19 cases, for instance due to virus mutations around the world,

Table 3 Results of further sensitivity analyses using brain activity as main predictor to explain stress burden.

covariate	ROI	Task / Assessment	Beta	T	p-value
Parenthood	Right IFG	Emotion Regulation / Baseline	-0.259	-2.681	.009
	Right MFG	Stop-Signal Task / Follow-Up	-0.333	-3.428	.001
	Right IFG	Stop-Signal Task / Follow-Up	-0.313	-3.288	.001
	Right IFG	Emotion Regulation / 2nd Follow-Up	-0.325	-3.203	.002
Income	Right IFG	Emotion Regulation / Baseline	-0.288	-2.997	.003
	Right MFG	Stop-Signal Task / Follow-Up	-0.351	-3.574	.001
	Right IFG	Stop-Signal Task / Follow-Up	-0.294	-2.973	.004
	Right IFG	Emotion Regulation / 2nd Follow-Up	-0.339	-3.404	.001
Current work status	Right IFG	Emotion Regulation / Baseline	-0.275	-2.853	.005
	Right MFG	Stop-Signal Task / Follow-Up	-0.375	-3.866	<.001
	Right IFG	Stop-Signal Task / Follow-Up	-0.319	-3.268	.001
Critical worker status	Right IFG	Emotion Regulation / 2nd Follow-Up	-0.311	-3.182	.002
	Right IFG	Emotion Regulation / Baseline	-0.271	-2.858	.005
	Right MFG	Stop-Signal Task / Follow-Up	-0.367	-3.765	<.001
	Right IFG	Stop-Signal Task / Follow-Up	-0.306	-3.113	.002
Workplace changes	Right IFG	Emotion Regulation / 2nd Follow-Up	-0.324	-3.221	.002
	Right IFG	Emotion Regulation / Baseline	-0.271	2.803	.006
	Right MFG	Stop-Signal Task / Follow-Up	-0.363	-3.702	<.001
	Right IFG	Stop-Signal Task / Follow-Up	-0.311	-3.148	.002
	Right IFG	Emotion Regulation / 2nd Follow-Up	-0.323	-3.225	.002

Fig. 2 Timeline and proposed stress model at different stages during the COVID-19 pandemic. An initial high load of emotional distress requires adequate affective coping of the IFG at the beginning of the first and the second wave of the COVID-19 pandemic. Prolonged socioeconomic uncertainties involve cognitive control of the IFG and MFG to overcome these challenges during the ongoing crisis.

Note: Weekly statistics on COVID-19 cases are freely available from the Robert Koch Institute, Berlin, Germany.

but also with unprecedented stressful events in general. Specifically, we argue that there is a particular need for prevention strategies which aim at improving the individual's affective and cognitive coping capacities in response to unforeseen events with a high stress load. As such, we suggest that stepped care neuromodulation might mitigate stress burden via gaining self-control over neural affective and cognitive regulation. Therefore, enhancing neuroplasticity through different methods, such as mindfulness or even neurofeedback in emotion regulation areas might offer the potential to acquire and foster primary preventive self-regulation skills when initially encountering stressful events, whereas learning self-control over cognitive regulation activity might rather serve as a secondary prevention tool during ongoing stressful events. In addition, ecological momentary assessment (EMA) might be a promising, affordable tool to further foster affective and cognitive control

strategies in response to real-time, real-life stressors on an individual's own smartphone; this could be addressed by future intervention studies.

Some limitations of our findings need to be addressed. Only a third of our initial sample was able to take part in all COVID-19 assessments. Moreover, despite the quick start, we were unable to include the first days of the lockdown, which might have exerted the greatest effects on perceived stress. However, since we were particularly interested in coping with stress burden caused by the crisis, this time period appears to be appropriate to capture stress burden given that previous studies found no differences when comparing well-being, anxiety and depression in the initial phase of the pandemic to four weeks later (Vindgaard and Benros, 2020). Finally, we concentrated on a more unspecific contrast of affective control, which not only captures brain activity related to emotion regulation, but also to neg-

Table 4 Additional sensitivity analyses after excluding one participant with COVID-19 infection.

ROI	Task / Assessment	Beta	T	p-value
Right IFG	Emotion Regulation / Baseline	-0.274	-2.916	.004
Right MFG	Stop-Signal Task / Follow-Up	-0.373	-3.833	<.001
Right IFG	Stop-Signal Task / Follow-Up	-0.343	-3.542	.001
Right IFG	Emotion Regulation / 2nd Follow-Up	-0.318	-3.120	.002

Table 2 Sample description and descriptive data.

Total <i>n</i> = 104			
	<i>N</i>	%	
Sex (female)	58	55.8	
Critical worker status	39	37.5	
Full-time employment	64	61.5	
Workplace changes due to COVID-19	69	66.3	
Parenthood	45	43.3	
Current mental disorder	28	26.9	
	<i>Mean</i>	<i>SD</i>	<i>range</i>
Age	33.31	0.54	32.25 - 34.25
COVID-19 impact (baseline)	12.13	5.92	4 - 25
COVID-19 impact (1st follow-up)	12.22	6.09	4 - 27
COVID-19 impact (2nd follow-up) ^A	14.39	5.68	4 - 26
Psychosocial risk factors at birth	1.80	1.90	0 - 7
Obstetric risk factors at birth	0.83	0.90	0 - 4
Income (in €)	4110	1816	450 - 8789
Life events prior to COVID-19 ^B	4.41	4.79	0 - 26
Life events during COVID-19 ^B (1st follow-up)	1.37	1.86	0 - 7
Life events during COVID-19 ^{A,B} (2nd follow-up)	1.76	1.75	0 - 8

Note: A. Sample size comprises participants who took part in the fMRI assessments and all follow-up questionnaires (*n* = 95). B. Higher scores of life events prior to COVID-19 resulted from retrospective assessment of a longer time period (i.e., time between regular assessment wave and baseline COVID-19 assessment) compared to life events during COVID-19 (time between baseline COVID-19 assessment and follow-up assessments).

active emotion processing. While this is not a standard approach, the contrast applied here might be more reflective of the affective challenges provided by the COVID-19 pandemic.

In the framework of an ongoing longitudinal study following at-risk participants since birth, we thus provide first evidence for the predictive value of neural underpinnings of emotion regulation during the first and second lockdown and cognitive regulation during the COVID-19 pandemic between both waves. These findings may inform future prevention strategies seeking to foster stress coping in unforeseen situations.

Contributors

M.M. conducted the MRI data collection, analyzed all data and wrote the paper. T.M.P. collected and analyzed the data and reviewed the paper. P.-M.A. collected the data and re-

viewed the paper. A.K. reviewed the paper. I.R. supervised all data analyses and reviewed the paper. A.H. contributed to the design of the study and reviewed the paper. A.M.-L. and D.B. contributed to the design of the study, supervised data analyses and reviewed the paper. T.B. obtained funding for the study, supervised data analyses and reviewed the paper. N.E.H. obtained funding for the study, designed the study, supervised all data analyses and wrote the paper.

Declaration of Competing Interest

T.B. served in an advisory or consultancy role for ADHS digital, Infectopharm, Lundbeck, Medice, Neurim Pharmaceuticals, Oberberg GmbH, Roche, and Takeda. He received conference support or speaker's fee by Medice and Takeda. He received royalties from Hogrefe, Kohlhammer, CIP Medien, Oxford University Press; the present work is unrelated to these relationships. D.B. serves as an unpaid scientific consultant for an EU-funded neurofeedback trial unrelated to the present work. A.M.-L. has received consultant fees from the American Association for the Advancement of Science, Atheneum Partners, Blueprint Partnership, Boehringer Ingelheim, Daimler und Benz Stiftung, Elsevier, F. Hoffmann-La Roche, ICARE Schizophrenia, K. G. Jebsen Foundation, L.E.K Consulting, Lundbeck International Foundation (LINF), R. Adamczak, Roche Pharma, Science Foundation, Sumitomo Dainippon Pharma, Synapsis Foundation - Alzheimer Research Switzerland, System Analytics, and has received lectures fees including travel fees from Boehringer Ingelheim, Fama Public Relations, Institut d'investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Janssen-Cilag, Klinikum Christophsbad, Göppingen, Lilly Deutschland, Luzerner Psychiatrie, LVR Klinikum Düsseldorf, LWL Psychiatrie Verbund Westfalen-Lippe, Otsuka Pharmaceuticals, Reunions i Ciencia S. L., Spanish Society of Psychiatry, Südwestrundfunk Fernsehen, Stern TV, and Vitos Klinikum Kurhessen. All other authors declare that they have no conflicts of interest.

Role of Funding Source

NH and TB gratefully acknowledge grant support from the [German Research Foundation](#) (grant numbers DFG HO 5674/2-1, GRK2350/1) and the Ministry of Science, Research and the Arts of the State of Baden-Württemberg, Germany (Sonderfördermaßnahme SARS CoV-2 pandemic to NH). The funding source had no role in study design, collection, analysis, and interpretation of data, in the writing of the report, and in the decision to submit the article for publication.

Acknowledgements

The authors gratefully thank Manfred Laucht (1946–2020), who was one of the founders of the Mannheim Study of Children at Risk and who continuously acted as an inspiring and supporting mentor giving impulses for innovative research projects. M.M. and N.H. had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. The authors thank

Sibylle Heinzel, Rafaela Gehr, Christin Loebel and Cäcilia Pracht for conducting and supporting the assessments.

References

- Appelhans, B.M., Thomas, A.S., Roisman, G.I., Booth-LaForce, C., Bleil, M.E., 2021. Preexisting executive function deficits and change in health behaviors during the COVID-19 pandemic. *Int. J. Behav. Med.* doi:[10.1007/s12529-021-09974-0](https://doi.org/10.1007/s12529-021-09974-0).
- Aron, A.R., 2007. The neural basis of inhibition in cognitive control. *Neuroscientist* 13 (3), 214–228. doi:[10.1177/1073858407299288](https://doi.org/10.1177/1073858407299288).
- Aron, A.R., Fletcher, P.C., Bullmore, E.T., Sahakian, B.J., Robbins, T.W., 2003. Stop-signal inhibition disrupted by damage to right inferior frontal gyrus in humans. *Nat. Neurosci.* 6 (2), 115–116. doi:[10.1038/nn1003](https://doi.org/10.1038/nn1003).
- Aron, A.R., Robbins, T.W., Poldrack, R.A., 2004. Inhibition and the right inferior frontal cortex. *Trends Cogn. Sci.* 8 (4), 170–177. doi:[10.1016/j.tics.2004.02.010](https://doi.org/10.1016/j.tics.2004.02.010).
- Bauerle, A., Teufel, M., Musche, V., Weismuller, B., Kohler, H., Hetkamp, M., ... Skoda, E.M., 2020. Increased generalized anxiety, depression and distress during the COVID-19 pandemic: a cross-sectional study in Germany. *J. Public Health* 42 (4), 672–678. doi:[10.1093/pubmed/fdaa106](https://doi.org/10.1093/pubmed/fdaa106).
- Beesdo-Baum, K., Zaudig, M., Wittchen, H.-U., 2019. *SCID-5-CV : Strukturiertes Klinisches Interview für DSM-5-Störungen - Klinische Version*. Hogrefe, Göttingen In (1. Auflage ed.).
- Benke, C., Autenrieth, L.K., Asselmann, E., Pané-Farré, C.A., 2020a. Lockdown, quarantine measures, and social distancing: associations with depression, anxiety and distress at the beginning of the COVID-19 pandemic among adults from Germany. *Psychiatry Res.* 293. doi:[10.1016/j.psychres.2020.113462](https://doi.org/10.1016/j.psychres.2020.113462).
- Benke, C., Autenrieth, L.K., Asselmann, E., Pané-Farré, C.A., 2020b. Stay-at-home orders due to the COVID-19 pandemic are associated with elevated depression and anxiety in younger, but not older adults: results from a nationwide community sample of adults from Germany. *Psychol. Med.* 1–2. doi:[10.1017/S0033291720003438](https://doi.org/10.1017/S0033291720003438).
- Berretz, G., Packheiser, J., Kumsta, R., Wolf, O.T., Ocklenburg, S., 2021. The brain under stress—a systematic review and activation likelihood estimation meta-analysis of changes in BOLD signal associated with acute stress exposure. *Neurosci. Biobehav. Rev.* 124, 89–99. doi:[10.1016/j.neubiorev.2021.01.001](https://doi.org/10.1016/j.neubiorev.2021.01.001).
- Breaux, R., Dvorsky, M.R., Marsh, N.P., Green, C.D., Cash, A.R., Shroff, D.M., ... Becker, S.P., 2021. Prospective impact of COVID-19 on mental health functioning in adolescents with and without ADHD: protective role of emotion regulation abilities. *J. Child Psychol. Psychiatry* doi:[10.1111/jcpp.13382](https://doi.org/10.1111/jcpp.13382).
- Brehl, A.-K., Schene, A., Kohn, N., Fernández, G., 2021. Maladaptive emotion regulation strategies in a vulnerable population predict increased anxiety during the Covid-19 pandemic: a pseudo-prospective study. *J. Affect. Disord. Rep.* 4. doi:[10.1016/j.jadr.2021.100113](https://doi.org/10.1016/j.jadr.2021.100113).
- Brooks, S.K., Webster, R.K., Smith, L.E., Woodland, L., Wessely, S., Greenberg, N., Rubin, G.J., 2020. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 395 (10227), 912–920. doi:[10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8).
- Campbell-Sills, L., Cohan, S.L., Stein, M.B., 2006. Relationship of resilience to personality, coping, and psychiatric symptoms in young adults. *Behav. Res. Ther.* 44 (4), 585–599. doi:[10.1016/j.brat.2005.05.001](https://doi.org/10.1016/j.brat.2005.05.001).
- Caston, A., Mauss, I., 2011. Resilience in the face of stress: emotion regulation as a protective factor. *Resil. Ment. Health* doi:[10.1017/CBO9780511994791.004](https://doi.org/10.1017/CBO9780511994791.004).
- Cavanagh, S.R., Fitzgerald, E.J., Urry, H.L., 2014. Emotion reactivity and regulation are associated with psychological functioning following the 2011 earthquake, tsunami, and nuclear crisis in Japan. *Emotion* 14 (2), 235–240. doi:[10.1037/a0035422](https://doi.org/10.1037/a0035422).
- Cutler, D.M., Summers, L.H., 2020. The COVID-19 Pandemic and the \$16 Trillion Virus. *JAMA* 324 (15), 1495–1496. doi:[10.1001/jama.2020.19759](https://doi.org/10.1001/jama.2020.19759).
- Davidson, R.J., Putnam, K.M., Larson, C.L., 2000. Dysfunction in the neural circuitry of emotion regulation—a possible prelude to violence. *Science* 289, 591–594. doi:[10.1126/science.289.5479.591](https://doi.org/10.1126/science.289.5479.591).
- Dubey, S., Biswas, P., Ghosh, R., Chatterjee, S., Dubey, M.J., Chatterjee, S., ... Lavie, C.J., 2020. Psychosocial impact of COVID-19. *Diabetes Metab. Syndr.* 14 (5), 779–788. doi:[10.1016/j.dsx.2020.05.035](https://doi.org/10.1016/j.dsx.2020.05.035).
- Feder, A., Fred-Torres, S., Southwick, S.M., Charney, D.S., 2019. The biology of human resilience: opportunities for enhancing resilience across the life span. *Biol. Psychiatry* 86 (6), 443–453. doi:[10.1016/j.biopsych.2019.07.012](https://doi.org/10.1016/j.biopsych.2019.07.012).
- Feder, A., Nestler, E.J., Charney, D.S., 2009. Psychobiology and molecular genetics of resilience. *Nat. Rev. Neurosci.* 10 (6), 446–457. doi:[10.1038/nrn2649](https://doi.org/10.1038/nrn2649).
- Fegert, J.M., Vitiello, B., Plener, P.L., Clemens, V., 2020. Challenges and burden of the Coronavirus 2019 (COVID-19) pandemic for child and adolescent mental health: a narrative review to highlight clinical and research needs in the acute phase and the long return to normality. *Child Adolesc. Psychiatry Ment. Health* 14, 20. doi:[10.1186/s13034-020-00329-3](https://doi.org/10.1186/s13034-020-00329-3).
- Fletcher, D., Sarkar, M., 2013. Psychological resilience: a review and critique of definitions, concepts, and theory. *Eur Psychol* 18 (1), 12–23. doi:[10.1027/1016-9040/a000124](https://doi.org/10.1027/1016-9040/a000124).
- Gavazzi, G., Giovannelli, F., Curro, T., Mascalchi, M., Viggiano, M.P., 2020. Contiguity of proactive and reactive inhibitory brain areas: a cognitive model based on ALE meta-analyses. *Brain Imaging Behav.* doi:[10.1007/s11682-020-00369-5](https://doi.org/10.1007/s11682-020-00369-5).
- Gloria, C.T., Steinhardt, M.A., 2016. Relationships among positive emotions, coping, resilience and mental health. *Stress Health* 32 (2), 145–156. doi:[10.1002/smi.2589](https://doi.org/10.1002/smi.2589).
- Golde, S., Wingenfeld, K., Riepenhausen, A., Schroter, N., Fleischer, J., Prussner, J., ... Otte, C., 2020. Healthy women with severe early life trauma show altered neural facilitation of emotion inhibition under acute stress. *Psychol. Med.* 50 (12), 2075–2084. doi:[10.1017/S0033291719002198](https://doi.org/10.1017/S0033291719002198).
- Grissom, N., Bhatnagar, S., 2009. Habituation to repeated stress: get used to it. *Neurobiol. Learn. Mem.* 92 (2), 215–224. doi:[10.1016/j.nlm.2008.07.001](https://doi.org/10.1016/j.nlm.2008.07.001).
- Groarke, J.M., Berry, E., Graham-Wisener, L., McKenna-Plumley, P.E., McGlinchey, E., Armour, C., 2020. Loneliness in the UK during the COVID-19 pandemic: cross-sectional results from the COVID-19 psychological wellbeing study. *PLoS One* 15 (9), e0239698. doi:[10.1371/journal.pone.0239698](https://doi.org/10.1371/journal.pone.0239698).
- Gross, J.J., 2015. Emotion Regulation: current Status and Future Prospects. *Psychol. Inq.* 26, 1–26. doi:[10.1080/1047840x.2014.940781](https://doi.org/10.1080/1047840x.2014.940781).
- Hermann, A., Kress, L., Stark, R., 2017. Neural correlates of immediate and prolonged effects of cognitive reappraisal and distraction on emotional experience. *Brain Imaging Behav.* 11 (5), 1227–1237. doi:[10.1007/s11682-016-9603-9](https://doi.org/10.1007/s11682-016-9603-9).
- Holley, S.R., Ewing, S.T., Stiver, J.T., Bloch, L., 2017. The relationship between emotion regulation, executive functioning, and aggressive behaviors. *J. Interpers. Violence* 32 (11), 1692–1707. doi:[10.1177/0886260515592619](https://doi.org/10.1177/0886260515592619).
- Joormann, J., Gotlib, I.H., 2010. Emotion regulation in depression: relation to cognitive inhibition. *Cogn. Emot.* 24 (2), 281–298. doi:[10.1080/02699930903407948](https://doi.org/10.1080/02699930903407948).
- Kanske, P., Heissler, J., Schonfelder, S., Bongers, A., Wessa, M., 2011. How to regulate emotion? Neural networks for reappraisal and distraction. *Cereb. Cortex* 21 (6), 1379–1388. doi:[10.1093/cercor/bhq216](https://doi.org/10.1093/cercor/bhq216).
- Lang, P.J., Bradley, M.M., Cuthbert, B.N., 1999. *International af-*

- fective picture system (IAPS): instruction manual and affective ratings. The Center For Research in Psychophysiology. University of Florida.
- Laucht, M., Esser, G., Baving, L., Gerhold, M., Hoesch, I., Ihle, W., ... Schmidt, M.H., 2000. Behavioral sequelae of perinatal insults and early family adversity at 8 years of age. *J. Am. Acad. Child Adolesc. Psychiatry* 39 (10), 1229–1237.
- Low, R., Overall, N., Chang, V., Henderson, A., 2020. Emotion Regulation and Psychological and Physical Health during a Nationwide COVID-19 Lockdown.
- Maier-Diewald, W., 1983. *Die Münchner Ereignisliste: MEL*. Max-Planck-Institut für Psychiatrie.
- Maldjian, J.A., Laurienti, P.J., Kraft, R.A., Burdette, J.H., 2003. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *Neuroimage* 19 (3), 1233–1239. doi:10.1016/s1053-8119(03)00169-1.
- McEwen, B.S., Gianaros, P.J., 2010. Central role of the brain in stress and adaptation: links to socioeconomic status, health, and disease. *Ann. N. Y. Acad. Sci.* 1186, 190–222. doi:10.1111/j.1749-6632.2009.05331.x.
- Min, J.A., Yu, J.J., Lee, C.U., Chae, J.H., 2013. Cognitive emotion regulation strategies contributing to resilience in patients with depression and/or anxiety disorders. *Compr. Psychiatry* 54 (8), 1190–1197. doi:10.1016/j.comppsych.2013.05.008.
- Nikolaidis, A., Paksarian, D., Alexander, L., DeRosa, J., Dunn, J., Nielson, D.M., ... Merikangas, K.R. (2020). The Coronavirus Health and Impact Survey (CRISIS) reveals reproducible correlates of pandemic-related mood states across the Atlantic. *medRxiv*. doi:10.1101/2020.08.24.20181123
- Ochsner, K.N., Bunge, S.A., Gross, J.J., Gabrieli, J.D.E., 2002. Rethinking feelings: an fMRI study of the cognitive regulation of emotion. *J. Cogn. Neurosci.* 14 (8), 1215–1229. doi:10.4324/9780203496190.
- Odriozola-Gonzalez, P., Planchuelo-Gomez, A., Irujo, M.J., de Luis-Garcia, R., 2020. Psychological effects of the COVID-19 outbreak and lockdown among students and workers of a Spanish university. *Psychiatry Res.* 290, 113108. doi:10.1016/j.psychres.2020.113108.
- Panayiotou, G., Panteli, M., Leonidou, C., 2021. Coping with the invisible enemy: the role of emotion regulation and awareness in quality of life during the COVID-19 pandemic. *J. Contextual. Behav. Sci.* 19, 17–27. doi:10.1016/j.jcbs.2020.11.002.
- Pedrosa, A.L., Bitencourt, L., Froes, A.C.F., Cazumba, M.L.B., Campos, R.G.B., de Brito, S., Simoes, E.S.A.C., 2020. Emotional, Behavioral, and Psychological Impact of the COVID-19 Pandemic. *Front. Psychol.* 11, 566212. doi:10.3389/fpsyg.2020.566212.
- Quinlan, E.B., Cattrell, A., Jia, T., Artiges, E., Banaschewski, T., Barker, G., ... Consortium, I., 2017. Psychosocial stress and brain function in adolescent psychopathology. *Am. J. Psychiatry* 174 (8), 785–794. doi:10.1176/appi.ajp.2017.16040464.
- Restubog, S.L.D., Ocampo, A.C.G., Wang, L., 2020. Taking control amidst the chaos: emotion regulation during the COVID-19 pandemic. *J. Vocat. Behav.* 119. doi:10.1016/j.jvb.2020.103440, 103440–103440.
- Roos, L.E., Knight, E.L., Beauchamp, K.G., Berkman, E.T., Faraday, K., Hyslop, K., Fisher, P.A., 2017. Acute stress impairs inhibitory control based on individual differences in parasympathetic nervous system activity. *Biol. Psychol.* 125, 58–63. doi:10.1016/j.biopsycho.2017.03.004.
- Rosenbaum, D., Thomas, M., Hilsendegen, P., Metzger, F., Häußinger, F., Nuerk, H.-C., ... Ehlis, A.-C., 2018. Stress-related dysfunction of the right inferior frontal cortex in high ruminators: an fNIRS study. *Neuroimage: Clin.* 18. doi:10.1016/j.nicl.2018.02.022.
- Rubia, K., Smith, A.B., Brammer, M.J., Taylor, E., 2003. Right inferior prefrontal cortex mediates response inhibition while mesial prefrontal cortex is responsible for error detection. *Neuroimage* 20 (1), 351–358. doi:10.1016/s1053-8119(03)00275-1.
- Rutter, M., Quinton, D., 1977. Psychiatric disorder: ecological factors and concepts of causation. In: McGurk, M. (Ed.), *Ecological Factors in Human Development*. Noord-Holland, Amsterdam, pp. 173–187.
- Shapero, B.G., Stange, J.P., McArthur, B.A., Abramson, L.Y., Alloy, L.B., 2019. Cognitive reappraisal attenuates the association between depressive symptoms and emotional response to stress during adolescence. *Cogn. Emot.* 33 (3), 524–535. doi:10.1080/02699931.2018.1462148.

- Sheffler, J.L., Piazza, J.R., Quinn, J.M., Sachs-Ericsson, N.J., Stanley, I.H., 2019. Adverse childhood experiences and coping strategies: identifying pathways to resiliency in adulthood. *Anxiety Stress Coping* 32 (5), 594-609. doi:[10.1080/10615806.2019.1638699](https://doi.org/10.1080/10615806.2019.1638699).
- Undheim, A.M., Sund, A.M., 2017. Associations of stressful life events with coping strategies of 12-15-year-old Norwegian adolescents. *Eur. Child Adolesc. Psychiatry* 26 (8), 993-1003. doi:[10.1007/s00787-017-0979-x](https://doi.org/10.1007/s00787-017-0979-x).
- Verbruggen, F., Logan, G.D., 2008. Response inhibition in the stop-signal paradigm. *Trends Cogn. Sci.* 12 (11), 418-424. doi:[10.1016/j.tics.2008.07.005](https://doi.org/10.1016/j.tics.2008.07.005).
- Vindegaard, N., Benros, M.E., 2020. COVID-19 pandemic and mental health consequences: systematic review of the current evidence. *Brain Behav. Immun.* 89, 531-542. doi:[10.1016/j.bbi.2020.05.048](https://doi.org/10.1016/j.bbi.2020.05.048).
- Vinkers, C.H., van Amelsvoort, T., Bisson, J.I., Branchi, I., Cryan, J.F., Domschke, K., ... van der Wee, N.J.A., 2020. Stress resilience during the coronavirus pandemic. *Eur. Neuropsychopharmacol.* 35, 12-16. doi:[10.1016/j.euroneuro.2020.05.003](https://doi.org/10.1016/j.euroneuro.2020.05.003).
- Weissman, D., Rodman, A., Rosen, M., Kasperek, S., Mayes, M., Sheridan, M., ... Meltzoff, A., 2021. Contributions of emotion regulation and brain structure and function to adolescent internalizing problems and stress vulnerability during the COVID-19 pandemic: a longitudinal study. *Biol. Psychiatry Glob. Open Sci.* doi:[10.1016/j.bpsgos.2021.06.001](https://doi.org/10.1016/j.bpsgos.2021.06.001).
- Yang, Y., Liu, K., Li, S., Shu, M., 2020. Social media activities, emotion regulation strategies, and their interactions on people's mental health in COVID-19 pandemic. *Int. J. Environ. Res. Public Health* (23) 17. doi:[10.3390/ijerph17238931](https://doi.org/10.3390/ijerph17238931).