



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2000

A finiteness result for associated primes of local cohomology modules

Brodmann, M ; Faghani, A

DOI: <https://doi.org/10.1090/S0002-9939-00-05328-4>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-22074>

Journal Article

Originally published at:

Brodmann, M; Faghani, A (2000). A finiteness result for associated primes of local cohomology modules. Proceedings of the American Mathematical Society, 128(10):2851-2853.

DOI: <https://doi.org/10.1090/S0002-9939-00-05328-4>

A FINITENESS RESULT FOR ASSOCIATED PRIMES OF LOCAL COHOMOLOGY MODULES

M. P. BRODMANN AND A. LASHGARI FAGHANI

(Communicated by Wolmer V. Vasconcelos)

ABSTRACT. We show that the first non-finitely generated local cohomology module $H_{\mathfrak{a}}^i(M)$ of a finitely generated module M over a noetherian ring R with respect to an ideal $\mathfrak{a} \subseteq R$ has only finitely many associated primes.

1. INTRODUCTION

Apparently very little is known about the finiteness of the set $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ of associated primes of the local cohomology module $H_{\mathfrak{a}}^i(M)$ of a finitely generated module M over a noetherian ring R with respect to an ideal \mathfrak{a} of R . So, in [M-S] the authors ask whether the sets $\text{Ass}_R(\text{Ext}_R^i(R/\mathfrak{a}^n, M))$ become stable for $n \gg 0$. An affirmative answer to this would imply that the set $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is always finite. On the other hand it seems not to be known whether the set $\text{Ass}_R(H_{\mathfrak{a}}^2(M))$ is finite in general even if \mathfrak{a} is generated by two elements only. Moreover, if for two elements $a, b \in R$ the set $S_M(a, b) := \bigcup_{n \in \mathbb{N}} \text{Ass}_R(M/(a^n, b^n)M)$ is finite, then $\text{Ass}_R(H_{(a,b)}^2(M))$ is finite. But Katzman [K] has shown that $S_M(a, b)$ need not be finite. In [B-R-Sh] we have shown that $S_M(a, b)$ is finite under certain conditions. These conditions imply that $H_{(a,b)}^1(M)$ is finitely generated and so the resulting finiteness of $\text{Ass}_R(H_{(a,b)}^2(M))$ is not surprising, as $\text{Ass}_R(H_{\mathfrak{a}}^2(M))$ is finite whenever $H_{\mathfrak{a}}^1(M)$ is finitely generated (see [B-R-Sh, (2.4), (2.5)]).

Finally, let us mention that the finiteness of the sets $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is related to the *local-global-principle for finiteness dimensions* due to Faltings [F] (cf. [B-Sh, (9.6.2)]) and also to the open problem of whether such a principle holds for the annihilation of local cohomology modules (see [R], [B-R-Sh]).

The aim of this note is to show that the set $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is finite, whenever the modules $H_{\mathfrak{a}}^1(M), \dots, H_{\mathfrak{a}}^{i-1}(M)$ are finitely generated. This generalizes the corresponding result which is shown in [B-R-Sh] for the special case $i \leq 2$ and which was already mentioned above.

Throughout this note, let R be a noetherian ring, let $\mathfrak{a} \subseteq R$ be an ideal and let M be a finitely generated R -module. If $i \in \mathbb{N}_0$, we write $H_{\mathfrak{a}}^i(M)$ for the i -th local cohomology module of M with respect to the ideal \mathfrak{a} . For convenience we write

Received by the editors September 8, 1998 and, in revised form, November 9, 1998.

2000 *Mathematics Subject Classification*. Primary 13D45; Secondary 13E99.

Key words and phrases. Local cohomology, associated primes.

$H_{\mathfrak{a}}^j(M) = 0$, whenever j is a negative integer. For the unexplained terminology we refer to [B-Sh].

2. THE FINITENESS RESULTS

Proposition 2.1. *Let $i \in \mathbb{N}_0$ be such that $H_{\mathfrak{a}}^j(M)$ is finitely generated for all $j < i$ and let $N \subseteq H_{\mathfrak{a}}^i(M)$ be a finitely generated submodule. Then, the set $\text{Ass}_R(H_{\mathfrak{a}}^i(M)/N)$ is finite.*

Proof. We proceed by induction on i . The case $i = 0$ is obvious as $H_{\mathfrak{a}}^0(M)$ is finitely generated. So, let $i > 0$ and set $\overline{M} := M/\Gamma_{\mathfrak{a}}(M)$. As $H_{\mathfrak{a}}^0(\overline{M}) = 0$ and in view of the natural isomorphisms $H_{\mathfrak{a}}^k(\overline{M}) \cong H_{\mathfrak{a}}^k(M)$ for all $k \in \mathbb{N}$, we may replace M by \overline{M} and hence assume that $\Gamma_{\mathfrak{a}}(M) = 0$. We thus find an M -regular element $y \in \mathfrak{a}$. By our choice of N , there is some $n \in \mathbb{N}$ with $y^n N = 0$.

We set $x := y^n$ and apply cohomology to the exact sequence $0 \rightarrow M \xrightarrow{x} M \rightarrow M/xM \rightarrow 0$. It follows that $H_{\mathfrak{a}}^l(M/xM)$ is finitely generated for all $l < i - 1$. Moreover, we get the following commutative diagram with exact rows and columns in which δ is the connecting homomorphism and in which ε and ϱ are the natural maps:

$$\begin{array}{ccccccc}
 (*) & H_{\mathfrak{a}}^{i-1}(M) & \xrightarrow{\varepsilon} & H_{\mathfrak{a}}^{i-1}(M/xM) & \xrightarrow{\delta} & H_{\mathfrak{a}}^i(M) & \xrightarrow{x} & H_{\mathfrak{a}}^i(M) \\
 & & & \downarrow & & \downarrow \varrho & & \downarrow \parallel \\
 & 0 & \longrightarrow & H_{\mathfrak{a}}^{i-1}(M/xM)/\delta^{-1}(N) & \xrightarrow{\overline{\delta}} & H_{\mathfrak{a}}^i(M)/N & \xrightarrow{\overline{x}} & H_{\mathfrak{a}}^i(M) \\
 & & & \downarrow & & \downarrow & & \\
 & & & 0 & & 0 & &
 \end{array}$$

As $\text{Ker}(\delta) = \varepsilon(H_{\mathfrak{a}}^{i-1}(M))$ and N are both finitely generated, so is $\delta^{-1}(N)$. Therefore, by induction

$$T := H_{\mathfrak{a}}^{i-1}(M/xM)/\delta^{-1}(N)$$

has only finitely many associated primes. It thus suffices to show the inclusion

$$(**) \quad \text{Ass}_R(H_{\mathfrak{a}}^i(M)/N) \subseteq \text{Ass}_R(T) \cup \text{Ass}_R(N) .$$

So, let $\mathfrak{p} \in \text{Ass}_R(H_{\mathfrak{a}}^i(M)/N) \setminus \text{Ass}_R(T)$. With an appropriate $h \in H_{\mathfrak{a}}^i(M)$ we may write $\mathfrak{p} = N \underset{R}{:} h$, hence $\mathfrak{p} = 0 \underset{R}{:} \varrho(h)$. As $\mathfrak{p} \notin \text{Ass}_R(T)$, the last equality and the second row of $(*)$ show that $\mathfrak{p} \in \text{Ass}_R(\overline{x}(\varrho(h))R) = \text{Ass}_R(xhR)$. This allows us to write $\mathfrak{p} = 0 \underset{R}{:} xsh$ for some $s \in R$.

As xsh is annihilated by some power of x , we have $x \in \mathfrak{p}$. By our choice of h this means that $xsh \in N$. This implies that $\mathfrak{p} \in \text{Ass}_R(N)$ and hence proves the inclusion $(**)$ and thus our result. □

Now, the announced result follows easily.

Theorem 2.2. *Let $i \in \mathbb{N}_0$ be such that $H_{\mathfrak{a}}^j(M)$ is finitely generated for all $j < i$. Then the set $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is finite.*

Proof. Apply Proposition (2.1) with $N = 0$. □

Next, let us introduce the \mathfrak{a} -finiteness dimension of M (see [B-Sh, (9.1.3)]):

$$f_{\mathfrak{a}}(M) := \min\{j \in \mathbb{N}_0 \mid H_{\mathfrak{a}}^j(M) \text{ not finitely generated}\}.$$

Using this notation we may write Theorem (2.2) in the form

Corollary 2.3. *If $i \leq f_{\mathfrak{a}}(M)$, then $\text{Ass}_R(H_{\mathfrak{a}}^i(M))$ is a finite set.* \square

Corollary 2.4 (see [B-R-Sh, (2.2)]). *The set $\text{Ass}_R(H_{\mathfrak{a}}^{\text{grade}_M(\mathfrak{a})}(M))$ is finite.*

Proof. Follows from Corollary (2.3) as $\text{grade}_M(\mathfrak{a}) \leq f_{\mathfrak{a}}(M)$. \square

REFERENCES

- [B-R-Sh] Brodmann, M., Rotthaus, Ch., Sharp, R., *On annihilators and associated primes of local cohomology modules*, to appear in Journal of Pure and Applied Algebra
- [B-Sh] Brodmann, M, Sharp, R., *Local cohomology – an algebraic introduction with geometric applications*, Cambridge studies in advanced mathematics No. 60, Cambridge University Press, 1998 CMP 98:10
- [F] Faltings, G., *Der Endlichkeitssatz in der lokalen Kohomologie*, Mathematische Annalen 255 (1981), 45-56 MR **82f**:13003
- [K] Katzman, M., *Finiteness of $\cup_e \text{Ass} F^e(M)$ and its connections to tight closure*; Illinois Journal of Mathematics 40 (1996), 330-337 MR **97i**:13003
- [M-S] Melkersson, L., Schenzel, P., *Asymptotic prime ideals related to derived functors*; Proc. AMS 117, No 4 (1993), 935- 938 MR **93e**:13027
- [R] Raghavan, K.N., *Local-global principle for annihilation of local cohomology*; Contemporary Math. 159 (1994), 329-331 MR **95c**:13018

MATHEMATICAL INSTITUTE, UNIVERSITY OF ZURICH, ZURICH, SWITZERLAND

E-mail address: brodmann@math.unizh.ch

E-mail address: lashagari@math.unizh.ch