



University of
Zurich^{UZH}

Zurich Open Repository and
Archive

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

Simpson type inequalities and applications

Awan, Muhammad Uzair ; Javed, Muhammad Zakria ; Rassias, Michael Th ; Noor, Muhammad Aslam ; Noor, Khalida Inayat

DOI: <https://doi.org/10.1007/s41478-021-00319-4>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-221304>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Awan, Muhammad Uzair; Javed, Muhammad Zakria; Rassias, Michael Th; Noor, Muhammad Aslam; Noor, Khalida Inayat (2021). Simpson type inequalities and applications. *Journal of Analysis*, 29(4):1403-1419.

DOI: <https://doi.org/10.1007/s41478-021-00319-4>



Simpson type inequalities and applications

Muhammad Uzair Awan¹ · Muhammad Zakria Javed¹ ·
Michael Th. Rassias² · Muhammad Aslam Noor³ · Khalida Inayat Noor³

Received: 19 December 2020 / Accepted: 1 April 2021 / Published online: 19 April 2021
© The Author(s) 2021

Abstract

A new generalized integral identity involving first order differentiable functions is obtained. Using this identity as an auxiliary result, we then obtain some new refinements of Simpson type inequalities using a new class called as strongly (s, m) -convex functions of higher order of $\sigma > 0$. We also discuss some interesting applications of the obtained results in the theory of means. In last we present applications of the obtained results in obtaining Simpson-like quadrature formula.

Keywords Convex · Strongly · Higher Order · Simpson · Inequality

Mathematics Subject Classification 26A51 · 26D15 · 65K15

Communicated by Samy Ponnusamy.



Michael Th. Rassias
michail.rassias@math.uzh.ch

Muhammad Uzair Awan
awan.uzair@gmail.com

Muhammad Zakria Javed
zakriajaved071@gmail.com

Muhammad Aslam Noor
noormaslam@gmail.com

Khalida Inayat Noor
khalidan@gmail.com

¹ Department of Mathematics, Government College University, Faisalabad, Pakistan

² Institute of Mathematics, University of Zurich, 8057 Zurich, Switzerland

³ Department of Mathematics, COMSATS University Islamabad, Islamabad, Pakistan

1 Introduction and preliminaries

The following inequality is known as Simpson's inequality.

Theorem 1.1 *Let $\mathcal{F} : [a_1, a_2] \rightarrow \mathbb{R}$ be a four times continuously differentiable function on (a_1, a_2) and $\|\mathcal{F}^4\|_\infty = \sup_{x \in (a_1, a_2)} |\mathcal{F}^4(x)| < \infty$, then*

$$\left| \frac{1}{3} \left[\frac{\mathcal{F}(a_1) + \mathcal{F}(a_2)}{2} + 2\mathcal{F}\left(\frac{a_1 + a_2}{2}\right) \right] - \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} \mathcal{F}(x) dx \right| \leq \frac{1}{2880} \|\mathcal{F}^4\|_\infty (a_2 - a_1)^4.$$

Simpson's inequality plays a significant role in modern analysis [4]. In past it has been extended and generalized in different directions using novel and innovative approaches. For example, Dragomir et al. [4] obtained the following version of the Simpson's inequality.

Theorem 1.2 *Let $\mathcal{F} : [a_1, a_2] \rightarrow \mathbb{R}$ be a differentiable function whose derivative is continuous on (a_1, a_2) and $\mathcal{F}' \in L[a_1, a_2]$, then*

$$\left| \frac{1}{3} \left[\frac{\mathcal{F}(a_1) + \mathcal{F}(a_2)}{2} + 2\mathcal{F}\left(\frac{a_1 + a_2}{2}\right) \right] - \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} \mathcal{F}(x) dx \right| \leq \frac{a_2 - a_1}{3} \|\mathcal{F}'\|_1,$$

where $\|\mathcal{F}'\|_1 = \int_{a_1}^{a_2} |\mathcal{F}'(x)| dx$.

Sarikaya et al. [13] obtained Simpson's inequality using differentiable convex functions.

Theorem 1.3 *Let $\mathcal{F} : [a_1, a_2] \rightarrow \mathbb{R}$ be a differentiable function on (a_1, a_2) such that $\mathcal{F}' \in L[a_1, a_2]$. If $|\mathcal{F}'|^q$ is convex, then*

$$\left| \frac{1}{6} \left[\mathcal{F}(a_1) + 4\mathcal{F}\left(\frac{a_1 + a_2}{2}\right) + \mathcal{F}(a_2) \right] - \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} \mathcal{F}(x) dx \right| \leq \frac{a_2 - a_1}{12} \left(\frac{1 + 2^{p+1}}{3(p+1)} \right)^{\frac{1}{p}} \left\{ \left(\frac{|\mathcal{F}'(a_1)|^q + 3|\mathcal{F}'(a_2)|^q}{4} \right)^{\frac{1}{q}} + \left(\frac{3|\mathcal{F}'(a_1)|^q + |\mathcal{F}'(a_2)|^q}{4} \right)^{\frac{1}{q}} \right\},$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Dragomir et al. [4] has written a very informative paper regarding the developments and applications of Simpson's inequality.

Fractional calculus although is very old but in recent decades has received special attention. It has emerged as an interdisciplinary field of mathematics. Due to its great utility and applications mathematicians have shown their keen interest in it.

The classical Riemann-Liouville fractional integrals are defined as:

Definition 1.1 ([7]) Let $\mathcal{F} \in L_1[a_1, a_2]$. Then the Riemann-Liouville integrals $J_{a_1^+}^\alpha \mathcal{F}$ and $J_{a_2^-}^\alpha \mathcal{F}$ of order $\alpha > 0$ with $a_1 \geq 0$ are defined by

$$J_{a_1^+}^\alpha \mathcal{F}(x) = \frac{1}{\Gamma(\alpha)} \int_{a_1}^x (x-v)^{\alpha-1} \mathcal{F}(v) dv, \quad x > a_1,$$

and

$$J_{a_2^-}^\alpha \mathcal{F}(x) = \frac{1}{\Gamma(\alpha)} \int_x^{a_2} (v-x)^{\alpha-1} \mathcal{F}(v) dv, \quad x < a_2,$$

where

$$\Gamma(x) = \int_0^\infty e^{-v} v^{x-1} dv, \quad x > 0$$

is the well known Gamma function.

Diaz and Pariguan [3] introduced the notion of generalized κ -gamma function. The integral form of Γ_κ is given by:

$$\Gamma_\kappa(x) = \int_0^\infty v^{x-1} e^{-\frac{v^\kappa}{\kappa}} dv, \quad \operatorname{Re}(x) > 0.$$

Note that

$$\Gamma_\kappa(x) = \kappa^{\frac{x}{\kappa}-1} \Gamma\left(\frac{x}{\kappa}\right).$$

κ -Beta function is defined as:

$$B_\kappa(x, y) = \frac{1}{\kappa} \int_0^1 v^{\frac{x}{\kappa}-1} (1-v)^{\frac{y}{\kappa}-1} dv.$$

Obviously

$$B_\kappa(x, y) = \frac{1}{\kappa} \beta\left(\frac{x}{\kappa}, \frac{y}{\kappa}\right).$$

Mubeen and Habibullah [9] were the first to extend the notion of classical Riemann-Liouville fractional integrals to κ -Riemann-Liouville fractional integrals. Sarikaya et al. [12] independently extended the notion of Riemann-Liouville fractional

integrals to κ -Riemann-Liouville fractional integrals and discussed some of its interesting properties.

To be more precise let \mathcal{F} be piecewise continuous on $I^* = (0, \infty)$ and integrable on any finite subinterval of $I = [0, \infty]$. Then for $\nu > 0$, we consider κ -Riemann-Liouville fractional integral of \mathcal{F} of order α

$${}_k J_{a_1}^\alpha \mathcal{F}(x) = \frac{1}{\kappa \Gamma_\kappa(\alpha)} \int_{a_1}^x (x - \nu)^{\frac{\alpha}{\kappa} - 1} \mathcal{F}(\nu) d\nu, \quad x > a_1, \kappa > 0.$$

For some recent studies involving κ -fractional calculus, see [1, 2, 5, 10, 11].

The main objective of this paper is to derive some new generalized variants of Simpson's like inequalities for the functions belonging to Lebesgue L_1 , L_q and L_∞ spaces. We essentially derive our results using a new class of convexity called as strongly (s, m) -convex functions of higher order of $\sigma > 0$. We also discuss some special cases of the obtained results. In order to show the significance of the obtained results, we also discuss some interesting applications of the main results. We hope that the ideas and the techniques of this paper will inspire interested readers working in this filed.

2 Main results

2.1 New definition

We now define the class of strongly (s, m) -convex functions of higher order of $\sigma > 0$.

Definition 2.1 A function $\mathcal{F} : [0, \infty) \rightarrow \mathbb{R}$ is said to be strongly (s, m) -convex functions of higher order of $\sigma > 0$, for certain fixed $(s, m) \in (0, 1]^2$, if

$$\begin{aligned} \mathcal{F}(\tau a_1 + m(1 - \tau)a_2) &\leq \tau^s \mathcal{F}(a_1) + m(1 - \tau)^s \mathcal{F}(a_2) \\ &\quad - \mu[\tau^\sigma(1 - \tau) + \tau(1 - \tau)^\sigma] \|a_2 - a_1\|^\sigma, \end{aligned}$$

where $\mu, \sigma > 0$ and $\tau \in [0, 1]$.

It is worth to point out here that, if $\sigma = 0$, then we have the class of (s, m) -convex functions introduced and studied by Eftkhari [6]. And if we take $\sigma = 2$, then the class of strongly (s, m) -convex functions of higher order of $\sigma > 0$ reduces to the class of strongly (s, m) -convex functions of order 2. To the best of our knowledge this is also new in the literature.

Definition 2.2 A function $\mathcal{F} : [0, \infty) \rightarrow \mathbb{R}$ is said to be strongly (s, m) -convex functions of order 2, for certain fixed $(s, m) \in (0, 1]^2$, if

$$\mathcal{F}(\tau a_1 + m(1 - \tau)a_2) \leq \tau^s \mathcal{F}(a_1) + m(1 - \tau)^s \mathcal{F}(a_2) - \mu\tau(1 - \tau) \|a_2 - a_1\|^2,$$

where $\mu > 0$ and $\tau \in [0, 1]$.

2.2 A key lemma

In this section, we derive a new integral identity which will be used as an auxiliary result in obtaining results of the subsequent sections.

Lemma 2.1 *Let $0 < m \leq 1$, $\alpha > 0$, $n \in \mathbb{N}$, and $\mathcal{F} : J = [ma_1, \frac{a_2}{m}] \rightarrow \mathbb{R}$ be a differentiable function with $a_1 < a_2$. If $\mathcal{F} \in L_1[J]$, then*

$$\begin{aligned} &\Omega(a_1, a_2; m, \alpha, w) \\ &= \frac{(w - ma_1)^2}{(n + 1)(a_2 - ma_1)} \int_0^1 \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n + 1} - \frac{1}{n + 4} \right) \mathcal{F}' \left(\frac{1 - \tau}{n + 1} ma_1 + \frac{n + \tau}{n + 1} w \right) d\tau \\ &\quad + \frac{(a_2 - w)^2}{(n + 1)(a_2 - ma_1)} \int_0^1 \left(\frac{1}{n + 4} - \frac{\tau^{\frac{\alpha}{\kappa}}}{n + 1} \right) \mathcal{F}' \left(\frac{n + \tau}{n + 1} w + \frac{1 - \tau}{n + 1} a_2 \right) d\tau, \end{aligned}$$

where

$$\begin{aligned} &\Omega(a_1, a_2; m, \alpha, w) \\ &= \frac{3}{(n + 1)(n + 4)(a_2 - ma_1)} \mathcal{F}(w) \\ &\quad + \frac{1}{(n + 4)(a_2 - ma_1)} \left[(w - ma_1) \mathcal{F} \left(\frac{(nw + ma_1)}{(n + 1)} \right) \right. \\ &\quad \left. + (a_2 - w) \mathcal{F} \left(\frac{(nw + a_2)}{(n + 1)} \right) \right] \\ &\quad - \frac{(n + 1)^{\frac{\alpha}{\kappa} - 1} \Gamma_{\kappa}(\alpha + \kappa)}{(a_2 - ma_1)(a_2 - w)^{\frac{\alpha}{\kappa} - 1}} {}_k J_{w^+}^{\alpha} \mathcal{F} \left(\frac{(nw + a_2)}{n + 1} \right) \\ &\quad - \frac{(n + 1)^{\frac{\alpha}{\kappa} - 1} \Gamma_{\kappa}(\alpha + \kappa)}{(a_2 - ma_1)(w - ma_1)^{\frac{\alpha}{\kappa} - 1}} {}_k J_{w^-}^{\alpha} \mathcal{F} \left(\frac{(nw + ma_1)}{n + 1} \right), \end{aligned}$$

and $w = (1 - \beta)ma_1 + \beta a_2$, $\beta \in [0, 1]$.

Proof Consider

$$\begin{aligned} I &= \frac{(w - ma_1)^2}{(n + 1)(a_2 - ma_1)} \int_0^1 \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n + 1} - \frac{1}{n + 4} \right) \mathcal{F}' \left(\frac{1 - \tau}{n + 1} ma_1 + \frac{n + \tau}{n + 1} w \right) d\tau \\ &\quad + \frac{(a_2 - w)^2}{(n + 1)(a_2 - ma_1)} \int_0^1 \left(\frac{1}{n + 4} - \frac{\tau^{\frac{\alpha}{\kappa}}}{n + 1} \right) \mathcal{F}' \left(\frac{n + \tau}{n + 1} w + \frac{1 - \tau}{n + 1} a_2 \right) d\tau \quad (2.1) \\ &= \frac{(w - ma_1)^2}{(n + 1)(a_2 - ma_1)} I_1 + \frac{(a_2 - w)^2}{(n + 1)(a_2 - ma_1)} I_2. \end{aligned}$$

Now

$$\begin{aligned}
 I_1 &= \int_0^1 \left(\frac{\tau^{\frac{\alpha}{k}}}{n+1} - \frac{1}{n+4} \right) \mathcal{F}' \left(\frac{1-\tau}{n+1} ma_1 + \frac{n+\tau}{n+1} w \right) d\tau \\
 &= \frac{3}{(n+4)(w-ma_1)} \mathcal{F}(w) + \frac{n+1}{(n+4)(w-ma_1)} \mathcal{F} \left(\frac{(nw+ma_1)}{n+1} \right) \\
 &\quad - \frac{(n+1)^{\frac{\alpha}{k}} \Gamma_{\kappa}(\alpha+\kappa)}{(w-ma_1)^{\frac{\alpha}{k}+1}} {}_k J_{w^-}^{\alpha} \mathcal{F} \left(\frac{(nw+ma_1)}{n+1} \right).
 \end{aligned} \tag{2.2}$$

Similarly

$$\begin{aligned}
 I_2 &= \int_0^1 \left(\frac{1}{n+4} - \frac{\tau^{\frac{\alpha}{k}}}{n+1} \right) \mathcal{F}' \left(\frac{n+\tau}{n+1} w + \frac{1-\tau}{n+1} a_2 \right) d\tau \\
 &= \frac{3}{(n+4)(a_2-w)} \mathcal{F}(w) + \frac{n+1}{(n+4)(a_2-w)} \mathcal{F} \left(\frac{(nw+a_2)}{n+1} \right) \\
 &\quad - \frac{(n+1)^{\frac{\alpha}{k}} \Gamma_{\kappa}(\alpha+\kappa)}{(a_2-w)^{\frac{\alpha}{k}+1}} {}_k J_{w^+}^{\alpha} \mathcal{F} \left(\frac{(nw+a_2)}{n+1} \right).
 \end{aligned} \tag{2.3}$$

Using the values of I_1 and I_2 in (2.1), we get the required result. □

2.3 Simpson’s like inequalities

We now derive some new Simpson’s like inequalities using the class of strongly (s, m) convex functions of higher order $\sigma > 0$.

Theorem 2.2 *Under all the assumption of Lemma 2.1, if $|\mathcal{F}'|$ is bounded, that is $\|\mathcal{F}'\|_{\infty} = \sup_{x \in [ma_1, a_2]} |\mathcal{F}'(x)| < \infty$, then*

$$|\Omega(a_1, a_2; m, \alpha, w)| \leq \phi(\tau) \left[\frac{(w-ma_1)^2 + (a_2-w)^2}{(n+1)(a_2-ma_1)} \right] \|\mathcal{F}'\|_{\infty}.$$

Proof Using Lemma 2.1, boundedness $|\mathcal{F}'|$ and properties of absolute value, we have

$$\begin{aligned}
 &|\Omega(a_1, a_2; m, \alpha, w)| \\
 &\leq \frac{(w-ma_1)^2}{(n+1)(a_2-ma_1)} \int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{k}}}{n+1} - \frac{1}{n+4} \right) \right| \left| \mathcal{F}' \left(\frac{1-\tau}{n+1} ma_1 + \frac{n+\tau}{n+1} w \right) \right| d\tau \\
 &\quad + \frac{(a_2-w)^2}{(n+1)(a_2-ma_1)} \int_0^1 \left| \left(\frac{1}{n+4} - \frac{\tau^{\frac{\alpha}{k}}}{n+1} \right) \right| \left| \mathcal{F}' \left(\frac{n+\tau}{n+1} w + \frac{1-\tau}{n+1} a_2 \right) \right| d\tau \\
 &\leq \left[\frac{(w-ma_1)^2 + (a_2-w)^2}{(n+1)(a_2-ma_1)} \right] \|\mathcal{F}'\|_{\infty} \int_0^1 \left| \frac{\tau^{\frac{\alpha}{k}}}{n+1} - \frac{1}{n+4} \right| d\tau \\
 &= \phi(\tau) \left[\frac{(w-ma_1)^2 + (a_2-w)^2}{(n+1)(a_2-ma_1)} \right] \|\mathcal{F}'\|_{\infty},
 \end{aligned}$$

where

$$\begin{aligned} & \int_0^1 \left| \left(\frac{1}{n+4} - \frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} \right) \right| \\ &= \frac{\alpha}{(n+1)(\alpha+\kappa)} \left(\frac{n+1}{n+4} \right)^{\frac{\kappa}{\alpha}} + \frac{\kappa}{(n+1)(\alpha+\kappa)} \left(1 - \left(\frac{n+1}{n+4} \right)^{\frac{\kappa}{\alpha}+1} \right) \\ & \quad - \frac{1}{n+4} \left(1 - \left(\frac{n+1}{n+4} \right)^{\frac{\kappa}{\alpha}} \right) := \phi(\tau) \end{aligned}$$

This completes the proof. □

Theorem 2.3 Under all the assumption of Lemma 2.1, if $\mathcal{F}' \in L_1[ma_1, a_2]$, then

$$|\Omega(a_1, a_2; m, \alpha, w)| \leq \frac{3}{(n+1)(n+4)} \|\mathcal{F}'\|_1,$$

where $\|\mathcal{F}'\|_1 = \int_{ma_1}^{a_2} |\mathcal{F}'(x)| dx$.

Proof Using Lemma 2.1, boundedness $|\mathcal{F}'|$ and properties of absolute value, we have

$$\begin{aligned} & |\Omega(a_1, a_2; m, \alpha, w)| \\ & \leq \frac{(w - ma_1)^2}{(n+1)(a_2 - ma_1)} \int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right| \left| \mathcal{F}' \left(\frac{1-\tau}{n+1} ma_1 + \frac{n+\tau}{n+1} w \right) \right| d\tau \\ & \quad + \frac{(a_2 - w)^2}{(n+1)(a_2 - ma_1)} \int_0^1 \left| \left(\frac{1}{n+4} - \frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} \right) \right| \left| \mathcal{F}' \left(\frac{n+\tau}{n+1} w + \frac{1-\tau}{n+1} a_2 \right) \right| d\tau \\ & \leq \frac{3}{(n+1)(n+4)} \left[\frac{w - ma_1}{a_2 - ma_1} \int_{\frac{ma_1+nw}{n+1}}^w |\mathcal{F}'(x)| dx + \right. \\ & \quad \left. \frac{a_2 - w}{a_2 - ma_1} \int_w^{\frac{a_2+nw}{n+1}} |\mathcal{F}'(x)| dx \right] \\ & \leq \frac{3}{(n+1)(n+4)} \left[\int_{\frac{ma_1+nw}{n+1}}^w |\mathcal{F}'(x)| dx + \int_w^{\frac{mw+a_2}{n+1}} |\mathcal{F}'(x)| dx \right] \\ & \leq \frac{3}{(n+1)(n+4)} \int_{\frac{ma_1+nw}{n+1}}^{\frac{mw+a_2}{n+1}} |\mathcal{F}'(x)| dx \\ & \leq \frac{3}{(n+1)(n+4)} \int_{ma_1}^{a_2} |\mathcal{F}'(x)| dx \\ & = \frac{3}{(n+1)(n+4)} \|\mathcal{F}'\|_1, \end{aligned}$$

where we have used the fact that $\left| \frac{1}{n+4} - \frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} \right| \leq \frac{3}{(n+1)(n+4)}$, for $\tau \in [0, 1]$ and $\alpha > 0$. This completes the proof. □

Theorem 2.4 Under all the assumption of Lemma 2.1, if $\mathcal{F}' \in L_q[ma_1, a_2]$ with $1 < q < \infty$, then

$$|\Omega(a_1, a_2; m, \alpha, w)| \leq N^{\frac{1}{p}} \left((n+1)^{\frac{1}{q}} (a_2 - ma_1)^{\frac{1}{p}} \right) \|\mathcal{F}'\|_p,$$

where

$$T = \frac{\kappa(n+1)^{\frac{\kappa}{\alpha}} \left((n+1)^{\frac{-\kappa}{\alpha}} - (n+4)^{\frac{-\kappa}{\alpha}} \right)^{\frac{2p}{\kappa} + 1}}{\alpha p + \kappa} + \frac{\kappa(n+1)^{\frac{\kappa}{\alpha}} (n+4)^{-p \frac{\kappa}{\alpha}}}{\alpha p + \kappa},$$

and $\|\mathcal{F}'\|_q = \left(\int_{ma_1}^{a_2} |\mathcal{F}'(x)|^q dx \right)^{\frac{1}{q}} < \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$.

Proof Using Lemma 2.1, boundedness $|\mathcal{F}'|$ and properties of absolute value, we have

$$\begin{aligned} & |\Omega(a_1, a_2; m, \alpha, w)| \\ & \leq \frac{(w - ma_1)^2}{(n+1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \right)^{\frac{1}{p}} \\ & \quad \left(\int_0^1 \left| \mathcal{F}' \left(\frac{1-\tau}{n+1} ma_1 + \frac{n+\tau}{n+1} w \right) \right|^q d\tau \right)^{\frac{1}{q}} \\ & \quad + \frac{(a_2 - w)^2}{(n+1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{1}{n+4} - \frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} \right) \right|^p d\tau \right)^{\frac{1}{p}} \\ & \quad \left(\int_0^1 \left| \mathcal{F}' \left(\frac{n+\tau}{n+1} w + \frac{1-\tau}{n+1} a_2 \right) \right|^q d\tau \right)^{\frac{1}{q}} \\ & \leq \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \right)^{\frac{1}{p}} \\ & \quad \left[\frac{(w - ma_1)^{2 - \frac{1}{q}}}{(n+1)^{1 - \frac{1}{q}} (a_2 - ma_1)} \left(\int_{\frac{ma_1+nw}{n+1}}^w |\mathcal{F}'(x)|^q dx \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \frac{(a_2 - w)^{2 - \frac{1}{q}}}{(n+1)^{1 - \frac{1}{q}} (a_2 - ma_1)} \left(\int_w^{\frac{a_2+nw}{n+1}} |\mathcal{F}'(x)|^q dx \right)^{\frac{1}{q}} \right] \\ & \leq \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \right)^{\frac{1}{p}} [(n+1)^{\frac{1}{q}-1} (a_2 - ma_1)^{\frac{1}{p}}] \\ & \quad \times \left[\left(\int_{\frac{ma_1+nw}{n+1}}^w |\mathcal{F}'(x)|^q dx \right)^{\frac{1}{q}} + \left(\int_w^{\frac{a_2+nw}{n+1}} |\mathcal{F}'(x)|^q dx \right)^{\frac{1}{q}} \right] \\ & \leq \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \right)^{\frac{1}{p}} [(n+1)^{\frac{1}{q}} (a_2 - ma_1)^{\frac{1}{p}}] \\ & \quad \times \left[\left(\int_{\frac{ma_1+nw}{n+1}}^w |\mathcal{F}'(x)|^q dx + \int_w^{\frac{a_2+nw}{n+1}} |\mathcal{F}'(x)|^q dx \right)^{\frac{1}{q}} \right] \\ & \leq \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \right)^{\frac{1}{p}} [(n+1)^{\frac{1}{q}} (a_2 - ma_1)^{\frac{1}{p}}] \left(\int_{ma_1}^{a_2} |\mathcal{F}'(x)|^q dx \right)^{\frac{1}{q}} \\ & = T^{\frac{1}{p}} (n+1)^{\frac{1}{q}} (a_2 - ma_1)^{\frac{1}{p}} \|\mathcal{F}'\|_p, \end{aligned}$$

where

$$\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \leq \int_0^1 \left| \left(\frac{\tau}{(n+4)^{\frac{\kappa}{\alpha}}} - \frac{1}{(n+4)^{\frac{\kappa}{\alpha}}} \right) \right|^{\frac{p\kappa}{\alpha}} d\tau$$

$$= \frac{\kappa(n+1)^{\frac{\kappa}{\alpha}} \left((n+1)^{-\frac{\kappa}{\alpha}} - (n+4)^{-\frac{\kappa}{\alpha}} \right)^{\frac{p\kappa}{\alpha} + 1}}{\alpha p + \kappa} + \frac{\kappa(n+1)^{\frac{\kappa}{\alpha}} (n+4)^{-p - \frac{\kappa}{\alpha}}}{\alpha p + \kappa} := N$$

which follows from $|A^\theta - B^\theta| \leq |A - B|^\theta$ for any $A, B \geq 1$ and $\theta \in (0, 1]$. This completes the proof. □

Theorem 2.5 *Let $\alpha > 0, \mu > 0, n \in \mathbb{N}$ and $\mathcal{F} : [ma_1, \frac{a_2}{m}] \rightarrow \mathbb{R}$ be a differentiable function with $a_1 < a_2$ such that $\mathcal{F} \in L_1[ma_1, \frac{a_2}{m}]$. If $|\mathcal{F}'|^q$ is strongly (s, m) -convex function of higher order of $\sigma > 0$ with $(s, m) \in (0, 1]^2$ for $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$, then for $w \in [ma_1, a_2]$ and $\alpha \in (0, 1]$, we have*

$$\begin{aligned} & \Omega(a_1, a_2; m, \alpha, w) \\ & \leq T^{\frac{1}{p}} \left[\frac{(w - ma_1)^2}{(n+1)(a_2 - ma_1)} \left(\frac{m}{(n+1)^s(s+1)} |\mathcal{F}'(a_1)|^q \right. \right. \\ & \quad \left. \left. + \frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} |\mathcal{F}'(w)|^q - \frac{2\mu}{(\sigma+1)(\sigma+2)} \|w - a_1\|^\sigma \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \frac{(a_2 - w)^2}{(n+1)(a_2 - ma_1)} \left(\frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} |\mathcal{F}'(w)|^q \right. \right. \\ & \quad \left. \left. + \frac{m}{(n+1)^s(s+1)} \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right|^q - \frac{2\mu}{(\sigma+1)(\sigma+2)} \left\| \frac{a_2}{m} - w \right\|^\sigma \right)^{\frac{1}{q}} \right], \end{aligned}$$

where T is defined in Theorem 2.4.

Proof Using Lemma 2.1 and the fact that $|\mathcal{F}'|^q$ is strongly (s, m) -convex function of higher order of $\sigma > 0$, we have

$$\begin{aligned} & |\Omega(a_1, a_2; m, \alpha, w)| \\ & \leq \frac{(w - ma_1)^2}{(n+1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \right)^{\frac{1}{p}} \\ & \quad \left(\int_0^1 \left| \mathcal{F}' \left(\frac{1-\tau}{n+1} ma_1 + \frac{n+\tau}{n+1} w \right) \right|^q d\tau \right)^{\frac{1}{q}} \\ & \quad + \frac{(a_2 - w)^2}{(n+1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{1}{n+4} - \frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} \right) \right|^p d\tau \right)^{\frac{1}{p}} \\ & \quad \left(\int_0^1 \left| \mathcal{F}' \left(\frac{n+\tau}{n+1} w + \frac{1-\tau}{n+1} a_2 \right) \right|^q d\tau \right)^{\frac{1}{q}} \\ & \leq \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\alpha}{\kappa}}}{n+1} - \frac{1}{n+4} \right) \right|^p d\tau \right)^{\frac{1}{p}} \\ & \quad \left[\frac{(w - ma_1)^2}{(n+1)(a_2 - ma_1)} \left(\int_0^1 \left(m \left(\frac{1-\tau}{n+1} \right)^s |\mathcal{F}'(a_1)|^q + \left(\frac{n+\tau}{n+1} \right)^s |\mathcal{F}'(w)|^q \right) d\tau \right. \right. \\ & \quad \left. \left. - \mu(\tau^\sigma(1-\tau) + \tau(1-\tau)^\sigma) \|w - a_1\|^\sigma \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \frac{(a_2 - w)^2}{(n+1)(a_2 - ma_1)} \left(\int_0^1 \left(\left(\frac{n+\tau}{n+1} \right)^s |\mathcal{F}'(w)|^q + m \left(\frac{1-\tau}{n+1} \right)^s \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right|^q \right) d\tau \right. \right. \\ & \quad \left. \left. - \mu(\tau^\sigma(1-\tau) + \tau(1-\tau)^\sigma) \left\| \frac{a_2}{m} - w \right\|^\sigma \right)^{\frac{1}{q}} \right], \end{aligned}$$

To obtain our required result, we use the following:

$$\int_0^1 \left(\frac{n + \tau}{n + 1}\right)^s d\tau = \frac{(n + 1)^{s+1} - n^{s+1}}{(n + 1)^s(s + 1)},$$

$$\int_0^1 \left(\frac{1 - \tau}{n + 1}\right)^s d\tau = \frac{1}{(n + 1)^s(s + 1)}$$

and

$$\int_0^1 \tau^\sigma(1 - \tau)d\tau = \frac{1}{(\sigma + 1)(\sigma + 2)}.$$

□

We now discuss some special cases.

Corollary 2.1 *If $w = ma_1$ and $\mu = 0$ in Theorem 2.5, we get*

$$\begin{aligned} &|\Omega(a_1, a_2; m, \alpha, ma_1)| \\ &\leq T^{\frac{1}{p}} \frac{(a_2 - ma_1)}{(n + 1)} \left[\left(\frac{(n + 1)^{s+1} - n^{s+1}}{(n + 1)^s(s + 1)} |\mathcal{F}'(ma_1)|^q \right. \right. \\ &\quad \left. \left. + \frac{m}{(n + 1)^s(s + 1)} \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right|^q \right)^{\frac{1}{q}} \right]. \end{aligned} \tag{2.4}$$

Corollary 2.2 *If $w = a_2$ and $\mu = 0$ in Theorem 2.5, we get*

$$\begin{aligned} &|\Omega(a_1, a_2; m, \alpha, a_2)| \\ &\leq T^{\frac{1}{p}} \frac{(a_2 - ma_1)}{(n + 1)} \left[\left(\frac{m}{(n + 1)^s(s + 1)} |\mathcal{F}'(a_1)|^q \right. \right. \\ &\quad \left. \left. + \frac{(n + 1)^{s+1} - n^{s+1}}{(n + 1)^s(s + 1)} |\mathcal{F}'(a_2)|^q \right)^{\frac{1}{q}} \right]. \end{aligned} \tag{2.5}$$

Corollary 2.3 *If $w = \frac{ma_1+a_2}{n+1}$ and $\mu = 0$ in Theorem 2.5, we get*

$$\begin{aligned} &|\Omega(a_1, a_2; m, \alpha, \frac{ma_1 + a_2}{n + 1})| \\ &\leq T^{\frac{1}{p}} \left[\left(\frac{(a_2 - mna)^2}{(n + 1)^3(a_2 - ma_1)} \right) \left(\frac{m}{(n + 1)^s(s + 1)} \right)^{\frac{1}{q}} |\mathcal{F}'(a_1)| \right. \\ &\quad \left. \left(\frac{(n + 1)^{s+1} - n^{s+1}}{(n + 1)^s(s + 1)} \right)^{\frac{1}{q}} \left\{ \left(\frac{(a_2 - mna)^2}{(n + 1)^3(a_2 - ma_1)} \right) + \left(\frac{(nb - ma_1)^2}{(n + 1)^3(a_2 - ma_1)} \right) \right\} \right. \\ &\quad \left. \left| \mathcal{F}'\left(\frac{ma_1 + a_2}{n + 1}\right) \right| + \left(\frac{m}{(n + 1)^s(s + 1)} \right)^{\frac{1}{q}} \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right| \right]. \end{aligned}$$

Corollary 2.4 Combining (2.4) and (2.5), we have

$$\begin{aligned}
 & \left| \frac{3}{(n+1)(n+4)} [\mathcal{F}(a_2) + \mathcal{F}(ma_1)] \right. \\
 & \quad + \frac{1}{(n+4)} \left[\mathcal{F}\left(\frac{nb+ma_1}{n+1}\right) + \mathcal{F}\left(\frac{nma+a_2}{n+1}\right) \right] \\
 & \quad - \frac{(n+1)^{\frac{\alpha}{\kappa}-1} \Gamma_{\kappa}(\alpha+\kappa)}{(a_2-ma_1)^{\frac{\alpha}{\kappa}}} {}_k J_{(ma_1)^+}^{\alpha} \mathcal{F}\left(\frac{nma+a_2}{n+1}\right) \\
 & \quad \left. - \frac{(n+1)^{\frac{\alpha}{\kappa}-1} \Gamma_{\kappa}(\alpha+\kappa)}{(a_2-ma_1)^{\frac{\alpha}{\kappa}}} {}_k J_{a_2^-}^{\alpha} \mathcal{F}\left(\frac{nb+ma_1}{n+1}\right) \right| \\
 & \leq \frac{(a_2-ma_1)^2}{(n+1)} T_p^{\frac{1}{p}} \left[\left(\frac{m}{(n+1)^s(s+1)} |\mathcal{F}'(a_1)|^q \right. \right. \\
 & \quad \left. \left. + \frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} |\mathcal{F}'(a_2)|^q \right)^{\frac{1}{q}} \right. \\
 & \quad \left. \left(\frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} |\mathcal{F}'(ma_1)|^q + \frac{m}{(n+1)^s(s+1)} \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right|^q \right)^{\frac{1}{q}} \right]
 \end{aligned} \tag{2.6}$$

Corollary 2.5 By taking $\alpha = \kappa = m = 1$ in (2.6), we have

$$\begin{aligned}
 & \left| \frac{3(a_2-a_1)}{(n+1)(n+4)} [\mathcal{F}(a_1) + \mathcal{F}(a_2)] \right. \\
 & \quad + \frac{a_2-a_1}{(n+4)} \left[\mathcal{F}\left(\frac{nb+a_1}{n+1}\right) + \mathcal{F}\left(\frac{na+a_2}{n+1}\right) \right] \\
 & \quad \left. - \left[\int_{a_1}^{\frac{na+a_2}{n+1}} \mathcal{F}(x) dx + \int_{\frac{a_1+nb}{n+1}}^{a_2} \mathcal{F}(x) dx \right] \right| \\
 & \leq \frac{(a_2-a_1)^2}{(n+1)} \left[\frac{(n+1)[3^{p+1} - (n+1)^{p+1}]}{(n+1)^{p+1}(n+4)^{p+1}(p+1)} \right]^{\frac{1}{p}} \\
 & \quad \times \left[\left(\frac{m}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} + \left(\frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} \right] [|\mathcal{F}'(a_1)| + |\mathcal{F}'(a_2)|].
 \end{aligned} \tag{2.7}$$

Theorem 2.6 Let $\alpha > 0$, $n \in \mathbb{N}$ and $\mathcal{F} : [ma_1, \frac{a_2}{m}] \rightarrow \mathbb{R}$ be a differentiable function with $a_1 < a_2$ such that $\mathcal{F} \in L_1[ma_1, \frac{a_2}{m}]$. If $|\mathcal{F}'|^q$ is strongly (s, m) -convex function of higher order of $\sigma > 0$ with $(s, m) \in (0, 1]^2$ for $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$, then for $w \in [ma_1, a_2]$ and $\alpha \in (0, 1]$, we have

$$\begin{aligned}
 & |\Omega(a_1, a_2; m, \alpha, w)| \\
 & \leq T^{\frac{1}{p}} \left[\frac{(w - ma_1)^2}{(n + 1)(a_2 - ma_1)} \left(m \left(\frac{1}{(n + 1)^{qs}(qs + 1)} \right)^{\frac{1}{q}} |\mathcal{F}'(a_1)| \right. \right. \\
 & \quad \left. \left. + \left(\frac{(n + 1)^{qs+1} - n^{qs+1}}{(n + 1)^{qs}(qs + 1)} \right)^{\frac{1}{q}} |\mathcal{F}'(w)| \right) \right. \\
 & \quad \left. - 2\mu \|w - a_1\|^{\sigma} (kB_{\kappa}(\kappa(q\sigma + 1), \kappa(q + 1)))^{\frac{1}{q}} \right) \\
 & \quad + \frac{(a_2 - w)^2}{(n + 1)(a_2 - ma_1)} \left(\left(\frac{(n + 1)^{qs+1} - n^{qs+1}}{(n + 1)^{qs}(qs + 1)} \right)^{\frac{1}{q}} |\mathcal{F}'(w)| \right. \\
 & \quad \left. + m \left(\frac{1}{(n + 1)^{qs}(qs + 1)} \right)^{\frac{1}{q}} \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right| \right. \\
 & \quad \left. - 2\mu \left\| \frac{a_2}{m} - w \right\|^{\sigma} (kB_{\kappa}(\kappa(q\sigma + 1), \kappa(q + 1)))^{\frac{1}{q}} \right) \right], \tag{2.8}
 \end{aligned}$$

where T is defined in Theorem 2.4.

Proof Using Lemma 2.1 and convexity of $|\mathcal{F}'|^q$ with $p, q > 1$, we have

$$\begin{aligned}
 & |\Omega(a_1, a_2; m, \alpha, w)| \\
 & \leq \frac{(w - ma_1)^2}{(n + 1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\kappa}{q}}}{n + 1} - \frac{1}{n + 4} \right) \right| \left| \mathcal{F}' \left(\frac{1 - \tau}{n + 1} ma_1 + \frac{n + \tau}{n + 1} w \right) \right| d\tau \right) \\
 & \quad + \frac{(a_2 - w)^2}{(n + 1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{1}{n + 4} - \frac{\tau^{\frac{\kappa}{q}}}{n + 1} \right) \right| \left| \mathcal{F}' \left(\frac{n + \tau}{n + 1} w + \frac{1 - \tau}{n + 1} a_2 \right) \right| d\tau \right) \\
 & \leq \frac{(w - ma_1)^2}{(n + 1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\kappa}{q}}}{n + 1} - \frac{1}{n + 4} \right) \right| \right. \\
 & \quad \left. \left(m \left(\frac{1 - \tau}{n + 1} \right)^s |\mathcal{F}'(a_1)| + \left(\frac{n + \tau}{n + 1} \right)^s |\mathcal{F}'(w)| \right) \right. \\
 & \quad \left. - \mu (\tau^{\sigma}(1 - \tau) + \tau(1 - \tau)^{\sigma}) \|w - a_1\|^{\sigma} d\tau \right) \\
 & \quad + \frac{(a_2 - w)^2}{(n + 1)(a_2 - ma_1)} \left(\int_0^1 \left| \left(\frac{1}{n + 4} - \frac{\tau^{\frac{\kappa}{q}}}{n + 1} \right) \right| \right. \\
 & \quad \left. \left(\left(\frac{n + \tau}{n + 1} \right)^s |\mathcal{F}'(w)| + m \left(\frac{1 - \tau}{n + 1} \right)^s \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right| \right) \right. \\
 & \quad \left. - \mu (\tau^{\sigma}(1 - \tau) + \tau(1 - \tau)^{\sigma}) \left\| \frac{a_2}{m} - w \right\|^{\sigma} d\tau \right) \\
 & \leq \left(\int_0^1 \left| \left(\frac{\tau^{\frac{\kappa}{q}}}{n + 1} - \frac{1}{n + 4} \right) \right|^p d\tau \right)^{\frac{1}{p}} \\
 & \quad \left[\frac{(w - ma_1)^2}{(n + 1)(a_2 - ma_1)} \left(\left(\int_0^1 \left(m \left(\frac{1 - \tau}{n + 1} \right)^{qs} d\tau \right)^{\frac{1}{q}} |\mathcal{F}'(a_1)| \right. \right. \right. \\
 & \quad \left. \left. + \left(\int_0^1 \left(\frac{n + \tau}{n + 1} \right)^{qs} d\tau \right)^{\frac{1}{q}} |\mathcal{F}'(w)| \right. \right. \\
 & \quad \left. \left. - \mu \|w - a_1\|^{\sigma} \left(\left(\int_0^1 \tau^{q\sigma} (1 - \tau)^q d\tau \right)^{\frac{1}{q}} + \left(\int_0^1 \tau^q (1 - \tau)^{q\sigma} d\tau \right)^{\frac{1}{q}} \right) \right) \right. \\
 & \quad \left. + \frac{(a_2 - w)^2}{(n + 1)(a_2 - ma_1)} \left(\left(\left(\int_0^1 \left(\frac{n + \tau}{n + 1} \right)^{qs} d\tau \right)^{\frac{1}{q}} |\mathcal{F}'(w)| \right. \right. \right. \\
 & \quad \left. \left. + m \left(\int_0^1 \left(\frac{1 - \tau}{n + 1} \right)^{qs} d\tau \right)^{\frac{1}{q}} \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right| \right. \right. \\
 & \quad \left. \left. - \mu \left\| \frac{a_2}{m} - w \right\|^{\sigma} \left(\left(\int_0^1 \tau^{q\sigma} (1 - \tau)^q d\tau \right)^{\frac{1}{q}} + \left(\int_0^1 \tau^q (1 - \tau)^{q\sigma} d\tau \right)^{\frac{1}{q}} \right) \right) \right) \right].
 \end{aligned}$$

By using these integrals in above expression, we obtain our required result

$$\int_0^1 \left(\frac{n + \tau}{n + 1}\right)^{qs} d\tau = \frac{(n + 1)^{qs+1} - n^{qs+1}}{(n + 1)^{qs}(qs + 1)},$$

$$\int_0^1 \left(\frac{1 - \tau}{n + 1}\right)^{qs} d\tau = \frac{1}{(n + 1)^{qs}(qs + 1)}$$

and

$$\int_0^1 \tau^q (1 - \tau)^{q\sigma} d\tau = \kappa B_{\kappa}(\kappa(q\sigma + 1), \kappa(q + 1)).$$

This completes the proof. □

Corollary 2.6 *If we put $w = \frac{ma_1+a_2}{n+1}$ in Theorem 2.6, then*

$$\begin{aligned} & \left| \Omega(a_1, a_2; m, \alpha, \frac{ma_1 + a_2}{n + 1}) \right| \\ & \leq T^{\frac{1}{p}} \left[\left(\frac{(a_2 - mna)^2}{(n + 1)^3(a_2 - ma_1)} \right) \left(\frac{m}{(n + 1)^{qs}(qs + 1)} \right)^{\frac{1}{q}} |\mathcal{F}'(a_1)| \right. \\ & \quad \left. + \left(\frac{(n + 1)^{qs+1} - n^{qs+1}}{(n + 1)^{qs}(qs + 1)} \right)^{\frac{1}{q}} \left\{ \left(\frac{(a_2 - mna)^2}{(n + 1)^3(a_2 - ma_1)} \right) + \left(\frac{(nb - ma_1)^2}{(n + 1)^3(a_2 - ma_1)} \right) \right\} \right. \\ & \quad \left| \mathcal{F}'\left(\frac{ma_1 + a_2}{n + 1}\right) \right| \\ & \quad \left. + \left(\frac{(nb - ma_1)^2}{(n + 1)^3(a_2 - ma_1)} \right) \left(\frac{m}{(n + 1)^{qs}(qs + 1)} \right)^{\frac{1}{q}} \left| \mathcal{F}'\left(\frac{a_2}{m}\right) \right| \right]. \end{aligned} \tag{2.9}$$

3 Applications

3.1 Applications to means

In this section, we give some applications to means. First of all we recall the definitions of some special means.

For $0 < a_1 < a_2$ and $n \in \mathbb{N}$, we have

1. The arithmetic mean is defined as: $A(a_1, a_2) = \frac{a_1+a_2}{n+1}$,
2. The β -logarithmic mean is defined as: $L_{\beta}(a_1, a_2) = \left[\frac{a_2^{\beta+1} - a_1^{\beta+1}}{(\beta+1)(a_2 - a_1)} \right]^{\frac{1}{\beta}}$ where $\beta \in \mathbb{Z} - \{0, 1\}$.

We now discuss main results of this section.

Proposition 3.1 *Let $0 < a_1 < a_2$, $0 < s < 1$ and $\frac{1}{p} + \frac{1}{q} = 1$ with $q > 1$, then*

$$\begin{aligned} & \left| \frac{3q}{(n+1)(n+4)(a_2-a_1)(s+q)} A^{\frac{s}{q}+1}(a_1, a_2) \frac{q}{(n+4)(a_2-a_1)(s+q)} \right. \\ & \left[\left(\frac{a_2-na}{n+1} \right) A^{\frac{s}{q}+1} \left(\frac{(2n+1)a_1}{n+1}, \frac{nb}{n+1} \right) \right. \\ & \left. + \left(\frac{nb-a_1}{n+1} \right) A^{\frac{s}{q}+1} \left(\frac{na}{n+1}, \frac{(2n+1)a_2}{n+1} \right) \right] \\ & \left. - \frac{q}{(s+q)} \left[I_{\frac{s}{q}+1}^{\frac{s}{q}+1} \left(\frac{(2n+1)a_1+nb}{(n+1)^2}, \frac{na+(2n+1)a_2}{(n+1)^2} \right) \right] \right| \\ & \leq \left[\frac{(n+1)(3^{p+1}+(n+1)^{p+1})}{(n+1)^{p+1}(n+4)^{p+1}(p+1)} \right]^{\frac{1}{p}} \left[\left(\frac{(a_2-na)^2}{(n+1)^3(a_2-a_1)} \right) \right. \\ & \left. \left(\frac{1}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} a_1^{\frac{s}{q}} \left(\frac{(n+1)^{s+1}-n^{s+1}}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} \right. \\ & \left. \left\{ \left(\frac{(a_2-na)^2}{(n+1)^3(a_2-a_1)} \right) + \left(\frac{(nb-a_1)^2}{(n+1)^3(a_2-a_1)} \right) \right\} A^{\frac{s}{q}}(a_1, a_2) \right. \\ & \left. + \left(\frac{(nb-a_1)^2}{(n+1)^3(a_2-a_1)} \right) \left(\frac{1}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} a_2^{\frac{s}{q}} \right]. \end{aligned}$$

Proof Using $\alpha = \kappa = m = 1$ and $\mathcal{F}(x) = \frac{x^{\frac{s}{q}+1}}{\frac{s}{q}+1}$ in Corollary 2.3 completes the proof. □

Proposition 3.2 Let $0 < a_1 < a_2$, $0 < s < 1$ and $\frac{1}{p} + \frac{1}{q} = 1$ with $q > 1$, then

$$\begin{aligned} & \left| \frac{3q}{(n+1)(n+4)(a_2-a_1)(s+q)} A^{\frac{s}{q}+1}(a_1, a_2) \frac{q}{(n+4)(a_2-a_1)(s+q)} \right. \\ & \left[\left(\frac{a_2-na}{n+1} \right) A^{\frac{s}{q}+1} \left(\frac{(2n+1)a_1}{n+1}, \frac{nb}{n+1} \right) \right. \\ & \left. + \left(\frac{nb-a_1}{n+1} \right) A^{\frac{s}{q}+1} \left(\frac{na}{n+1}, \frac{(2n+1)a_2}{n+1} \right) \right] \\ & \left. - \frac{q}{(s+q)} \left[I_{\frac{s}{q}+1}^{\frac{s}{q}+1} \left(\frac{(2n+1)a_1+nb}{(n+1)^2}, \frac{na+(2n+1)a_2}{(n+1)^2} \right) \right] \right| \\ & \leq \left[\frac{(n+1)(3^{p+1}+(n+1)^{p+1})}{(n+1)^{p+1}(n+4)^{p+1}(p+1)} \right]^{\frac{1}{p}} \\ & \left[\left(\frac{(a_2-na)^2}{(n+1)^3(a_2-a_1)} \right) \left(\frac{1}{(n+1)^{qs}(qs+1)} \right)^{\frac{1}{q}} a_1^{\frac{s}{q}} \right. \\ & \left. \left(\frac{(n+1)^{qs+1}-n^{qs+1}}{(n+1)^{qs}(qs+1)} \right)^{\frac{1}{q}} \left\{ \left(\frac{(a_2-na)^2}{(n+1)^3(a_2-a_1)} \right) \right. \right. \\ & \left. \left. + \left(\frac{(nb-a_1)^2}{(n+1)^3(a_2-a_1)} \right) \right\} A^{\frac{s}{q}}(a_1, a_2) \right. \\ & \left. + \left(\frac{(nb-a_1)^2}{(n+1)^3(a_2-a_1)} \right) \left(\frac{1}{(n+1)^{qs}(qs+1)} \right)^{\frac{1}{q}} a_2^{\frac{s}{q}} \right]. \end{aligned}$$

Proof Using $\alpha = \kappa = m = 1$ and $\mathcal{F}(x) = \frac{x^{\frac{s}{q}+1}}{\frac{s}{q}+1}$ in Corollary 2.6 completes the proof. □

3.2 Simpson-like quadrature formula

Let P be the partition of the interval $[a_1, a_2]$, that is $P : a_1 = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = a_2$. Consider the quadrature formula

$$\int_{a_1}^{a_2} \mathcal{F}(x)dx = {}_sA(\mathcal{F}, P) + {}_sR(\mathcal{F}, P),$$

where

$$\begin{aligned} {}_sA(\mathcal{F}, P) &= \frac{3}{10} \sum_{i=1}^{n-1} [\mathcal{F}(x_i) + \mathcal{F}(x_{i+1})](x_{i+1} - x_i) \\ &+ \frac{2}{5} \sum_{i=0}^{n-1} \mathcal{F}\left(\frac{x_i + x_{i+1}}{2}\right)(x_{i+1} - x_i), \end{aligned}$$

for Simpson-Like formula and ${}_sR(\mathcal{F}, P)$ denotes the relative approximation error of the integral $\int_{a_1}^b f(x)dx$.

Proposition 3.3 *Suppose that all the assumptions of Theorem 2.5 are satisfied. then the Simpson-like error estimate exists*

$$\begin{aligned} &|{}_sR(\mathcal{F}, P)| \\ &\leq \left[\frac{(n+1)[3^{p+1} + (n+1)^{p+1}]}{(n+1)^{p+1}(n+4)^{p+1}(p+1)} \right]^{\frac{1}{p}} \\ &\times \left[\left(\frac{1}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} + \left(\frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} \right] \\ &\sum_{i=0}^{n-1} \frac{(x_{i+1} - x_i)^2}{(n+1)} [|\mathcal{F}'(x_i)| + |\mathcal{F}'(x_{i+1})|]. \end{aligned} \tag{3.1}$$

Proof Considering the subintervals of $[a_1, a_2]$, that is $[x_i, x_{i+1}] \subset [a_1, a_2]$ and using in Corollary 2.5, we obtain

$$\begin{aligned}
& \left| \frac{3(x_{i+1} - x_i)}{(n+1)(n+4)} [\mathcal{F}(x_i) + \mathcal{F}(x_{i+1})] \right. \\
& \quad + \frac{x_{i+1} - x_i}{(n+4)} \left[\mathcal{F}\left(\frac{(nx_{i+1} + x_i)}{(n+1)}\right) + \mathcal{F}\left(\frac{(nx_i + x_{i+1})}{(n+1)}\right) \right] \\
& \quad \left. - \left[\int_{x_i}^{\frac{nx_i + x_{i+1}}{n+1}} \mathcal{F}(x) dx + \int_{\frac{x_i + nx_{i+1}}{n+1}}^{x_{i+1}} \mathcal{F}(x) dx \right] \right| \\
& \leq \left[\frac{(n+1)[3^{p+1} + (n+1)^{p+1}]}{(n+1)^{p+1}(n+4)^{p+1}(p+1)} \right]^{\frac{1}{p}} \\
& \quad \times \left[\left(\frac{1}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} + \left(\frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} \right] \\
& \quad \frac{(x_{i+1} - x_i)^2}{(n+1)} [|\mathcal{F}'(x_i)| + |\mathcal{F}'(x_{i+1})|].
\end{aligned}$$

By taking sum over i from 0 to $n-1$, we get

$$\begin{aligned}
|{}_sR(\mathcal{F}, P)| &= \sum_{i=0}^{n-1} \left| \frac{3(x_{i+1} - x_i)}{(n+1)(n+4)} [\mathcal{F}(x_i) + \mathcal{F}(x_{i+1})] \right. \\
& \quad + \frac{x_{i+1} - x_i}{(n+4)} \left[\mathcal{F}\left(\frac{(nx_{i+1} + x_i)}{(n+1)}\right) + \mathcal{F}\left(\frac{(nx_i + x_{i+1})}{(n+1)}\right) \right] \\
& \quad \left. - \left[\int_{x_i}^{\frac{nx_i + x_{i+1}}{n+1}} \mathcal{F}(x) dx + \int_{\frac{x_i + nx_{i+1}}{n+1}}^{x_{i+1}} \mathcal{F}(x) dx \right] \right| \\
& \leq \left[\frac{(n+1)[3^{p+1} + (n+1)^{p+1}]}{(n+1)^{p+1}(n+4)^{p+1}(p+1)} \right]^{\frac{1}{p}} \\
& \quad \times \left[\left(\frac{1}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} + \left(\frac{(n+1)^{s+1} - n^{s+1}}{(n+1)^s(s+1)} \right)^{\frac{1}{q}} \right] \\
& \quad \sum_{i=0}^{n-1} \frac{(x_{i+1} - x_i)^2}{(n+1)} [|\mathcal{F}'(x_i)| + |\mathcal{F}'(x_{i+1})|].
\end{aligned}$$

This completes the proof. \square

Author Contributions All the authors contributed equally for the preparation of the present paper.
Funding Open Access funding provided by Universität Zürich. Not applicable

Declarations

Competing interests The authors declare that they do not have any competing interests.

Conflicts of interest All authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Awan, M. U., Talib, S., Y. M. Chu, M. A. Noor, and K. I. Noor. (2020) Some new refinements of Hermite–Hadamard-type inequalities involving ψ_k -Riemann–Liouville fractional integrals and applications, *Math. Probl. Eng.*, 2020, Article ID 3051920.
2. Chu, Y.-M., M.U. Awan, S. Talib, M.A. Noor, and K.I. Noor. 2020. Generalizations of Hermite–Hadamard like inequalities involving χ_k -Hilfer fractional integrals. *Adv. Diff. Equ.* 2020: 594.
3. Diaz, R., and E. Pariguan. 2007. On hypergeometric functions and Pochhammer k -symbol. *Divulg. Math.* 15: 179–192.
4. Dragomir, S.S., R.P. Agarwal, and P. Cerone. 2000. On Simpson's inequality and applications. *J. Inequal. Appl.* 5: 533–579.
5. Du, T. S., Awan, M. U., A. Kashuri, and S. S. Zhao, Some k -fractional extensions of the trapezium inequalities through generalized relative semi- (m, h) -preinvexity, *Appl. Anal.*, <https://doi.org/10.1080/00036811.2019.1616083>.
6. Eftekhari, N. 2014. Some remarks on (s, m) -convexity in the second sense. *J. Math. Inequal.* 8 (3): 489–495.
7. Kilbas, A., H.M. Srivastava, and J.J. Trujillo. 2006. *Theory and applications of fractional differential equations*. Amsterdam, Netherlands: Elsevier B.V.
8. Luo, C.Y., and T.-S. Du. 2020. Generalized Simspon-like inequalities involving Riemann–Liouville fractional integrals and their applications. *Filomat* 34 (3): 751–760.
9. Mubeen, S., and G.M. Habibullah. 2012. k -fractional integrals and application. *Int. J. Contemp. Math. Sci.* 7 (2): 89–94.
10. Rehman, G., and S. Mubeen. 2014. Some inequalities involving k -gamma and k -beta functions with applications-II. *J. Inequal. Appl.* 2014: 445.
11. Rahman, G., S. Mubeen, and K.S. Nisar. 2020. On generalized k -fractional derivative operator. *AIMS Math.* 5 (3): 1936–1945.
12. Sarikaya, M.Z., and A. Karaca. 2014. On the k -Riemann–Liouville fractional integral and applications. *Int. J. Stat. Math.* 1 (3): 33–43.
13. Sarikaya, M. Z., Set, E., and E. Ozdemir, On new inequalities of Simpson's type for convex functions, RGMIA Research Report Collection, 13(2), article 2, (2010).

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.