



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
Main Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 1998

---

**On the numerical analysis of some variational problems with  
nonhomogeneous boundary conditions**

Chipot, M ; Elfanni, A

Abstract: The goal of this note is to expose a new techniques to get energy estimates for nonconvex problems with nonlinear boundary conditions in term of the mesh size of a Lagrange finite elements method.

DOI: <https://doi.org/10.1007/BF03167317>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-22186>

Journal Article

Published Version

Originally published at:

Chipot, M; Elfanni, A (1998). On the numerical analysis of some variational problems with nonhomogeneous boundary conditions. *Japan Journal of Industrial and Applied Mathematics*, 15(3):345-361.

DOI: <https://doi.org/10.1007/BF03167317>

## On the Numerical Analysis of Some Variational Problems with Nonhomogeneous Boundary Conditions

M. CHIPOT<sup>†</sup> and A. ELFANNI<sup>††</sup>

<sup>†</sup>*Institut für Angewandte Mathematik, Universität Zürich,  
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

<sup>††</sup>*Centre d'Analyse Non Linéaire, Université de Metz,  
URA CNRS 399, Ile du Saulcy, 57045 Metz Cedex 01, France*

Received November 14, 1996

The goal of this note is to expose a new techniques to get energy estimates for nonconvex problems with nonlinear boundary conditions in term of the mesh size of a Lagrange finite elements method.

*Key words:* approximation, non-convex, calculus of variations, finite elements

### 1. Introduction

Let  $\Omega$  be a bounded polyhedral domain of  $\mathbf{R}^n$ ,  $n \geq 2$ , of boundary  $\Gamma$ . Let  $w_i \in \mathbf{R}^n$ ,  $i = 1, \dots, p$ ,  $p \geq 2$ , and consider a function  $\varphi : \mathbf{R}^n \rightarrow \mathbf{R}$  such that

$$\varphi(w_i) = 0 \quad \forall i = 1, \dots, p, \quad (1.1)$$

$$\varphi(w) > 0 \quad \forall w \neq w_i, \quad i = 1, \dots, p, \quad (1.2)$$

$$w_i - w_p, \quad i = 1, \dots, p-1 \quad \text{are linearly independent.} \quad (1.3)$$

For instance, in a physical setting,  $\varphi$  could be some stored energy density that vanishes at wells  $w_i$ 's. These wells stand for natural states with low or no energy. Let us denote by  $\Psi$  a nonnegative continuous function such that for some  $q > 0$ ,  $c > 0$ ,

$$0 \leq \Psi(\xi) \leq c|\xi|^q \quad \forall \xi \in (-1, 1). \quad (1.4)$$

If  $A(x)$  is a Lipschitz continuous function i.e. if

$$A \in W^{1,\infty}(\Omega) \quad (1.5)$$

we denote by  $W_A^{1,\infty}(\Omega)$  the set

$$W_A^{1,\infty}(\Omega) = \{v \in W^{1,\infty}(\Omega) / v(x) = A(x) \text{ on } \Gamma\}. \quad (1.6)$$

Then, we would like to consider the problems

$$\inf_{v \in W_A^{1,\infty}(\Omega)} \int_{\Omega} \varphi(\nabla v(x)) dx, \quad (1.7)$$

$$\inf_{v \in W_A^{1,\infty}(\Omega)} \int_{\Omega} \Psi(v(x) - A(x)) + \varphi(\nabla v(x)) dx, \tag{1.8}$$

$$\inf_{v \in W^{1,\infty}(\Omega)} \int_{\Omega} \Psi(v(x) - A(x)) + \varphi(\nabla v(x)) dx, \tag{1.9}$$

More precisely we would like to investigate the case when

$$\nabla A(x) \in \text{Co}(w_i) \text{ a.e. } x \in \Omega \tag{1.10}$$

( $\text{Co}(w_i)$  denotes the convex hull of the  $w_i$ 's).

As we mentioned before, such minimization problems arise in various physical settings (see for instance the connection with continuum mechanics in [1], [2], [21], [23], [24], [25], [27], [28], [29], [30], [31]) and it is of importance to know with what accuracy a finite element method will provide an estimate of the energy level. Moreover, our theoretical analysis provides minimizing sequences for the problems at hand that describe of course the limit behavior of the system. However, we will not insist here on the fact that these problems have no minimizers and are described by the mean of these minimizing sequences. The interested reader is referred for instance to [8] or [11] for details. The relationship between (1.7), (1.9) is exposed in [3], we refer for instance to [26] for notation on Sobolev spaces.

For our purpose, let  $(\mathcal{T}_h)_{h>0}$  be a family of triangulations of  $\Omega$  (see [33]), that is to say satisfying

$$\forall h > 0 \begin{cases} \forall K \in \mathcal{T}_h, K \text{ is a } n\text{-simplex,} \\ \max_{K \in \mathcal{T}_h} (h_K) = h, \\ \exists \nu > 0 \text{ such that } \forall K \in \mathcal{T}_h \frac{h_K}{\rho_K} \leq \nu. \end{cases} \tag{1.11}$$

where  $h_K$  is the diameter of the  $n$ -simplex and  $\rho_K$  its roundness (i.e. the largest diameter of the balls that could fit in  $K$ ). If  $P_1(K)$  is the space of polynomials of degree 1 on  $K$ , set

$$V_h = \{v : \Omega \rightarrow \mathbf{R} \text{ continuous, } v|_K \in P_1(K), \forall K \in \mathcal{T}_h\}. \tag{1.12}$$

Moreover, let us denote by  $\widehat{A}$  the interpolate of  $A$  on  $\mathcal{T}_h$ -i.e.  $\widehat{A}$  is the unique function of  $V_h$  that agrees with  $A$  on the vertices of  $\mathcal{T}_h$ .

REMARK 1.1. If  $\nabla A(x) \in \text{Co}(w_i)$ , note that one does not have necessarily

$$\nabla \widehat{A} \in \text{Co}(w_i).$$

Then set

$$V_h^{\widehat{A}} = \{v : \Omega \rightarrow \mathbf{R} \text{ continuous, } v|_K \in P_1(K), \forall K \in \mathcal{T}_h, v = \widehat{A} \text{ on } \Gamma\}. \tag{1.13}$$

Our goal is to obtain estimates for the infima

$$\inf_{v \in V_h^{\widehat{A}}} \int_{\Omega} \varphi(\nabla v(x)) dx \tag{1.14}$$

$$\inf_{v \in V_h^{\widehat{A}}} \int_{\Omega} \Psi(v(x) - A(x)) + \varphi(\nabla v(x)) dx \tag{1.15}$$

$$\inf_{v \in V_h} \int_{\Omega} \Psi(v(x) - A(x)) + \varphi(\nabla v(x)) dx \tag{1.16}$$

in terms of the mesh size  $h$ . Moreover, in doing so, we will exhibit sequences providing approximations for (1.7)–(1.9).

REMARK 1.2. In the case where  $A$  is affine on each face of  $\Gamma$  then  $\widehat{A} = A$  on  $\Gamma$ . Note that no assumption is needed on  $\varphi$  for (1.14)–(1.16) to be defined. For (1.7)–(1.9) one might require  $\varphi$  to be a Borel function.

Problems of this type have been first considered by C. Collins, D. Kinderlehrer and M. Luskin in one dimension (see [16], [17]–[18], [15]). In higher dimension estimates were also obtained in [7], [8], [10] in the case where  $\nabla A$  is a constant and in [11] in the vectorial case. Sharp estimates were obtained in [14]. In the case when  $A$  is nonlinear, estimates were obtained in [9] and [13]. However, in these papers the estimates are not optimal and using new test functions we are able to improve them and get in term of  $h$  the same power than in the linear case i.e. when  $\nabla A$  is constant (see [8], [10]).

First, remark that there is no loss of generality in assuming that

$$0 \in \text{ri}(\text{Co}(w_i)) \tag{1.17}$$

where  $\text{ri}(\text{Co}(w_i))$  denotes the relative interior of  $\text{Co}(w_i)$ -i.e. its interior in the space spanned by the  $w_i - w_p$ 's. Indeed, let  $w \in \text{ri}(\text{Co}(w_i))$ . Then, we can look for a  $v$  of the form

$$v = u + w.x \tag{1.18}$$

where  $w.x$  denotes the scalar product of  $w$  and  $x$ . Then we are led to minimize

$$\int_{\Omega} \Psi(u(x) - (A(x) - w.x)) + \varphi(w + \nabla u(x)) dx \tag{1.19}$$

over some space. If we set

$$\tilde{\varphi}(\xi) = \varphi(\xi + w), \quad \tilde{A}(x) = A(x) - w.x \tag{1.20}$$

we end up to deal with the same problem with the wells  $w_i - w$  that satisfy

$$0 \in \text{ri}(\text{Co}(w_i - w)), \quad \nabla \tilde{A} \in \text{Co}(w_i - w) \text{ a.e. } x \in \Omega. \tag{1.21}$$

So, in what follows we will always assume that (1.17) holds. In particular, by (1.3) there exist  $\alpha_i$ 's,  $i = 1, \dots, p$ , such that

$$0 = \sum_{i=1}^p \alpha_i w_i, \quad \alpha_i \in (0, 1). \tag{1.22}$$

Moreover the space  $W$  spanned by the  $w_i - w_p$ 's coincides with the space spanned by the  $w_i$ 's.

## 2. Energy Estimates

Our main theorem is the following:

**THEOREM 2.1.** *Let us assume that  $\Omega$  is convex and that  $\varphi$  is a function bounded on bounded subsets of  $\mathbf{R}^n$  satisfying (1.1), (1.2). Moreover, assume that  $\Psi$  is a continuous function satisfying (1.4). Then, if  $A \in W^{1,\infty}(\Omega)$  satisfies (1.10), and (1.3) holds, there exists a constant  $C$ , independent of  $h \in (0, 1)$ , such that*

$$E_h^1 = \inf_{v \in V_h^A} \int_{\Omega} \varphi(\nabla v(x)) dx \leq Ch^{1/2} \tag{2.1}$$

$$E_h^2 = \inf_{v \in V_h^A} \int_{\Omega} \Psi(v(x) - A(x)) + \varphi(\nabla v(x)) dx \leq Ch^{r/(r+1)} \tag{2.2}$$

$$E_h^3 = \inf_{v \in V_h} \int_{\Omega} \Psi(v(x) - A(x)) + \varphi(\nabla v(x)) dx \leq Ch^{q/(q+1)} \tag{2.3}$$

where  $r = q \wedge 1$ ,  $\wedge$  denotes the infimum of two numbers.

**REMARK 2.1.** Assuming for instance that  $\lim_{|\xi| \rightarrow +\infty} \varphi(\xi) = +\infty$  or  $\lim_{|\xi| \rightarrow +\infty} \Psi(\xi) = +\infty$  one can show via an easy compactness argument that the infima (2.1)–(2.3) are in fact achieved (see [8] or [10] for a proof).

In order to prove Theorem 2.1 we will need some preparatory lemmas.

First:

**LEMMA 2.1.** *Assume  $A \in W^{1,\infty}(\Omega)$  satisfies (1.10). Then if the segment  $[x, x']$  is included in  $\Omega$  one has:*

$$\bigwedge_{i=1}^p w_{i \cdot} (x - x') \leq A(x) - A(x') \leq \bigvee_{i=1}^p w_{i \cdot} (x - x') \tag{2.4}$$

(we denote by  $\wedge$  the infimum of numbers, by  $\vee$  the supremum).

*Proof.* It is enough to apply the mean value theorem after regularization (see [13]). ■

REMARK 2.2. An immediate consequence of this lemma is that

$$A(x) - A(x') = 0$$

on each segment  $[x, x']$  such that  $x - x' \in W^\perp$ , where  $W^\perp$  denotes the orthogonal of  $W$  that could be, of course, reduced to 0 when the  $w_i$ 's are spanning the whole space.

Let us denote by  $v_1, \dots, v_{p-1}$  the dual basis of  $w_i - w_p$  i.e. the basis such that

$$(w_i - w_p).v_j = \delta_{ij} \quad \forall i, j = 1, \dots, p - 1. \tag{2.5}$$

Denote also by  $x_z$  the points of the lattice of size  $h^\alpha$ ,  $\alpha \in (0, 1)$  will be chosen later on, spanned by the  $v_i$ 's-i.e. for any  $z = (z_1, \dots, z_{p-1})$  set

$$x_z = \sum_{i=1}^{p-1} z_i h^\alpha v_i. \tag{2.6}$$

Then, let us define the function  $A$  by

$$A(x) = \bigvee_{z \in \mathbb{Z}^{p-1}} \left( \bigwedge_{i=1}^p w_i.(x - x_z) + \tilde{A}(x_z) \right) \tag{2.7}$$

where  $\tilde{A}$  is the extension of  $A$  (see the Appendix) that satisfies

$$\nabla \tilde{A}(x) \in \text{Co}(w_i) \quad \text{a.e. } x \in \mathbf{R}^n.$$

Note that  $A$  is constant in the  $W^\perp$  direction, (see Remark 2.2), and we will consider it as a function of  $x \in W$  or  $x \in \mathbf{R}^n$  as well. By a unit cell of the lattice spanned by the  $h^\alpha v_i$  we mean a set of the type

$$C_z = x_z + \left\{ \sum_{i=1}^{p-1} \beta_i h^\alpha v_i / \beta_i \in [0, 1] \right\}$$

where  $x_z$  is defined by (2.6). Then one has:

LEMMA 2.2. *Let us assume that  $\Omega$  is convex. Under the above assumptions, denote by  $C_{z_0}$  a unit cell spanned by  $h^\alpha v_i$ 's and by  $E$  the set*

$$E = \{z \in \mathbb{Z}^{p-1} / z_i = 0 \text{ or } 1, \forall i = 1, \dots, p - 1\},$$

*then one has*

$$A(x) = \bigvee_{z' \in z_0 + E} \left( \bigwedge_{i=1}^p w_i.(x - x_{z'}) + \tilde{A}(x_{z'}) \right) \quad \forall x \in C_{z_0}. \tag{2.8}$$

*i.e. in (2.7) instead of taking the supremum on a number of  $z$  that could be unbounded when  $h \rightarrow 0$  it is enough to take it on a fixed finite number.*

*Proof.* Assume that  $x \in C_{z_0}$ , and fix  $z \in \mathbb{Z}^{p-1}$ . Then, for some  $l = 1, \dots, p$  one has

$$\bigwedge_{i=1}^p w_i.(x - x_z) = w_l.(x - x_z). \quad (2.9)$$

We claim that there exists  $z' \in z_0 + E$  such that

$$\bigwedge_{i=1}^p w_i.(x - x_z) = \bigwedge_{i=1}^p w_i.(x - x_{z'}) + \bigwedge_{i=1}^p w_i.(x_{z'} - x_z). \quad (2.10)$$

To prove that let us show that there exists a  $z' \in z_0 + E$  such that

$$\begin{cases} w_i.(x - x_{z'}) \geq w_l.(x - x_{z'}) & \forall i = 1, \dots, p, i \neq l, \\ w_i.(x_{z'} - x_z) \geq w_l.(x_{z'} - x_z) & \forall i = 1, \dots, p, i \neq l. \end{cases} \quad (2.11)$$

Indeed, if (2.11) holds, by (2.9) one would have

$$\begin{aligned} \bigwedge_{i=1}^p w_i.(x - x_z) &= w_l.(x - x_z) \\ &= w_l.(x - x_{z'}) + w_l.(x_{z'} - x_z) \\ &= \bigwedge_{i=1}^p w_i.(x - x_{z'}) + \bigwedge_{i=1}^p w_i.(x_{z'} - x_z) \end{aligned} \quad (2.12)$$

and (2.10) will follow. Assume that

$$x = \sum_{k=1}^{p-1} (\beta_k + z_{0k}) h^\alpha v_k, \quad x_z = \sum_{k=1}^{p-1} z_k h^\alpha v_k, \quad x_{z'} = \sum_{k=1}^{p-1} z'_k h^\alpha v_k. \quad (2.13)$$

Then, clearly (2.9) reads

$$(w_i - w_l).x \geq (w_i - w_l).x_z \quad \forall i = 1, \dots, p, i \neq l$$

or

$$\sum_{k=1}^{p-1} \{\beta_k + z_{0k}\} (w_i - w_l).v_k \geq \sum_{k=1}^{p-1} z_k (w_i - w_l).v_k \quad \forall i = 1, \dots, p, i \neq l. \quad (2.14)$$

Similarly (2.11) reads

$$\begin{aligned} \sum_{k=1}^{p-1} \{\beta_k + z_{0k}\} (w_i - w_l).v_k &\geq \sum_{k=1}^{p-1} z'_k (w_i - w_l).v_k \quad \forall i = 1, \dots, p, i \neq l \\ \sum_{k=1}^{p-1} z'_k \{w_i - w_l\}.v_k &\geq \sum_{k=1}^{p-1} z_k \{w_i - w_l\}.v_k \quad \forall i = 1, \dots, p, i \neq l. \end{aligned} \quad (2.15)$$

One has (see (2.5)), when  $l \neq p$

\* if  $i = p$ ,

$$(w_i - w_l).v_k = -\delta_{lk}. \tag{2.16}$$

\* if  $i \neq p$ ,

$$\begin{aligned} (w_i - w_l).v_k &= [(w_i - w_p) + (w_p - w_l)].v_k \\ &= (w_i - w_p).v_k + (w_p - w_l).v_k \\ &= \delta_{ik} - \delta_{lk}. \end{aligned}$$

Hence, in this second case:

$$(w_i - w_l).v_k = \begin{cases} 0 & \text{when } k \neq i, l, \\ 1 & \text{when } k = i, \\ -1 & \text{when } k = l. \end{cases} \tag{2.17}$$

When  $l = p$

$$(w_i - w_l).v_k = \delta_{ik}. \tag{2.18}$$

Let us assume first that  $l \neq p$ , then (2.14), and (2.15) can be written respectively

$$\begin{cases} -(\beta_l + z_{0l}) \geq -z_l \\ (\beta_i + z_{0i}) - (\beta_l + z_{0l}) \geq z_i - z_l \quad \forall i = 1, \dots, p-1, i \neq l \end{cases} \tag{2.19}$$

$$\begin{cases} -(\beta_l + z_{0l}) \geq -z'_l \\ (\beta_i + z_{0i}) - (\beta_l + z_{0l}) \geq z'_i - z'_l \quad \forall i = 1, \dots, p-1, i \neq l \end{cases} \tag{2.20}$$

$$\begin{cases} -z'_l \geq -z_l \\ z'_i - z'_l \geq z_i - z_l \quad \forall i = 1, \dots, p-1, i \neq l. \end{cases} \tag{2.21}$$

Thus, knowing (2.19), we have to find  $z' \in z_0 + E$  such that

$$\begin{cases} -(\beta_l + z_{0l}) \geq -z'_l \geq -z_l \\ (\beta_i + z_{0i}) - (\beta_l + z_{0l}) \geq z'_i - z'_l \geq z_i - z_l \quad \forall i = 1, \dots, p-1, i \neq l. \end{cases} \tag{2.22}$$

If  $\beta_l \in ]0, 1]$ , we choose

$$z'_l = z_{0l} + 1, \begin{cases} z'_i = z_{0i} + 1 & \text{if } \beta_i \geq \beta_l, \\ z'_i = z_{0i} & \text{if } \beta_i < \beta_l, \quad i = 1, \dots, p-1, i \neq l. \end{cases} \tag{2.23}$$



In the case when  $\beta_i \geq \beta_l$ , we have  $0 \leq \beta_i - \beta_l < 1$ , since  $z_l, z_i - z_l$  are integers satisfying (2.19) one has

$$\begin{cases} -(\beta_l + z_{0l}) \geq -(z_{0l} + 1) \geq -z_l \\ (\beta_i + z_{0i}) - (\beta_l + z_{0l}) \geq (z_{0i} + 1) - (z_{0l} + 1) \geq z_i - z_l \end{cases} \quad (2.24)$$

$\forall i = 1, \dots, p-1, i \neq l.$

which is (2.22) in this case. When  $\beta_i < \beta_l$ , since  $-1 \leq \beta_i - \beta_l < 0$  we have

$$\begin{cases} -(\beta_l + z_{0l}) \geq -(z_{0l} + 1) \geq -z_l \\ (\beta_i + z_{0i}) - (\beta_l + z_{0l}) \geq z_{0i} - (z_{0l} + 1) \geq z_i - z_l \end{cases} \quad (2.25)$$

$\forall i = 1, \dots, p-1, i \neq l.$

which is (2.22). If  $\beta_l = 0$ , we choose

$$z'_l = z_{0l}, \begin{cases} z'_i = z_{0i} & \text{if } \beta_i \in [0, 1[, \\ z'_i = z_{0i} + 1 & \text{if } \beta_i = 1, \quad i = 1, \dots, p-1, i \neq l. \end{cases}$$

Then clearly (2.22) holds. So, it remains only to treat the case where  $l = p$ . In this case, the system to be solved is (see (2.14), (2.15), (2.18)):  
find  $z'$  such that

$$\beta_i + z_{0i} \geq z'_i \geq z_i \quad \forall i = 1, \dots, p-1, \quad (2.26)$$

knowing that

$$\beta_i + z_{0i} \geq z_i \quad \forall i = 1, \dots, p-1.$$

We choose

$$\begin{cases} z'_i = z_{0i} & \text{if } \beta_i \in [0, 1[, \\ z'_i = z_{0i} + 1 & \text{if } \beta_i = 1, \quad i = 1, \dots, p-1. \end{cases}$$

Then clearly we have (2.22). This proves (2.10).

To complete the proof of Lemma 2.2, consider a point  $x \in C_{z_0}$ . If  $x_z$  is arbitrary by (2.10), there exists a point  $z' \in z_0 + E$  such that

$$\bigwedge_{i=1}^p w_i \cdot (x - x_z) = \bigwedge_{i=1}^p w_i \cdot (x - x_{z'}) + \bigwedge_{i=1}^p w_i \cdot (x_{z'} - x_z).$$

Using Lemma 2.1 we deduce that

$$\bigwedge_{i=1}^p w_i \cdot (x - x_z) \leq \bigwedge_{i=1}^p w_i \cdot (x - x_{z'}) + \tilde{A}(x_{z'}) - \tilde{A}(x_z)$$

i.e.

$$\bigwedge_{i=1}^p w_i \cdot (x - x_z) + \tilde{A}(x_z) \leq \bigwedge_{i=1}^p w_i \cdot (x - x_{z'}) + \tilde{A}(x_{z'})$$

and this completes the proof. ■

We have also:

LEMMA 2.3. *Let us assume that  $\Omega$  is convex. Under the preceding assumptions one has for some positive constant  $C$  independent of  $h$*

$$A(x) - Ch^\alpha \leq \Lambda(x) \leq A(x) \quad \forall x \in \Omega. \tag{2.27}$$

*Proof.* Using Lemma 2.1 we get for  $x \in \Omega$  :

$$\bigwedge_{i=1}^p w_i \cdot (x - x_z) + \tilde{A}(x_z) \leq A(x) \quad \forall x_z.$$

Hence

$$\Lambda(x) \leq A(x).$$

Denote by  $P_W(\Omega)$  the orthogonal projection of  $\Omega$  onto  $W$ . Let  $x' \in P_W(\Omega)$  be the component of  $x$  in  $P_W(\Omega)$ . There exists a  $z_0$  such that  $x' \in C_{z_0}$ , so that

$$\begin{aligned} A(x) &\geq \bigwedge_{i=1}^p w_i \cdot (x - x_{z_0}) + \tilde{A}(x_{z_0}) \\ &= \bigwedge_{i=1}^p w_i \cdot (x' - x_{z_0}) + \tilde{A}(x_{z_0}) \\ &\geq -Ch^\alpha + \tilde{A}(x_{z_0}) - A(x') + A(x) \\ &\geq -Ch^\alpha + A(x) \end{aligned}$$

we used the fact that  $A(x) = A(x')$ , see Remark 2.2. This completes the proof. ■

*Proof of Theorem 2.1.*

-*Proof of (2.3).* Define  $\hat{u}_h$  as the interpolate of the function  $\Lambda(x)$  that has been introduced in (2.7). Clearly,  $\hat{u}_h \in V_h$ . Then, first notice that

$$\nabla \Lambda(x) = w_i \quad \text{a.e. } x \in \Omega,$$

so that  $\Lambda$  is a Lipschitz continuous function with a Lipschitz constant uniformly bounded. It then results that the same holds for  $\hat{u}_h$  (see for instance [4], [5], [6]). Then, by the mean value theorem

$$|\Lambda(x) - \hat{u}_h(x)| \leq Ch$$

for some constant  $C$ . Since  $\alpha \in (0, 1)$  we deduce from (2.27)

$$\begin{aligned} |A(x) - \widehat{u}_h(x)| &\leq |A(x) - A(x)| + |A(x) - \widehat{u}_h(x)| \\ &\leq Ch^\alpha + Ch \leq Ch^\alpha \end{aligned} \tag{2.28}$$

if one assumes  $h < 1$ . In particular this leads to

$$\int_{\Omega} \Psi(\widehat{u}_h - A(x)) dx \leq C|\Omega|h^{q\alpha} \tag{2.29}$$

(note that (1.4) holds on any bounded subset of  $\mathbf{R}$ ,  $|\Omega|$  denotes the measure of  $\Omega$ .)

Next we need to estimate

$$\int_{\Omega} \varphi(\nabla \widehat{u}_h(x)) dx. \tag{2.30}$$

First, notice that

$$\nabla \widehat{u}_h = w_i$$

except maybe on the set  $I$  composed of the simplices where interpolation occurred. Since on this set  $\nabla \widehat{u}_h$  remains bounded one has

$$\int_{\Omega} \varphi(\nabla \widehat{u}_h(x)) dx = \int_I \varphi(\nabla \widehat{u}_h(x)) dx \leq C|I|$$

where  $|I|$  denotes the measure of  $I$ . Now, interpolation occurs on a  $h$ -neighborhood of the ridge of  $A(x)$ -i.e. a  $h$ -neighborhood of the set where  $A(x)$  has a discontinuity in its gradient. Clearly  $A(x)$  has a jump in its gradient on a unit cell of the lattice spanned by  $h^\alpha v_i$  when one of the function

$$w_i \cdot (x - x_z) + \widetilde{A}(x_z)$$

is equal to an other. These two functions are then equal on a set of  $p-2$ -dimensional measure bounded by  $C(h^\alpha)^{p-2}$  (this is the intersection of a hyperplane and a cell of diameter bounded by  $Ch^\alpha$ ). Since in (2.7) the supremum is taken on a finite number of functions it is clear that

$$|I| \leq C(h^\alpha)^{p-2} \cdot h \cdot N(h^\alpha) \tag{2.31}$$

where  $N(h^\alpha)$  is the number of cells of size  $h^\alpha$  included in  $P_W(\Omega)$  (Recall that  $P_W$  denotes the projection of  $\Omega$  onto  $W$ ). Clearly,

$$N(h^\alpha) \cdot (h^\alpha)^{p-1} \leq C.$$

Hence (2.31) reads

$$|I| \leq Ch^{1-\alpha}. \tag{2.32}$$

Collecting (2.29)–(2.32) we get

$$\int_{\Omega} \Psi(\widehat{u}_h - A(x)) + \varphi(\nabla \widehat{u}_h(x)) \leq C[h^{\alpha q} + h^{1-\alpha}].$$

The power is the best in this inequality when  $1 - \alpha = \alpha q$ , i.e. when  $\alpha = \frac{1}{1+q}$ . This completes the proof of (2.3).

REMARK 2.3. The above estimate is sharp (see for instance [14]).

-*Proof of (2.1), (2.2).* Define  $\Lambda$  as before. Now, this function is not necessarily equal to  $A(x)$  on the boundary of  $\Omega$ . So, its interpolate will not be in  $V_h^A$ . To correct that, set

$$u_h(x) = \Lambda(x) \vee (A(x) - \text{dist}(x, \Gamma))$$

where  $\text{dist}(x, \Gamma)$  denotes the distance from  $x$  to the boundary  $\Gamma$ . Due to (2.27) one has

$$|A(x) - \Lambda(x)| \leq Ch^\alpha.$$

So, if

$$\text{dist}(x, \Gamma) \geq Ch^\alpha,$$

then,

$$u_h(x) = A(x).$$

Now, if

$$u_h(x) = A(x) - \text{dist}(x, \Gamma),$$

one has necessarily

$$|u_h(x) - A(x)| = \text{dist}(x, \Gamma) \leq Ch^\alpha.$$

It results that in all cases one has

$$|u_h(x) - A(x)| \leq Ch^\alpha.$$

Since - when  $\widehat{u}_h$  denotes the interpolate of  $u_h$

$$|u_h(x) - \widehat{u}_h(x)| \leq Ch$$

one deduces -say for  $h \leq 1$

$$|\widehat{u}_h(x) - A(x)| \leq Ch^\alpha. \tag{2.33}$$

So, we get (see (1.4), (2.33))

$$\int_{\Omega} \Psi(\widehat{u}_h(x) - A(x)) dx \leq Ch^{\alpha q}. \quad (2.34)$$

Next, as in part 1

$$\int_{\Omega} \varphi(\nabla \widehat{u}_h(x)) dx \leq C|I| \quad (2.35)$$

where  $I$  denotes the set where we are interpolating. Recall that when

$$\text{dist}(x, \Gamma) \geq Ch^{\alpha},$$

one has

$$u_h(x) = A(x).$$

So, when  $\text{dist}(x, \Gamma) \geq Ch^{\alpha} + h$  one has

$$\widehat{u}_h = \text{the interpolate of } A(x)$$

and, by the estimate we developed in part 1, if  $I_1 = \{x/\text{dist}(x, \Gamma) \geq Ch^{\alpha} + h\}$ , we obtain

$$\int_{I_1} \varphi(\nabla \widehat{u}_h(x)) dx \leq Ch^{1-\alpha}. \quad (2.36)$$

Next we have to estimate

$$\int_{I \setminus I_1} \varphi(\nabla \widehat{u}_h(x)) dx. \quad (2.37)$$

But since  $I \setminus I_1$  is a  $h^{\alpha}$ -neighborhood of  $\Gamma$ . (note that  $h \leq h^{\alpha}$ ), and since the gradient of  $\widehat{u}_h$  is uniformly bounded we get

$$\int_{I \setminus I_1} \varphi(\nabla \widehat{u}_h(x)) dx \leq Ch^{\alpha}. \quad (2.38)$$

Combining (2.34)–(2.38) we obtain

$$\begin{aligned} \int_{\Omega} \Psi(\widehat{u}_h(x) - A(x)) + \varphi(\nabla \widehat{u}_h(x)) dx &\leq C[h^{\alpha q} + h^{1-\alpha} + h^{\alpha}] \\ &\leq [h^{\alpha r} + h^{1-\alpha}] \end{aligned} \quad (2.39)$$

and

$$\int_{\Omega} \varphi(\nabla \widehat{u}_h(x)) dx \leq C[h^{1-\alpha} + h^{\alpha}] \quad (2.40)$$

In (2.39) the estimate is the best when  $1 - \alpha = \alpha r$ , i.e. when  $\alpha = \frac{1}{r+1}$ . This gives (2.2). In (2.40) the estimate is optimal for  $\alpha = \frac{1}{2}$  which gives (2.1). This completes the proof of Theorem (2.1). ■

REMARK 2.4. In the case where  $\Omega$  is not convex, when  $q \geq 1$ , and under the assumptions of Theorem 2.1 one has for some constant  $C$

$$E_h^i \leq Ch^{1/2} \quad \forall i = 1, 2, 3. \tag{2.41}$$

Indeed, thanks to (1.4) one has

$$E_h^1 \leq E_h^2$$

and since  $V_h^{\hat{A}} \subset V_h$ ,

$$E_h^3 \leq E_h^2.$$

So, it is enough to prove (2.41) when  $i = 2$ . But clearly a polyhedral domain can be decomposed into polyhedral domains  $\Omega_i, i = 1, \dots, N$  that are convex. Then, if on each of these domains we construct  $u_h$  as in part 2 of the proof of Theorem 2.1, the different  $u_h$  are matching on the boundary of the different  $\Omega_i$ . Then, we call  $\hat{u}_h$  the interpolate of these functions on  $\mathcal{T}_h$ . The estimate of

$$\int_{\Omega} \Psi(\hat{u}_h(x) - A(x)) + \varphi(\nabla \hat{u}_h(x)) dx = \sum_{i=1}^N \int_{\Omega_i} \Psi(\hat{u}_h(x) - A(x)) + \varphi(\nabla \hat{u}_h(x)) dx$$

goes the same way than in part 2 above and the result follows.

In the case  $q < 1$ , one would get similarly

$$E_h^2, E_h^3 \leq Ch^{q/(q+1)}.$$

### Appendix

We would like to prove here an extension Theorem that we have used in Section 2. Let  $\Omega$  be a domain of  $\mathbf{R}^n$ . Let  $w_1, \dots, w_p \in \mathbf{R}^n$ , and consider a function  $A$  defined in  $\Omega$ . We have:

THEOREM A.1. *Assume that  $\Omega$  is convex. If  $A \in W^{1,\infty}(\Omega)$  satisfies*

$$\nabla A(x) \in \text{Co}(w_i) \quad \text{a.e. } x \in \Omega \tag{A.1}$$

*then there exists a Lipschitz continuous function  $\tilde{A}$  defined in  $\mathbf{R}^n$  such that:*

$$\begin{aligned} \tilde{A} &= A \quad \text{in } \Omega, \\ \nabla \tilde{A}(x) &\in \text{Co}(w_i) \quad \text{a.e. } x \in \mathbf{R}^n. \end{aligned}$$

Before proving this theorem, remark that when  $A$  satisfies (A.1) then

$$A(x) - A(x') \geq \bigwedge_{i=1}^p w_i \cdot (x - x') \quad \forall x, x' \in \Omega.$$

In the following lemma we would like to establish that conversely:

LEMMA A.1. *If  $\Omega$  is a domain of  $\mathbf{R}^n$  and  $A$  satisfies*

$$A(x) - A(x') \geq \bigwedge_{i=1}^p w_i \cdot (x - x') \quad \forall x, x' \in \Omega$$

*then*

$$\nabla A(x) \in \text{Co}(w_i) \quad \text{for a.e. } x \in \Omega.$$

*Proof.* We know that there exists  $u_h \in W^{1,\infty}(\Omega)$  (cf. [9]) such that

$$\begin{aligned} u_h &\longrightarrow A \quad \text{uniformly,} \\ \nabla u_h &= w_i, \quad i = 1, \dots, p. \end{aligned}$$

By extracting eventually a subsequence we have:

$$\nabla u_h \rightarrow \nabla A \quad \text{in } L^\infty(\Omega) \text{ weak } *. \tag{A.2}$$

Let us now denote by  $B$  an arbitrary ball included in  $\Omega$ . We have

$$\frac{1}{|B|} \int_B \nabla u_h(x) dx \in \text{Co}(w_i).$$

On the other hand we have by (A.2)

$$\lim_{h \rightarrow 0} \frac{1}{|B|} \int_B \nabla u_h(x) dx = \frac{1}{|B|} \int_B \nabla A(x) dx.$$

Since  $\text{Co}(w_i)$  is closed

$$\frac{1}{|B|} \int_B \nabla A(x) dx \in \text{Co}(w_i).$$

Using the Lebesgue differentiation theorem we get

$$\nabla A(x) \in \text{Co}(w_i) \quad \text{a.e. } x \in \Omega.$$

This completes the proof in the lemma. ■

We are now able to prove Theorem A.1. Define the function  $\tilde{A}$  on  $\mathbf{R}^n$  by setting

$$\tilde{A}(x) = \inf_{e \in \Omega} \left\{ A(e) - \bigwedge_{i=1}^p w_i \cdot (e - x) \right\} \quad (\text{A.3})$$

First, let us show that  $\tilde{A} = A$  in  $\Omega$ . Let  $x \in \Omega$ . It is clear that  $\tilde{A}(x) \leq A(x)$ . Moreover, see Lemma 2.1,

$$\forall e \in \Omega, \quad A(e) - A(x) \geq \bigwedge_{i=1}^p w_i \cdot (e - x)$$

or

$$\forall e \in \Omega, \quad A(e) - \bigwedge_{i=1}^p w_i \cdot (e - x) \geq A(x)$$

i.e.

$$\tilde{A}(x) \geq A(x).$$

Hence

$$\tilde{A}(x) = A(x) \quad \forall x \in \Omega.$$

Finally we have:

$$\forall x, y \in \mathbf{R}^n, \quad \tilde{A}(x) - \tilde{A}(y) \geq \bigwedge_{i=1}^p w_i \cdot (x - y).$$

Indeed:

$$\begin{aligned} \tilde{A}(x) &= \inf_{e \in \Omega} \left\{ A(e) - \bigwedge_{i=1}^p w_i \cdot (e - x) \right\} \\ &= \inf_{e \in \Omega} \left\{ A(e) - \bigwedge_{i=1}^p w_i \cdot (e - y) + \bigwedge_{i=1}^p w_i \cdot (e - y) - \bigwedge_{i=1}^p w_i \cdot (e - x) \right\} \\ &\geq \inf_{e \in \Omega} \left\{ A(e) - \bigwedge_{i=1}^p w_i \cdot (e - y) \right\} + \bigwedge_{i=1}^p w_i \cdot (x - y) \\ &\quad \left( \text{since } \bigwedge_{i=1}^p w_i \cdot (e - y) - \bigwedge_{i=1}^p w_i \cdot (e - x) \geq \bigwedge_{i=1}^p w_i \cdot (x - y) \right) \\ &= \tilde{A}(y) + \bigwedge_{i=1}^p w_i \cdot (x - y). \end{aligned}$$

Taking  $y \in \Omega$  this shows in particular that the infimum in (A.3) is finite. Moreover, by Lemma A.1 it follows that:

$$\nabla \tilde{A}(x) \in \text{Co}(w_i) \quad \text{for a.e. } x \in \mathbf{R}^n$$



This completes the proof of the Theorem A.1.

*Acknowledgments.* We would like to thank the referee for his remarks that help to improve the earlier version of this work.

### References

- [ 1 ] J.M. Ball and R.D. James, Fine phase mixtures as minimizers of energy. Arch. Rational Mech. Anal., **100** (1987), 13–52.
- [ 2 ] J.M. Ball and R.D. James, Proposed experimental tests of a theory of fine microstructures. Phil. Trans. Roy. Soc. London A, **338** (1992), 350–389.
- [ 3 ] J.M. Ball and F. Murat,  $W^{1,p}$  quasiconvexity and variational problems for multiples integrals. J. Funct. Anal., **58** (1984), 225–253.
- [ 4 ] B. Brighi, Sur quelques problèmes de calcul des variations et l’approximation de leur fonctionnelles relaxées. Thesis, University of Metz, 1991.
- [ 5 ] B. Brighi and M. Chipot, Approximation in nonconvex problems. Proceedings of the first European Conference on Elliptic and Parabolic Problems, Pont-à-Mousson, June 1991. Pitman research notes in mathematics, #267, 1992, 150–157.
- [ 6 ] B. Brighi and M. Chipot, Approximated convex envelope of a function. SIAM J. Numer. Anal., **31**, 1 (1994), 128–148.
- [ 7 ] M. Chipot, Hyperelasticity for crystals. Eur. J. Appl. Math., **1** (1990), 113–129.
- [ 8 ] M. Chipot, Numerical analysis of oscillations in nonconvex problems. Numerische Mathematik, **59** (1991), 747–767.
- [ 9 ] M. Chipot, Energy estimates for variational problems with nonhomogeneous boundary conditions. Nonlinear mathematical problems in industry, Gakuto International Series, Mathematical Sciences and Applications, **2**, 1993, 473–487.
- [10] M. Chipot and C. Collins, Numerical approximation in variational problems with potential wells. SIAM J. Numer. Anal., **29**, 4 (1993), 473–487.
- [11] M. Chipot, C. Collins and D. Kinderlehrer, Numerical analysis of oscillations in multiple well problems. Numerische Mathematik, **70** (1995), 259–282.
- [12] M. Chipot and D. Kinderlehrer, Equilibrium configurations of crystals. Arch. Rational Mech. Anal., **103** (1988), 237–277.
- [13] M. Chipot and W. Li, Variational problems with potential wells and nonhomogeneous boundary conditions. Calculus of variations, homogenization and continuum mechanics (eds. G. Bouchitté, G. Buttazzo and P. Suquet), Series on Advances in Mathematics for Applied Sciences, **18**, World Scientific, 1994, 149–168.
- [14] M. Chipot and S. Müller, Sharp energy estimates to finite element approximations for nonconvex problems (to appear).
- [15] C. Collins, Thesis, University of Minnesota, 1990.
- [16] C. Collins, D. Kinderlehrer and M. Luskin, Numerical approximation of the solution of a variational problem with a double well potential. SIAM J. Numer. Anal., **28** (1991), 321–332.
- [17] C. Collins and M. Luskin, The computation of austenetic-martensitic phase transition. Partial Differential Equations and Continuum Models of Phase Transitions (eds. M. Rasle, D. Serre and M. Slemrod), Lecture Notes in Physics, #344, Springer-Verlag, Berlin, New York, 1989, 34–50.
- [18] C. Collins and M. Luskin, Computational results for phase transitions in shape memory materials. Smart Materials, Structure, and Mathematical Issues (ed. C. Rogers), Technomic Publishing Co., Lancaster, Pennsylvania, 1989, 198–215.
- [19] C. Collins and M. Luskin, Numerical modeling of the microstructure of crystals with symmetry-related variants. Proceedings of the ARO US-Japan Workshop on Smart/Intelligent Materials and Systems, Technomic Publishing Co., Lancaster, Pennsylvania, 1990, 309–318.
- [20] C. Collins and M. Luskin, Optimal order error estimates for the finite element approximation of the solution of a nonconvex variational problem. Math. Comp., **57** (1991), 621–637.
- [21] B. Dacorogna, Weak continuity and weak lower semicontinuity of non linear functionals. Springer Lectures Notes, #922, 1982.
- [22] A. Elfanni, Sur quelques questions d’analyse numérique relatives à des problèmes non con-

- vexes. Thesis, University of Metz, 1996.
- [23] J.L. Ericksen, Some constrained elastic crystals. *Material Instabilities in Continuum Mechanics and Related Problems* (ed. J.M. Ball), Oxford University Press, Oxford, 1987, 119–137.
  - [24] I. Fonseca, Variational methods for elastic crystals. *Arch. Rational Mech. Anal.*, **97** (1985), 189–220.
  - [25] I. Fonseca, The lower quasiconvex envelope of stored energy function for an elastic crystal. *J. Math. Pures et Appl.*, **67** (1988), 175–195.
  - [26] D. Gilbarg and N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*. Springer Verlag, 1985.
  - [27] R.D. James, Basic principles for the improvement of shape-memory and related materials. *Smart Materials, Structure, and Mathematical Issues* (ed. C. Rogers), Technomic Publishing Co., Lancaster, Pennsylvania, PA, 1989.
  - [28] R.D. James, Microstructure and weak convergence. *Material Instabilities in Continuum Mechanics and Related Problems* (ed. J.M. Ball), Oxford University Press, Oxford, 1987, 175–196.
  - [29] R.D. James and D. Kinderlehrer, Theory of diffusionless phase transitions. *Partial Differential Equations and Continuum Models of Phase Transitions* (eds. M. Rascle, D. Serre and M. Slemrod), *Lecture Notes in Physics*, #344, Springer-Verlag, Berlin, New York, 1989, 51–84.
  - [30] D. Kinderlehrer, Remarks about equilibrium configurations of crystals. *Material Instabilities in Continuum Mechanics and Related Problems* (ed. J.M. Ball), Oxford University Press, Oxford, 1987, 217–242.
  - [31] R. Kohn, The relationship between linear and nonlinear variational models of coherent phase transitions. *Proceedings of seventh Army Conference on Applied Mathematics and Computing*, West Point, June 1989.
  - [32] W. Li, Thesis, University of Metz, 1993.
  - [33] P.A. Raviart and J.M. Thomas, *Introduction à l'analyse numérique des équations aux dérivées partielles*. Masson, Paris, 1988.