



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

The cohomological crepant resolution conjecture for $\mathbb{P}(1, 3, 4, 4)$

Boissière, S ; Mann, E ; Perroni, F

DOI: <https://doi.org/10.1142/S0129167X09005479>

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-23480>
Journal Article

Originally published at:

Boissière, S; Mann, E; Perroni, F (2009). The cohomological crepant resolution conjecture for $\mathbb{P}(1, 3, 4, 4)$. *International Journal of Mathematics*, 20(6):791-801.

DOI: <https://doi.org/10.1142/S0129167X09005479>

THE COHOMOLOGICAL CREPANT RESOLUTION CONJECTURE FOR $\mathbb{P}(1, 3, 4, 4)$

SAMUEL BOISSIÈRE, ÉTIENNE MANN AND FABIO PERRONI

ABSTRACT. We prove the cohomological crepant resolution conjecture of Ruan for the weighted projective space $\mathbb{P}(1, 3, 4, 4)$. To compute the quantum corrected cohomology ring we combine the results of Coates-Corti-Iritani-Tseng on $\mathbb{P}(1, 1, 1, 3)$ and our previous results.

1. INTRODUCTION

The *Cohomological Crepant Resolution Conjecture*, as proposed by Ruan, predicts the existence of an isomorphism between the Chen-Ruan cohomology ring of a complex Gorenstein orbifold \mathcal{X} and the quantum corrected cohomology ring of any crepant resolution $\rho : Z \rightarrow X$ of the coarse moduli space X of \mathcal{X} [Rua06]. The quantum corrected cohomology ring of Z is the ring obtained from the small quantum cohomology of Z after specialization of the quantum parameters corresponding to the rational exceptional curves to c_1, \dots, c_m and the remaining parameters to zero, it is denoted by $H_\rho^*(Z)(c_1, \dots, c_m)$ (see Sec. 2.3). It is an important issue (see [Rua06]) to determine the c_1, \dots, c_m . Examples suggest that there can be different choices (see e.g. [BGP08], [CCIT07], [Per07]); however, if one assumes the validity of Conjecture 4.1 in [CR], then Ruan's conjecture follows (Theorem 7.2 in [CR]) and the c_1, \dots, c_m acquire a precise meaning (see section 11 in [CR]).

In this paper we consider the weighted projective space $\mathcal{X} = \mathbb{P}(1, 3, 4, 4)$. The quantum corrected cohomology ring depends on four quantum parameters: q_1, q_2, q_3 and q_4 ; the first three correspond to the components of the exceptional divisor over the transverse singularity, while the fourth corresponds to the component of the exceptional divisor over the isolated singularity (see Sec. 2.3). Our main result, Theorem 4.1, states that, for $(c_1, \dots, c_4) \in \{(i, i, i, 1), (-i, -i, -i, 1)\}$, the quantum corrected cohomology ring of Z is isomorphic to the Chen-Ruan cohomology ring of $\mathbb{P}(1, 3, 4, 4)$. These values of the quantum parameters are relevant for the cohomological crepant resolution conjecture and its generalization, Conj. 4.1 in [CR]. In the previous paper [BMPa], we proved that $H_\rho^*(Z)(q_1, \dots, q_4)$ becomes isomorphic to $H_{\text{CR}}^*(\mathbb{P}(1, 3, 4, 4))$ when we set the quantum parameters to $(i, i, i, 0)$ or $(-i, -i, -i, 0)$, which we found strange in regards to the conjecture,

2000 *Mathematics Subject Classification.* Primary 14M25; Secondary 14A20.

Thanks. This research was partially supported by SNF, No 200020-107464/1.

where the value $c_4 = 0$ is not considered. A motivation for this work was that to clarify this point.

To prove the theorem, we compute explicitly a presentation of $H_\rho^*(Z)(c_1, \dots, c_4)$ with quantum parameters equal to $(i, i, i, 1)$ and $(-i, -i, -i, 1)$, then we give an explicit isomorphism $H_\rho^*(Z; \mathbb{C})(c_1, \dots, c_4) \rightarrow H_{\text{CR}}^*(\mathbb{P}(1, 3, 4, 4))$. The quantum corrections coming from the exceptional divisor over the transverse singularity have been computed in [BMPa]. To compute the quantum corrections coming from the component of the exceptional divisor over the isolated singularity we use results from [CCIT].

The paper is organized as follows. In Sec. 2 we collect some background material: we first give a presentation of the Chen-Ruan cohomology ring of $\mathbb{P}(1, 3, 4, 4)$; then we describe the crepant resolution of $|\mathbb{P}(1, 3, 4, 4)|$; finally we write a presentation of $H_\rho^*(Z)(c_1, \dots, c_4)$ with $(c_1, \dots, c_4) = (i, i, i, c_4)$, here c_4 is a generic complex number and we assume the convergence of the power series (6). The fact that (6) converges in a neighborhood of the origin to an analytic function that admits analytic continuation in $1 \in \mathbb{C}$ is proved in Sec. 3; more precisely we show that (6) is equal to a constant structure of the small quantum cohomology of the crepant resolution of $|\mathbb{P}(1, 1, 1, 3)|$ and then use results from [CCIT]. In the last section 4 we prove Theorem 4.1.

2. BACKGROUND

The weighted projective space $\mathbb{P}(1, 3, 4, 4)$ is the quotient stack $[\mathbb{C}^4 - \{0\}/\mathbb{C}^*]$, where \mathbb{C}^* acts diagonally with weights $w_0 = 1, w_1 = 3, w_2 = 4$ and $w_3 = 4$, it will be denoted by \mathcal{X} . The coarse moduli space $X := |\mathbb{P}(1, 3, 4, 4)|$ is a projective variety whose singular locus is the disjoint union of the curve $C := \{[0 : 0 : x_2 : x_3]\} \subset X$ and the isolated point $P := [0 : 1 : 0 : 0] \in X$. Along C , X has transverse A_3 singularities (see [Per07]), the point P is a singularity of type $\frac{1}{3}(1, 1, 1)$, according to Reid's notation [Rei87].

2.1. The Chen-Ruan cohomology. To compute the Chen-Ruan cohomology ring $H_{\text{CR}}^*(\mathcal{X}; \mathbb{C})$ we follow [BMPb]. The twisted sectors are indexed by the set $\mathbb{T} := \{\exp(2\pi i \gamma) \mid \gamma \in \{0, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}\}\}$. For any $g \in \mathbb{T}$, $\mathcal{X}_{(g)}$ is a weighted projective space: set $\mathbb{I}(g) := \{i \in \{0, 1, 2, 3\} \mid g^{w_i} = 1\}$, then $\mathcal{X}_{(g)} = \mathbb{P}(w_{\mathbb{I}(g)})$, where $(w_{\mathbb{I}(g)}) = (w_i)_{i \in \mathbb{I}(g)}$. The inertia stack is the disjoint union of the twisted sectors:

$$\mathbb{I}\mathcal{X} = \sqcup_{g \in \mathbb{T}} \mathbb{P}(w_{\mathbb{I}(g)}).$$

As a vector space, the Chen-Ruan cohomology is the cohomology of the inertia stack; the graded structure is obtained by shifting the degree of the cohomology of any twisted sector by twice the corresponding age, $\text{age}(g)$. We have

$$\begin{aligned} (1) \quad H_{\text{CR}}^p(\mathcal{X}; \mathbb{C}) &= \bigoplus_{g \in \mathbb{T}} H^{p-2\text{age}(g)}(\mathbb{P}(w_{\mathbb{I}(g)}); \mathbb{C}) \\ &= H^p(\mathbb{P}(1, 3, 4, 4); \mathbb{C}) \oplus H^{p-2}(\mathbb{P}(3); \mathbb{C}) \oplus H^{p-4}(\mathbb{P}(3); \mathbb{C}) \oplus \\ &\quad H^{p-2}(\mathbb{P}(4, 4); \mathbb{C}) \oplus H^{p-2}(\mathbb{P}(4, 4); \mathbb{C}) \oplus H^{p-2}(\mathbb{P}(4, 4); \mathbb{C}). \end{aligned}$$

A basis of $H_{\text{CR}}^*(\mathcal{X}; \mathbb{C})$ is easily obtained in the following way: set

$$H, E_1, E_2, E_3, E_4 \in H_{\text{CR}}^2(\mathcal{X}; \mathbb{C})$$

be the image of $c_1(\mathcal{O}_{\mathcal{X}}(1)) \in H^2(\mathcal{X}; \mathbb{C})$, $1 \in H^0(\mathcal{X}_{(\exp(\pi i/2))}; \mathbb{C})$, $1 \in H^0(\mathcal{X}_{(\exp(\pi i))}; \mathbb{C})$, $1 \in H^0(\mathcal{X}_{(\exp(\pi i/3))}; \mathbb{C})$ and $1 \in H^0(\mathcal{X}_{(\exp(2\pi i/3))}; \mathbb{C})$ respectively, under the inclusion $H^*(\mathbb{P}(w_{1(g)})) \rightarrow H_{\text{CR}}^*(\mathcal{X})$ determined by the decomposition (1). As a commutative \mathbb{C} -algebra, the Chen-Ruan cohomology ring is generated by H, E_1, E_2, E_3, E_4 with relations (see [BMPb]):

$$\begin{aligned} HE_4, E_1E_1 - 3HE_2, E_1E_2 - 3HE_3, E_1E_3 - 3H^2, \\ E_2E_2 - 3H^2, E_2E_3 - HE_1, E_3E_3 - HE_2, 16H^3 - E_4^3, \\ H^2E_1, H^2E_2, H^2E_3, E_1E_4, E_2E_4, E_3E_4. \end{aligned}$$

We see that the following elements form a basis of $H_{\text{CR}}^*(\mathcal{X}; \mathbb{C})$ which we fix for the rest of the paper:

$$(2) \quad 1, H, E_1, E_2, E_3, E_4, H^2, HE_1, HE_2, HE_3, E_4^2, H^3.$$

REMARK 2.1. Note that the elements of our basis are different from those used in [BMPb] by a combinatorial factor.

Other methods are suitable in order to compute the Chen-Ruan cup product of weighted projective spaces, here are a few: the results in [BCS05] provide a presentation of the Chen-Ruan cohomology ring for a general toric Deligne-Mumford stack; results from [Man08] and from [CCLT].

2.2. The crepant resolution. We study some properties of the crepant resolution of $X := |\mathbb{P}(1, 3, 4, 4)|$. We begin with the following

PROPOSITION 2.2. *The variety $X = |\mathbb{P}(1, 3, 4, 4)|$ has a unique crepant resolution $\rho : Z \rightarrow X$, up to isomorphism.*

PROOF. This is a direct consequence of the following facts: the 3-fold singularity $\frac{1}{3}(1, 1, 1)$ has a unique crepant resolution (see e.g. [FN04]) and any variety with transverse ADE singularities has a unique crepant resolution ([Per07], Prop. 4.2), up to isomorphism. \square

An explicit model for the crepant resolution $\rho : Z \rightarrow X$ can be constructed using methods from toric geometry, we follow [Ful93]. The toric variety X is associated to the lattice \mathbb{Z}^3 and the fan Σ , where Σ is the fan whose cones are generated by proper subsets of $\{v_0 := (-3, -4, -4), v_1 := (1, 0, 0), v_2 = (0, 1, 0), v_3 = (0, 0, 1)\} \subset \mathbb{Z}^3$. The 3-dimensional cones which correspond to singular open affine subvarieties of X are: $\sigma_1 := \langle v_0, v_2, v_3 \rangle$, $\sigma_2 := \langle v_0, v_1, v_3 \rangle$, $\sigma_3 := \langle v_0, v_1, v_2 \rangle$. More precisely, σ_1 gives rise to the isolated singularity,

while σ_2 and σ_3 to the transverse one. To resolve the isolated singularity, we subdivide σ_1 by inserting the ray generated by

$$Q_4 := (-1, -1, -1) = \frac{1}{3}v_0 + \frac{1}{3}v_2 + \frac{1}{3}v_3.$$

To resolve the transverse singularity, we subdivide σ_2 and σ_3 by inserting the rays generated by

$$Q_1 := (0, -1, -1) = \frac{1}{4}v_0 + \frac{3}{4}v_1,$$

$$Q_2 := (-1, -2, -2) = \frac{1}{2}v_0 + \frac{1}{2}v_1,$$

$$\text{and } Q_3 := (-2, -3, -3) = \frac{3}{4}v_0 + \frac{1}{4}v_1.$$

The subdivision Σ' is shown in Figure 1. Then we set Z to be the toric variety associated




FIGURE 1. Polar polytopes of $|\mathbb{P}(1, 3, 4, 4)|$ and crepant resolution

to \mathbb{Z}^3 and Σ' , and $\rho : Z \rightarrow X$ to be the morphism associated to the identity $\mathbb{Z}^3 \rightarrow \mathbb{Z}^3$.

LEMMA 2.3. *The morphism $\rho : Z \rightarrow X$ defined above is a crepant resolution.*

PROOF. Since any cone of Σ' is generated by a part of a basis of the lattice, it follows that Z is smooth, [Ful93] Sec. 2.1. The crepancy of ρ follows from the existence of a continuous Σ -piecewise linear function $h' : \mathbb{R}^3 \rightarrow \mathbb{R}$ such that $h'(Q_i) = h'(v_j) = -1$, for any $i \in \{1, 2, 3, 4\}$ and $j \in \{0, 1, 2, 3\}$, [Ful93] Sec. 3.4. \square

Observe that Σ' is the maximal projective subdivision of the polytope of $|\mathbb{P}(1, 3, 4, 4)|$ (see [CK99] or [Voi96]).

2.3. Quantum corrected cohomology ring of Z . Let us denote by b_j (resp. e_i) the first Chern class of the holomorphic line bundle associated to the torus invariant divisor in Z corresponding to the ray generated by v_j (resp. Q_i), for any $j \in \{0, 1, 2, 3\}$ ($i \in \{1, 2, 3, 4\}$ resp.). Set $h = \frac{1}{12} \left(\sum_{i=0}^3 b_i + \sum_{j=1}^4 e_j \right)$, then $H^*(Z; \mathbb{C}) \cong \mathbb{C}[h, e_1, e_2, e_3, e_4]/I$ where I is the ideal generated by (see [BMPa] for more details)

$$\begin{aligned} & 3he_4, e_1e_3, e_1e_4, e_2e_4, e_3e_4, \\ & e_1^2 - 10he_1 - 4he_2 - 2he_3 + 24h^2, \\ & e_1e_2 + 3he_1 + 2he_2 + he_3 - 12h^2, \\ & e_2^2 - 6he_1 - 12he_2 - 2he_3 + 24h^2, \\ & e_2e_3 + 3he_1 + 6he_2 + he_3 - 12h^2, \\ & e_3^2 - 6he_1 - 12he_2 - 14he_3 + 24h^2, \\ & 16h^2e_1, 16h^2e_2, 16h^2e_3, 16h^3 - \frac{1}{27}e_4^3. \end{aligned}$$

Let us fix the following basis of $H^*(Z; \mathbb{C})$:

$$(3) \quad \begin{aligned} \psi_0 &= 1, \psi_1 = h, \psi_2 = e_1, \psi_3 = e_2, \psi_4 = e_3, \psi_5 = e_4, \\ \psi_6 &= h^2, \psi_7 = he_1, \psi_8 = he_2, \psi_9 = he_3, \psi_{10} = e_4^2, \psi_{11} = h^3, \end{aligned}$$

with dual basis:

$$\begin{aligned} \psi^0 &= 48h^3, \psi^1 = 48h^2, \psi^2 = -3he_1 - 2he_2 - he_3, \psi^3 = -2he_1 - 4he_2 - 2he_3, \\ \psi^4 &= -he_1 - 2he_2 - 3he_3, \psi^5 = \frac{1}{9}e_4^2, \psi^6 = 48h, \psi^7 = -3e_1 - 2e_2 - e_3, \\ \psi^8 &= -2e_1 - 4e_2 - 2e_3, \psi^9 = -e_1 - 2e_2 - 3e_3, \psi^{10} = \frac{1}{9}e_4, \psi^{11} = 48 \cdot 1. \end{aligned}$$

Let $M_\rho(Z) \subset A_1(Z; \mathbb{Z})$ be the cone of effective 1-cycles in Z which are contracted by ρ . It is freely generated by $\Gamma_i := \text{PD}(4he_i)$ for $i \in \{1, 2, 3\}$ and $\Gamma_4 := \text{PD}\left(-\frac{1}{3}e_4^2\right)$, where PD means Poincaré dual (see e.g. [BMPa]). Let q_1, q_2, q_3, q_4 be formal variables and let $\Lambda := \mathbb{C}[[q_1, \dots, q_4]]$ be the ring of formal power series in q_1, \dots, q_4 . We have an associative product on $H^*(Z; \Lambda)$ defined as:

$$(4) \quad \psi_i \star_\rho \psi_j := \sum_{\Gamma \in M_\rho(Z)} \sum_{\ell=0}^{11} \langle \psi_i, \psi_j, \psi^\ell \rangle_{0,3,\Gamma}^Z \psi_\ell q^\Gamma,$$

where $\langle \dots \rangle_{0,3,\Gamma}^Z$ is the Gromov-Witten invariant of Z of genus zero, three marked points, homology class Γ and $q^\Gamma := q_1^{d_1} \cdots q_4^{d_4}$ for $\Gamma = d_1\Gamma_1 + \dots + d_4\Gamma_4 \in M_\rho(Z)$.

In [BMPa] we computed explicitly the product (4) whenever $\psi_i \neq e_4$ or $\psi_j \neq e_4$; as a result it follows that, in these cases, the power series involved in (4) converge in a

neighborhood of the origin. We will see in Section 3 that also $e_4 \star_\rho e_4$ is convergent in a neighborhood of the origin. Therefore the expression (4) defines a family of \mathbb{C} -algebras over the vector space $H^*(Z; \mathbb{C})$ whose structure constants are analytic functions defined in some region of the complex space \mathbb{C}^4 . For any c_1, \dots, c_4 in this region, the algebra obtained by setting $q_i = c_i$ is the *quantum corrected cohomology ring* of Z with quantum parameters specialized to c_1, \dots, c_4 and is denoted by $H^*(Z; \mathbb{C})(c_1, \dots, c_4)$ [Rua06].

In particular we have the following presentation for $H_\rho^*(Z)(i, i, i, c_4)$ (see [BMPa] for more details):

$$\begin{aligned}
(5) \quad & \alpha_1 \star_\rho \alpha_2 = \alpha_1 \cup \alpha_2, \quad \text{if } \deg(\alpha_1) \neq 2, \\
& h \star_\rho \alpha = h \cup \alpha, \quad \text{for any } \alpha, \\
& e_1 \star_\rho e_1 = -24h^2 + (-2 + 6i)he_1 - 4he_2 + (-2 - 2i)he_3, \\
& e_1 \star_\rho e_2 = 12h^2 + (-1 - 4i)he_1 + (2 - 4i)he_2 + he_3, \\
& e_1 \star_\rho e_3 = -2ihe_1 - 2ihe_3, \\
& e_2 \star_\rho e_2 = -24h^2 + (2 + 2i)he_1 + 8ihe_2 + (-2 + 2i)he_3, \\
& e_2 \star_\rho e_3 = 12h^2 - he_1 + (-2 - 4i)he_2 + (1 - 4i)he_3, \\
& e_3 \star_\rho e_3 = -24h^2 + (2 - 2i)he_1 + 4he_2 + (2 + 6i)he_3, \\
& e_4 \star_\rho e_4 = \epsilon(c_4)e_4^2,
\end{aligned}$$

where \cup is the usual cup product, and

$$\begin{aligned}
(6) \quad \epsilon(q) & := 1 + \frac{1}{9} \left(\int_{\Gamma_4} e_4 \right)^3 \sum_{a=1}^{\infty} a^3 \deg [\overline{\mathcal{M}}_{0,0}(Z, a\Gamma_4)]^{\text{vir}} q^a \\
& = 1 - 3 \sum_{a=1}^{\infty} a^3 \deg [\overline{\mathcal{M}}_{0,0}(Z, a\Gamma_4)]^{\text{vir}} q^a.
\end{aligned}$$

A similar presentation holds for $H^*(Z)(-i, -i, -i, c_4)$, also in this case $e_4 \star_\rho e_4 = \epsilon(c_4)e_4^2$ with $\epsilon(q)$ defined in (6).

3. RELATIONS WITH THE QUANTUM COHOMOLOGY OF \mathbb{F}_3

To state our main result (Theorem 4.1), we need to know that the series (6) converges in a neighborhood of the origin to an analytic function that admits analytic continuation at the point $1 \in \mathbb{C}$. To this aim, we show in this section that (6) appears as a structure constant of the small quantum cohomology of \mathbb{F}_3 , the crepant resolution of $|\mathbb{P}(1, 1, 1, 3)|$, so that we can use the results of [CCIT] and [CR]. For reader's convenience, we stick to the notation of [CCIT].

Let us consider the coarse moduli space of the weighted projective space $\mathbb{P}(1, 1, 1, 3)$, it has an isolated singularity of type $\frac{1}{3}(1, 1, 1)$. Let $\mathbb{F}_3 := \mathbb{P}(\mathcal{O}_{\mathbb{P}^2}(-3) \oplus \mathcal{O}_{\mathbb{P}^2})$, then there exists a morphism $\chi : \mathbb{F}_3 \rightarrow |\mathbb{P}(1, 1, 1, 3)|$ which is a crepant resolution. Following the

notation of [CCIT], let $p_1 \in H^2(\mathbb{F}_3; \mathbb{C})$ be the class Poincaré dual to the preimage in \mathbb{F}_3 of a hyperplane in \mathbb{P}^2 ; let $p_2 \in H^2(\mathbb{F}_3; \mathbb{C})$ the class Poincaré dual to the infinity section. The cohomology ring of \mathbb{F}_3 has the following presentation

$$H^*(\mathbb{F}_3; \mathbb{C}) = \mathbb{C}[p_1, p_2] / \langle p_1^3, p_2^2 - 3p_1p_2 \rangle.$$

Let us fix the following basis of $H^*(\mathbb{F}_3; \mathbb{C})$ (as in [CCIT])

$$\phi_0 = 1, \phi_1 = \frac{p_2}{3}, \phi_2 = \frac{p_1p_2}{3}, \phi_3 = \frac{p_2 - 3p_1}{3}, \phi_4 = -\frac{p_1(p_2 - 3p_1)}{3}, \phi_5 = \frac{p_1^2p_2}{3},$$

then its dual is

$$\phi^0 = p_1^2p_2, \phi^1 = p_1p_2, \phi^2 = p_2, \phi^3 = -p_1(p_2 - 3p_1), \phi^4 = p_2 - 3p_1, \phi^5 = 3.$$

The Poincaré dual to the exceptional divisor is $\mathbf{p} := p_2 - 3p_1$ (in [CCIT] \mathbf{p} is denoted by \mathbf{p}_1). We want to compute the product $\mathbf{p} \circ_{\hat{q}} \mathbf{p}$ in the small quantum cohomology of \mathbb{F}_3 with $(\hat{q}_1, \hat{q}_2) = (1, 0)$ (see (8)).

By definition, we have (see [CCIT], sec. 2.4)

$$(7) \quad \mathbf{p} \circ_{\hat{q}} \mathbf{p} = \sum_{\gamma \in M(\mathbb{F}_3)} \sum_{\ell=0}^5 \langle \mathbf{p}, \mathbf{p}, \phi_\ell \rangle_{0,3,\gamma}^{\mathbb{F}_3} \hat{q}^\gamma \phi^\ell,$$

where $M(\mathbb{F}_3) \subset H_2(\mathbb{F}_3; \mathbb{Z})$ is the cone of the classes of effective curves, for $\gamma \in M(\mathbb{F}_3)$ $\hat{q}^\gamma : \frac{H^2(\mathbb{F}_3; \mathbb{C})}{2\pi i H^2(\mathbb{F}_3; \mathbb{Z})} \rightarrow \mathbb{C}^*$ is the function $[\tau] \mapsto \exp\left(\int_\gamma \tau\right)$, $\langle \dots \rangle_{0,3,\gamma}^{\mathbb{F}_3}$ is the Gromov-Witten invariant of \mathbb{F}_3 of genus zero, three marked points and class γ .

Since p_1, p_2 form a basis of $H^2(\mathbb{F}_3; \mathbb{C})$, any $[\tau]$ can be written as $[\tau] = [\tau^1 p_1 + \tau^2 p_2]$, therefore

$$(8) \quad \hat{q}^\gamma([\tau]) = \hat{q}_1^{\int_\gamma p_1} \hat{q}_2^{\int_\gamma p_2},$$

where $\hat{q}_i := \exp(\tau^i)$, $i \in \{1, 2\}$. Let us consider the classes $\gamma_1 := \text{PD}(-\frac{1}{3}\mathbf{p}^2 = p_1(p_2 - 3p_1))$ and $\gamma_2 := \text{PD}(p_2^2)$, they form a basis for $M(\mathbb{F}_3)$ such that $\int_{\gamma_i} p_j = \delta_{ij}$, $i, j \in \{1, 2\}$. It follows that the product $\mathbf{p} \circ_{\hat{q}} \mathbf{p}$ restricted to $\hat{q} = (\hat{q}_1, 0)$ is given by:

$$\mathbf{p} \circ_{(\hat{q}_1, 0)} \mathbf{p} = \sum_{a \geq 0} \sum_{\ell=0}^5 \langle \mathbf{p}, \mathbf{p}, \phi_\ell \rangle_{0,3,a\gamma_1}^{\mathbb{F}_3} \hat{q}_1^a \phi^\ell.$$

Using the degree axiom, the divisor axiom and $\int_{\gamma_1} p_2 = 0$, we deduce that

$$\mathbf{p} \circ_{(\hat{q}_1, 0)} \mathbf{p} = \sum_{a=0}^{\infty} \langle \mathbf{p}, \mathbf{p}, \frac{1}{3}\mathbf{p} \rangle_{0,3,a\gamma_1}^{\mathbb{F}_3} \cdot (-p_1(p_2 - 3p_1))^a \hat{q}_1^a.$$

Finally, since $\mathbf{p}^2 := \mathbf{p} \cup \mathbf{p} = -3p_1(p_2 - 3p_1)$, using again the divisor axiom we rewrite the previous expression as

$$(9) \quad \mathbf{p} \circ_{(\hat{q}_1, 0)} \mathbf{p} = \left(1 - 3 \sum_{a=1}^{\infty} a^3 \text{deg} [\overline{\mathcal{M}}_{0,0}(\mathbb{F}_3, a\gamma_1)]^{\text{vir}} \hat{q}_1^a \right) \mathbf{p}^2.$$

Then we have:

LEMMA 3.1. *The power series in (9) is equal to the power series ϵ (see Formula (6)).*

PROOF. From the uniqueness of the crepant resolution of the singularity $\frac{1}{3}(1, 1, 1)$, there exists an isomorphism between a neighborhood in Z of the component of the exceptional divisor over $[0 : 1 : 0 : 0] \in |\mathbb{P}(1, 3, 4, 4)|$ and a neighborhood in \mathbb{F}_3 of the exceptional divisor of χ . The isomorphism between these neighborhoods induces isomorphisms of stacks (see e.g. [Per07], Lemma 7.1):

$$(10) \quad \overline{\mathcal{M}}_{0,0}(Z, a\Gamma_4) \cong \overline{\mathcal{M}}_{0,0}(\mathbb{F}_3, a\gamma_1) \quad \forall a \in \mathbb{N} - \{0\}.$$

Since the virtual fundamental classes of the above stacks depend only on neighborhoods of the exceptional divisors, it follows that

$$\deg [\overline{\mathcal{M}}_{0,0}(Z, a\Gamma_4)]^{\text{vir}} = \deg [\overline{\mathcal{M}}_{0,0}(\mathbb{F}_3, a\gamma_1)]^{\text{vir}} \quad \forall a \in \mathbb{N} - \{0\}.$$

Hence the result follows. \square

As a consequence we have:

COROLLARY 3.2. *The series (6) converges in a neighborhood of the origin to an analytic function $f(q)$ that has analytic continuation in $1 \in \mathbb{C}$. Moreover the value of $f(q)$ at $q = 1$ is*

$$f(1) = \frac{2\pi\beta_1}{9\beta_2^2} = \frac{(2\pi)^6}{27\Gamma(\frac{1}{3})^9},$$

where $\beta_i = \frac{2\pi}{9\Gamma(\frac{i}{3})^3}$, $i \in \{1, 2\}$.

PROOF. It follows from [CCIT] (see also [CR] Theorem 7.2) that the statement is true for (9) and Lemma 3.1 implies that it is also true for (6).

To compute $f(1)$, consider the map $\Theta(q) : H_{\text{CR}}^*(\mathbb{P}(1, 1, 1, 3)) \rightarrow H^*(\mathbb{F}_3)$ defined in [CCIT] (pag. 56) and set $q = 0$. We consider $H^*(\mathbb{F}_3)$ with the algebra structure coming from the quantum cohomology when we set the quantum parameter \hat{q} at $(1, 0)$ (see (8)). $\Theta(0)$ is a morphism of algebras, so we have [CCIT]¹:

$$\frac{2\pi}{3}\beta_1\mathbf{p}^2 = \Theta(0)(\mathbf{1}_{\frac{1}{3}}) = \Theta(0)(\mathbf{1}_{\frac{2}{3}}) \circ_{(1,0)} \Theta(0)(\mathbf{1}_{\frac{2}{3}}) = 3\beta_2^2\mathbf{p} \circ_{(1,0)} \mathbf{p},$$

therefore

$$\mathbf{p} \circ_{(1,0)} \mathbf{p} = f(1)\mathbf{p}^2 = \frac{2\pi\beta_1}{9\beta_2^2}\mathbf{p}^2 \quad \text{and} \quad f(1) = \frac{2\pi\beta_1}{9\beta_2^2}.$$

The second equality follows from standard identities of the Γ -function. \square

¹The careful reader will notice that, in [CCIT], $\mathbb{P}(1, 1, 1, 3)$ is defined as the quotient $[\mathbb{C}^4 - \{0\}/\mathbb{C}^*]$ where the action has negative weights.

4. THE MAIN RESULT

From the results in [BMPa] and Cor. 3.2 it follows that the power series in (4) converge in a neighborhood of the origin of \mathbb{C}^4 to analytic functions that admit analytic continuations in $(i, i, i, 1)$ and in $(-i, -i, -i, 1)$, moreover the following result holds.

THEOREM 4.1. *For $(c_1, c_2, c_3, c_4) \in \{(i, i, i, 1), (-i, -i, -i, 1)\}$ there is a ring isomorphism*

$$H_\rho^*(Z; \mathbb{C})(c_1, c_2, c_3, c_4) \xrightarrow{\cong} H_{\text{CR}}^*(\mathbb{P}(1, 3, 4, 4); \mathbb{C})$$

which is an isometry with respect to the Poincaré pairings.

PROOF. Let us first consider the case where $(c_1, \dots, c_4) = (i, i, i, 1)$. Let

$$(11) \quad \Xi : H_\rho^*(Z; \mathbb{C})(i, i, i, 1) \longrightarrow H_{\text{CR}}^*(\mathbb{P}(1, 3, 4, 4); \mathbb{C})$$

be the linear map defined by the following matrix with respect to the basis (2) and (3),

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\sqrt{2} & -2i & \sqrt{2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i\sqrt{2} & 2i & -i\sqrt{2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sqrt{2} & -2i & -\sqrt{2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \alpha & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\sqrt{2} & -2i & \sqrt{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -i\sqrt{2} & 2i & -i\sqrt{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sqrt{2} & -2i & -\sqrt{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

where α and β are complex numbers to be determined. Note that Ξ coincides with the map (4.3) in [BMPa] except possibly for the image of e_4 and e_4^2 .

Clearly Ξ is an isomorphism of vector spaces if and only if $\alpha \cdot \beta \neq 0$. We have that $\Xi(e_i \star_\rho e_j) = \Xi(e_i) \cup_{\text{CR}} \Xi(e_j)$ for $i, j \in \{1, 2, 3\}$ [BMPa], therefore it remains to find α and β such that $\alpha \cdot \beta \neq 0$, $\Xi(e_4 \star_\rho e_4) = \Xi(e_4) \cup_{\text{CR}} \Xi(e_4)$ and $\Xi(e_4 \star_\rho e_4^2) = \Xi(e_4) \cup_{\text{CR}} \Xi(e_4^2)$. This is equivalent to the equations:

$$(12) \quad \alpha^3 = 27f(1) \quad \text{and} \quad \alpha \cdot \beta = 27.$$

The existence of α and β verifying (12) follows from Corollary 3.2. The fact that Ξ is an isometry is then a direct consequence of [BMPa] and (12).

The case where $(c_1, \dots, c_4) = (-i, -i, -i, 1)$ is similar to the previous one, the linear map $H_\rho^*(Z; \mathbb{C})(-i, -i, -i, 1) \rightarrow H_{\text{CR}}^*(\mathbb{P}(1, 3, 4, 4); \mathbb{C})$ coincides with (4.4) in [BMPa] on $1, e_1, e_2, e_3, h^2, he_1, he_2, he_3, h^3$ and coincides with (11) on e_4 and e_4^2 . Also in this case the

fact that it is a ring isomorphism and that preserves the metrics follows from [BMPa] and (12). This concludes the proof. \square

REFERENCES

- [BCS05] Lev A. Borisov, Linda Chen, and Gregory G. Smith, *The orbifold Chow ring of toric Deligne-Mumford stacks*, J. Amer. Math. Soc. **18** (2005), no. 1, 193–215.
- [BGP08] Jim Bryan, Tom Graber, and Rahul Pandharipande, *The orbifold quantum cohomology of $\mathbb{C}^2/\mathbb{Z}_3$ and Hurwitz-Hodge integrals*, J. Algebraic Geom. **17** (2008), 1–28.
- [BMPa] Samuel Boissiere, Etienne Mann, and Fabio Perroni, *Crepant resolutions of weighted projective spaces and quantum deformations*, math/0610617v2.
- [BMPb] ———, *A model for the orbifold Chow ring of weighted projective spaces*, arXiv:0709.4559.
- [CCIT] Tom Coates, Alessio Corti, Hiroshi Iritani, and Hsian Hua Tseng, *Wall-crossings in toric Gromov-Witten theory I: Crepant examples*, math.AG/0611550.
- [CCIT07] ———, *The crepant resolution conjecture for type A surface singularities*, arXiv:0704.2034.
- [CCLT] Tom Coates, Alessio Corti, Yuan Pin Lee, and Hsian Hua Tseng, *The quantum orbifold cohomology of weighted projective space*, arXiv:math.AG/0608481.
- [CK99] David A. Cox and Sheldon Katz, *Mirror symmetry and algebraic geometry*, Mathematical Surveys and Monographs, vol. 68, American Mathematical Society, Providence, RI, 1999.
- [CR] Tom Coates and Yongbin Ruan, *Quantum cohomology and crepant resolutions: A conjecture*, arXiv:0710.5901.
- [FN04] Baohua Fu and Yoshinori Namikawa, *Uniqueness of crepant resolutions and symplectic singularities*, Ann. Inst. Fourier (Grenoble) **54** (2004), no. 1, 1–19.
- [Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry.
- [Man08] Etienne Mann, *Orbifold quantum cohomology of weighted projective spaces*, J. Algebraic Geom. **17** (2008), 137–166.
- [Per07] Fabio Perroni, *Chen-Ruan cohomology of ADE-singularities*, International Journal of Mathematics **18** (2007), 1–51.
- [Rei87] Miles Reid, *Young person’s guide to canonical singularities*, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), Proc. Sympos. Pure Math., vol. 46, Amer. Math. Soc., Providence, RI, 1987, pp. 345–414.
- [Rua06] Yongbin Ruan, *The cohomology ring of crepant resolutions of orbifolds*, Gromov-Witten theory of spin curves and orbifolds, Contemp. Math., vol. 403, Amer. Math. Soc., Providence, RI, 2006, pp. 117–126.
- [Voi96] Claire Voisin, *Symétrie miroir*, Panoramas et Synthèses [Panoramas and Syntheses], vol. 2, Société Mathématique de France, Paris, 1996.

LABORATOIRE J.A.DIEUDONNÉ UMR CNRS 6621
 UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS
 PARC VALROSE, 06108 NICE
 FRANCE
 E-mail address: sb@math.unice.fr

DÉPARTEMENT DE MATHÉMATIQUES
 UNIVERSITÉ DE MONTPELLIER 2

PLACE EUGÈNE BATAILLON
F-34 095 MONTPELLIER CEDEX 5
E-mail address: emann@math.univ-montp2.fr

INSTITUT FÜR MATHEMATIK
UNIVERSITÄT ZÜRICH
WINTERTHURERSTRASSE 190
8057 ZÜRICH
SWITZERLAND
E-mail address: fabio.perroni@math.unizh.ch