



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2023

Renormalization of twist-two operators in covariant gauge to three loops in QCD

Gehrmann, Thomas ; von Manteuffel, Andreas ; Yang, Tong-Zhi

DOI: [https://doi.org/10.1007/jhep04\(2023\)041](https://doi.org/10.1007/jhep04(2023)041)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-239998>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Gehrmann, Thomas; von Manteuffel, Andreas; Yang, Tong-Zhi (2023). Renormalization of twist-two operators in covariant gauge to three loops in QCD. *Journal of High Energy Physics*, 2023(4):41.

DOI: [https://doi.org/10.1007/jhep04\(2023\)041](https://doi.org/10.1007/jhep04(2023)041)

Renormalization of twist-two operators in covariant gauge to three loops in QCD

Thomas Gehrmann,^a Andreas von Manteuffel^b and Tong-Zhi Yang^{a,b}

^a*Physik-Institut, Universität Zürich,
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

^b*Department of Physics and Astronomy, Michigan State University,
East Lansing, MI 48824, U.S.A.*

E-mail: thomas.gehrmann@uzh.ch, vmante@msu.edu, toyang@physik.uzh.ch

ABSTRACT: The leading short-distance contributions to hadronic hard-scattering cross sections in the operator product expansion are described by twist-two quark and gluon operators. The anomalous dimensions of these operators determine the splitting functions that govern the scale evolution of parton distribution functions. In massless QCD, these anomalous dimensions can be determined through the calculation of off-shell operator matrix elements, typically performed in a covariant gauge, where the physical operators mix with gauge-variant operators of the same quantum numbers. We derive a new method to systematically extract the counterterm Feynman rules resulting from these gauge-variant operators. As a first application of the new method, we rederive the unpolarized three-loop singlet anomalous dimensions, independently confirming previous results obtained with other methods. Employing a general covariant gauge, we observe the explicit cancellation of the gauge parameter dependence in these results.

KEYWORDS: Higher-Order Perturbative Calculations, Renormalization and Regularization, Deep Inelastic Scattering or Small-x Physics

ARXIV EPRINT: [2302.00022](https://arxiv.org/abs/2302.00022)

Contents

1	Introduction	1
2	Notation and formalism	2
2.1	Lagrangian density in massless QCD	2
2.2	Twist-two operators	3
2.3	Renormalization of the twist-two operators	4
2.4	Operator matrix elements	8
3	Computation of bare OMEs up to three loops	11
4	GV counterterm Feynman rules from renormalization conditions	14
4.1	General formulae to determine the Feynman rules for the O_{ABC} operators	15
4.2	Determining the counterterm Feynman rules for $[ZO]_g^{\text{GV},(2)}$ and beyond	17
5	Computation of Feynman rules for the GV operators	19
5.1	Lorentz structures for the Feynman rules of a general twist-two operator	20
5.2	Feynman rules for O_{ABC} up to four legs	22
5.3	Two-loop counterterm Feynman rules for $[ZO]_g^{\text{GV}}$	27
5.4	Comparison with previous fixed n results	34
6	Three-loop splitting functions from operator insertions	36
7	Summary and conclusions	40
A	Feynman rules for physical operators	42
A.1	Feynman rules for the operator O_q	43
A.2	Feynman rules for the operator O_g	44
B	Feynman rules for all GV operators at lowest multiplicity	45
C	Renormalization constants Z_{qA} and Z_{gA} to three loops	46
D	Standard QCD renormalization constants	54
E	Instructions for supplementary material	56

1 Introduction

The operator product expansion (OPE) [1, 2] provides an elegant method to separate short-distance from long-distance contributions in quantum field theory. Its early application to deeply inelastic lepton-nucleon scattering processes [3] in quantum chromodynamics (QCD) successfully predicted the violation of Bjorken scaling [4, 5], thereby enabling the development of the QCD-improved parton model [6]. The anomalous dimensions of

quark and gluon operators in the OPE are directly related to the Altarelli-Parisi splitting functions [6–8] of the QCD-improved parton model by an inverse Mellin transformation.

The splitting functions determine the scale evolution of the parton distributions, which are an essential ingredient to all quantitative predictions for high-energy hadron collider processes. The precise determination of parton distributions requires the iterated comparison of highly accurate experimental data for a multitude of processes with theoretical predictions at a comparable level of precision. These predictions require higher order perturbative corrections [9] to the underlying hard scattering processes as well as to the splitting functions.

Splitting functions are currently known to three-loop order in QCD [10, 11], which enables a consistent description of hadron collider processes at next-to-next-to-leading order (NNLO) in perturbative QCD. Despite its early success and its computational simplicity, the OPE method has not played a significant role in this progress towards precision QCD for collider observables. Its applicability at higher loop orders is limited by the currently incomplete understanding of the renormalization of singlet quark and gluon operators, which involves the mixing with so-called gauge-variant (GV) operators, that are unphysical operators resulting from the gauge fixing in QCD. Although the existence of such operators has been known [12, 13] since the initial applications of the OPE in QCD, it has not been possible to determine the number and the form of these GV operators — or even only the renormalization counterterms that result from them — beyond what is required for two-loop calculations [14].

In this paper, we revisit the long-standing question of the renormalization of the leading-twist quark and gluon operators, whose anomalous dimensions determine the scale evolution of parton distribution functions. We devise a new method to extract the Feynman rules for renormalization counterterms that result from GV operators, through the computation of multi-leg operator matrix elements. We apply the newly developed method to determine all counterterms required in the OPE up to three loops in QCD. Section 2 establishes our notation and briefly summarizes the OPE of QCD. We describe the calculation of operator matrix elements (OMEs) up to three loops in section 3. Our method for the determination of Feynman rules for operators and counterterms is developed in section 4 and, in section 5, applied to compute all counterterms that are required for the three-loop renormalization. These counterterms are used in section 6 to rederive the three-loop anomalous dimensions, thereby rigorously establishing their independence of the QCD gauge parameter. We conclude with an extensive summary of our method and key results in section 7.

2 Notation and formalism

2.1 Lagrangian density in massless QCD

The dynamics of quarks and gluons are controlled by the QCD Lagrangian. The classical gauge invariant Lagrangian of QCD is given by

$$\mathcal{L}_c = \bar{\psi}_i \dot{\gamma}^\mu (D_\mu)_{ij} \psi_j - \frac{1}{4} G_{\mu\nu}^a G_a^{\mu\nu}, \quad (2.1)$$

where ψ_i is the quark field in the fundamental representation of SU(3) gauge group and i is the color index. Further, $(D_\mu)_{ij}$ is the gauge-covariant derivative in the fundamental

representation,

$$(D_\mu)_{ij} = \partial_\mu \delta_{ij} - ig_s (T^a)_{ij} A_\mu^a, \quad (2.2)$$

and $G_{\mu\nu}^a$ is the gluon field strength tensor,

$$G_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g_s f^{abc} A_\mu^b A_\nu^c, \quad (2.3)$$

with gluon fields A_μ^a in the adjoint representation of the SU(3) gauge group.

The classical QCD Lagrangian is invariant under the following infinitesimal gauge transformations,

$$A_\mu^a \rightarrow A_\mu^a + D_\mu^{ab} \theta^b, \quad (2.4)$$

where the covariant derivative D_μ^{ab} in the adjoint representation is given by

$$D_\mu^{ab} = \partial_\mu \delta^{ab} - g_s f^{abc} A_\mu^c. \quad (2.5)$$

The gauge freedom causes difficulties when trying to quantize gauge field theories. One way to deal with it is the introduction of a gauge fixing term in the Lagrangian to eliminate the problematic gauge freedom. This procedure leads to a ghost term, which cancels the unphysical longitudinal degrees of the gauge field. The commonly used gauge-fixing and ghost terms in covariant gauge are

$$\mathcal{L}_{\text{GF+ghost}} = -\frac{1}{2\xi} (\partial^\mu A_\mu^a)^2 - \bar{c}^a \partial^\mu D_\mu^{ab} c^b, \quad (2.6)$$

where c^a is the ghost field, \bar{c}^a is the anti-ghost field and ξ is the gauge parameter with $\xi = 1$ being the 't Hooft-Feynman gauge. The full QCD Lagrangian is thus

$$\mathcal{L}_{\text{QCD}} = \mathcal{L}_c + \mathcal{L}_{\text{GF+ghost}}, \quad (2.7)$$

in covariant gauge.

2.2 Twist-two operators

To study the collinear behavior of QCD, we consider the operator product expansion. In the following, we focus entirely on twist-two operators, which encode the collinear physics at leading power. According to the flavour group, the twist-two operators are divided into non-singlet and singlet parts. The non-singlet operators of spin n and twist two are given by

$$O_{q,k}^{\mu_1 \cdots \mu_n} = \frac{i^{n-1}}{2} \mathcal{S} \left[\bar{\psi}_{i_1} \gamma^{\mu_1} D_{i_1 i_2}^{\mu_2} D_{i_2 i_3}^{\mu_3} \cdots D_{i_{n-1} i_n}^{\mu_n} \frac{\lambda_k}{2} \psi_{i_n} - \text{trace terms} \right], \quad k = 3, \cdots, n_f^2 - 1, \quad (2.8)$$

where $\lambda_k/2$ is a diagonal generator of the flavour group SU(n_f) and \mathcal{S} denotes the symmetrization of Lorentz indices $\mu_1 \cdots \mu_n$. Since only the traceless part of the operator is of relevance, trace terms are subtracted to render the operator definition traceless.

There are two operators in the flavor singlet case. The twist-two singlet quark and gluon operators that are obtained from the OPE are

$$\begin{aligned} O_q^{\mu_1 \cdots \mu_n} &= \frac{i^{n-1}}{2} \mathcal{S} \left[\bar{\psi}_{i_1} \gamma^{\mu_1} D_{i_1 i_2}^{\mu_2} D_{i_2 i_3}^{\mu_3} \cdots D_{i_{n-1} i_n}^{\mu_n} \psi_{i_n} - \text{trace terms} \right], \\ O_g^{\mu_1 \cdots \mu_n} &= -\frac{i^{n-2}}{2} \mathcal{S} \left[G_{a_1 \mu}^{\mu_1} D_{a_1 a_2}^{\mu_2} \cdots D_{a_{n-2} a_{n-1}}^{\mu_{n-1}} G_{a_{n-1} a_n}^{\mu_n} - \text{trace terms} \right], \end{aligned} \quad (2.9)$$

where the covariant derivatives in the quark and gluon operators are defined in (2.2) and (2.5) respectively. No momentum transfer is associated with the operators.

Since the operators are traceless, we can extract the information of interest by contracting the operators with an external source

$$J_{\mu_1 \dots \mu_n} = \Delta_{\mu_1} \Delta_{\mu_2} \dots \Delta_{\mu_n} \tag{2.10}$$

where Δ is a light-like vector $\Delta^2 = 0$. We thus define the following spin- n twist-two operators

$$\begin{aligned} O_{q,k} &= O_{q,k}^{\mu_1 \dots \mu_n} J_{\mu_1 \dots \mu_n}, \\ O_q &= O_q^{\mu_1 \dots \mu_n} J_{\mu_1 \dots \mu_n}, \\ O_g &= O_g^{\mu_1 \dots \mu_n} J_{\mu_1 \dots \mu_n}. \end{aligned} \tag{2.11}$$

Here and in the following, the dependence of $O_{q,k}$, O_q and O_g on n is understood.

To compute matrix elements of these operators for a given set of external parton states, one needs the corresponding operator Feynman rules, which can be obtained from (2.9) by a functional variation. In addition to the primary $q\bar{q}$ and gg states for O_q and O_g respectively, the covariant derivatives in (2.9) lead to Feynman rules for an arbitrary number of additional gluons. Since each additional gluon contributes a factor g_s , only a finite number of these operator Feynman rules need to be considered at a given order in perturbation theory. These Feynman rules can be cast into an all- n form [15], as outlined in section 3 below.

2.3 Renormalization of the twist-two operators

The non-singlet operator $O_{q,k}$ is distinguished from singlet operators by quark flavour, and can therefore be renormalized separately. The renormalized non-singlet operator is defined by

$$O_{q,k}^R = Z_{\text{ns}} O_{q,k}^B, \tag{2.12}$$

where Z_{ns} is a multiplicative renormalization constant. Here and in the following, the superscripts R and B denote the renormalized and bare operators, respectively.

The two singlet operators belong to the same irreducible representation and mix with each other under renormalization. Due to the absence of further physical operators in the same representation, one could naively expect the following operator renormalization to hold:

$$\begin{pmatrix} O_q \\ O_g \end{pmatrix}^{\text{R,naive}} = \begin{pmatrix} Z_{qq} & Z_{qg} \\ Z_{gq} & Z_{gg} \end{pmatrix} \begin{pmatrix} O_q \\ O_g \end{pmatrix}^B. \tag{2.13}$$

As we will discuss further below, this picture needs to be extended to include further, non-physical operators.

First let us note that, for non-singlet and singlet operators, the anomalous dimension γ can be extracted from the renormalization constant according to

$$\frac{dZ}{d \ln \mu} = -2\gamma \cdot Z. \tag{2.14}$$

For the non-singlet case, we have $Z = Z_{\text{ns}}$ and $\gamma = \gamma_{\text{ns}}$. For the singlet case, both Z and γ are two-by-two matrices due to operator mixing:

$$Z = \begin{pmatrix} Z_{qq} & Z_{qg} \\ Z_{gq} & Z_{gg} \end{pmatrix}, \quad \gamma = \begin{pmatrix} \gamma_{qq} & \gamma_{qg} \\ \gamma_{gq} & \gamma_{gg} \end{pmatrix}. \quad (2.15)$$

We expand the renormalization constants perturbatively according to

$$Z = \sum_{i=0}^{\infty} a_s^i Z^{(i)}, \quad (2.16)$$

where we have defined $a_s = \alpha_s/(4\pi)$. For the anomalous dimensions, we use

$$\gamma = \sum_{i=0}^{\infty} a_s^{i+1} \gamma^{(i)}. \quad (2.17)$$

By using the definition of the d -dimensional QCD β function

$$\beta(a_s, \epsilon) = \frac{da_s}{d \ln \mu} = -2\epsilon a_s - 2a_s \sum_{i=0}^{\infty} a_s^{i+1} \beta_i, \quad (2.18)$$

with ϵ being the dimensional regulator $\epsilon = (4 - d)/2$, we can express the renormalization constant Z in terms of the anomalous dimension. In non-singlet case, the renormalization constant is

$$\begin{aligned} Z_{\text{ns}} = & 1 + a_s \frac{\gamma_{\text{ns}}^{(0)}}{\epsilon} + a_s^2 \left(\frac{\gamma_{\text{ns}}^{(1)}}{2\epsilon} + \frac{1}{2\epsilon^2} \left[-\beta_0 \gamma_{\text{ns}}^{(0)} + (\gamma_{\text{ns}}^{(0)})^2 \right] \right) \\ & + a_s^3 \left(\frac{1}{3\epsilon} \gamma_{\text{ns}}^{(2)} + \frac{1}{6\epsilon^2} \left[-2\beta_1 \gamma_{\text{ns}}^{(0)} - 2\beta_0 \gamma_{\text{ns}}^{(1)} + 3\gamma_{\text{ns}}^{(0)} \gamma_{\text{ns}}^{(1)} \right] \right. \\ & \left. + \frac{1}{6\epsilon^3} \left[2\beta_0^2 \gamma_{\text{ns}}^{(0)} - 3\beta_0 (\gamma_{\text{ns}}^{(0)})^2 + (\gamma_{\text{ns}}^{(0)})^3 \right] \right) + \mathcal{O}(a_s^4). \end{aligned} \quad (2.19)$$

In the singlet case, one has

$$\begin{aligned} Z_{ij} = & \delta_{ij} + a_s \frac{\gamma_{ij}^{(0)}}{\epsilon} + a_s^2 \left(\frac{\gamma_{ij}^{(1)}}{2\epsilon} + \frac{1}{2\epsilon^2} \left[-\beta_0 \gamma_{ij}^{(0)} + \sum_{k=q,g} \gamma_{ik}^{(0)} \gamma_{kj}^{(0)} \right] \right) \\ & + a_s^3 \left(\frac{1}{3\epsilon} \gamma_{ij}^{(2)} + \frac{1}{6\epsilon^2} \left[-2\beta_1 \gamma_{ij}^{(0)} - 2\beta_0 \gamma_{ij}^{(1)} + 2 \sum_{k=q,g} \gamma_{ik}^{(1)} \gamma_{kj}^{(0)} + \sum_k \gamma_{ik}^{(0)} \gamma_{kj}^{(1)} \right] \right. \\ & \left. + \frac{1}{6\epsilon^3} \left[2\beta_0^2 \gamma_{ij}^{(0)} - 3\beta_0 \sum_{k=q,g} \gamma_{ik}^{(0)} \gamma_{kj}^{(0)} + \sum_{k=q,g} \sum_{l=q,g} \gamma_{ik}^{(0)} \gamma_{kl}^{(0)} \gamma_{lj}^{(0)} \right] \right) + \mathcal{O}(a_s^4), \end{aligned} \quad (2.20)$$

where $i, j = q, g$. By extracting the anomalous dimensions, one can subsequently determine the splitting functions, see the end of section 6.

It was already pointed out by Gross and Wilczek in the first calculation of the one-loop singlet anomalous dimensions [3] that the quark and gluon operators may mix with further

GV operators under renormalization. These GV operators originate from the interplay of gauge-fixing at the QCD Lagrangian level and the OPE, with the twist-two operator basis (2.9) being obtained from the classical gauge-invariant QCD Lagrangian (2.1) prior to gauge fixing.

The derivation of a consistent OPE on the basis of the gauge-fixed QCD Lagrangian (2.7) has been investigated extensively in the literature. In the seminal work of Dixon and Taylor [12], the set of GV operators relevant at order g_s was constructed explicitly. These results demonstrated that the naive renormalization of the singlet operators is consistent at the one-loop level, and, subsequently, enabled the construction of the correct singlet renormalization at two loops [14], thereby resolving earlier inconsistencies [16, 17]. All-order renormalization conditions for the gauge-invariant operators O_q and O_g were derived by Joglekar and Lee [18], confirming the results of [12] but lacking a procedure for the construction of GV operators at higher orders in g_s . An independent approach to the renormalization of the singlet operators was based on the renormalization of the QCD energy-momentum tensor [19, 20], equally yielding [21] the GV operators at order g_s . This order is sufficient for the extraction of the anomalous dimension matrix of the singlet operators up to two loops. Previous calculations of the three-loop anomalous dimensions did not employ the OPE, but used the forward scattering amplitude in deep inelastic scattering [11, 22], inclusive hadron collider cross sections [23, 24] or beam functions [25–29] to determine the three-loop mass factorization counterterms of parton distributions, which contain the required anomalous dimensions.

Most recently, Falcioni and Herzog [30] translated the conditions formulated by Joglekar and Lee [18] into a set of constraint equations, that allow to infer the GV operators and their associated renormalization constants at fixed n , order-by-order in the number of loops and powers of the strong coupling constant. They demonstrated their method in the derivation of the three-loop anomalous dimensions for $n \leq 6$ and the four-loop anomalous dimensions for $n \leq 4$.

In the following, we propose to overcome the current lack of understanding of the full structure of the GV operators by devising a procedure for the direct extraction of the all- n counterterm Feynman rules resulting from these operators. We start from the generic form of the renormalization of the singlet operators, including their mixing with the GV operators.

The most general form of the renormalization for O_q and O_g can be written as follows,

$$\begin{aligned}
 O_q^R &= Z_{qq} O_q^B + Z_{qg} O_g^B + \sum_{i=1}^{\infty} \sum_{j=1}^{N_i} \left(Z_{qA_{i,j}} O_{A_{i,j}}^B + Z_{qB_{i,j}} O_{B_{i,j}}^B + Z_{qC_{i,j}} O_{C_{i,j}}^B \right), \\
 O_g^R &= Z_{gq} O_q^B + Z_{gg} O_g^B + \sum_{i=1}^{\infty} \sum_{j=1}^{N_i} \left(Z_{gA_{i,j}} O_{A_{i,j}}^B + Z_{gB_{i,j}} O_{B_{i,j}}^B + Z_{gC_{i,j}} O_{C_{i,j}}^B \right). \quad (2.21)
 \end{aligned}$$

In the above equations, the GV operators carry subscripts A , B and C to distinguish three kinds of operators: $O_{A_{i,j}}$ involve gluon fields only, $O_{B_{i,j}}$ two quark fields plus gluon fields, and $O_{C_{i,j}}$ two ghost fields plus gluon fields. The index i assigned to the operators is used to indicate the order in the strong coupling constant at which each operator starts to

contribute to the renormalization of the physical quark and gluon operators, such that the renormalization constants $Z_{qA_{i,j}}, Z_{qB_{i,j}}, Z_{qC_{i,j}}$ start at $\mathcal{O}(a_s^{i+1})$, and $Z_{gA_{i,j}}, Z_{gB_{i,j}}, Z_{gC_{i,j}}$ start at $\mathcal{O}(a_s^i)$. In principle, the index i extends to infinity. However, we only need operators up to finite i for practical computations at a given fixed loop order. For example, we need $i \leq 2$ to extract three-loop splitting functions and $i \leq 3$ to extract four-loop splitting functions. We also assign another index j to enumerate N_i different operator structures, which contribute to the same index i but require their own (n dependent) renormalization constant. We note that the number N_i is in general not known and could a priori be infinite.

Our key idea is to directly extract the counterterm Feynman rules resulting from a linear combination of operators instead of determining the operators themselves or determining the Feynman rules resulting from each operator separately. From explicit computations described in section 5 below, we find that $N_1 = 1$ and $N_2 > 1$; it even seems possible that $N_{\geq 2} = \infty$. In other words, we can disentangle the operator from its corresponding renormalization constant for $i = 1$, but not for $i \geq 2$. Furthermore, we find that the type A, B, and C renormalization constants for $i = 1$ are identical at the lowest order in a_s . We expect this to hold to all orders, such that the following relations are fulfilled:

$$\begin{aligned} Z_{qA_{1,1}} &= Z_{qB_{1,1}} = Z_{qC_{1,1}} \equiv Z_{qA}, \\ Z_{gA_{1,1}} &= Z_{gB_{1,1}} = Z_{gC_{1,1}} \equiv Z_{gA}. \end{aligned} \tag{2.22}$$

It is then sufficient to consider the following combination of $i = 1$ type operators,

$$O_{ABC} = O_A + O_B + O_C, \tag{2.23}$$

where we have abbreviated $O_{A_{1,1}} \equiv O_A$, $O_{B_{1,1}} \equiv O_B$, and $O_{C_{1,1}} \equiv O_C$. It is possible to prove the relations (2.22) in the context of BRST symmetry, noting that only the combination shown in (2.23) is compatible with the requirement of transversality of the physical gluon fields. Given the simplicity of the $i = 1$ type operators, as compared to all $i \geq 2$ type operators, it is natural to treat them separately. By defining the following counterterm operators as the linear combination of all $i \geq 2$ type GV operators,

$$\begin{aligned} [ZO]_q^{\text{GV}} &= \sum_{i=2}^{\infty} \sum_{j=1}^{N_i} \left(Z_{qA_{i,j}} O_{A_{i,j}}^{\text{B}} + Z_{qB_{i,j}} O_{B_{i,j}}^{\text{B}} + Z_{qC_{i,j}} O_{C_{i,j}}^{\text{B}} \right), \\ [ZO]_g^{\text{GV}} &= \sum_{i=2}^{\infty} \sum_{j=1}^{N_i} \left(Z_{gA_{i,j}} O_{A_{i,j}}^{\text{B}} + Z_{gB_{i,j}} O_{B_{i,j}}^{\text{B}} + Z_{gC_{i,j}} O_{C_{i,j}}^{\text{B}} \right), \end{aligned} \tag{2.24}$$

equation (2.21) can be simplified as follows:

$$O_q^{\text{R}} = Z_{qq} O_q^{\text{B}} + Z_{qg} O_g^{\text{B}} + Z_{qA} \left(O_A^{\text{B}} + O_B^{\text{B}} + O_C^{\text{B}} \right) + [ZO]_q^{\text{GV}}, \tag{2.25}$$

$$O_g^{\text{R}} = Z_{gq} O_q^{\text{B}} + Z_{gg} O_g^{\text{B}} + Z_{gA} \left(O_A^{\text{B}} + O_B^{\text{B}} + O_C^{\text{B}} \right) + [ZO]_g^{\text{GV}}. \tag{2.26}$$

We used symbols $[ZO]_q^{\text{GV}}$ and $[ZO]_g^{\text{GV}}$ for the reason that we can not disentangle the renormalization constant from the corresponding operator, otherwise, we should write them

as $Z_{qV} O_V$ and $Z_{gV} O_V$, where V stands for any $i \geq 2$ type GV operator. The counterterm operators can be further decomposed as follows according to the number of loops,

$$[ZO]_q^{\text{GV}} = \sum_{l=3}^{\infty} a_s^l [ZO]_q^{\text{GV},(l)}, \quad [ZO]_g^{\text{GV}} = \sum_{l=2}^{\infty} a_s^l [ZO]_g^{\text{GV},(l)}, \quad (2.27)$$

where the expansions apply to the renormalization constants, for example

$$[ZO]_g^{\text{GV},(2)} = \sum_{j=1}^{N_2} \left(Z_{gA_{2,j}}^{(2)} O_{A_{2,j}} + Z_{gB_{2,j}}^{(2)} O_{B_{2,j}} + Z_{gC_{2,j}}^{(2)} O_{C_{2,j}} \right). \quad (2.28)$$

As we stated above, the renormalization of physical operators mixes with GV operators. However, the renormalization of GV operators can not mix with physical operators, as shown by Joglekar and Lee [18]. It means that the eigenvalues of the corresponding mixing matrix factorize into physical and non-physical parts. In our context, it implies that the inclusion of GV operators affects only the determination of physical renormalization constants Z_{kj} with $k, j = q$ or g , but not the extraction of physical anomalous dimensions from the renormalization constants, such that (2.20) remains valid even with the presence of GV operators. By also considering the renormalization of the $i = 1$ type GV operators, the equations (2.25) and (2.26) are written as the following matrix form,

$$\begin{pmatrix} O_q \\ O_g \\ O_{ABC} \end{pmatrix}^{\text{R}} = \begin{pmatrix} Z_{qq} & Z_{qg} & Z_{qA} \\ Z_{gq} & Z_{gg} & Z_{gA} \\ 0 & 0 & Z_{AA} \end{pmatrix} \begin{pmatrix} O_q \\ O_g \\ O_{ABC} \end{pmatrix}^{\text{B}} + \begin{pmatrix} [ZO]_q^{\text{GV}} \\ [ZO]_g^{\text{GV}} \\ [ZO]_A^{\text{GV}} \end{pmatrix}^{\text{B}}, \quad (2.29)$$

where we have introduced another counterterm operator $[ZO]_A^{\text{GV}}$ renormalizing the operator O_{ABC} defined in (2.23).

We emphasize again that the distinction between operators of type $i = 1$ and of types $i \geq 2$ in (2.29) is a choice made by us for the sake of computational simplicity, and will be justified in detail in section 5 below. Alternatively, one could choose not to single out the $i = 1$ operator O_{ABC} and include its contributions in the counterterm operators used for $i \geq 2$.

2.4 Operator matrix elements

To extract the renormalization constants as well as to derive the counterterm Feynman rules of the GV operators, we need to introduce the concept of OMEs which are defined as correlation functions or matrix elements with an operator insertion

$$A_{ij} = \langle j(p) | O_i | j(p) \rangle, \quad (2.30)$$

where O_i is a twist-two operator, j denotes a quark, gluon or ghost state, and p is the momentum of j . Up to three-loop order, the results for the above OMEs can be conveniently expressed in terms of Casimir invariants C_A , C_F , and $d_{abc} d^{abc}$, and the number of massless

quark flavours N_f . For an $SU(N_c)$ gauge group, the Casimir invariants or color factors read

$$\begin{aligned} C_A &= N_c, \\ C_F &= \frac{N_c^2 - 1}{2N_c}, \\ d_{abc}d^{abc} &= 4 \text{Tr}[(T_a T_b + T_b T_a) T_c] \text{Tr}[(T^a T^b + T^b T^a) T^c] = \frac{(N_c^2 - 1)(N_c^2 - 4)}{N_c}, \end{aligned} \quad (2.31)$$

with $N_c = 3$ for QCD. We use the normalization

$$\text{Tr}(T^a T^b) = T_F \delta^{ab} = \frac{1}{2} \delta^{ab}. \quad (2.32)$$

The OMEs with two-quark external states can be decomposed into their physical part and an equation-of-motion (EOM) part, which are described by form factors A_{iq}^{Phy} and A_{iq}^{EOM} , respectively,

$$A_{iq} = \langle q(p) | O_i | q(p) \rangle = A_{iq}^{\text{Phy}} (\Delta \cdot p)^{n-1} \not{\Delta} + A_{iq}^{\text{EOM}} (\Delta \cdot p)^n \not{p}. \quad (2.33)$$

The OMEs with two-ghost external states involve only a single form factor,

$$A_{ic} = \langle c(p) | O_i | c(p) \rangle = A_{ic,1} (\Delta \cdot p)^n, \quad (2.34)$$

where we use c to represent a ghost state and the index 1 in $A_{ic,1}$ is used to distinguish $A_{ic,1}$ from A_{ic} . Similarly, the decomposition of OMEs with two-gluon external states is given in terms of four form factors,

$$A_{ig}^{\mu\nu} = \langle g(p) | O_i | g(p) \rangle^{\mu\nu} = \sum_{k=1}^4 A_{ig,k} T_k^{\mu\nu}, \quad (2.35)$$

with the four tensor structures

$$\begin{aligned} T_1^{\mu\nu} &= \frac{1 + (-1)^n}{2} \left[(\Delta \cdot p)^2 g^{\mu\nu} - \Delta \cdot p (p^\mu \Delta^\nu + \Delta^\mu p^\nu) + \Delta^\mu \Delta^\nu p^2 \right] (\Delta \cdot p)^{n-2}, \\ T_2^{\mu\nu} &= \frac{1 + (-1)^n}{2} \left[\frac{p^\mu p^\nu}{p^2} (\Delta \cdot p)^2 - \Delta \cdot p (p^\mu \Delta^\nu + \Delta^\mu p^\nu) + \Delta^\mu \Delta^\nu p^2 \right] (\Delta \cdot p)^{n-2}, \\ T_3^{\mu\nu} &= \frac{1 + (-1)^n}{2} \left[-\Delta \cdot p (p^\mu \Delta^\nu + \Delta^\mu p^\nu) + 2\Delta^\mu \Delta^\nu p^2 \right] (\Delta \cdot p)^{n-2}, \\ T_4^{\mu\nu} &= \frac{1 + (-1)^n}{2} \left[\Delta \cdot p (p^\mu \Delta^\nu + \Delta^\mu p^\nu) + 2\Delta^\mu \Delta^\nu p^2 \right] (\Delta \cdot p)^{n-2}, \end{aligned} \quad (2.36)$$

where $n \geq 2$. The above tensor structures satisfy the relations

$$p_\mu T_i^{\mu\nu} = 0 \quad (i = 1, 2), \quad p_\mu T_i^{\mu\nu} \neq 0 \quad (i = 3, 4), \quad (2.37)$$

$$p_\mu p_\nu T_i^{\mu\nu} = 0 \quad (i = 1, 2, 3), \quad p_\mu p_\nu T_i^{\mu\nu} \neq 0 \quad (i = 4). \quad (2.38)$$

Once we determined all GV counterterm Feynman rules for the renormalization of the physical twist-two operators, the physical, EOM, or non-physical form factors in (2.33)

and (2.35) should be renormalized independently. For the purpose of extracting the splitting functions, it is sufficient to consider the renormalization of the following form factors,

$$\begin{aligned}
 \mathcal{F}_{iq} &= \frac{1}{2N_c} \cdot \frac{\text{Tr}(\not{p}A_{iq})}{(\Delta \cdot p)^n}, \\
 \mathcal{F}_{ig} &= \frac{1}{(d-2)(N_c^2-1)} \cdot \frac{(-g_{\mu\nu}A_{ig}^{\mu\nu})}{(\Delta \cdot p)^n}, \\
 \mathcal{F}_{ic} &= \frac{1}{(d-2)(N_c^2-1)} \cdot \frac{A_{ic}}{(\Delta \cdot p)^n},
 \end{aligned}
 \tag{2.39}$$

where we averaged and summed over spins and colors of incoming and outgoing states, respectively. We employ the perturbative expansion

$$\mathcal{F} = \sum_{l=0}^{\infty} \mathcal{F}^{(l)} a_s^l
 \tag{2.40}$$

for the form factors. With the above definitions, both \mathcal{F}_{qq} and \mathcal{F}_{gg} are normalized to unity at lowest order,

$$\mathcal{F}_{qq}^{(0)} = \mathcal{F}_{gg}^{(0)} = 1.
 \tag{2.41}$$

If the external state j in (2.30) is on-shell, the OMEs of the GV operators do not mix under renormalization with those of the physical operators and the naive renormalization procedure (2.13) remains valid. Likewise, the GV operators do not mix with the polarized analogues of the singlet operators (2.9), which involve the difference of the quark or gluon spin states instead of their sums. This can be understood from the structure of the Fock space in a covariant gauge, where the ghost degrees of freedom mix only with the unphysical gauge field polarization states. In unpolarized OMEs, all polarization states (including the unphysical ones) are summed over, while polarized OMEs are constructed as differences between the two physical polarization states, thereby decoupling the unphysical sector of the Fock space. Consequently, the renormalization of polarized operators also fulfils (2.13).

Another approach to avoid GV operators is to perform the quantization in an axial gauge, which requires to introduce a light-like reference direction n^μ for the quantization. In an axial gauge, the Fock spaces of physical and unphysical polarizations decouple completely, again allowing for the naive operator renormalization (2.13).

The form factors appearing on the right-hand sides of (2.33), (2.34), and (2.35) are Lorentz scalars. By dimensional analysis, one finds that they vanish trivially in massless QCD for external on-shell states if dimensional regularization is applied. To generate a mass scale, one either needs to insert an internal mass scale or to consider off-shell external states $p^2 < 0$. Working with an internal quark mass, three-loop on-shell OME have been used to extract heavy-flavour contributions to the three-loop anomalous dimensions [31–35], as well as for the computation of a subset of the genuine massless anomalous dimensions at three loops [36].

In terms of computational simplicity, the calculations of purely massless off-shell OMEs are preferable over on-shell OMEs with an internal mass, since the underlying all-massless




Figure 1. Representative 3-loop Feynman diagrams with physical operator insertions to extract 3-loop splitting functions.

Feynman integrals are considerably simpler than corresponding integrals involving an internal mass scale.

Massless off-shell OMEs have been computed up to three-loop order and used for the determination of the three-loop non-singlet anomalous dimensions [37] and for the polarized singlet anomalous dimensions at two loops [38, 39] and three loops [40]. In all these cases, the GV operators do not contribute.

The first successful extraction of the two-loop singlet anomalous dimensions [41–43] was based on performing QCD quantization in an axial gauge (where the GV operators decouple [44]) and computing all-massless off-shell OMEs. Owing to the presence of the gauge vector n^μ , the resulting Feynman integrals are considerably more complicated than in a covariant gauge, such that the use of an axial gauge is not a viable option at the higher loop orders.

3 Computation of bare OMEs up to three loops

Before working out the Feynman rules for the GV operators, we first explain how to compute the bare OMEs for the physical operators O_q and O_g to three-loop order. The Feynman rules resulting from these operators are summarized in appendix A.

As the first step, we generate all relevant Feynman diagrams to three-loop order using QGRAF [45]. Some representative diagrams are shown in figure 1. To translate the diagrams into expressions, one needs the standard QCD Feynman rules as well as the Feynman rules for the twist-two operators. As shown in the appendix A, the Feynman rules for the twist-two operators are non-standard, in the sense that they involve terms like $(\Delta \cdot p)^n$ with n being an arbitrary non-negative integer. If n is fixed to a specific integer, as adopted in [30, 46], the Feynman rules become standard and the relevant Feynman integrals are the two-point integrals.

Here, we want to keep n arbitrary, in order to directly obtain the all- n splitting functions. However, it is not immediately obvious how to perform the integration-by-parts (IBP) reductions [47] with terms like $(\Delta \cdot p)^n$ present. In the following, we adopt a method first proposed in [15, 31]. The method sums terms proportional to $(\Delta \cdot p)^n$ to linear

propagators with the help of a tracing parameter x , for example,

$$\begin{aligned}
 (\Delta \cdot p)^n &\rightarrow \sum_{n=0}^{\infty} x^n (\Delta \cdot p)^n = \frac{1}{1-x\Delta \cdot p}, \\
 \sum_{j=0}^{n-3} (\Delta \cdot p_1)^{n-3-j} (\Delta \cdot p_2)^j &\rightarrow \sum_{n=3}^{\infty} x^n \sum_{j=0}^{n-3} (\Delta \cdot p_1)^{n-3-j} (\Delta \cdot p_2)^j = \frac{x^3}{(1-x\Delta \cdot p_1)(1-x\Delta \cdot p_2)}.
 \end{aligned}
 \tag{3.1}$$

The above method translates the non-standard terms depending on n into standard linear propagators depending on x , which can be easily handled by standard IBP algorithms. To revert back to n space, we symbolically extract the coefficient of x^n from the results in the x -parameter space. It should be noted that the limit $x \rightarrow 0$ trivializes the linear propagators and that the corresponding Feynman integrals converge to massless two-point functions in this limit. The auxiliary parameter x should not be confused with the Bjorken- x variable, which relates to n by an inverse Mellin transformation.

We work in parameter- x space throughout, starting at the level of the Feynman rules, which are all transformed from n space to parameter- x space. Mathematica is used to substitute the standard QCD Feynman rules and the effective Feynman rules in parameter x -space into the Feynman diagrams. The Dirac and color algebra is performed with FORM [48]. Subsequently, the Feynman integrals are classified into different integral families with an in-house code invoking Reduze 2 [49] and the latest version of FeynCalc [50, 51]. During the topology classification, partial fractions with respect to the Feynman propagators are also needed, and they are performed using Apart [52].

For the singlet operator insertions, we find 1, 2, and 7 integral families at one, two, and three loop order, respectively. Here, an integral family is defined as a complete set of quadratic and linear denominator polynomials, such that any scalar product of a loop momentum can be expressed in terms of them. In general, one integral family may describe more than one “top-level topology”. As an example, we give the 2 integral families at two-loop order,

$$\left(\begin{array}{l} 1 - x\Delta \cdot l_1, \quad 1 - x\Delta \cdot l_2, l_1^2, (l_1 - p)^2, l_2^2, (l_2 - p)^2, (l_1 - l_2)^2 \\ 1 - x\Delta \cdot (l_2 - l_1), 1 - x\Delta \cdot l_2, l_1^2, (l_1 - p)^2, l_2^2, (l_2 - p)^2, (l_1 - l_2)^2 \end{array} \right), \tag{3.2}$$

where l_1, l_2 are the loop momenta. Each integral family contains 7 propagators with the first two being the linear propagators obtained through (3.1). We perform the IBP reduction using the combination of LiteRed [53] and FIRE6 [54], Reduze 2 [49] and Kira [55], which are based on implementations of the Laporta algorithm [56].

To directly compute the master integrals in parameter- x space, we derive differential equations (DEs) [57] with respect to x for the master integrals. The DE system is turned into canonical form [58] using a combination of CANONICA [59, 60] and Libra [61, 62]. Due to (3.1), the master integrals are regular in the limit $x \rightarrow 0$ and reduce to two-point integrals. The two-point integrals are known to four-loop order [63, 64] and serve as the boundary conditions for the DEs. In this way, we manage to express all master integrals up to three

loops in terms of harmonic polylogarithms (HPLs) [65], which are defined by

$$\begin{aligned}
 H(a_1, a_2, \dots, a_m; x) &= \int_0^x dt f_{a_1}(t) H(a_2, \dots, a_m; t), \\
 H(\vec{0}_m; x) &= \frac{\ln^n x}{n!}, \\
 H(x) &= 1,
 \end{aligned}
 \tag{3.3}$$

where a_i is 0, 1, or -1 , and the kernel $f_a(t)$ is defined as

$$f_1(t) = \frac{1}{1-t}, \quad f_0(t) = \frac{1}{t}, \quad f_{-1}(t) = \frac{1}{1+t}.
 \tag{3.4}$$

Inserting IBP relations and the solutions for the master integrals into the integrand, we obtain the final expressions for the bare OMEs.

As the last step before renormalization, we turn the results from the parameter- x space into n space. The HPLs can be expanded around $x = 0$,

$$H(a_1, a_2, \dots; x) = \sum_{n=0}^{\infty} b_n x^n,
 \tag{3.5}$$

with b_n being composed of the harmonic sums [66, 67] of argument n , which are defined recursively by

$$\begin{aligned}
 S_{\pm m_1, m_2, \dots, m_d}(n) &= \sum_{j=1}^n (\pm 1)^j j^{-m_1} S_{m_2, \dots, m_d}(j) \quad (m_i \in \mathbb{N}), \\
 S_{\emptyset}(n) &= 1.
 \end{aligned}
 \tag{3.6}$$

We use the Mathematica package `HarmonicSums` [68–73] to perform the expansion (3.5) of the HPLs. In order to extract the relevant coefficients of x^n , one needs to take into account also the rational prefactors multiplying the HPLs. In our non-singlet computation, we find powers of x , $1/x$, and $1/(1-x)$ after partial fraction decomposition. For the singlet case, we also encounter the factor $1/(1+x)$, which can, however, always be mapped to $1/(1-x)$ due to the fact that only even Mellin moments contribute. At this stage, an OME $G(x)$ in parameter- x space can be written as

$$G(x) = \sum_{m=0}^{m_{\max}} F_m(x),
 \tag{3.7}$$

with $F_m(x)$ being defined as

$$F_m(x) = \frac{B_m(x)}{(1-x)^m},
 \tag{3.8}$$

where $B_m(x)$ is a linear combination of HPLs with coefficients involving powers of x only, and m is a non-negative integer. Prior to simplifications, we find $m_{\max} = 3$ for individual contributions. We expand both $F_m(x)$ and $B_m(x)$ of (3.8),

$$F_m(x) = \sum f_n x^n = \frac{1}{(1-x)^m} \left[\sum_{j_1=0}^{\alpha-1} h_{j_1} x^{j_1} + \sum_{j_1=\alpha}^{\infty} c_{j_1} x^{j_1} \right],
 \tag{3.9}$$

where f_n is the desired result in Mellin space and α is a small integer that is used to separate $B(x)$ into two parts. The coefficients h_{j_1} in the first part are stated explicitly for each value of j_1 (since for them typically no closed form for symbolic j_1 can be obtained), while c_{j_1} can be written in terms of rational functions or harmonic sums depending on the symbol j_1 . For example, if $B_0(x)$ equals $(1+x)\ln(1-x)$, we can express it as follows,

$$(1+x)\ln(1-x) = \sum_{j_1=0}^{\alpha-1} h_{j_1} x^{j_1} + \sum_{j_1=\alpha}^{\infty} c_{j_1} x^{j_1}, \quad (3.10)$$

with

$$\alpha = 2, \quad h_0 = 0, \quad h_1 = -1, \quad c_{j_1} = \frac{-1}{j_1} + \frac{-1}{j_1 - 1},$$

where c_0 and c_1 are not well defined. Taking into account the expansion of the $1/(1-x)^m$ factor, the resulting f_n for generic n in (3.9) can be written down recursively,

$$\begin{aligned} f_n|_{m=1} &= f_\alpha + \sum_{j_1=\alpha+1}^n c_{j_1}, \\ f_n|_{m=2} &= f_\alpha + \sum_{j_1=\alpha+1}^n \left[-f_\alpha + f_{\alpha+1} + \sum_{j_2=\alpha+2}^{j_1} c_{j_2} \right], \\ f_n|_{m=3} &= \left(f_n|_{m=2} \right) \Big|_{c_{j_2} \rightarrow f_\alpha - 2f_{\alpha+1} + f_{\alpha+2} + \sum_{j_3=\alpha+3}^{j_2} c_{j_3}}, \\ &\vdots \\ f_n|_m &= \left(f_n|_{m-1} \right) \Big|_{c_{j_{m-1}} \rightarrow \sum_{k=0}^{m-1} \binom{k}{m-1} (-1)^k f_{\alpha-k+m-1} + \sum_{j_m=\alpha+m}^{j_{m-1}} c_{j_m}}. \end{aligned} \quad (3.11)$$

The multiple sums appearing on the right side of the above formula can be again transformed to harmonic sums by `HarmonicSums`. We note that, after simplification, B_m in (3.8) is non-zero only for $m = 0, 1$ for all partonic channels at three loops. Following the steps outlined above, we are able to get all bare three-loop OMEs of the singlet quark and gluon operators in terms of harmonic sums.

4 GV counterterm Feynman rules from renormalization conditions

We are now ready to derive the counterterm Feynman rules resulting from the GV operators. Several ingredients are needed: the renormalization equations in (2.25), (2.26), the OMEs in (2.30) generalized to any number of external states, and the renormalization conditions.

First, we consider equations (2.25), (2.26). For simplicity, we restrict our discussion to (2.26), which can be used to determine the Feynman rules for operators O_{ABC} as well as for the counterterm operator $[ZO]_g^{\text{GV}}$. Feynman rules for the other counterterm operator $[ZO]_q^{\text{GV}}$ can be extracted from (2.25) in a similar way. The main point is to consider the off-shell one-particle-irreducible (1PI) OMEs for both sides of equation (2.26) and impose the renormalization conditions.

To determine the vertex Feynman rules for the GV operators, it is sufficient to consider the following off-shell 1PI OMEs with external states consisting of two particles of type j and m gluons,

$$\begin{aligned} \langle j|O_g|j+m g\rangle_{1\text{PI}}^{\mu_1\cdots\mu_m, \text{R}} &= Z_j(\sqrt{Z_g})^m \left[\langle j|(Z_{gq}O_q + Z_{gg}O_g)|j+m g\rangle_{1\text{PI}}^{\mu_1\cdots\mu_m, \text{B}} \right] \\ &+ Z_j(\sqrt{Z_g})^m \left[Z_{gA} \langle j|O_{ABC}|j+m g\rangle_{1\text{PI}}^{\mu_1\cdots\mu_m, \text{B}} \right. \\ &\quad \left. + \langle j|[ZO]_g^{\text{GV}}|j+m g\rangle_{1\text{PI}}^{\mu_1\cdots\mu_m, \text{B}} \right]. \end{aligned} \quad (4.1)$$

Here, j could denote quarks(q), gluons(g), or ghosts(c), $\sqrt{Z_j}$ is the corresponding field renormalization constant, and the renormalization of the strong coupling constant is implicitly understood on the right-hand side. To make the extraction of counterterm Feynman rules from the above equation transparent, we expand the OMEs according to the number of loops and legs,

$$\langle j|O|j+m g\rangle^{\mu_1\cdots\mu_m} = \sum_{l=1}^{\infty} \left[\langle j|O|j+m g\rangle^{\mu_1\cdots\mu_m, (l), (m)} \right] a_s^l g_s^m, \quad (4.2)$$

where O denotes a generic operator. For fixed m , we can compute the off-shell OMEs in the first line of the right-hand side of (4.1) order-by-order in a loop expansion. Since the left-hand side of (4.1) is an ultraviolet renormalized and infrared finite quantity, the sum of terms on the right-hand side should also be finite. If the GV operators were not to contribute, the divergences should cancel within the OMEs of the physical operators. Otherwise, the remaining divergences should be absorbed into the contribution from GV operators. Requiring the finiteness of the right-hand side of (4.1), we can extract the counterterm Feynman rules from the GV operators (or even the Feynman rules for the GV operators themselves) order-by-order in a loop expansion.

4.1 General formulae to determine the Feynman rules for the O_{ABC} operators

Due to the hierarchy of GV operators, the counterterm operators $[ZO]_j^{\text{GV}}$ contribute to the renormalization of the OMEs only from two loops onwards. Consequently, the Feynman rules for the O_{ABC} operators can be extracted from the evaluation of one-loop off-shell 1PI OMEs.

To extract the Feynman rules for O_C , we consider the one-loop OMEs in (4.1) with two ghosts, $j = c$, and m gluons in the external states. In this case, by employing the renormalization conditions, equation (4.1) simplifies to the form

$$Z_{gA}^{(1)} \langle c|O_C|c+m g\rangle_{1\text{PI}}^{\mu_1\cdots\mu_m, (0), (m)} = - \left[\langle c|O_g|c+m g\rangle_{1\text{PI}}^{\mu_1\cdots\mu_m, (1), (m), \text{B}} \right]_{\text{div}}, \quad (4.3)$$

where the subscript ‘div’ denotes the divergent contribution, that is, the poles in ϵ . We notice that the left-hand side of the above equation is the multiplication of a renormalization constant and the vertex Feynman rules for the O_C operator. By evaluating the right-hand

side for $m = 0$ and factorizing the dependence on kinematics, $Z_{gA}^{(1)}$ can be determined up to an overall m -independent constant

$$Z_{gA}^{(1)} = -\frac{1}{\epsilon} \frac{C_A}{n(n-1)}. \quad (4.4)$$

For $m = 1$, a similar method was used to fix the unknown coefficients in an ansatz constrained by a generalized BRST symmetry in [14].

To determine the Feynman rule for the O_B operator, we consider the OMEs with two quarks and m gluon external states. Setting j to a quark state and expanding to one-loop order, (4.1) becomes

$$Z_{gA}^{(1)} \langle q|O_B|q + m g\rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (0), (m)} = -\left\{ Z_{gq}^{(1)} \left[\langle q|O_q|q + m g\rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (0), (m)} \right] + \left[\langle q|O_g|q + m g\rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} \right]_{\text{div}} \right\}, \quad (4.5)$$

with $Z_{gq}^{(1)}$ being the first order renormalization constant for the $q \rightarrow g$ transition

$$Z_{gq}^{(1)} = \frac{1}{\epsilon} \left[\frac{4}{n} - \frac{2}{n+1} - \frac{4}{n-1} \right] C_F. \quad (4.6)$$

The results of the OMEs $\left[\langle q|O_g|q + m g\rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} \right]_{\text{div}}$ contain both C_F and C_A color factors. Since $Z_{gA}^{(1)}$ is proportional to C_A , the C_F from the one-loop OMEs must cancel against the one from $Z_{gq}^{(1)}$ in the above equation.

Finally, to determine the Feynman rule for O_A operator, we consider the OMEs with $m + 2$ gluon states in a similar way. Setting j to a gluon state and expanding to one-loop order, (4.1) becomes

$$Z_{gA}^{(1)} \langle g|O_A|g + m g\rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (0), (m)} = -\left\{ \left[\langle g|O_g|g + m g\rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (1), (m), \text{B}} \right]_{\text{div}} + \left[Z_{gg}^{(1)} - \frac{m}{2\epsilon} \beta_0 + \frac{m+2}{2} Z_g^{(1)} \right] \langle g|O_g|g + m g\rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (0), (m)} \right\}, \quad (4.7)$$

where β_0 is the one-loop QCD beta function

$$\beta_0 = \frac{11C_A}{3} - \frac{2N_f}{3}, \quad (4.8)$$

$Z_g^{(1)}$ is the first-order gluon field renormalization constant

$$Z_g^{(1)} = \frac{1}{\epsilon} \left[\frac{13C_A}{6} - \frac{C_A \xi}{2} - \frac{2N_f}{3} \right], \quad (4.9)$$

and $Z_{gg}^{(1)}$ is the first order renormalization constant for the $g \rightarrow g$ transition.

$$Z_{gg}^{(1)} = \frac{1}{\epsilon} \left[C_A \left(4S_1(n) - \frac{4}{n-1} + \frac{4}{n} - \frac{4}{n+1} + \frac{4}{n+2} - \frac{11}{3} \right) + \frac{2N_f}{3} \right]. \quad (4.10)$$

For a one-loop, purely gluonic scattering process involving the operator O_g , one expects that no N_f factor appears. Indeed, through a simple calculation, we found that the N_f factor cancels out in the combination $Z_{gg}^{(1)} - \frac{m}{2\epsilon}\beta_0 + \frac{m+2}{2}Z_g^{(1)}$ of equation (4.7).

Through equations (4.3), (4.5) and (4.7), deriving the Feynman rules for the unknown operators O_{ABC} is equivalent to the computation of the divergent contributions to one-loop off-shell 1PI OMEs with an operator insertion of O_g . Since the Feynman rules resulting from O_g are known to all multiplicities, as shown in (2.9), the Feynman rules for O_{ABC} could in principle also be calculated to all multiplicities.

4.2 Determining the counterterm Feynman rules for $[ZO]_g^{\text{GV},(2)}$ and beyond

In the last subsection, the Feynman rules for the O_{ABC} operators were extracted from one-loop OMEs in (4.1). Similarly, the l -loop corrections of (4.1) can be used to extract the counterterm Feynman rules for $[ZO]_g^{\text{GV},(l)}$ with $l \geq 2$. Firstly, we work out the explicit formulae for the Feynman rules of $[ZO]_g^{\text{GV},(2)}$. Considering two ghost plus m gluon external states, to two-loop order, (4.1) reads

$$\begin{aligned}
 & \langle c | [ZO]_g^{\text{GV},(2)} | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (0), (m)} = \\
 & - \left\{ \left[\langle c | O_g | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (2), (m), \text{B}} \right. \right. \\
 & \quad + \left(Z_c^{(1)} + \frac{m Z_g^{(1)}}{2} + Z_{gg}^{(1)} - \frac{\beta_0(m+2)}{2\epsilon} \right) \langle c | O_g | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} \\
 & \quad + \left(Z_c^{(1)} Z_{gA}^{(1)} + \frac{1}{2} m Z_g^{(1)} Z_{gA}^{(1)} - \frac{\beta_0 m Z_{gA}^{(1)}}{2\epsilon} + Z_{gA}^{(2)} \right) \langle c | O_C | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (0), (m), \text{B}} \\
 & \quad + Z_{gA}^{(1)} \langle c | O_{AC} | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} \\
 & \quad \left. \left. + Z_g^{(1)} \sum_{t=1}^s \xi^t t \langle c | O_g | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), (t), \text{B}} \right] \right\}_{\text{div}} \Big|_{\xi^{\text{B}} \rightarrow \xi}, \quad (4.11)
 \end{aligned}$$

where the last term originates from the gauge parameter renormalization

$$\xi^{\text{B}} = Z_\xi \xi, \quad (4.12)$$

with $Z_\xi = Z_g$ in covariant gauge. We further decompose the OMEs according to the power of the gauge parameter,

$$\langle c | O_g | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} = \sum_{t=0}^s (\xi^{\text{B}})^t \langle c | O_g | c + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), (t), \text{B}}, \quad (4.13)$$

where s is a small positive integer.

For the two quark plus m gluon external states, to two-loop order, (4.1) becomes

$$\begin{aligned}
 & \langle q | [ZO]_g^{\text{GV}, (2)} | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (0), (m)} = \\
 & - \left\{ \left[\langle q | O_g | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (2), (m), \text{B}} \right. \right. \\
 & \quad + \left(\frac{m Z_g^{(1)}}{2} + Z_{gg}^{(1)} + Z_q^{(1)} - \frac{\beta_0(m+2)}{2\epsilon} \right) \langle q | O_g | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} \\
 & \quad + \left(\frac{1}{2} m Z_g^{(1)} Z_{gA}^{(1)} - \frac{\beta_0 m Z_{gA}^{(1)}}{2\epsilon} + Z_{gA}^{(1)} Z_q^{(1)} + Z_{gA}^{(2)} \right) \langle q | O_B | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (0), (m), \text{B}} \\
 & \quad + \left(\frac{1}{2} m Z_g^{(1)} Z_{gq}^{(1)} - \frac{\beta_0 m Z_{gq}^{(1)}}{2\epsilon} + Z_{gq}^{(1)} Z_q^{(1)} + Z_{gq}^{(2)} \right) \langle q | O_q | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (0), (m), \text{B}} \\
 & \quad + Z_{gA}^{(1)} \langle q | O_{AB} | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} + Z_{gq}^{(1)} \langle q | O_q | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), \text{B}} \\
 & \quad \left. \left. + Z_g^{(1)} \sum_{t=1}^s \xi^t t \langle q | O_g | q + m g \rangle_{\text{1PI}}^{\mu_1 \cdots \mu_m, (1), (m), (t), \text{B}} \right]_{\text{div}} \right\} \Big|_{\xi^{\text{B}} \rightarrow \xi}. \tag{4.14}
 \end{aligned}$$

Similarly, for $m + 2$ gluon external states, to two-loop order, (4.1) reads

$$\begin{aligned}
 & \langle g | [ZO]_g^{\text{GV}, (2)} | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (0), (m)} = \\
 & - \left\{ \left[\langle g | O_g | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (2), (m), \text{B}} \right. \right. \\
 & \quad + \left(\frac{1}{2} (m+2) Z_g^{(1)} + Z_{gg}^{(1)} - \frac{\beta_0(m+2)}{2\epsilon} \right) \langle g | O_g | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (1), (m), \text{B}} \\
 & \quad + \left(\frac{1}{2} (m+2) Z_g^{(1)} Z_{gA}^{(1)} - \frac{\beta_0 m Z_{gA}^{(1)}}{2\epsilon} + Z_{gA}^{(2)} \right) \langle g | O_A | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (0), (m), \text{B}} \\
 & \quad + Z_{gA}^{(1)} \langle g | O_{ABC} | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (1), (m), \text{B}} + Z_{gq}^{(1)} \langle g | O_q | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (1), (m), \text{B}} \\
 & \quad + \left(\frac{-\frac{1}{4} \beta_0 m (m+2) Z_g^{(1)} - \frac{1}{2} \beta_0 m Z_{gg}^{(1)} - \frac{\beta_1 m}{4}}{\epsilon} + \frac{1}{2} (m+2) (Z_g^{(1)} Z_{gg}^{(1)} + Z_g^{(2)}) + Z_{gg}^{(2)} \right) \\
 & \quad + \frac{1}{8} m (m+2) (Z_g^{(1)})^2 + \frac{\beta_0^2 m (m+2)}{8\epsilon^2} \left. \langle g | O_g | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (0), (m), \text{B}} \right. \\
 & \quad \left. \left. + Z_g^{(1)} \sum_{t=1}^s \xi^t t \langle g | O_g | g + m g \rangle_{\text{1PI}}^{\mu\nu\mu_1 \cdots \mu_m, (1), (m), (t), \text{B}} \right]_{\text{div}} \right\} \Big|_{\xi^{\text{B}} \rightarrow \xi}. \tag{4.15}
 \end{aligned}$$

Here, we introduced some short-hand notations, $O_{AB} = O_A + O_B$, $O_{AC} = O_A + O_C$. The above equations express the Feynman rules for the counterterm operator $[ZO]_g^{\text{GV}, (2)}$ through the divergent terms of two-loop off-shell OMEs with O_g insertion plus contributions from lower-loop OMEs with other operator insertions. We noticed that the one-loop OMEs need to be evaluated to order ϵ^0 .

In addition to the renormalization constants appearing in (4.3), (4.5), (4.7), we need several further renormalization constants in the above equations, including the first order quark and ghost field renormalization constants $Z_q^{(1)}$, $Z_c^{(1)}$, the second order gluon field renormalization constant $Z_g^{(2)}$, the 2-loop QCD beta function β_1 as well as the second order renormalization constants $Z_{gA}^{(2)}$, $Z_{gq}^{(2)}$, $Z_{gg}^{(2)}$. The field renormalization constants and

beta function are known from the computations of diagrams without a twist-two operator insertion, they are summarized in appendix D. However, $Z_{gA}^{(2)}$, $Z_{gq}^{(2)}$, $Z_{gg}^{(2)}$ are supposed to be extracted from two-point two-loop OMEs in the above equations for $m = 0$, and thus should be regarded unknown. Therefore, it seems that the equations (4.11), (4.14) and (4.15) can not be used to determine the Feynman rules for $[ZO]_g^{\text{GV},(2)}$.

However, as shown in appendix B, the two-point vertex Feynman rules ($m = 0$) for all GV operators except O_A and O_C are zero. Consequently,

$$\begin{aligned} \langle c | [ZO]_g^{\text{GV},(2)} | c \rangle_{\text{1PI}}^{(0),(0)} &= 0, \\ \langle q | [ZO]_g^{\text{GV},(2)} | q \rangle_{\text{1PI}}^{(0),(0)} &= 0, \\ \langle g | [ZO]_g^{\text{GV},(2)} | g \rangle_{\text{1PI}}^{\mu\nu,(0),(0)} &= 0. \end{aligned} \tag{4.16}$$

These equations allow us to easily determine $Z_{gA}^{(2)}$, $Z_{gq}^{(2)}$, $Z_{gg}^{(2)}$ by separately evaluating (4.11), (4.14) and (4.15) for $m = 0$. With all renormalization constants determined, the counterterm Feynman rules for $[ZO]_g^{\text{GV},(2)}$ can, in principle, be determined to an arbitrary number of legs from (4.11), (4.14) and (4.15), similarly to the case of O_{ABC} . Since the vertex Feynman rules with two legs for $[ZO]_q^{\text{GV},(3)}$ and $[ZO]_g^{\text{GV},(3)}$ are zero, the counterterm operators $[ZO]_q^{\text{GV},(3)}$ and $[ZO]_g^{\text{GV},(3)}$ contribute to the splitting functions only from four loops onwards. We leave them for future study.

5 Computation of Feynman rules for the GV operators

We demonstrated in the above section that, in order to derive the Feynman rules of GV operators, one needs to work with general kinematics and keep all relevant Lorentz structures when computing the multi-loop, multi-particle OMEs. This task is in general non-trivial. For a k particle scattering process with O_g insertion, the external kinematics introduces a reference vector Δ and $k - 1$ independent momenta p_1, \dots, p_{k-1} of the scattering particles. Therefore, the total number of mass scales is

$$\frac{1}{2}(k-1)(k+2). \tag{5.1}$$

For example, for a three-particle scattering, one has the 5 scales

$$p_1 \cdot p_2, \quad p_1^2, \quad p_2^2, \quad \Delta \cdot p_1, \quad \Delta \cdot p_2. \tag{5.2}$$

For four-particle and five-particle scatterings, we need 9 and 14 scales respectively. The presence of many mass scales renders the computations complicated, and the number of independent Lorentz structures increases quickly especially for pure gluon scatterings. Before considering the symmetries, the 5 Lorentz structures for 2-gluon scattering can be listed as

$$\Delta^\mu \Delta^\nu, \quad \Delta^\mu p^\nu, \quad p^\mu \Delta^\mu, \quad g^{\mu\nu}, \quad p^\mu p^\nu. \tag{5.3}$$

For a 3-gluon scattering, we divide the Lorentz structures into different categories according to the number of Δ ,

$$\begin{aligned}
 3 \Delta &: \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} \\
 2 \Delta &: \Delta^{\mu_1} \Delta^{\mu_2} p_1^{\mu_3}, \Delta^{\mu_1} \Delta^{\mu_2} p_2^{\mu_3}, \dots \\
 1 \Delta &: \Delta^{\mu_1} g^{\mu_2 \mu_3}, \Delta^{\mu_1} p_1^{\mu_2} p_1^{\mu_3}, \Delta^{\mu_1} p_1^{\mu_2} p_2^{\mu_3}, \Delta^{\mu_1} p_2^{\mu_2} p_2^{\mu_3}, \dots \\
 0 \Delta &: g^{\mu_1 \mu_2} p_1^{\mu_3}, g^{\mu_1 \mu_2} p_2^{\mu_3}, p_1^{\mu_1} p_1^{\mu_2} p_1^{\mu_3}, p_1^{\mu_1} p_1^{\mu_2} p_2^{\mu_3}, p_1^{\mu_1} p_2^{\mu_2} p_2^{\mu_3}, p_2^{\mu_1} p_2^{\mu_2} p_2^{\mu_3}, \dots
 \end{aligned} \tag{5.4}$$

where we have eliminated p_3 using momentum conservation $p_3 = -p_1 - p_2$, and the dots represent Lorentz structures that can be obtained from the permutations of the Lorentz indices μ_1, μ_2, μ_3 . In total, we have 36 Lorentz structures for 3-gluon scattering, 353 Lorentz structures for 4-gluon scattering, and 4400 for 5-gluon scattering. A naive projection method will make the computations extremely complicated. However, we observe that many Lorentz structures do not appear in the Feynman rules of the physical operators O_q and O_g , as shown in appendix A. Since the GV operators are also twist-two operators, the observation may be used to simplify the computations. To make things explicit, we work out all allowed Lorentz structures for a general twist-two operator in the following.

5.1 Lorentz structures for the Feynman rules of a general twist-two operator

For the sake of simplicity and without loss of generality, we focus on the Lorentz structures of Feynman rules for GV operators O_{ABC} . The same conclusion also applies to a general twist-two operator. Our guiding principles are the properties of twist-two operators and the fact that no inverse mass scale can be generated in the Feynman rule of a vertex.

We first analyze the Lorentz structures of m -gluon Feynman rules for O_A . For the twist-two O_A operator, the mass dimension is $n + 2$ and the spin is n . By factoring out m gluon fields, the mass dimension becomes $n + 2 - m$. If we denote the number of Δ^μ , p_i^μ and $g^{\mu\nu}$ appearing in a Lorentz structure by n_1, n_2, n_3 respectively, then they should sum up to the total number of Lorentz indices m ,

$$m = n_1 + n_2 + 2n_3. \tag{5.5}$$

Next, we count the mass dimension for the coefficient of a Lorentz structure. The coefficient is composed of Lorentz scalar products formed by the reference vector Δ and the momentum p_i of scattering particles. The dependence of the coefficients on the scalar products is of polynomial form, in particular, propagator-type terms (for example $1/(p_1 \cdot p_2)$) can not appear in the Feynman rule of a vertex. Each monomial is further divided into two factors, the first one involves the contractions of Δ and the momenta p_i (for example $\Delta \cdot p_1$), and the second one includes the contractions within the momenta p_i (for example $p_1 \cdot p_2$). The mass dimension from the first factor must be $n - n_1$ since the operator should always include a total number n of Δ vectors. We set the mass dimension of the second factor to a non-negative integer $2n_4$.

By adding up the mass dimensions from the different ingredients, we should recover the total mass dimensions,

$$(n - n_1) + n_2 + 2n_4 = n + 2 - m. \tag{5.6}$$

Substituting (5.5) into (5.6) gives us

$$n_2 + n_3 + n_4 = 1. \quad (5.7)$$

Since all n_2, n_3, n_4 should be non-negative, the above equation tells us that each of them should not be larger than 1 and that their sum must be 1. The equation drastically constrains the allowed Lorentz structures. For example, the structure $\Delta^{\mu_1} p_1^{\mu_2} p_1^{\mu_3}$ in (5.4) can not appear since $n_2 = 2 > 1$. As another example, $g^{\mu_1 \mu_2} p_1^{\mu_3}$ can not appear due to $n_2 = 1, n_3 = 1$.

With the constraint in (5.7), it is easy to write down the general ansatz for the 3-gluon Feynman rule of O_A ,

$$\begin{aligned} & \left[\langle g | O_A | gg \rangle_{\text{1PI}}^{\mu_1 \mu_2 \mu_3, (0), (1)} \right] = \\ & a_1 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} + a_2 \Delta^{\mu_1} \Delta^{\mu_2} p_1^{\mu_3} + a_3 \Delta^{\mu_1} \Delta^{\mu_3} p_1^{\mu_2} \\ & + a_4 \Delta^{\mu_2} \Delta^{\mu_3} p_1^{\mu_1} + a_5 \Delta^{\mu_1} \Delta^{\mu_2} p_2^{\mu_3} + a_6 \Delta^{\mu_1} \Delta^{\mu_3} p_2^{\mu_2} + a_7 \Delta^{\mu_2} \Delta^{\mu_3} p_2^{\mu_1} \\ & + a_8 \Delta^{\mu_1} g^{\mu_2 \mu_3} + a_9 \Delta^{\mu_2} g^{\mu_1 \mu_3} + a_{10} \Delta^{\mu_3} g^{\mu_1 \mu_2}. \end{aligned} \quad (5.8)$$

The number of Lorentz structures is reduced from 36 to 10 for the 3-gluon Feynman rule in the above equation. Similarly, the general ansatz for the 4-gluon Feynman rule of O_A is obtained as

$$\begin{aligned} & \left[\langle g | O_A | ggg \rangle_{\text{1PI}}^{\mu_1 \mu_2 \mu_3 \mu_4, (0), (2)} \right] = \\ & b_1 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} + b_2 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} p_1^{\mu_4} \\ & + b_3 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_4} p_1^{\mu_3} + b_4 \Delta^{\mu_1} \Delta^{\mu_3} \Delta^{\mu_4} p_1^{\mu_2} + b_5 \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} p_1^{\mu_1} \\ & + b_6 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} p_2^{\mu_4} + b_7 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_4} p_2^{\mu_3} + b_8 \Delta^{\mu_1} \Delta^{\mu_3} \Delta^{\mu_4} p_2^{\mu_2} + b_9 \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} p_2^{\mu_1} \\ & + b_{10} \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} p_3^{\mu_4} + b_{11} \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_4} p_3^{\mu_3} + b_{12} \Delta^{\mu_1} \Delta^{\mu_3} \Delta^{\mu_4} p_3^{\mu_2} + b_{13} \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} p_3^{\mu_1} \\ & + b_{14} \Delta^{\mu_1} \Delta^{\mu_2} g^{\mu_3 \mu_4} + b_{15} \Delta^{\mu_1} \Delta^{\mu_3} g^{\mu_2 \mu_4} + b_{16} \Delta^{\mu_1} \Delta^{\mu_4} g^{\mu_2 \mu_3} + b_{17} \Delta^{\mu_2} \Delta^{\mu_3} g^{\mu_1 \mu_4} \\ & + b_{18} \Delta^{\mu_2} \Delta^{\mu_4} g^{\mu_1 \mu_3} + b_{19} \Delta^{\mu_3} \Delta^{\mu_4} g^{\mu_1 \mu_2}, \end{aligned} \quad (5.9)$$

where we need only 19 out of the initial list of 353 Lorentz structures.

We also analyze the dependence of the scalar coefficients of Lorentz tensors in (5.8) and (5.9) on the kinematic invariants. For term $a_1 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3}$, $n_2 = n_3 = 0$, therefore, n_4 must be 1. It indicates that the coefficient a_1 must be linear in p_1^2, p_2^2 , or $p_1 \cdot p_2$. Except for $\Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3}$ and $\Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4}$, other tensor structures have either $n_2 = 1$ or $n_3 = 1$. Therefore, their corresponding coefficients a_i, b_i with $i > 1$ can be only constructed from scalar products involving Δ . Exploiting the above properties allows for a particularly efficient determination of the coefficients with modern finite-field and function reconstruction techniques [74, 75].

Since Lorentz indices can not be carried by the quark fields and ghost fields, the Lorentz structures appearing in the Feynman rules for the operators O_B and O_C are much more constrained. Regarding a vertex involving two quarks plus m gluons, or two ghosts plus m gluons, (5.5) is still valid by using the same convention as for the pure m -gluon vertex.

The difference is the extra mass dimension from quark fields or ghost fields. Regarding the operator O_C , (5.6) should be modified to

$$(n - n_1) + n_2 + 2n_4 = n + 2 - [\bar{c}c] - m = n + 2 - 2 - m, \quad (5.10)$$

where $[\bar{c}c] = 2$ stands for the mass dimension of a pair of ghost and anti-ghost fields. For the operator O_B , one always needs a factor \not{A} to compensate the extra mass dimension from $\bar{\psi}\psi$ compared with $\bar{c}c$. Correspondingly, (5.6) is modified to

$$(n - n_1 - 1) + n_2 + 2n_4 = n + 2 - [\bar{\psi}\psi] - m = n + 2 - 3 - m, \quad (5.11)$$

where $[\bar{\psi}\psi] = 3$ stands for the mass dimension of a pair of quark and anti-quark fields. The above equation is equivalent to the equation (5.10). Solving (5.5) and (5.10), the result reads,

$$n_2 + n_3 + n_4 = 0. \quad (5.12)$$

The above equation indicates, for both Feynman rules of O_B and O_C , the Lorentz tensors can be composed of Δ^μ only,

$$\begin{aligned} \left[\langle c | O_C | c + m g \rangle_{\text{1PI}}^{\mu_1 \dots \mu_m, (0), (m)} \right] &= c_m \Delta^{\mu_1} \Delta^{\mu_2} \dots \Delta^{\mu_m}, \\ \left[\langle q | O_B | q + m g \rangle_{\text{1PI}}^{\mu_1 \dots \mu_m, (0), (m)} \right] &= d_m \Delta^{\mu_1} \Delta^{\mu_2} \dots \Delta^{\mu_m}, \end{aligned} \quad (5.13)$$

where the coefficients c_m and d_m have the same property as the coefficients a_i, b_i with $i > 1$ in (5.8) and (5.9).

Since the counterterm operators $[ZO]_q^{\text{GV}}$ and $[ZO]_g^{\text{GV}}$ are also twist-two operators, the above considerations remain valid if we replace O_A, O_B, O_C in (5.8), (5.9) and (5.13) by $[ZO]_q^{\text{GV}}$ or $[ZO]_g^{\text{GV}}$. To compute the multi-loop, multi-particle OMEs, we project out the coefficients of the corresponding Lorentz structures in (5.8), (5.9) and (5.13). In general, the Lorentz structures that are allowed in the Feynman rules do not form a complete basis of the vector space, such that we need extra Lorentz structures in the construction of the projectors. However, the number of independent projectors is the same as the number of required Lorentz structures in corresponding equations. For example, a single projector $p_{1, \mu_1} p_{1, \mu_2} \dots p_{1, \mu_m}$ is enough to project out c_m and d_m in (5.13).

5.2 Feynman rules for O_{ABC} up to four legs

From (4.3), (4.5) and (4.7), we observe that the Feynman rules for O_{ABC} are either proportional to the divergent terms of the one-loop OMEs, or expressed as a linear combination of one-loop OMEs and the Feynman rules for physical operators O_q and O_g . Therefore, the same ansatz as in (5.8), (5.9) and (5.13) also applies to the divergent terms of the one-loop OMEs. Having worked out the simplified ansatz and projectors, we are ready to compute the one-loop, multi-particle OMEs by adopting the method in section 3. We work directly in parameter- x space. Since we need just the divergent terms, with generic kinematics, only




Figure 2. Sample diagrams to determine Feynman rules with $gggg$ vertex for O_A (left), $q\bar{q}gg$ vertex for O_B (middle) and $c\bar{c}gg$ vertex for O_C (right). All Feynman diagrams contain an insertion of the physical operator O_g .

two kinds of master integrals contribute. One is the bubble integral, the other one is the bubble integral with a linear propagator insertion,

$$\begin{aligned}
 I_1 &= (\mu^2)^\epsilon \int \frac{d^d l}{i\pi^{d/2}} \frac{1}{(l - q_1)^2 l^2}, \\
 I_2 &= (\mu^2)^\epsilon \int \frac{d^d l}{i\pi^{d/2}} \frac{1}{(l - q_1)^2 l^2 (1 - x\Delta \cdot (l + q_2))},
 \end{aligned}
 \tag{5.14}$$

where q_1, q_2 are a linear combination of momenta p_1, \dots, p_{k-1} and μ is the 't Hooft scale. The two master integrals can be computed easily, and the only special function that appears in their divergent parts is the logarithm. In particular, we encounter x -dependent logarithms from the master integral I_2 ,

$$I_2 = \frac{1}{\epsilon} \left[\frac{\ln(1 - x\Delta \cdot q_1 - x\Delta \cdot q_2) - \ln(1 - x\Delta \cdot q_2)}{-x\Delta \cdot q_1} \right] + \mathcal{O}(\epsilon^0).
 \tag{5.15}$$

The spurious poles appearing in the results for the OMEs in parameter- x space can be eliminated using the `MultivariateApart` [76] and `Singular_pfd` [77] packages implementing multivariate partial fraction algorithms. Using the method in section 3 and the following type of replacement,

$$\begin{aligned}
 \frac{x^3}{(1 - x\Delta \cdot p_1)(1 - x\Delta \cdot p_2)} &= \sum_{n=3}^{\infty} x^n \sum_{j_1=0}^{n-3} (\Delta \cdot p_1)^{n-3-j_1} (\Delta \cdot p_2)^{j_1} \\
 &\rightarrow \sum_{j_1=0}^{n-3} (\Delta \cdot p_1)^{n-3-j_1} (\Delta \cdot p_2)^{j_1},
 \end{aligned}
 \tag{5.16}$$

we manage to express the divergent terms of one-loop OMEs in terms of a single harmonic sum $S_1(n)$ and multiple summations of the form shown in the last line of the above equation.

We compute the divergent terms of the one-loop OMEs in (4.3), (4.5) and (4.7) for $m = 0, 1, 2$ with general ξ dependence. The computation of the $m = 3$ case is also straightforward, but becomes relevant only for the 4-loop splitting functions. In all cases, the right-hand sides are found to be proportional to $Z_{gA}^{(1)}$. The observation verifies our statement right above (2.22). Explicitly for $m = 0, 1, 2$, the OMEs involving ghosts have no dependence on $S_1(n)$ and are directly proportional to $Z_{gA}^{(1)}$. The OMEs involving quarks also have no dependence on $S_1(n)$ and their dependence on C_F cancels against the $Z_{gq}^{(1)}$ terms.

The dependence on $S_1(n)$ of OMEs involving only gluons cancels against the $Z_{gg}^{(1)}$ terms. By dividing out the common factor $Z_{gA}^{(1)}$ in the left-hand and right-hand side of (4.3), (4.5) and (4.7), we manage to obtain all- n Feynman rules up to 4 legs for the operators O_A , O_B and O_C . We emphasize that all Feynman rules are found to be ξ independent.

Up to three legs, the Feynman rules for the operators O_A and O_C were already available in [14], we find full agreement with them upon corrections of typographical errors according to [78]. The Feynman rule up to 3 legs for O_B was given in [79], and we find full agreement in that case as well. Our all- n Feynman rules for operators O_{ABC} with four legs are new. For completeness, we present all Feynman rules up to 4 legs in the following with the convention of all momenta flowing into the vertices.

The all- n Feynman rules for the O_B operator with up to 4 legs are given by


$$\begin{array}{c}
 \begin{array}{ccc}
 \xrightarrow{p_1, i_1} & \bullet & \xrightarrow{p_2, i_2} \\
 & & \\
 & \rightarrow 0, &
 \end{array} \\
 \end{array} \tag{5.17}$$

$$\begin{array}{c}
 \begin{array}{ccc}
 \xrightarrow{p_1, i_1} & \bullet & \xrightarrow{p_2, i_2} \\
 & \downarrow \text{red wavy} & \\
 & p_3, \mu_3, a_3 &
 \end{array} \\
 \rightarrow -\frac{1 + (-1)^n}{2} g_s \Delta^{\mu_3} T_{i_2 i_1}^{a_3} \not{\Delta} (\Delta \cdot (p_1 + p_2))^{n-2}, \\
 \end{array} \tag{5.18}$$


$$\begin{array}{c}
 \begin{array}{ccc}
 \xrightarrow{p_1, i_1} & \bullet & \xrightarrow{p_2, i_2} \\
 \downarrow \text{red wavy} & & \downarrow \text{red wavy} \\
 p_3, \mu_3, a_3 & & p_4, \mu_4, a_4
 \end{array} \\
 \rightarrow -\frac{1}{4} \frac{1 + (-1)^n}{2} g_s^2 \Delta^{\mu_3} \Delta^{\mu_4} (T^{a_3} T^{a_4} - T^{a_4} T^{a_3})_{i_2 i_1} \not{\Delta} \sum_{j_1=0}^{n-3} \left(3 (\Delta \cdot (p_1 + p_2))^{-j_1+n-3} \right. \\
 \left. \times \left[(-\Delta \cdot p_3)^{j_1} - (-\Delta \cdot p_4)^{j_1} \right] - (-\Delta \cdot p_4)^{j_1} (\Delta \cdot p_3)^{-j_1+n-3} \right). \\
 \end{array} \tag{5.19}$$

The all- n Feynman rules for the O_C operator with up to 4 legs are given by

$$\begin{array}{c}
 \begin{array}{ccc}
 \dashrightarrow p_1, a_1 & \bullet & \dashrightarrow p_2, a_2 \\
 & & \\
 & \rightarrow \frac{1 + (-1)^n}{2} \delta^{a_1 a_2} (\Delta \cdot p_1)^n, &
 \end{array} \\
 \end{array} \tag{5.20}$$




$$\begin{aligned}
 &\rightarrow \frac{-i}{4} \frac{1+(-1)^n}{2} \Delta^{\mu_3} g_s f^{a_1 a_2 a_3} \left(3\Delta \cdot p_1 \Delta \cdot p_2 \sum_{j_1=0}^{n-3} \left((-\Delta \cdot p_2)^{j_1} (\Delta \cdot p_1)^{-j_1+n-3} \right) \right. \\
 &\quad \left. + (\Delta \cdot p_1 - \Delta \cdot p_2) (\Delta \cdot (p_1 + p_2))^{n-2} - (\Delta \cdot p_1)^{n-1} + (\Delta \cdot p_2)^{n-1} \right), \tag{5.21}
 \end{aligned}$$




$$\begin{aligned}
 &\rightarrow \frac{1}{24} \frac{1+(-1)^n}{2} g_s^2 \Delta^{\mu_3} \Delta^{\mu_4} \left\{ f^{a_1 a_3 a} f^{a_2 a_4 a} \left(6(-\Delta \cdot p_4)^{n-2} + 6(\Delta \cdot p_3)^{n-2} \right. \right. \\
 &\quad \left. \left. + 6(\Delta \cdot (p_1 + p_3))^{n-2} + 6(\Delta \cdot (p_2 + p_3))^{n-2} - \sum_{j_1=0}^{n-2} \left[\right. \right. \right. \\
 &\quad \left. \left. \left. + [(-\Delta \cdot p_3)^{j_1} + (-\Delta \cdot p_4)^{j_1}] [3(\Delta \cdot p_1)^{n-j_1-2} \right. \right. \right. \\
 &\quad \left. \left. \left. + 3(\Delta \cdot p_2)^{n-j_1-2} + (\Delta \cdot (p_1 + p_2))^{n-j_1-2} \right] \right. \right. \\
 &\quad \left. \left. + 9[(\Delta \cdot p_1)^{n-j_1-2} + (-\Delta \cdot p_2)^{n-j_1-2}] [(\Delta \cdot (-p_2 - p_3))^{j_1} + (\Delta \cdot (p_1 + p_3))^{j_1}] \right) \right. \\
 &\quad \left. + 13 \sum_{j_1=0}^{n-2} \sum_{j_2=0}^{j_1} \left[(-\Delta \cdot p_2)^{j_1-j_2} (\Delta \cdot p_1)^{n-j_1-2} [(\Delta \cdot (-p_2 - p_3))^{j_2} + (\Delta \cdot (p_1 + p_3))^{j_2}] \right] \right) \\
 &\quad + f^{a_1 a_2 a} f^{a_3 a_4 a} \left(-6(\Delta \cdot p_3)^{n-2} - 6(\Delta \cdot (p_2 + p_3))^{n-2} \right. \\
 &\quad \left. + \sum_{j_1=0}^{n-2} \left[3(-\Delta \cdot p_4)^{j_1} (\Delta \cdot p_1)^{n-j_1-2} \right. \right. \\
 &\quad \left. \left. + 3(-\Delta \cdot p_3)^{j_1} (\Delta \cdot p_2)^{n-j_1-2} + [5(-\Delta \cdot p_3)^{j_1} - 4(-\Delta \cdot p_4)^{j_1}] (\Delta \cdot (p_1 + p_2))^{n-j_1-2} \right. \right. \\
 &\quad \left. \left. + 9[(\Delta \cdot p_1)^{n-j_1-2} + (-\Delta \cdot p_2)^{n-j_1-2}] (\Delta \cdot (-p_2 - p_3))^{j_1} \right] - 3\Delta \cdot p_2 \sum_{j_1=0}^{n-3} \left[\right. \right. \\
 &\quad \left. \left. 3[(-\Delta \cdot p_3)^{j_1} - (-\Delta \cdot p_4)^{j_1}] (\Delta \cdot (p_1 + p_2))^{n-j_1-3} - (-\Delta \cdot p_4)^{j_1} (\Delta \cdot p_3)^{n-j_1-3} \right] \right) \\
 &\quad \left. + \sum_{j_1=0}^{n-2} \sum_{j_2=0}^{j_1} \left[(-\Delta \cdot p_2)^{j_1-j_2} (\Delta \cdot p_1)^{n-j_1-2} [(\Delta \cdot (p_1 + p_3))^{j_2} - 14(\Delta \cdot (p_1 + p_4))^{j_2}] \right] \right)
 \end{aligned}$$

$$\begin{aligned}
& + \frac{6d_A^{a_1 a_2 a_3 a_4}}{C_A} \left(- \sum_{j_1=0}^{n-2} \left[[(-\Delta \cdot p_3)^{j_1} + (-\Delta \cdot p_4)^{j_1}] (\Delta \cdot (p_1 + p_2))^{-j_1 + n - 2} \right] \right. \\
& \left. + \sum_{j_1=0}^{n-2} \sum_{j_2=0}^{j_1} \left[(-\Delta \cdot p_2)^{j_1 - j_2} (\Delta \cdot p_1)^{-j_1 + n - 2} [(\Delta \cdot (p_1 + p_4))^{j_2} + (\Delta \cdot (p_1 + p_3))^{j_2}] \right] \right) \Big\}.
\end{aligned} \tag{5.22}$$


Finally, the all- n Feynman rules for the operator O_A and up to 4 legs are given by



$$\rightarrow \delta^{a_1 a_2} \frac{1 + (-1)^n}{2} \left[-\Delta \cdot p_1 (p_1^{\mu_1} \Delta^{\mu_2} + \Delta^{\mu_1} p_1^{\mu_2}) + 2\Delta^{\mu_1} \Delta^{\mu_2} p_1 \cdot p_1 \right] (\Delta \cdot p_1)^{n-2}, \tag{5.23}$$



$$\begin{aligned}
& \rightarrow \frac{-i}{4} \frac{1 + (-1)^n}{2} g_s f^{a_1 a_2 a_3} \left(-4\Delta^{\mu_3} g^{\mu_1 \mu_2} \Delta \cdot p_1 (\Delta \cdot (p_1 + p_2))^{n-2} \right. \\
& - 3\Delta^{\mu_1} \Delta^{\mu_3} p_2^{\mu_2} \sum_{j_1=0}^{n-2} \left((-\Delta \cdot p_2)^{j_1} (\Delta \cdot p_1)^{-j_1 + n - 2} \right) \\
& + 2\Delta^{\mu_1} \Delta^{\mu_2} (4p_2^{\mu_3} + p_3^{\mu_3}) (\Delta \cdot p_1)^{n-2} - \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} (p_1 \cdot p_1 - p_1 \cdot p_2 + p_2 \cdot p_2) \\
& \left. \times \sum_{j_1=0}^{n-3} \left((-\Delta \cdot p_2)^{j_1} (\Delta \cdot p_1)^{-j_1 + n - 3} \right) \right) + \text{permutations}, \tag{5.24}
\end{aligned}$$



$$\begin{aligned}
& \rightarrow \frac{1}{48} \frac{1 + (-1)^n}{2} g_s^2 \left\{ 3\Delta^{\mu_1} \Delta^{\mu_2} g^{\mu_3 \mu_4} \left(8(f^{aa_1 a_3} f^{aa_2 a_4} - f^{aa_1 a_2} f^{aa_3 a_4}) (\Delta \cdot p_1)^{n-2} \right. \right. \\
& - f^{aa_1 a_2} f^{aa_3 a_4} (\Delta \cdot p_1 + \Delta \cdot p_2 + 2\Delta \cdot p_3) \\
& \left. \times \sum_{j_1=0}^{n-3} \left[[6(\Delta \cdot (-p_1 - p_2))^{j_1} + (\Delta \cdot p_2)^{j_1}] (-\Delta \cdot p_1)^{-j_1 + n - 3} \right] \right) \\
& + \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} \left((p_1 \cdot p_1 + p_2 \cdot p_2 + p_3 \cdot p_3 + p_4 \cdot p_4) f^{aa_1 a_3} f^{aa_2 a_4} \right. \\
& \left. + (13p_1 \cdot p_1 - 5p_2 \cdot p_2 + 13p_3 \cdot p_3 - 5p_4 \cdot p_4) f^{aa_1 a_2} f^{aa_3 a_4} \right)
\end{aligned}$$

$$\begin{aligned}
 & \times \sum_{j_1=0}^{n-4} \sum_{j_2=0}^{j_1} \left((-\Delta \cdot p_3)^{j_2} (\Delta \cdot (p_1 + p_2))^{j_1-j_2} (\Delta \cdot p_1)^{-j_1+n-4} \right) \\
 & - \Delta^{\mu_2} \Delta^{\mu_4} (f^{aa_1 a_3} f^{aa_2 a_4} + 13 f^{aa_1 a_2} f^{aa_3 a_4}) (\Delta^{\mu_1} p_3^{\mu_3} - \Delta^{\mu_3} p_1^{\mu_1}) \\
 & \times \sum_{j_1=0}^{n-3} \sum_{j_2=0}^{j_1} \left((\Delta \cdot (-p_1 - p_2))^{j_1-j_2} (\Delta \cdot p_3)^{j_2} (-\Delta \cdot p_1)^{-j_1+n-3} \right) + 3 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_4} \\
 & \times (4 p_1^{\mu_3} + p_3^{\mu_3}) f^{aa_1 a_3} f^{aa_2 a_4} \sum_{j_1=0}^{n-3} \left[[4 (\Delta \cdot (p_1 + p_3))^{j_1} + (\Delta \cdot p_4)^{j_1}] (-\Delta \cdot p_2)^{-j_1+n-3} \right. \\
 & \left. + 2 (\Delta \cdot p_4)^{j_1} (\Delta \cdot (-p_1 - p_3))^{-j_1+n-3} \right] + \frac{12 d_A^{a_1 a_2 a_3 a_4}}{C_A} \left(\right. \\
 & \left. - \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} p_1^{\mu_1} \sum_{j_1=0}^{n-3} \sum_{j_2=0}^{j_1} \left[(-\Delta \cdot p_2)^{j_1-j_2} (\Delta \cdot (p_1 + p_4))^{j_2} (\Delta \cdot p_1)^{-j_1+n-3} \right] \right. \\
 & \left. + \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} (p_1 \cdot p_1 + p_3 \cdot p_3) \right. \\
 & \left. \times \sum_{j_1=0}^{n-4} \sum_{j_2=0}^{j_1} \left[(-\Delta \cdot p_2)^{j_1-j_2} (\Delta \cdot (-p_2 - p_3))^{j_2} (\Delta \cdot p_1)^{-j_1+n-4} \right] \right) \Bigg\} + \text{permutations}, \tag{5.25}
 \end{aligned}$$

where plus *permutations* indicates the summation over all the external gluon indices (simultaneous permutation of μ_i, a_i, p_i). In the above equations, the fully symmetric color structure $d_A^{a_1 a_2 a_3 a_4}$ in the adjoint representation is defined by

$$d_A^{a_1 a_2 a_3 a_4} = \frac{1}{4!} \left[\text{Tr} (T_A^{a_1} T_A^{a_2} T_A^{a_3} T_A^{a_4}) + \text{symmetric permutations} \right], \tag{5.26}$$

with $(T_A)_{a_2 a_3}^{a_1} = -i f^{a_1 a_2 a_3}$.

5.3 Two-loop counterterm Feynman rules for $[ZO]_g^{\text{GV}}$

As discussed in subsection 4.2, the second order renormalization constants $Z_{gA}^{(2)}$, $Z_{gq}^{(2)}$ and $Z_{gg}^{(2)}$ can be derived from (4.11), (4.14) and (4.15) for $m = 0$. In practice, this can be done following the same computational method as described in section 3. As a reference, we write down the result for $Z_{gA}^{(2)}$ extracted from (4.11),

$$\begin{aligned}
 Z_{gA}^{(2)} &= \frac{1}{\epsilon^2 n(n-1)} \left[C_A^2 \left(\frac{56n^4 + 115n^3 - 8n^2 - 85n + 30}{12(n-1)n(n+1)(n+2)} - \frac{11S_1(n)}{4} \right) - \frac{2C_A N_f}{3} \right] \\
 &+ \frac{1}{\epsilon n(n-1)} \left[C_A^2 \left(-\frac{5}{4} S_{1,1}(n) + \frac{(35n^2 - 11n - 12) S_1(n)}{12(n-1)n} + S_{-2}(n) + 3S_2(n) \right. \right. \\
 &\left. \left. - \frac{301n^8 + 1195n^7 + 539n^6 - 2102n^5 - 1747n^4 + 1159n^3 + 1555n^2 - 72n - 180}{36(n-1)^2 n^2 (n+1)^2 (n+2)^2} \right) \right. \\
 &\left. + (1 - \xi) C_A^2 \left(\frac{2n^3 + n^2 - 1}{8(n-1)n(n+1)} - \frac{S_1(n)}{8} \right) + C_A N_f \left(\frac{16}{9} - \frac{2S_1(n)}{3} \right) \right], \tag{5.27}
 \end{aligned}$$

where two color structures, C_A^2 and $C_A N_f$, appear in the result. The three-loop correction to Z_{gA} is documented in appendix C. Unlike the physical renormalization constants Z_{ij} with $i, j = q$ or g , Z_{gA} depends on the gauge parameter ξ .

In the following, we focus on equations (4.11), (4.14) and (4.15) with $m = 1$, especially on the computation of the off-shell, two-loop, three-leg OMEs. We follow closely the method described in section 3 and work directly in parameter- x space. However, the computations are much more involved compared to the two-leg case, due to the larger number of scales:

$$p_1 \cdot p_1, p_2 \cdot p_2, p_1 \cdot p_2, \Delta \cdot p_1 = 1, \Delta \cdot p_2 = z_1, \quad (5.28)$$

where we consider an off-shell, three-particle interaction with all momenta incoming, and eliminate p_3 using momentum conservation $p_3 = -p_1 - p_2$. We also set $\Delta \cdot p_1$ to 1 and $\Delta \cdot p_2$ to z_1 .

According to the analysis in subsection 5.1, for a twist-two operator, the Feynman rules involving quarks or ghosts do not depend on the Mandelstam variables. Further, the Feynman rules containing only gluons are linear in the Mandelstam variables. Therefore, we can perform an IBP reduction by setting $p_1 \cdot p_1, p_2 \cdot p_2, p_1 \cdot p_2$ to some random non-zero rational numbers. A single numeric sample is then sufficient to determine the Feynman rules involving quarks or ghosts. Another sample is used as a cross-check. For the case involving only gluons, we also need only one numeric sample to fix the single unknown parameter, due to the symmetry constraint resulting from gluons obeying Bose statistics.

Another simplification is due to the application of simplified Lorentz structures in the Feynman rules of a twist-two operator, as discussed in subsection 5.1. In contrast to the last subsection, the application to the present case requires us to consider each of the right-hand sides of equations (4.11), (4.14) and (4.15) as a whole. To explain the reason, we analyze the Lorentz structures of each term in (4.11) for the case $m = 1$. The same analysis also applies to (4.14) and (4.15). The left-hand side of (4.11) is composed of one Lorentz structure only:

$$\langle c | [ZO]_g^{\text{GV},(2)} | c g \rangle_{\text{1PI}}^{\mu_1, (0), (1)} = d_1 \Delta^{\mu_1}. \quad (5.29)$$

The sum on the right-hand side of (4.11) also yields the same Lorentz structure, which is however not true for each term individually. For example, the following two-loop OMEs depend on three Lorentz structures

$$\left[\langle c | O_g | c g \rangle_{\text{1PI}}^{\mu_1, (2), (1), \text{B}} \right]_{\text{div}} = c_1 \Delta^{\mu_1} + c_2 p_1^{\mu_1} + c_3 p_2^{\mu_1}, \quad (5.30)$$

where c_2 and c_3 are in general non-zero, but they cancel with the corresponding structures from the $\mathcal{O}(\epsilon^0)$ part of the one-loop OMEs. Even if we do not care about the values of c_2 or c_3 , three projectors are still needed to determine the value of c_1 . To resolve this problem, we postpone the evaluation of c_1 and only evaluate the right-hand side of (4.11) as a whole. Explicitly, we contract p_{1, μ_1} with each term in the right-hand side of (4.11) and sum them up. The contributions from the last two Lorentz structures in (5.30) cancel in the result. Therefore, d_1 in (5.29) is determined by dividing the result by $\Delta \cdot p_1$, i.e.,

$$d_1 = \frac{1}{\Delta \cdot p_1} p_{1, \mu_1} \cdot [\text{right-hand side of (4.11) for } m = 1]. \quad (5.31)$$




Figure 3. Sample 2-loop Feynman diagrams to determine the counterterm Feynman rules with 3 legs stemming from $[ZO]_g^{\text{GV},(2)}$.

With the above two simplifications, the computations become much more feasible. Since they otherwise follow a standard chain of steps described in section 3, we do not need to go into much further detail. We stress that the IBP reductions and the constructions of DEs are done by setting $p_1 \cdot p_1$, $p_2 \cdot p_2$, $p_1 \cdot p_2$ to three different non-zero prime integers. About 500 master integrals appear in the result, for which we derive only the DEs with respect to x , not with respect to z_1 . For the solutions of the DEs, we fix the boundary conditions in the limit of $x \rightarrow 0$, where the integrals coincide with the all-off-shell, two-loop, 3-leg integrals without operator insertion [80]. Only a few master integrals are needed to fix the boundary conditions, but their solutions are in terms of rather complicated Goncharov multiple polylogarithms (GPLs) or Bloch-Wigner functions. From subsection 5.1, we know that Feynman rules should not contain GPLs with Mandelstam variables as their arguments. Moreover, as two-loop counterterm Feynman rules, the results are free from transcendental numbers for a fixed n , similarly to (5.27). Thus we can simplify the boundary conditions by only keeping the transcendental-weight zero contributions, which involve the following three types of master integrals only,

$$\begin{aligned}
 J_1 &= (\mu^2)^{2\epsilon} \int \frac{d^d l_1}{i\pi^{d/2}} \frac{d^d l_2}{i\pi^{d/2}} \frac{1}{l_1^2 l_2^2 (l_1 + l_2 + p_1)^2} = \frac{-p_1^2}{2\epsilon(1-2\epsilon)(2-3\epsilon)(1-3\epsilon)} + \mathcal{O}(\lambda), \\
 J_2 &= (\mu^2)^{2\epsilon} \int \frac{d^d l_1}{i\pi^{d/2}} \frac{d^d l_2}{i\pi^{d/2}} \frac{1}{l_1^2 (l_1 + p_1)^2 l_2^2 (l_2 + p_2)^2} = \frac{1}{\epsilon^2(1-2\epsilon)^2} + \mathcal{O}(\lambda), \\
 J_3 &= (\mu^2)^{2\epsilon} \int \frac{d^d l_1}{i\pi^{d/2}} \frac{d^d l_2}{i\pi^{d/2}} \frac{1}{l_1^2 (l_1 + p_1)^2 l_2^2 (l_2 - l_1 + p_2)^2} = \frac{1}{2\epsilon^2(1-2\epsilon)(1-3\epsilon)} + \mathcal{O}(\lambda),
 \end{aligned}
 \tag{5.32}$$

where λ^i indicates the contribution from polylogarithmic functions and numbers of transcendental weight i and μ is the 't Hooft scale. We do not try to solve the DEs in terms of special functions. Instead, we extract fixed Mellin moments of master integrals by solving the DEs in terms of a power series expansion in x . For each master integrals f_i we write

$$f_i(\epsilon, z_1) = \sum_{j=0}^{\infty} a_{ij}(\epsilon, z_1) x^j,
 \tag{5.33}$$

where it is straightforward to expand to high powers in x . We stress that we do not need to include $\ln(x)$ or x^ϵ terms due to our interpretation of x as a tracing parameter, see (3.1). In practice, we also expand in ϵ and set z_1 to some random prime numbers to speed up

the expansion. The reconstruction of the z_1 dependence is then performed after combining different contributions to a Feynman rule for fixed n .

For the one-loop OMEs appearing in (4.11), (4.14) and (4.15), we adopt the same method as described above, since they are required to be evaluated to order ϵ^0 . We obtain all counterterm Feynman rules with 3 legs for $[ZO]_g^{\text{GV},(2)}$ up to $n = 96$. In order to obtain all- n Feynman rules, we match our results against a heuristic ansatz. While we find that harmonic sums (3.6) are not sufficient to express the Feynman rules, we were successful in expressing them using also generalized harmonic sums [81] defined by

$$S_{\pm m}(y; n) = \sum_{j=1}^n j^{\mp m} y^j, \quad S_0(y; n) = \sum_{j=1}^n y^j,$$

$$S_{\pm m_1, m_2, \dots, m_d}(y_1, y_2, \dots, y_d; n) = \sum_{j=1}^n j^{\mp m_1} y_1^j S_{m_2, \dots, m_d}(y_2, \dots, y_d; j). \quad (5.34)$$

From the structure of the leading poles, we expect sums up to weight two with integer m_i and arguments

$$y_i \in \left\{ 1, \pm z_1, \pm(z_1 + 1), \pm \frac{1}{z_1}, \pm \frac{1}{z_1 + 1}, \pm \frac{z_1}{z_1 + 1} \right\}. \quad (5.35)$$

We reconstruct the all- n Feynman rules from fixed Mellin moments up to $n = 76$, and then cross-check our symbolic result against the remaining numerical data for moments up to $n = 96$. The package `FiniteFlow` [82] was used to speed up the reconstruction process.

As shown in (4.16), the two-leg counterterm Feynman rules for $[ZO]_g^{\text{GV},(2)}$ are zero,



$$\rightarrow 0, \quad (5.36)$$



$$\rightarrow 0, \quad (5.37)$$




$$\rightarrow 0. \quad (5.38)$$


The two-loop counterterm Feynman rules contributing to $[ZO]_g^{\text{GV}}$, i.e, the left-hand sides of (4.11), (4.14), (4.15), are listed in the following, with the convention of all momenta flowing into the vertices,



$$\rightarrow 0, \quad (5.39)$$



$$\begin{aligned} &\rightarrow 2ig_s C_A^2 f^{a_1 a_2 a_3} \frac{1 + (-1)^n}{256n(n-1)} (\Delta \cdot p_1)^{n-1} \Delta^{\mu_3} \left\{ \frac{F_{-2,0} + (1-\xi) F_{-2,1}}{\epsilon^2} \right. \\ &\quad \left. + \frac{F_{-1,0} + (1-\xi) F_{-1,1}}{\epsilon} \right\}, \end{aligned} \quad (5.40)$$



$$\begin{aligned} &\rightarrow 2ig_s C_A^2 f^{a_1 a_2 a_3} \frac{1 + (-1)^n}{256n(n-1)} \frac{(\Delta \cdot p_1)^{n-2}}{\Delta \cdot p_2} \left(-\Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} (p_1 \cdot p_1 + p_2 \cdot p_2 + p_3 \cdot p_3) \right. \\ &\quad \left. + \Delta^{\mu_2} \Delta^{\mu_3} p_1^{\mu_1} \Delta \cdot p_1 + \Delta^{\mu_1} \Delta^{\mu_3} p_2^{\mu_2} \Delta \cdot p_2 + \Delta^{\mu_1} \Delta^{\mu_2} p_3^{\mu_3} \Delta \cdot p_3 \right) \\ &\quad \times \left\{ \frac{F_{-2,0} + (1-\xi) F_{-2,1}}{\epsilon^2} + \frac{F_{-1,0} + (1-\xi) F_{-1,1}}{\epsilon} \right\}, \end{aligned} \quad (5.41)$$

where the quark-quark-gluon counterterm Feynman rule for $[ZO]_g^{\text{GV},(2)}$ is zero. The counterterm Feynman rules for 3-gluon and ghost-ghost-gluon vertices are proportional to each other, indicating they are related by a generalized BRST symmetry. The scalar form factors in (5.41) are as follows:

$$F_{-2,1} = -\frac{(z_1-1)(1+z_1)^n}{(z_1+1)^2} + \frac{(z_1+2)z_1^n}{z_1(z_1+1)} - \frac{(2z_1+1)1^n}{z_1+1}, \quad (5.42)$$

$$\begin{aligned} F_{-2,0} = z_1^n &\left[-\frac{2(z_1-2)}{z_1(z_1+1)} S_1\left(-\frac{1}{z_1}; n\right) + \frac{2(3z_1+2)}{z_1(z_1+1)} S_1\left(\frac{z_1+1}{z_1}; n\right) \right. \\ &\quad \left. - \frac{2(z_1+2)(2n^2 z_1 + 2n^2 - 2nz_1^2 - 9nz_1 - 9n + 2z_1^2 + 3z_1 + 3)}{(n-1)nz_1(z_1+1)^2} - \frac{6(z_1+2)}{z_1(z_1+1)} S_1(n) \right] \\ &+ (1+z_1)^n \left[\frac{2(z_1+3)}{(z_1+1)^2} S_1\left(\frac{1}{z_1+1}; n\right) - \frac{2(3z_1+1)}{(z_1+1)^2} S_1\left(\frac{z_1}{z_1+1}; n\right) \right. \\ &\quad \left. + \frac{2(z_1-1)(2n^2 z_1 + 2nz_1^2 - 5nz_1 + 2n - 2z_1^2 - z_1 - 2)}{(n-1)nz_1(z_1+1)^2} + \frac{6(z_1-1)}{(z_1+1)^2} S_1(n) \right] \\ &+ 1^n \left[-\frac{2(2z_1-1)}{z_1+1} S_1(-z_1; n) - \frac{2(2z_1+3)}{z_1+1} S_1(z_1+1; n) + \frac{6(2z_1+1)}{z_1+1} S_1(n) \right. \\ &\quad \left. + \frac{2(2z_1+1)(2n^2 z_1^2 + 2n^2 z_1 - 9nz_1^2 - 9nz_1 - 2n + 3z_1^2 + 3z_1 + 2)}{(n-1)nz_1(z_1+1)^2} \right], \end{aligned} \quad (5.43)$$

$$\begin{aligned}
 F_{-1,1} = z_1^n & \left[-\frac{(z_1+2)(n^2 z_1 + n^2 - n z_1^2 + 2n z_1 + 2n + z_1^2 - z_1 - 1)}{(n-1)n z_1 (z_1+1)^2} \right. \\
 & + \frac{S_1\left(-\frac{1}{z_1}; n\right)}{z_1} + \frac{S_1\left(\frac{z_1+1}{z_1}; n\right)}{z_1(z_1+1)} \left. \right] + (1+z_1)^n \left[\frac{S_1\left(\frac{z_1}{z_1+1}; n\right)}{(z_1+1)^2} - \frac{z_1 S_1\left(\frac{1}{z_1+1}; n\right)}{(z_1+1)^2} \right. \\
 & + \frac{(z_1-1)(n^2 z_1 + n z_1^2 + 4n z_1 + n - z_1^2 - 3z_1 - 1)}{(n-1)n z_1 (z_1+1)^2} \left. \right] + 1^n \left[-S_1(-z_1; n) \right. \\
 & \left. - \frac{z_1 S_1(z_1+1; n)}{z_1+1} + \frac{(2z_1+1)(n^2 z_1^2 + n^2 z_1 + 2n z_1^2 + 2n z_1 - n - z_1^2 - z_1 + 1)}{(n-1)n z_1 (z_1+1)^2} \right], \tag{5.44}
 \end{aligned}$$

$$\begin{aligned}
 F_{-1,0} = z_1^n & \left[-\frac{4n}{z_1(z_1+1)} S_2\left(\frac{1}{z_1}; n\right) + \frac{8n(z_1+2)}{z_1(z_1+1)} S_{-2}(n) - \frac{4(n+1)(z_1+2)}{z_1(z_1+1)} S_2(n) \right. \\
 & + \frac{6(z_1+2)}{z_1(z_1+1)} S_{1,1}(n) + \frac{4(n+z_1)}{z_1(z_1+1)} S_2\left(\frac{z_1+1}{z_1}; n\right) - \frac{2(3z_1+2)}{z_1(z_1+1)} S_{1,1}\left(1, \frac{z_1+1}{z_1}; n\right) \\
 & + \frac{4(z_1 n + n - z_1)}{z_1(z_1+1)} S_2\left(-\frac{1}{z_1}; n\right) - \frac{4n}{z_1} S_2\left(-\frac{z_1+1}{z_1}; n\right) - \frac{4n}{z_1} S_{1,1}\left(-\frac{1}{z_1}, 1; n\right) \\
 & - \frac{2(z_1+2)(6z_1 n^3 + 6n^3 - 4z_1^2 n^2 - 15z_1 n^2 - 15n^2 + 6z_1^2 n + 16z_1 n + 16n - 2z_1^2 - 3z_1 - 3)}{(n-1)n^2 z_1 (z_1+1)^2} \\
 & - \frac{2(z_1+2)(2z_1^2 n^2 + z_1 n^2 + n^2 - 2z_1^2 n - 7z_1 n - 7n + 2z_1 + 2)}{(n-1)n z_1 (z_1+1)^2} S_1(n) \\
 & - \frac{2(2nz_1^3 - nz_1^2 + 2z_1^2 - 5nz_1 + 2z_1 - 4n)}{nz_1(z_1+1)^2} S_1\left(\frac{z_1+1}{z_1}; n\right) + \frac{4n}{z_1} S_{1,1}\left(-\frac{1}{z_1}, -z_1; n\right) \\
 & + \frac{2(2nz_1^3 + 3nz_1^2 + 2z_1^2 + 7nz_1 + 2z_1 + 4n)}{nz_1(z_1+1)^2} S_1\left(-\frac{1}{z_1}; n\right) - \frac{4n}{z_1} S_{1,1}\left(-\frac{1}{z_1}, z_1+1; n\right) \\
 & - \frac{4n}{z_1(z_1+1)} S_{1,1}\left(\frac{z_1+1}{z_1}, 1; n\right) - \frac{4n S_{1,1}\left(\frac{z_1+1}{z_1}, \frac{1}{z_1+1}; n\right)}{z_1(z_1+1)} + \frac{4n S_{1,1}\left(\frac{z_1+1}{z_1}, \frac{z_1}{z_1+1}; n\right)}{z_1(z_1+1)} \\
 & + \frac{2(z_1-2)}{z_1(z_1+1)} S_{1,1}\left(1, -\frac{1}{z_1}; n\right) \left. \right] + (1+z_1)^n \left[-\frac{4(nz_1 - z_1 - 1)}{(z_1+1)^2} S_2\left(\frac{1}{z_1+1}; n\right) \right. \\
 & - \frac{2(z_1+3)}{(z_1+1)^2} S_{1,1}\left(1, \frac{1}{z_1+1}; n\right) + \frac{2(3z_1+1)}{(z_1+1)^2} S_{1,1}\left(1, \frac{z_1}{z_1+1}; n\right) - \frac{4n}{(z_1+1)^2} S_2\left(-\frac{1}{z_1+1}; n\right) \\
 & - \frac{2(z_1-1)(2z_1^2 n^2 + 3z_1 n^2 + 2n^2 - 2z_1^2 n + 3z_1 n - 2n - 2z_1)}{(n-1)n z_1 (z_1+1)^2} S_1(n) \\
 & + \frac{2(z_1-1)(6z_1 n^3 + 4z_1^2 n^2 - 7z_1 n^2 + 4n^2 - 6z_1^2 n + 4z_1 n - 6n + 2z_1^2 + z_1 + 2)}{(n-1)n^2 z_1 (z_1+1)^2} \\
 & + \frac{2(2nz_1^3 + 3nz_1^2 - 2z_1^2 + 7nz_1 - 2z_1 + 2n)}{nz_1(z_1+1)^2} S_1\left(\frac{1}{z_1+1}; n\right) + \frac{4nz_1}{(z_1+1)^2} S_2\left(-\frac{z_1}{z_1+1}; n\right) \\
 & - \frac{2(2nz_1^3 + 7nz_1^2 - 2z_1^2 + 3nz_1 - 2z_1 + 2n)}{nz_1(z_1+1)^2} S_1\left(\frac{z_1}{z_1+1}; n\right) - \frac{4n}{(z_1+1)^2} S_{1,1}\left(\frac{z_1}{z_1+1}, 1; n\right) \\
 & - \frac{4n}{(z_1+1)^2} S_{1,1}\left(\frac{z_1}{z_1+1}, -\frac{1}{z_1}; n\right) + \frac{4n S_{1,1}\left(\frac{z_1}{z_1+1}, \frac{z_1+1}{z_1}; n\right)}{(z_1+1)^2} + \frac{4(n-z_1-1)}{(z_1+1)^2} S_2\left(\frac{z_1}{z_1+1}; n\right) \\
 & - \frac{8n(z_1-1)}{(z_1+1)^2} S_{-2}(n) + \frac{4(n+1)(z_1-1)}{(z_1+1)^2} S_2(n) - \frac{6(z_1-1)}{(z_1+1)^2} S_{1,1}(n) \\
 & \left. + \frac{4nz_1 S_{1,1}\left(\frac{1}{z_1+1}, 1; n\right)}{(z_1+1)^2} + \frac{4nz_1 S_{1,1}\left(\frac{1}{z_1+1}, -z_1; n\right)}{(z_1+1)^2} - \frac{4nz_1 S_{1,1}\left(\frac{1}{z_1+1}, z_1+1; n\right)}{(z_1+1)^2} \right]
 \end{aligned}$$

$$\begin{aligned}
& +1^n \left[4nS_2(-z_1-1;n) + 4nS_{1,1}(-z_1,1;n) - 4nS_{1,1}\left(-z_1,-\frac{1}{z_1};n\right) \right. \\
& + 4nS_{1,1}\left(-z_1,\frac{z_1+1}{z_1};n\right) + \frac{2(2z_1-1)}{z_1+1}S_{1,1}(1,-z_1;n) - \frac{8n(2z_1+1)}{z_1+1}S_{-2}(n) \\
& + \frac{4(n+1)(2z_1+1)}{z_1+1}S_2(n) - \frac{6(2z_1+1)}{z_1+1}S_{1,1}(n) + \frac{2(2z_1+3)}{z_1+1}S_{1,1}(1,z_1+1;n) \\
& - \frac{4(nz_1+1)}{z_1+1}S_2(z_1+1;n) - \frac{4(z_1n+n-1)}{z_1+1}S_2(-z_1;n) \\
& + \frac{2(2z_1+1)(z_1^2n^2+z_1n^2+2n^2-7z_1^2n-7z_1n-2n+2z_1^2+2z_1)}{(n-1)nz_1(z_1+1)^2}S_1(n) \\
& + \frac{1}{(n-1)n^2z_1(z_1+1)^2} \left[2(2z_1+1)(6z_1^2n^3+6z_1n^3-15z_1^2n^2-15z_1n^2 \right. \\
& \left. - 4n^2+16z_1^2n+16z_1n+6n-3z_1^2-3z_1-2) \right] \\
& - \frac{2(4nz_1^3+5nz_1^2-2z_1^2+nz_1-2z_1-2n)}{nz_1(z_1+1)^2}S_1(z_1+1;n) + \frac{4nz_1}{z_1+1}S_{1,1}(z_1+1,1;n) \\
& - \frac{2(4nz_1^3+7nz_1^2+2z_1^2+3nz_1+2z_1+2n)}{nz_1(z_1+1)^2}S_1(-z_1;n) + \frac{4nz_1}{z_1+1}S_2(z_1;n) \\
& \left. - \frac{4nz_1S_{1,1}\left(z_1+1,\frac{1}{z_1+1};n\right)}{z_1+1} + \frac{4nz_1S_{1,1}\left(z_1+1,\frac{z_1}{z_1+1};n\right)}{z_1+1} \right], \tag{5.45}
\end{aligned}$$

where

$$z_1 = \frac{\Delta \cdot p_2}{\Delta \cdot p_1},$$

and the following 46 (generalized) harmonic sums up to weight 2 appear

$$\begin{aligned}
& \left\{ S_1(n), S_1\left(-\frac{1}{z_1};n\right), S_1(-z_1;n), S_1\left(\frac{1}{z_1+1};n\right), S_1\left(\frac{z_1}{z_1+1};n\right), S_1(z_1+1;n), \right. \\
& S_1\left(\frac{z_1+1}{z_1};n\right), S_{-2}(n), S_2(n), S_{1,1}(n), S_2(-z_1-1;n), S_2\left(-\frac{1}{z_1};n\right), S_2\left(\frac{1}{z_1};n\right), \\
& S_2(-z_1;n), S_2(z_1;n), S_2\left(-\frac{1}{z_1+1};n\right), S_2\left(\frac{1}{z_1+1};n\right), S_2\left(-\frac{z_1}{z_1+1};n\right), \\
& S_2\left(\frac{z_1}{z_1+1};n\right), S_2(z_1+1;n), S_2\left(-\frac{z_1+1}{z_1};n\right), S_2\left(\frac{z_1+1}{z_1};n\right), S_{1,1}\left(1,-\frac{1}{z_1};n\right), \\
& S_{1,1}\left(1,-z_1;n\right), S_{1,1}\left(1,\frac{1}{z_1+1};n\right), S_{1,1}\left(1,\frac{z_1}{z_1+1};n\right), S_{1,1}\left(1,z_1+1;n\right), \\
& S_{1,1}\left(1,\frac{z_1+1}{z_1};n\right), S_{1,1}\left(-\frac{1}{z_1},1;n\right), S_{1,1}\left(-\frac{1}{z_1},-z_1;n\right), S_{1,1}\left(-\frac{1}{z_1},z_1+1;n\right), \\
& S_{1,1}\left(-z_1,1;n\right), S_{1,1}\left(-z_1,-\frac{1}{z_1};n\right), S_{1,1}\left(-z_1,\frac{z_1+1}{z_1};n\right), S_{1,1}\left(\frac{1}{z_1+1},1;n\right), \\
& S_{1,1}\left(\frac{1}{z_1+1},-z_1;n\right), S_{1,1}\left(\frac{1}{z_1+1},z_1+1;n\right), S_{1,1}\left(\frac{z_1}{z_1+1},1;n\right), \\
& S_{1,1}\left(\frac{z_1}{z_1+1},\frac{z_1+1}{z_1};n\right), S_{1,1}\left(z_1+1,1;n\right), S_{1,1}\left(z_1+1,\frac{1}{z_1+1};n\right), \\
& S_{1,1}\left(z_1+1,\frac{z_1}{z_1+1};n\right), S_{1,1}\left(\frac{z_1+1}{z_1},1;n\right), S_{1,1}\left(\frac{z_1+1}{z_1},\frac{1}{z_1+1};n\right), \\
& \left. S_{1,1}\left(\frac{z_1}{z_1+1},-\frac{1}{z_1};n\right), S_{1,1}\left(\frac{z_1+1}{z_1},\frac{z_1}{z_1+1};n\right) \right\}. \tag{5.46}
\end{aligned}$$

We used `HarmonicSums` to evaluate these (generalized) harmonic sums throughout.

We stress that our counterterm Feynman rules are given in closed form with symbolic n dependence. Due to the appearance of the generalized harmonic sums, by keeping the all- n dependence, it is not possible to disentangle the renormalization constants (characterized by their independence of z_1 in this case) from the corresponding operators (which are independent of harmonic sums). Therefore, it appears impossible to determine the individual $i = 2$ operators in a closed form for symbolic n in this way.

It is, however, possible to work out the operator basis n -by- n , as considered in [30]. Typically, more and more operators are needed as n increases, and the number is expected to go to infinity when n tends to infinity. For a given value of n , only a finite number of operators are contributing through their operator Feynman rules. From our all- n counterterm Feynman rules, we can deduce the operator basis n -by- n by setting n to a specific number. For example, the scalar products in (5.45) for $n = 12$ are

$$\begin{aligned}
 F_{-2,1} \Big|_{n=12} &= -(z_1 - 1) z_1 (z_1 + 2) (2z_1 + 1) (z_1^2 + z_1 + 2) (2z_1^2 + z_1 + 1) (2z_1^2 + 3z_1 + 2) , \\
 F_{-2,0} \Big|_{n=12} &= \frac{1}{630} (z_1 - 1) z_1 (z_1 + 2) (2z_1 + 1) \left[34126z_1^6 + 102378z_1^5 + 215515z_1^4 \right. \\
 &\quad \left. + 260400z_1^3 + 215515z_1^2 + 102378z_1 + 34126 \right] , \\
 F_{-1,1} \Big|_{n=12} &= \frac{(z_1 - 1) z_1 (z_1 + 2) (2z_1 + 1)}{27720} \left[180899z_1^6 + 542697z_1^5 + 1137425z_1^4 \right. \\
 &\quad \left. + 1370355z_1^3 + 1137425z_1^2 + 542697z_1 + 180899 \right] , \\
 F_{-1,0} \Big|_{n=12} &= \frac{(z_1 - 1) z_1 (z_1 + 2) (2z_1 + 1)}{1164240} \left[9395264z_1^6 + 28185792z_1^5 \right. \\
 &\quad \left. + 53213569z_1^4 + 59450818z_1^3 + 53213569z_1^2 + 28185792z_1 + 9395264 \right] , \tag{5.47}
 \end{aligned}$$

where the results are polynomials in z_1 with constant coefficients, which makes it straightforward to separate the renormalization constants (rational numbers) and the Feynman rules of the corresponding operators. To infer the operators from the corresponding Feynman rules, we need to replace a momentum with the derivative of a field, for example, $(\Delta \cdot p_1(1 + z_1))^n = (-\Delta \cdot p_3)^n \rightarrow (-\Delta \cdot \partial)^n A_\mu^a$, where A_μ^a is a gluon field.

5.4 Comparison with previous fixed n results

In a recent work [30], Falcioni and Herzog have used gauge and BRST symmetry to derive a set of constraint equations for the GV operators. By applying these constraint equations to a general ansatz for the GV operators at fixed values of n , they were able to determine the required operator bases for low Mellin moments $n = 2, 4, 6$ and to compute the associated renormalization constants.

In the following, we compare our all- n counterterm Feynman rules evaluated at $n = 2, 4$ and 6 with their results. Since our approach does not allow to disentangle GV operators and renormalization constants, the comparison requires us to first determine the GV operator Feynman rules resulting from [30], which are subsequently multiplied with corresponding

renormalization constants in order to determine the counterterm Feynman rules. It can be easily verified from (5.40) and (5.41) that the counterterm Feynman rules from $[ZO]_g^{\text{GV}}$ are non-zero only for $n \geq 6$. Therefore, for $n = 2, 4$ the GV operator basis is contained in the operators O_{ABC} . We presented detailed comparisons for each n separately below. All comparisons are truncated up to four legs, and we find full agreement with [30].

For $n = 2$, the GV operators were expressed as $\mathcal{O}_1^{(2)}$ and $\mathcal{O}_2^{(2)}$ in [30], and we find that they are related to our results as follows:

$$\begin{aligned} O_g \Big|_{n=2} &= -\mathcal{O}_1^{(2)}, \\ Z_{gA} (O_{AC} \Big|_{\text{FR}}) \Big|_{n=2, N_f=0} &= -\delta Z_{12}^{(2)} \mathcal{O}_2^{(2)} \Big|_{\text{FR}}, \end{aligned} \tag{5.48}$$

where we use ‘FR’ to indicate the Feynman rules resulting from the corresponding operators. The renormalization constant $\delta Z_{12}^{(2)}$ in the above equation is given to order a_s^3 with full ξ dependence in [30]. Our result for $Z_{gA}|_{n=2, N_f=0}$, in (C.8), is in full agreement with $\delta Z_{12}^{(2)}$ presented in [30]. The overall minus sign in the above equation is due to different normalizations of O_g and \mathcal{O}_1 ,

$$O_g = -i^{n-2} \mathcal{O}_1. \tag{5.49}$$

For $n = 4$, the GV operator basis in [30] consists of three elements, which relate to our results as follows:

$$\begin{aligned} O_g \Big|_{n=4} &= \mathcal{O}_1^{(4)}, \\ Z_{gA} (O_{AC} \Big|_{\text{FR}}) \Big|_{n=4, N_f=0} &= \delta Z_{12}^{(4)} \mathcal{O}_2^{(4)} \Big|_{\text{FR}} + \delta Z_{13}^{(4)} \mathcal{O}_3^{(4)} \Big|_{\text{FR}}, \end{aligned} \tag{5.50}$$

where the operators $\mathcal{O}_2^{(4)}, \mathcal{O}_3^{(4)}$ are distinguished from each other by their different color structures with $\mathcal{O}_3^{(4)}$ being proportional to $d_A^{a_1 a_2 a_3 a_4}$ as defined in (5.26). In [30], the constant $\delta Z_{12}^{(4)}$ was given to a_s^3 with $a_s^3(1 - \xi)^2$ being dropped, and $\delta Z_{13}^{(4)}$ was given to a_s^1 with

$$\delta Z_{13}^{(4)} = a_s \frac{C_A}{24\epsilon} + \mathcal{O}(a_s^2). \tag{5.51}$$

We find the following relations between the renormalization constants:

$$\begin{aligned} Z_{gA} \Big|_{n=4, N_f=0} - \delta Z_{12}^{(4)} &= \mathcal{O}(a_s^3(1 - \xi)^2), \\ Z_{gA} \Big|_{n=4, N_f=0} + 2\delta Z_{13}^{(4)} &= \mathcal{O}(a_s^2), \end{aligned} \tag{5.52}$$

with $Z_{gA}|_{n=4}$ listed in (C.9). Given that these renormalization constants were truncated [83] at order $(1 - \xi)$ in [30], we thus find full agreement.

For $n = 6$, the comparison becomes more interesting, since the counterterm operator $[ZO]_g^{\text{GV},(2)}$ starts to contribute. In this case, we further decompose the comparison into three different categories, depending on the number of legs. As the first category, for counterterm Feynman rules with two legs, we find the following relation,

$$Z_{gA} (O_{AC} \Big|_{\text{FR}}) \Big|_{n=6, N_f=0} = -\delta Z_{12}^{(6)} \mathcal{O}_2^{(6)} \Big|_{\text{FR}}. \tag{5.53}$$

The above relation is expected to be satisfied to all orders. It is checked explicitly to order a_s^2 since $\delta Z_{12}^{(6)}$ is given to a_s^2 in [30]. For counterterm Feynman rules with three legs, the following relation holds,

$$Z_{gA} (O_{AC}|_{\text{FR}}) \Big|_{n=6, N_f=0} + \left[[ZO]_g^{\text{GV},(2)} |_{\text{FR}} \right] \Big|_{n=6} = - \left\{ \delta Z_{12}^{(6)} \mathcal{O}_2^{(6)} |_{\text{FR}} + \delta Z_{13}^{(6)} \mathcal{O}_3^{(6)} |_{\text{FR}} \right\} + \mathcal{O}(a_s^3). \quad (5.54)$$

Interestingly, both $\delta Z_{12}^{(6)}$ and $\delta Z_{13}^{(6)}$ start to contribute at order a_s , while only the first term in the left-hand side of the above equation contributes at order a_s . At order a_s^2 , also $[ZO]_g^{\text{GV},(2)}$ starts to contribute. The above relation is confirmed explicitly to order a_s^2 . For counterterm Feynman rules with four legs, the following relation holds¹ at order a_s^1 ,

$$Z_{gA} (O_{AC}|_{\text{FR}}) \Big|_{n=6, N_f=0} = - \left\{ \delta Z_{12}^{(6)} \mathcal{O}_2^{(6)} |_{\text{FR}} + \delta Z_{13}^{(6)} \mathcal{O}_3^{(6)} |_{\text{FR}} + \delta Z_{14}^{(6)} \mathcal{O}_4^{(6)} |_{\text{FR}} + \delta Z_{15}^{(6)} \mathcal{O}_5^{(6)} |_{\text{FR}} \right\} + \mathcal{O}(a_s^2), \quad (5.55)$$

where the four operators on the right-hand side are assembled into a single operator O_{AC} on the left-hand side. At order a_s^2 , one more operator $\mathcal{O}_6^{(6)}$ and one more counterterm operator $[ZO]_g^{\text{GV},(2)}$ will appear on the right-hand side and the left-hand side of this equation, respectively. We do not consider them in this paper, since they start to contribute to the OME renormalization first at the four-loop order.

6 Three-loop splitting functions from operator insertions

The final goal of this paper is the application of our framework to the computation of the three-loop splitting functions. Having worked out the Feynman rules for physical operators as well as the renormalization counterterms that originate from GV operators, the main remaining task is the computation of the two-point OMEs with different operator insertions, i.e. (2.30).

With the insertion of the operator O_q or O_g , the two-point OMEs need to be evaluated to three loops. Regarding the insertion of O_{ABC} , we need to compute the corresponding two-point OMEs to two loops. Finally, the two-point OMEs with the $[ZO]_g^{\text{GV},(2)}$ counterterm insertion are needed to one loop. In table 1, we list all required OMEs for the extraction of the three-loop splitting functions, including also the multi-leg OMEs used for the determination of the counterterms and specifying the respective loop order. Examples for two-point diagrams with counterterm insertions are depicted in figure 4.

For physical operators as well as for $i = 1$ GV operators O_{ABC} , it is straightforward to turn the corresponding Feynman rules from n -space into x -parameter space using linear propagators according to (3.1). However, the all- n Feynman rules for the counterterm $[ZO]_g^{\text{GV},(2)}$, as shown in (5.40), (5.41), leads to polylogarithms in x -parameter space. In

¹We thank Falcioni and Herzog for pointing out to us, that the term $+A^{a_1} \partial^2 A^{a_2} \partial c^{a_3}$ in the third line of equation (5.31) in [30] defining the operator $\mathcal{O}_3^{(6)}$ should be modified to $+4A^{a_1} \partial^2 A^{a_2} \partial c^{a_3}$ [83]. After correcting this minor typesetting issue, we find full agreement between their and our results.

Legs \ Loops	2	3	4	5
0		$[ZO]_g^{\text{GV},(2)}$	O_{ABC}	O_q, O_g
1	$[ZO]_g^{\text{GV},(2)}$	O_{ABC}	O_g	
2	O_{ABC}	O_g		
3	O_q, O_g			

Table 1. Summary of all OMEs entering the calculation of 3-loop splitting functions. OMEs with two legs have two quarks, two ghosts or two gluons in the external state. The determination of the counterterm Feynman rules requires also OMEs with additional external gluons. For each operator, the table denotes the maximal number of loops needed for OMEs with a given number of legs.




Figure 4. Sample diagrams with GV counterterm insertions entering the calculation of the 3-loop splitting functions.

practice, it is not clear how to perform IBP reductions of Feynman integrals involving polylogarithmic dependence on the loop kinematics. Therefore, the method of section 3 can only be used to compute two-point OMEs with an insertion of O_q , O_g or O_{ABC} , but not with $[ZO]_g^{\text{GV},(2)}$.

The contributions due to $[ZO]_g^{\text{GV},(2)}$ consists of three-gluon and ghost-ghost-gluon counterterms. In the following, we explain our procedure to compute the corresponding one-loop, two-gluon OMEs using the example of the ghost-ghost-gluon counterterm (5.40); the procedure for the three-gluon counterterm (5.41) follows identical steps. For a fixed value of n , the general structure of the ghost-ghost-gluon vertex (5.40) reads:

$$ig_s f^{a_1 a_2 a_3} \Delta^{\mu_3} \sum_{m=1}^{n-2} a_{mn} (\Delta \cdot p_1)^m (\Delta \cdot p_2)^{n-1-m}, \tag{6.1}$$

where a_{mn} can not be written as a closed rational form in (m, n) . If a_{mn} does not depend on m and n (this is the case for the Feynman rules of the operators O_q , O_g as well as O_{ABC}), or if it has a polynomial dependence on m and n , a single auxiliary parameter x as used in section 3 is sufficient to express the above equation in terms of linear propagators. For example, if $a_{mn} = m$, then

$$\sum_{n=3}^{\infty} x^n \sum_{m=1}^{n-2} m (\Delta \cdot p_1)^m (\Delta \cdot p_2)^{n-1-m} = \frac{x^2 \Delta \cdot p_1}{(1 - x \Delta \cdot p_1)^2} \frac{x \Delta \cdot p_2}{1 - x \Delta \cdot p_2}. \tag{6.2}$$

Without turning (6.1) into a linear propagator, it is still possible to directly evaluate the corresponding OMEs for low Mellin moments n . However, when n increases (typically

values of n required for an all- n reconstruction range into the hundreds or thousands), the numerator degrees of the resulting Feynman integrals become so large that an IBP reduction seems inefficient.

We thus employ an alternative method to bypass this complexity. In addition to x , we introduce one more auxiliary parameter t and replace a_{mn} in (6.1) with t^m ,

$$h(x, t) = \sum_{n=3}^{\infty} x^n \sum_{m=1}^{n-2} t^m (\Delta \cdot p_1)^m (\Delta \cdot p_2)^{n-1-m} = \frac{x t \Delta \cdot p_1}{1 - x t \Delta \cdot p_1} \frac{x^2 \Delta \cdot p_2}{1 - x \Delta \cdot p_2}. \quad (6.3)$$

In this way, we manage to turn (6.1) into linear propagators depending on two tracing parameters x and t . By considering the matrix elements with an insertion of the above linear propagators, we have

$$\langle g(p) | h(x, t) | g(p) \rangle = \sum_{n=3}^{\infty} x^n \sum_{m=1}^{n-2} c_{mn} t^m, \quad (6.4)$$

where in this way we can perform a standard IBP reduction with the corresponding Feynman integrals without encountering high numerator degrees (at the expense of an extra symbolic parameter t in the IBP relations). The c_{mn} can be determined from differential equations in x up to high values of n . Using the following formula, we can then read out the results of OMEs for multiple fixed values of n easily,

$$\langle g(p) | \sum_{m=1}^{n-2} a_{mn} (\Delta \cdot p_1)^m (\Delta \cdot p_2)^{n-1-m} | g(p) \rangle = \sum_{m=1}^{n-2} a_{mn} c_{mn}. \quad (6.5)$$

Using the all- n counterterm Feynman rules, we can thus efficiently compute the OMEs at fixed Mellin moments up to high values of n . By introducing the auxiliary parameter t , we avoid the growth in complexity for the IBP reduction at fixed numerical n with increasing n , and the IBP-reduced results are expressed as simple one-loop bubble integrals with a linear propagator insertion.

We compute all required two-point OMEs as listed in table 1 in a covariant gauge with full ξ dependence. The all- n results for OMEs with an operator insertion of O_q , O_g or O_{ABC} are computed using the method described in section 3. The 3-loop non-singlet splitting functions can be determined in the off-shell approach without the GV counterterms presented in this paper; the required OMEs have been computed in [37], and we find full agreement with them. The OMEs with the counterterm operator $[ZO]_g^{\text{GV},(2)}$ insertion are computed for fixed Mellin moments up to $n = 500$ with the procedure described above. These moments are then used for a reconstruction and cross-check of the all- n results to order ϵ^0 , with the following combination contributing to the 3-loop splitting functions,

$$\begin{aligned} & \frac{1}{(d-2)(N_c^2-1)} \left[-g_{\mu\nu} \langle g | [ZO]_g^{\text{GV},(2)} | g \rangle_{\text{1PI}}^{\mu\nu,(1),(0)} - 2 \langle c | [ZO]_g^{\text{GV},(2)} | c \rangle_{\text{1PI}}^{(1),(0)} \right] = \\ & \frac{(1-\xi)C_A^3}{128n(n-1)\epsilon^2} \left\{ -\frac{4(4n^2+13n-13)S_1(n)}{(n-1)n} + \frac{4(n+1)(8n^2-7n-3)}{(n-1)^2n^2} \right. \\ & \left. + (1-\xi) \left(\frac{2(n+1)(2n-3)}{(n-1)n} - 2S_1(n) \right) - 8S_{-2}(n) + 12S_2(n) + 12S_{1,1}(n) \right\} \end{aligned}$$

$$\begin{aligned}
& + \frac{(1-\xi)C_A^3}{128n(n-1)\epsilon} \left\{ (1-\xi)^2 \left[S(1,n) - \frac{(n+1)(2n-3)}{(n-1)n} \right] \right. \\
& + (1-\xi) \left[\frac{2(4n^2+18n-21)S(1,n)}{(n-1)n} + 8S(-2,n) - 8S(2,n) - 8S(1,1,n) \right. \\
& + \left. \left. \frac{2(6n^4-28n^3+16n^2+17n-3)}{(n-1)^2n^2} \right] + \frac{8(2n^2-n+1)S(-2,n)}{(n-1)n} \right. \\
& - \frac{8(n^2+6n-6)S(2,n)}{(n-1)n} + \frac{4(12n^2-31n+15)S(1,1,n)}{(n-1)n} \\
& - \frac{4(20n^4-24n^3-23n^2+44n-7)S(1,n)}{(n-1)^2n^2} + 4(6n-7)S(-3,n) \\
& - 8(2n-3)S(3,n) - 8(10n+1)S(-2,1,n) + 16S(1,-2,n) + 28S(1,2,n) \\
& \left. \left. - 8(n+1)S(2,1,n) + \frac{8(4n^5+2n^4-13n^3+12n^2-10n+3)}{(n-1)^3n^2} \right\} + \mathcal{O}(\epsilon^0). \tag{6.6}
\end{aligned}$$

The quantity in the above equation is zero in Feynman gauge with $\xi = 1$. It implies that the correct three-loop splitting functions can be extracted without considering the contribution from the counterterm operator $[ZO]_g^{\text{GV},(2)}$ if we work in Feynman gauge, which is exactly what we found in [84]. We point out that we compute all two-parton OMEs to one order higher in ϵ than what is needed for the extraction of the three-loop splitting functions, since those contributions will enter the renormalization at the four-loop level. That is, we compute the OMEs at three-loop, two-loop and one-loop order to ϵ^0 , ϵ^1 and ϵ^2 , respectively. We collect results of all two-parton OMEs in supplementary material attached to this paper and provide instructions for their usage in appendix E.

Having the results for all required OMEs at hand, it is straightforward to extract the physical renormalization constants according to the renormalization procedure in (2.25) and (2.26). We find that the ξ dependence indeed cancels for the physical renormalization constants, stressing that the inclusion of (6.6) is crucial for this cancellation. According to (2.20), we extract the physical singlet anomalous dimensions $\gamma_{ij}(n)$ to three loops. Our results are in full agreement with the results in [11]. Through a similar procedure, we also extract non-singlet anomalous dimension $\gamma_{\text{ns}}(n)$ to three loops. Separating even and odd moments, γ_{ns} can be decomposed as follows:

$$\gamma_{\text{ns}} = \frac{1+(-1)^n}{2} \gamma_{\text{ns}}^+ + \frac{1-(-1)^n}{2} (\gamma_{\text{ns}}^- + \gamma_{\text{ns}}^s), \tag{6.7}$$

where the definitions of γ_{ns}^\pm and γ_{ns}^s can be found for example in [10]. The anomalous dimension γ_{ns}^s starts to contribute at the three-loop level and is proportional to the color structure $d^{abc}d_{abc}$ at this loop order. Our results for these quantities agree with those in [10]. Through the equations

$$\gamma_{ij}(n) = - \int_0^1 dz z^{n-1} P_{ij}(z), \tag{6.8}$$

$$\gamma_{\text{ns}}^{\pm,s} = - \int_0^1 dz z^{n-1} P_{\text{ns}}^{\pm,s}(z), \tag{6.9}$$

the physical anomalous dimensions are related to the splitting functions in momentum fraction z space. Using `HarmonicSums`, the splitting functions $P_{ij}(z)$ ($P_{\text{ns}}^{\pm, s}(z)$) are obtained from γ_{ij} ($\gamma_{\text{ns}}^{\pm, s}$) by an inverse Mellin transformation. Also for the splitting functions, we find complete agreement with the results in [10, 11].

7 Summary and conclusions

The operator-product expansion (OPE) allows to systematically separate short-distance and long-distance contributions to hadron-induced processes in QCD. The anomalous dimensions of the resulting quark and gluon operators determine the scale evolution of parton distributions. More specifically, these anomalous dimensions are directly related to the Altarelli-Parisi splitting functions by a Mellin transformation.

The quantization of QCD in a covariant gauge enlarges the particle spectrum by ghosts and induces an a priori infinite number of gauge-variant (GV) operators, which contribute to the renormalization of the physical quark singlet and gluon operators (2.21). Only a finite number of these GV operators contribute to the renormalization of the quark and gluon operators for a fixed Mellin moment and at a given loop order. The fully general structure of the GV operators is only poorly understood at present, thus preventing the calculation of anomalous dimensions beyond the two-loop order in the OPE up to now.

In this setup, the anomalous dimensions of the quark and gluon operators are determined from the divergences of the simplest operator matrix elements (OMEs) with two external, off-shell partons. Compared to alternative methods to determine anomalous dimensions or splitting functions, the calculation of off-shell OMEs is generally considered to involve lower computational complexity at the level of amplitudes and Feynman integrals. It is therefore highly desirable to extend the applicability of the OPE to higher loop orders by developing a consistent renormalization framework for the quark and gluon operators.

In this paper, we developed a new approach to systematically determine the counterterm Feynman rules that result from the GV operators. These counterterm Feynman rules are sufficient to determine the renormalization of operator matrix elements (OMEs) for a fixed number of external partons at a given loop order, even without full knowledge of the GV operators themselves. As an example, the first line of table 1 summarizes which Feynman rules for operators or counterterms are required for the computation of the three-loop, two-parton OMEs and their renormalization. With increasing loop order, OMEs with an increasing number of additional gluons need to be considered, corresponding to vertices with increasing multiplicity entering the bare matrix elements.

Our approach is based on the extraction of the counterterm Feynman rules from the divergent parts of multi-leg OMEs in general off-shell kinematics. The renormalization of these OMEs for two partons and an arbitrary number of gluons is illustrated at two-loop order in (4.11), (4.14), (4.15). The central observation is that the computation of the divergent parts of the OME for a given external state at a given loop order can be used to infer the counterterm Feynman rules for this external state at this loop order.

A special role is played by the GV operators that contribute to the renormalization of the quark and gluon operators already at one loop. At this order, only three operators

contribute, and they are highly constrained in their structure. We collectively denote them by O_{ABC} , and their contribution to the one-loop renormalization of the physical operators is controlled by a single renormalization constant (4.4). It is thus possible to extract the operator Feynman rules for O_{ABC} for a fixed multiplicity of external partons. These turn into the respective counterterm Feynman rules only upon multiplication with the operator renormalization constant. The O_{ABC} operator Feynman rules had been computed previously [14, 78, 79] for external states with up to three partons. We extend their determination now to four partons in (5.19), (5.22), (5.25). The O_{ABC} operators are particularly special, since they already exhaust all allowed Lorentz structures in GV operators at the lowest multiplicities (two-gluon, two-ghost and two-quark-plus-gluon), as we demonstrate in appendix B. Consequently, the counterterm Feynman rules for the lowest multiplicities must be proportional to the O_{ABC} operator Feynman rules, multiplied with a higher-loop correction to the operator renormalization constant (5.27), which can be determined from a low-multiplicity OME calculation.

At two loops, an infinite number of GV operators besides O_{ABC} contribute to the operator renormalization. These other GV operators yield, however, only counterterm Feynman rules starting at higher multiplicities than O_{ABC} . By computing the respective three-parton OMEs in general kinematics, we explicitly determined the two-loop counterterm Feynman rules relevant to the renormalization of the gluon operator $[ZO]_g^{\text{GV},(2)}$ for the ghost-ghost-gluon (5.40) and three-gluon (5.41) operator vertices. From the structural properties of these counterterm Feynman rules (their dependence on n and on the external kinematics), we can infer that they can not originate from a single operator, but only from a linear combination of an unknown number of operators.

The computations of multi-leg OMEs are performed for fixed numerical values of the external kinematics. A finite number of these numerical samples is then sufficient to reconstruct the full kinematical dependence of the resulting counterterm Feynman rules, exploiting their symmetries and their dimensional scaling. The full n -dependence of the OMEs is usually retained through the introduction of resummed propagators (3.1). The two-loop counterterm Feynman rules can not be cast into this resummed form. Instead, OMEs with these counterterm insertions are evaluated repeatedly for multiple integer values of n (based on the all- n Feynman rules as well as a generalized resummed form (6.3), such that the computational effort does not increase with n), allowing subsequently their all- n reconstruction.

We applied our newly computed GV operator and counterterm Feynman rules to rederive the three-loop anomalous dimensions of the unpolarized quark singlet and gluon operators in a general covariant gauge using the OPE method. These three-loop anomalous dimensions (or the corresponding splitting functions) were previously computed with several other approaches [11, 23–29], and we find full agreement with the literature. Our result establishes the independence of the anomalous dimensions on the gauge parameter, which was expected and supported by calculations at fixed n , but not proven up to now.

A method for the systematic construction of GV operators and counterterms for the renormalization of the gluon operator based on BRST symmetry has recently been outlined in Reference [30]. The resulting constraint equations were, however, solved only for fixed

low values of n , and they show a substantial growth in complexity with increasing n . Our method enables the determination of the counterterm Feynman rules for symbolic n , thereby allowing the computation of counterterm OMEs for multiple values of n without an increase in complexity towards larger n . Our results reproduce the two-loop counterterm Feynman rules for low values of n that were obtained in [30].

Owing to the relative computational simplicity of two-parton OMEs at high loop orders, the OPE method holds the potential for computing the four-loop corrections to anomalous dimensions or splitting functions, which are of paramount importance to precision collider physics. While first results on the quark non-singlet [85] and most recently on low- n moments of the quark singlet and gluon anomalous dimensions [30, 46] were obtained, their all- n calculation still remains an outstanding challenge. The developments made in this paper lay out a strategy to determine the renormalization counterterms that are required in this context, thereby paving the way for a future derivation of the four-loop anomalous dimensions in the OPE method.

Acknowledgments

We thank Vasily Sotnikov for useful discussions and Kay Schönwald for helpful comments on the manuscript. We would like to thank the European Research Council (ERC) for funding of this work under the European Union’s Horizon 2020 research and innovation programme grant agreement 101019620 (ERC Advanced Grant TOPUP) and the National Science Foundation (NSF) for support under grant number 2013859.

A Feynman rules for physical operators

As outlined in subsection 2.2, the Feynman rules for physical operators can be obtained from (2.8) and (2.9) by a functional variation. Equivalently, one can read out the Feynman rules directly from the definitions of the operators by replacing a derivative of a field with $-i$ times the associated momentum. As an example, we derive the Feynman rule for one of the terms in O_g ,


$$\begin{aligned}
 & -\frac{i^{n-2}}{2} \mathcal{S} \left[\left(\Delta \cdot \partial A_\mu^{a_1} \right) (\Delta \cdot \partial)^{n-1} A^{\mu, a_2} \delta^{a_1 a_2} \right] \\
 & \rightarrow -\frac{i^{n-2}}{2} (-i)^n \delta^{a_1 a_2} (\Delta \cdot p_2)^{n-1} \Delta \cdot p_1 g^{\mu_1 \mu_2} + \{p_1 \leftrightarrow p_2, \mu_1 \leftrightarrow \mu_2, a_1 \leftrightarrow a_2\} \\
 & = -\delta^{a_1 a_2} (\Delta \cdot p_1)^n g^{\mu_1 \mu_2}, \tag{A.1}
 \end{aligned}$$

where in the last line we use momentum conservation $p_2 = -p_1$ to simplify the result. In the following, the Feynman rules for the physical operators O_q and O_g and up to 5 legs are listed for completeness, with the convention of all momenta flowing into the vertices.


A.1 Feynman rules for the operator O_q




$$\rightarrow \frac{1}{2} \not{\Delta} \delta_{i_1 i_2} (\Delta \cdot p_1)^{n-1}, \quad (\text{A.2})$$



$$\rightarrow \frac{1}{2} g_s \Delta^{\mu_3} T_{i_2 i_1}^{a_3} \not{\Delta} \sum_{j_1=0}^{n-2} \left((-\Delta \cdot p_2)^{j_1} (\Delta \cdot p_1)^{-j_1+n-2} \right), \quad (\text{A.3})$$



$$\rightarrow \frac{1}{2} g_s^2 \Delta^{\mu_3} \Delta^{\mu_4} \not{\Delta} \left\{ \sum_{j_1=0}^{n-3} \sum_{j_2=0}^{j_1} (-\Delta \cdot p_2)^{j_1-j_2} (\Delta \cdot p_1)^{-j_1+n-3} \right. \\ \left. \times \left[(T^{a_3} T^{a_4})_{i_2 i_1} (\Delta \cdot (p_1 + p_4))^{j_2} + (T^{a_4} T^{a_3})_{i_2 i_1} (\Delta \cdot (p_1 + p_3))^{j_2} \right] \right\}, \quad (\text{A.4})$$




$$\rightarrow \frac{1}{2} g_s^3 \Delta^{\mu_3} \Delta^{\mu_4} \Delta^{\mu_5} \not{\Delta} \left\{ \sum_{j_1=0}^{n-4} \sum_{j_2=0}^{j_1} \sum_{j_3=0}^{j_2} (\Delta \cdot p_1)^{-j_1+n-4} (-\Delta \cdot p_2)^{j_1-j_2} \right. \\ \times \left[(T^{a_3} T^{a_4} T^{a_5})_{i_2 i_1} (\Delta \cdot (-p_2 - p_3))^{j_2-j_3} (\Delta \cdot (p_1 + p_5))^{j_3} \right. \\ + (T^{a_3} T^{a_5} T^{a_4})_{i_2 i_1} (\Delta \cdot (-p_2 - p_3))^{j_2-j_3} (\Delta \cdot (p_1 + p_4))^{j_3} \\ + (T^{a_4} T^{a_3} T^{a_5})_{i_2 i_1} (\Delta \cdot (-p_2 - p_4))^{j_2-j_3} (\Delta \cdot (p_1 + p_5))^{j_3} \\ + (T^{a_4} T^{a_5} T^{a_3})_{i_2 i_1} (\Delta \cdot (-p_2 - p_4))^{j_2-j_3} (\Delta \cdot (p_1 + p_3))^{j_3} \\ + (T^{a_5} T^{a_3} T^{a_4})_{i_2 i_1} (\Delta \cdot (-p_2 - p_5))^{j_2-j_3} (\Delta \cdot (p_1 + p_4))^{j_3} \\ \left. \left. + (T^{a_5} T^{a_4} T^{a_3})_{i_2 i_1} (\Delta \cdot (-p_2 - p_5))^{j_2-j_3} (\Delta \cdot (p_1 + p_3))^{j_3} \right] \right\}. \quad (\text{A.5})$$


A.2 Feynman rules for the operator O_g




$$\rightarrow -\delta^{a_1 a_2} (\Delta \cdot p_1)^{n-2} \left[(\Delta \cdot p_1)^2 g^{\mu_1 \mu_2} - \Delta \cdot p_1 (p_1^{\mu_1} \Delta^{\mu_2} + \Delta^{\mu_1} p_1^{\mu_2}) + \Delta^{\mu_1} \Delta^{\mu_2} p_1 \cdot p_1 \right], \quad (\text{A.6})$$



$$\begin{aligned} \rightarrow & \frac{-i}{4} g_s f^{a_1 a_2 a_3} \left\{ (\Delta \cdot p_1)^{n-2} \left(-\Delta^{\mu_2} g^{\mu_1 \mu_3} \Delta \cdot p_1 \right. \right. \\ & + \Delta^{\mu_3} g^{\mu_1 \mu_2} \Delta \cdot p_1 - \Delta^{\mu_1} \Delta^{\mu_3} p_1^{\mu_2} + \Delta^{\mu_1} \Delta^{\mu_2} p_1^{\mu_3} \Big) \\ & + (-\Delta \cdot p_3)^{n-2} \left(-\Delta^{\mu_1} g^{\mu_2 \mu_3} \Delta \cdot p_3 + \Delta^{\mu_2} g^{\mu_1 \mu_3} \Delta \cdot p_3 - \Delta^{\mu_2} \Delta^{\mu_3} p_3^{\mu_1} + \Delta^{\mu_1} \Delta^{\mu_3} p_3^{\mu_2} \right) \\ & - 2\Delta^{\mu_2} \left(-g^{\mu_1 \mu_3} \Delta \cdot p_1 \Delta \cdot p_3 + \Delta^{\mu_3} p_3^{\mu_1} \Delta \cdot p_1 + \Delta^{\mu_1} p_3^{\mu_3} \Delta \cdot p_3 - \Delta^{\mu_1} \Delta^{\mu_3} p_1 \cdot p_3 \right) \\ & \times \sum_{j_1=0}^{n-3} \left((\Delta \cdot (p_1 + p_2))^{j_1} (\Delta \cdot p_1)^{-j_1+n-3} \right) \Big\} + \text{permutations}, \quad (\text{A.7}) \end{aligned}$$



$$\begin{aligned} \rightarrow & \frac{1}{8} g_s^2 f^{a a_1 a_2} f^{a a_3 a_4} \left\{ 2\Delta^{\mu_3} \left(\Delta^{\mu_1} g^{\mu_2 \mu_4} \Delta \cdot p_4 - \Delta^{\mu_2} g^{\mu_1 \mu_4} \Delta \cdot p_4 + \Delta^{\mu_2} \Delta^{\mu_4} p_4^{\mu_1} \right. \right. \\ & - \Delta^{\mu_1} \Delta^{\mu_4} p_4^{\mu_2} \Big) \sum_{j_1=0}^{n-3} \left((\Delta \cdot (p_1 + p_2 + p_3))^{j_1} (\Delta \cdot (p_1 + p_2))^{-j_1+n-3} \right) \\ & + 4\Delta^{\mu_2} \Delta^{\mu_3} \left(-g^{\mu_1 \mu_4} \Delta \cdot p_1 \Delta \cdot p_4 + \Delta^{\mu_4} p_4^{\mu_1} \Delta \cdot p_1 + \Delta^{\mu_1} p_1^{\mu_4} \Delta \cdot p_4 - \Delta^{\mu_1} \Delta^{\mu_4} p_1 \cdot p_4 \right) \\ & \times \sum_{j_1=0}^{n-4} \sum_{j_2=0}^{j_1} \left((\Delta \cdot (p_1 + p_2))^{j_1-j_2} (\Delta \cdot (p_1 + p_2 + p_3))^{j_2} (\Delta \cdot p_1)^{-j_1+n-4} \right) \\ & + \left(-\Delta^{\mu_2} \Delta^{\mu_4} g^{\mu_1 \mu_3} + \Delta^{\mu_1} \Delta^{\mu_4} g^{\mu_2 \mu_3} \right. \\ & + \Delta^{\mu_2} \Delta^{\mu_3} g^{\mu_1 \mu_4} - \Delta^{\mu_1} \Delta^{\mu_3} g^{\mu_2 \mu_4} \Big) (\Delta \cdot (p_1 + p_2))^{n-2} \\ & + 2\Delta^{\mu_2} \left(\Delta^{\mu_3} g^{\mu_1 \mu_4} \Delta \cdot p_1 - \Delta^{\mu_4} g^{\mu_1 \mu_3} \Delta \cdot p_1 + \Delta^{\mu_1} \Delta^{\mu_4} p_1^{\mu_3} - \Delta^{\mu_1} \Delta^{\mu_3} p_1^{\mu_4} \right) \\ & \times \sum_{j_1=0}^{n-3} \left((\Delta \cdot (p_1 + p_2))^{j_1} (\Delta \cdot p_1)^{-j_1+n-3} \right) \Big\} + \text{permutations}, \quad (\text{A.8}) \end{aligned}$$



$$\begin{aligned}
 &\rightarrow \frac{-i}{8} g_s^3 f^{aa_1 a_2} f^{aba_3} f^{ba_4 a_5} \left\{ -2\Delta^{\mu_2} \Delta^{\mu_3} \left(\Delta^{\mu_4} g^{\mu_1 \mu_5} \Delta \cdot p_1 - \Delta^{\mu_5} g^{\mu_1 \mu_4} \Delta \cdot p_1 + \Delta^{\mu_1} \Delta^{\mu_5} p_1^{\mu_4} \right. \right. \\
 &\quad - \Delta^{\mu_1} \Delta^{\mu_4} p_1^{\mu_5} \left. \right) \sum_{j_1=0}^{n-4} \sum_{j_2=0}^{j_1} \left((\Delta \cdot (p_1 + p_2))^{j_1 - j_2} (\Delta \cdot (p_1 + p_2 + p_3))^{j_2} (\Delta \cdot p_1)^{-j_1 + n - 4} \right) \\
 &\quad - 2\Delta^{\mu_3} \Delta^{\mu_4} \left(\Delta^{\mu_1} g^{\mu_2 \mu_5} \Delta \cdot p_5 - \Delta^{\mu_2} g^{\mu_1 \mu_5} \Delta \cdot p_5 + \Delta^{\mu_2} \Delta^{\mu_5} p_5^{\mu_1} - \Delta^{\mu_1} \Delta^{\mu_5} p_5^{\mu_2} \right) \\
 &\quad \times \sum_{j_1=0}^{n-4} \sum_{j_2=0}^{j_1} \left((\Delta \cdot (p_1 + p_2 + p_3))^{j_1 - j_2} (-\Delta \cdot p_5)^{j_2} (\Delta \cdot (p_1 + p_2))^{-j_1 + n - 4} \right) \\
 &\quad - 4\Delta^{\mu_2} \Delta^{\mu_3} \Delta^{\mu_4} \left(-g^{\mu_1 \mu_5} \Delta \cdot p_1 \Delta \cdot p_5 + \Delta^{\mu_5} p_5^{\mu_1} \Delta \cdot p_1 + \Delta^{\mu_1} p_1^{\mu_5} \Delta \cdot p_5 - \Delta^{\mu_1} \Delta^{\mu_5} p_1 \cdot p_5 \right) \\
 &\quad \times \sum_{j_1=0}^{n-5} \sum_{j_2=0}^{j_1} \sum_{j_3=0}^{j_2} \left[(\Delta \cdot (p_1 + p_2))^{j_1 - j_2} \right. \\
 &\quad \times (\Delta \cdot (p_1 + p_2 + p_3))^{j_2 - j_3} (-\Delta \cdot p_5)^{j_3} (\Delta \cdot p_1)^{-j_1 + n - 5} \left. \right] \\
 &\quad + \Delta^{\mu_3} \left(\Delta^{\mu_2} \Delta^{\mu_5} g^{\mu_1 \mu_4} - \Delta^{\mu_1} \Delta^{\mu_5} g^{\mu_2 \mu_4} - \Delta^{\mu_2} \Delta^{\mu_4} g^{\mu_1 \mu_5} + \Delta^{\mu_1} \Delta^{\mu_4} g^{\mu_2 \mu_5} \right) \\
 &\quad \times \sum_{j_1=0}^{n-3} \left((\Delta \cdot (p_1 + p_2 + p_3))^{j_1} (\Delta \cdot (p_1 + p_2))^{-j_1 + n - 3} \right) \left. \right\} + \text{permutations}, \quad (\text{A.9})
 \end{aligned}$$

where plus *permutations* indicates the summation over all the external gluon indices (simultaneous permutation of μ_i, a_i, p_i).

B Feynman rules for all GV operators at lowest multiplicity

As discussed at the end of subsection 4.2, the two-point vertex Feynman rules for all GV operators except O_A and O_C are zero. We prove this statement in the following. For the vertex Feynman rules with two quark or two ghost legs, the only possible forms are:

$$\langle q|O|q \rangle^{(0),(0)} = h_1 \not{\Delta} (\Delta \cdot p)^{n-1}, \quad (\text{B.1})$$

$$\langle c|O|c \rangle^{(0),(0)} = h_2 (\Delta \cdot p)^n, \quad (\text{B.2})$$

where O is an arbitrary twist-two operator, and h_1 and h_2 are constants. We recognize that the above two equations are exactly the Feynman rules for O_q and O_C respectively, as in (A.2) and (5.20). Therefore, the vertex Feynman rules with two quarks for all GV operators are zero, and the vertex Feynman rules with two ghosts for all GV operators except O_C are zero.

For the case of the vertex Feynman rules with two gluons, there are four possible tensor structures $T_i^{\mu\nu}$ with $i = 1 \dots 4$ as shown in (2.36). From subsection 5.1, $T_2^{\mu\nu}$ can not be the

Feynman rule of a twist-two operator. Moreover, the Feynman rules with two gluons are expected to satisfy the condition of being transverse, which is not the case for $T_4^{\mu\nu}$, i.e.,

$$p_\mu p_\nu T_4^{\mu\nu} \neq 0. \quad (\text{B.3})$$

The remaining tensor structures $T_1^{\mu\nu}$ and $T_3^{\mu\nu}$ are just the Feynman rules of O_g and O_A separately, as shown in (A.6) and (5.24). Therefore, all GV operators involving only two gluon fields except O_A are zero.

Based on the symmetry from the exchange of two quarks for a singlet twist-two operator, and from the Feynman rules in (A.3), (5.18) and (5.39), we conjecture but not prove the statement: the $q\bar{q}g$ vertex Feynman rules for all GV operators except O_B are zero.

C Renormalization constants Z_{qA} and Z_{gA} to three loops

We give the all- n results for the renormalization constants Z_{qA} and Z_{gA} to three loops, which can be extracted from the OMEs with two-ghost external states. Z_{qA} starts at a_s^2 and reads as follows,

$$\begin{aligned} Z_{qA} = & \frac{a_s^2 C_A N_f}{(n-1)n^2(n+1)(n+2)} \left\{ \frac{(n^2+n+2)}{\epsilon^2} - \frac{(n^2+5n+2)(n^3+n^2+n+2)}{n(n+1)(n+2)\epsilon} \right\} \\ & + \frac{a_s^3 N_f}{(n-1)n^2(n+1)(n+2)} \left\{ \frac{1}{\epsilon^3} \left[\frac{8}{9} (n^2+n+2) C_A N_f + C_A C_F \left(\frac{4}{3} (n^2+n+2) S_1(n) \right. \right. \right. \\ & - \left. \left. \frac{(n^2+n+2)(3n^2+3n+2)}{3n(n+1)} \right) + C_A^2 \left(\frac{11}{6} (n^2+n+2) S_1(n) \right. \right. \\ & \left. \left. - \frac{(n^2+n+2)(100n^4+203n^3-52n^2-173n+30)}{18(n-1)n(n+1)(n+2)} \right) \right] \\ & + \frac{1}{\epsilon^2} \left[C_A C_F \left(\frac{8}{3} (n^2+n+2) S_{1,1}(n) - \frac{8}{3} (n^2+n+2) S_2(n) \right. \right. \\ & \left. \left. - \frac{4(n^5+8n^4+20n^3+35n^2+36n+12) S_1(n)}{3n(n+1)(n+2)} \right. \right. \\ & \left. \left. + \frac{13n^7+71n^6+176n^5+309n^4+333n^3+182n^2+76n+24}{3n^2(n+1)^2(n+2)} \right) \right. \\ & \left. + C_A^2 \left(-\frac{11}{6} (n^2+n+2) S_{1,1}(n) - \frac{10}{3} (n^2+n+2) S_{-2}(n) - 2(n^2+n+2) S_2(n) \right. \right. \\ & \left. \left. + (1-\xi) \left(\frac{1}{12} (n^2+n+2) S_1(n) - \frac{(n^2+n+2)(2n^3+n^2-1)}{12(n-1)n(n+1)} \right) \right. \right. \\ & \left. \left. - \frac{(53n^6+219n^5+33n^4+77n^3+310n^2-284n-120) S_1(n)}{18(n-1)n(n+1)(n+2)} \right. \right. \\ & \left. \left. + \frac{1}{54(n-1)^2 n^2 (n+1)^2 (n+2)^2} \left(577n^{10} + 3302n^9 + 5318n^8 + 1049n^7 \right. \right. \right. \\ & \left. \left. - 4127n^6 - 3430n^5 - 3270n^4 + 291n^3 + 4742n^2 - 780n - 1080 \right) \right) \\ & \left. + C_A N_f \left(\frac{4}{9} (n^2+n+2) S_1(n) - \frac{4(11n^5+50n^4+80n^3+91n^2+68n+12)}{27n(n+1)(n+2)} \right) \right\} \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{\epsilon} \left[C_A^2 \left(\frac{1}{162(n-1)^3 n^3 (n+1)^3 (n+2)^3} \left(353n^{14} - 943n^{13} - 5075n^{12} + 1993n^{11} \right. \right. \right. \\
 & - 6788n^{10} - 54586n^9 - 38153n^8 + 57445n^7 + 61625n^6 - 49753n^5 - 42370n^4 \\
 & + 54772n^3 + 8808n^2 - 25344n - 8640 \Big) + \frac{2(9n^3 + 19n^2 + 24n + 12) S_{-3}(n)}{3(n+1)} \\
 & + \frac{2(6n^6 + 47n^5 + 105n^4 + 129n^3 + 137n^2 + 96n + 20) S_{-2}(n)}{3n(n+1)^2(n+2)} \\
 & - \frac{1}{54(n-1)n^2(n+1)^2(n+2)^2} \left(28n^9 - 144n^8 - 172n^7 + 1210n^6 \right. \\
 & - 379n^5 - 6950n^4 - 6539n^3 - 290n^2 + 1068n + 72 \Big) S_1(n) \\
 & + \frac{(5n^5 + 37n^4 + 18n^3 - 14n^2 + 40n + 8) S_2(n)}{3n(n+1)(n+2)} + \frac{(11n^3 + 20n^2 + 35n + 30) S_3(n)}{3(n+1)} \\
 & - \frac{8}{3} n(n+1) S_{-2,1}(n) + \frac{4(n^3 + n^2 + 2n + 4) S_{1,-2}(n)}{3(n+1)} \\
 & + \frac{(53n^6 + 219n^5 + 321n^4 + 221n^3 - 122n^2 - 284n - 120) S_{1,1}(n)}{18(n-1)n(n+1)(n+2)} \\
 & + (1-\xi) \left(- \frac{3n^9 - 7n^7 - 34n^6 - 50n^5 + 3n^4 + 26n^3 + 31n^2 - 8n - 12}{12(n-1)^2 n^2 (n+1)^2 (n+2)} \right. \\
 & + \frac{(n^5 + 5n^3 + 6n^2 - 12n - 8) S_1(n)}{12n(n+1)(n+2)} + \frac{1}{12} (-n^2 - n - 2) S_{1,1}(n) \\
 & + \frac{(11n^3 + 21n^2 + 32n + 24) S_{1,2}(n)}{3(n+1)} + \frac{n(n^2 + 3n + 4) S_{2,1}(n)}{3(n+1)} \\
 & - \frac{13}{6} (n^2 + n + 2) S_{1,1,1}(n) - 6(n^2 + n + 2) \zeta_3 \Big) \\
 & + N_f C_A \left(- \frac{4(13n^8 + 13n^7 + 45n^6 + 223n^5 - 51n^4 - 833n^3 - 718n^2 - 60n + 72)}{81n^2(n+1)^2(n+2)^2} \right. \\
 & + \frac{8(n-2)(n^4 + 4n^2 + 13n + 6) S_1(n)}{27n(n+1)(n+2)} - \frac{4}{9} (n^2 + n + 2) S_{1,1}(n) \Big) \\
 & + C_F C_A \left(- \frac{2}{3n^3(n+1)^3(n+2)^2} \left(11n^{10} + 101n^9 + 384n^8 + 875n^7 + 1523n^6 \right. \right. \\
 & + 2068n^5 + 1886n^4 + 1072n^3 + 448n^2 + 160n + 32 \Big) \\
 & - \frac{2(n^8 - 47n^6 - 202n^5 - 466n^4 - 726n^3 - 708n^2 - 360n - 80) S_1(n)}{3n^2(n+1)^2(n+2)^2} \\
 & + \frac{8(n^5 + 6n^4 + 11n^3 + 20n^2 + 24n + 8) S_2(n)}{3n(n+1)(n+2)} - \frac{4}{3} (n^2 + n + 2) S_3(n) \\
 & - \frac{4(2n^5 + 13n^4 + 22n^3 + 31n^2 + 36n + 12) S_{1,1}(n)}{3n(n+1)(n+2)} - \frac{8}{3} (n^2 + n + 2) S_{1,2}(n) \\
 & \left. + \frac{4}{3} (n^2 + n + 2) S_{1,1,1}(n) + 8(n^2 + n + 2) \zeta_3 \right) \Big] + \mathcal{O}(a_s^4). \tag{C.1}
 \end{aligned}$$

Z_{gA} starts at a_s^1 and we gave the first order $Z_{gA}^{(1)}$ and the second order $Z_{gA}^{(2)}$ in (4.4) and (5.27), respectively. The third order contribution to Z_{gA} is obtained as

$$\begin{aligned}
 Z_{gA}^{(3)} = & \frac{1}{\epsilon^3} \frac{1}{n(n-1)} \left[C_A^3 \left(-\frac{391}{48} S_{1,1}(n) + \frac{(2636n^4 + 5407n^3 - 698n^2 - 4051n + 1242) S_1(n)}{144(n-1)n(n+1)(n+2)} \right. \right. \\
 & - \frac{1}{144(n-1)^2 n^2 (n+1)^2 (n+2)^2} (2736n^8 + 11192n^7 + 9977n^6 \\
 & - 11572n^5 - 14586n^4 + 8032n^3 + 6901n^2 - 3020n + 1140) \\
 & + (1 - \xi) \left(\frac{S_1(n)}{96} - \frac{(n+1)(2n-3)}{96(n-1)n} \right) + \frac{1}{24} S_{-2}(n) + \frac{191 S_2(n)}{48} \Big) \\
 & + N_f C_A^2 \left(\frac{2(26n^4 + 53n^3 - 6n^2 - 39n + 14)}{9(n-1)n(n+1)(n+2)} - \frac{26 S_1(n)}{9} \right) \\
 & - \left. \frac{2(n^2 + n + 2)^2 C_A C_F N_f}{3(n-1)n^2(n+1)^2(n+2)} - \frac{4}{9} C_A N_f^2 \right] \\
 & + \frac{1}{\epsilon^2} \frac{1}{n(n-1)} \left[C_A^3 \left((1 - \xi)^2 \left(\frac{9n^3 + 6n^2 + n - 4}{192(n-1)n(n+1)} - \frac{S_1(n)}{32} \right) \right. \right. \\
 & + (1 - \xi) \left(-\frac{356n^7 + 863n^6 + 45n^5 - 912n^4 - 130n^3 + 619n^2 + 401n - 90}{288(n-1)^2 n^2 (n+1)^2 (n+2)} \right. \\
 & - \frac{1}{12} S_{-2}(n) + \frac{(343n^4 + 764n^3 + 59n^2 - 374n - 72) S_1(n)}{288(n-1)n(n+1)(n+2)} + \frac{5 S_2(n)}{16} - \frac{55}{96} S_{1,1}(n) \Big) \\
 & + \frac{1}{432(n-1)^3 n^3 (n+1)^3 (n+2)^3} (21598n^{12} + 130274n^{11} + 217650n^{10} - 97011n^9 \\
 & - 545042n^8 - 168389n^7 + 540328n^6 + 320207n^5 - 220384n^4 - 157997n^3 + 51298n^2 \\
 & - 9804n - 20520) + \frac{1}{4} (-n - 20) S_{-3}(n) - \frac{(63n^4 + 133n^3 + 102n^2 + 128) S_{-2}(n)}{12(n-1)n(n+1)(n+2)} \\
 & - \frac{1}{216(n-1)^2 n^2 (n+1)^2 (n+2)^2} (9787n^8 + 41377n^7 + 30596n^6 - 55169n^5 \\
 & - 61462n^4 + 36244n^3 + 57835n^2 - 852n - 6516) S_1(n) \\
 & - \frac{(385n^4 + 872n^3 + 158n^2 - 527n + 138) S_2(n)}{18(n-1)n(n+1)(n+2)} + \frac{1}{48} (8n - 477) S_3(n) \\
 & + \frac{1}{6} (5n + 14) S_{-2,1}(n) + \frac{47}{6} S_{1,-2}(n) \\
 & + \frac{(2818n^4 + 7001n^3 + 932n^2 - 4901n - 1170) S_{1,1}(n)}{144(n-1)n(n+1)(n+2)} \\
 & + \frac{187}{12} S_{1,2}(n) + \frac{1}{24} (2n + 377) S_{2,1}(n) - \frac{137}{16} S_{1,1,1}(n) \Big) + \left(\frac{32}{27} - \frac{4 S_1(n)}{9} \right) N_f^2 C_A \\
 & + \left(-\frac{1}{108(n-1)^2 n^2 (n+1)^2 (n+2)^2} (1791n^8 + 7217n^7 + 4685n^6 - 11071n^5 \right. \\
 & - 11004n^4 + 6286n^3 + 6040n^2 - 1880n - 768) \\
 & \left. + \frac{2}{3} S_{-2}(n) + (1 - \xi) \left(\frac{2n^3 + n^2 - 1}{18(n-1)n(n+1)} - \frac{S_1(n)}{18} \right) \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{(1205n^4 + 2575n^3 - 668n^2 - 2344n - 120) S_1(n) + \frac{61S_2(n)}{18} - \frac{71}{18} S_{1,1}(n)}{108(n-1)n(n+1)(n+2)} \Big) C_A^2 N_f \\
 & - \frac{2(n^2 + n + 2)(2n^7 + 10n^6 + 13n^5 - 12n^4 - 30n^3 - 15n^2 - 20n - 12) C_A C_F N_f}{3(n-1)n^3(n+1)^3(n+2)^2} \Big] \\
 & + \frac{1}{\epsilon} \frac{1}{n(n-1)} \left[C_A^3 \left(\left\{ - \frac{30n^6 - n^5 - 27n^4 - 31n^3 + 9n^2 + 20n - 8}{192(n-1)^2 n^2 (n+1)^2} \right. \right. \right. \\
 & + \left. \left. \frac{(15n^3 + 6n^2 - 7n - 6) S_1(n)}{192(n-1)n(n+1)} - \frac{S_2(n)}{192} - \frac{5}{192} S_{1,1}(n) \right\} (1-\xi)^2 \right. \\
 & + (1-\xi) \left\{ \frac{1}{864(n-2)(n-1)^3 n^3 (n+1)^3 (n+2)^2} (1589n^{12} + 4922n^{11} - 6625n^{10} \right. \\
 & - 26755n^9 - 7874n^8 + 40335n^7 + 31081n^6 - 35477n^5 - 16025n^4 + 15763n^3 \\
 & + 14n^2 - 8508n - 1512) - \frac{(39n^2 + 39n + 16) S_{-3}(n)}{96n(n+1)} \\
 & - \frac{(35n^6 - 11n^5 - 85n^4 - 61n^3 - 14n^2 + 8n + 32) S_{-2}(n)}{48(n-2)(n-1)n^2(n+1)^2} \\
 & - \frac{1}{864(n-1)^2 n^3 (n+1)^2 (n+2)^2} (1180n^9 + 6049n^8 + 8192n^7 + 1843n^6 - 730n^5 \\
 & - 2948n^4 - 5456n^3 + 4866n^2 + 4392n - 2808) S_1(n) \\
 & - \frac{(281n^4 + 345n^3 - 164n^2 + 306n - 234) S_2(n)}{288(n-1)n^2(n+1)} - \frac{(17n^2 + 17n - 8) S_3(n)}{48n(n+1)} \\
 & + \frac{1}{12} S_{-2,1}(n) + \frac{(21n^2 + 21n + 16) S_{1,-2}(n)}{48n(n+1)} \\
 & + \frac{(60n^4 + 201n^3 + 135n^2 - 82n - 152) S_{1,1}(n)}{96(n-1)n(n+1)(n+2)} \\
 & + \frac{(35n^3 + 71n^2 + 23n + 39) S_{1,2}(n)}{48n(n+1)^2} + \frac{(67n^3 + 132n^2 + 91n - 78) S_{2,1}(n)}{96n(n+1)^2} \\
 & - \left. \frac{35}{96} S_{1,1,1}(n) + \frac{(n^3 + 4n^2 - 31n + 70) \zeta_3}{16n(n+1)^2} \right\} \\
 & - \frac{1}{1296(n-2)(n-1)^4 n^4 (n+1)^4 (n+2)^4} (57083n^{17} + 341102n^{16} + 177962n^{15} \\
 & - 2219135n^{14} - 3448638n^{13} + 3989078n^{12} + 11054452n^{11} - 1139746n^{10} \\
 & - 17966031n^9 - 7357576n^8 + 14022462n^7 + 10545037n^6 \\
 & - 4729434n^5 - 5907736n^4 + 1571152n^3 + 1706640n^2 - 477216n - 359424) \\
 & + \frac{(12n^6 + 869n^5 + 3303n^4 + 6881n^3 + 5709n^2 + 1154n - 504) S_{-3}(n)}{144(n-1)n(n+1)^2(n+2)} \\
 & + \frac{1}{36(n-2)(n-1)^2 n^2 (n+1)^3 (n+2)^2} (629n^{10} + 1839n^9 - 2772n^8 - 9854n^7 \\
 & - 1227n^6 + 12939n^5 + 8026n^4 - 7548n^3 - 7536n^2 + 2336n + 1440) S_{-2}(n) \\
 & + \frac{1}{648(n-1)^3 n^3 (n+1)^3 (n+2)^3} (17788n^{12} + 127947n^{11} + 281430n^{10} + 30563n^9
 \end{aligned}$$

$$\begin{aligned}
& -617061n^8 - 517548n^7 + 589151n^6 + 1063392n^5 + 160500n^4 - 539150n^3 \\
& - 192804n^2 + 82008n + 44496) S_1(n) + \frac{1}{16}(6n + 197)S_{-4}(n) \\
& + \frac{1}{432(n-1)^2n^2(n+1)^2(n+2)^2} (19838n^8 + 80657n^7 + 42469n^6 - 130108n^5 \\
& - 109184n^4 + 98303n^3 + 133133n^2 - 1440n - 11196) S_2(n) \\
& + \frac{(197n^5 + 1036n^4 + 2214n^3 + 1526n^2 + 177n + 154) S_3(n)}{48(n-1)n(n+1)^2(n+2)} + \frac{1}{24}(257 - 6n)S_4(n) \\
& + \frac{1}{6}(-8n - 41)S_{-3,1}(n) + \frac{(3n^2 - 10n - 17) S_{-2,-2}(n)}{6(n+1)} \\
& + \frac{(3n^5 - 52n^4 - 185n^3 - 296n^2 - 34n + 276) S_{-2,1}(n)}{18(n-1)n(n+1)(n+2)} + \frac{1}{6}(n - 22)S_{-2,2}(n) \\
& - \frac{(48n^2 + 937n + 913) S_{1,-3}(n)}{48(n+1)} \\
& - \frac{(767n^5 + 2949n^4 + 2771n^3 - 933n^2 - 3274n - 2136) S_{1,-2}(n)}{72(n-1)n(n+1)^2(n+2)} \\
& - \frac{1}{432(n-1)^2n^2(n+1)^2(n+2)^2} (8524n^8 + 38614n^7 + 37691n^6 - 28388n^5 \\
& - 48148n^4 + 12310n^3 + 47149n^2 + 12780n + 468) S_{1,1}(n) \\
& - \frac{(1085n^5 + 4209n^4 + 3803n^3 - 1713n^2 - 3340n - 1236) S_{1,2}(n)}{72(n-1)n(n+1)^2(n+2)} \\
& + \frac{(12n^2 - 325n - 321) S_{1,3}(n)}{24(n+1)} + \frac{(3n^2 - 123n - 130) S_{2,-2}(n)}{12(n+1)} \\
& + \frac{(12n^6 - 2213n^5 - 8175n^4 - 7625n^3 + 1995n^2 + 6238n + 1992) S_{2,1}(n)}{144(n-1)n(n+1)^2(n+2)} \\
& + \frac{(n^2 - 119n - 122) S_{2,2}(n)}{6(n+1)} - \frac{(5n^2 + 347n + 346) S_{3,1}(n)}{24(n+1)} + \frac{1}{3}(5 - n)S_{-2,1,1}(n) \\
& + \frac{2}{3}(3n + 10)S_{1,-2,1}(n) + \frac{(135n + 143)S_{1,1,-2}(n)}{24(n+1)} \\
& + \frac{(989n^4 + 2725n^3 + 373n^2 - 2161n - 1278) S_{1,1,1}(n)}{144(n-1)n(n+1)(n+2)} + \frac{(173n + 165)S_{1,1,2}(n)}{24(n+1)} \\
& + \frac{(389n + 381)S_{1,2,1}(n)}{48(n+1)} - \frac{(4n^2 - 389n - 417) S_{2,1,1}(n)}{48(n+1)} - \frac{7}{3}S_{1,1,1,1}(n) \\
& + \left(\frac{19n^5 + 12n^4 - 27n^3 + 18n^2 - 30n - 4}{4(n-1)n(n+1)^2(n+2)} - \frac{(n+3)S_1(n)}{n+1} \right) \zeta_3 \\
& + \left(\frac{32S_1(n)}{27} - \frac{4}{9}S_{1,1}(n) - \frac{16}{27} \right) N_f^2 C_A \\
& + C_A^2 N_f \left(\frac{1}{648(n-1)^3n^3(n+1)^3(n+2)^3} (8065n^{12} + 48004n^{11} + 65499n^{10} \right. \\
& \left. - 81506n^9 - 224797n^8 - 10924n^7 + 311121n^6 + 187594n^5 - 180704n^4 - 169280n^3 \right.
\end{aligned}$$

$$\begin{aligned}
& + 19376n^2 + 38496n + 4608) - \frac{8}{9}S_{-3}(n) \\
& - \frac{(29n^4 + 58n^3 - 21n^2 - 50n - 64) S_{-2}(n)}{9(n-1)n(n+1)(n+2)} \\
& - \frac{1}{324(n-1)^2n^2(n+1)^2(n+2)^2} (3701n^8 + 16697n^7 + 13876n^6 - 23791n^5 \\
& - 34295n^4 + 3380n^3 + 21920n^2 + 6432n - 144) S_1(n) \\
& - \frac{(236n^3 + 27n^2 - 278n - 9) S_2(n)}{27(n-1)n(n+1)} - \frac{S_3(n)}{2} + \frac{8}{9}S_{-2,1}(n) + \frac{16}{9}S_{1,-2}(n) \\
& + (1 - \xi) \left\{ - \frac{14n^6 + 16n^5 + n^4 - 23n^3 - 6n^2 + 7n + 3}{54(n-1)^2n^2(n+1)^2} \right. \\
& + \frac{(19n^3 + 6n^2 - 7n - 6) S_1(n)}{54(n-1)n(n+1)} \\
& + \left. \frac{S_2(n)}{9} - \frac{1}{6}S_{1,1}(n) \right\} + \frac{(547n^4 + 1445n^3 + 116n^2 - 1172n - 576) S_{1,1}(n)}{108(n-1)n(n+1)(n+2)} \\
& + \frac{(28n^2 + 59n + 25) S_{1,2}(n)}{9(n+1)^2} + \frac{(29n^2 + 55n + 32) S_{2,1}(n)}{9(n+1)^2} - \frac{25}{18}S_{1,1,1}(n) \\
& + \frac{2(2n^2 + 5n + 1) \zeta_3}{(n+1)^2} + C_F C_A N_f \left[\frac{1}{9(n-1)n^4(n+1)^4(n+2)^3} (67n^{12} + 603n^{11} \right. \\
& + 2168n^{10} + 3846n^9 + 3021n^8 - 387n^7 - 3292n^6 - 4758n^5 - 4748n^4 - 3384n^3 \\
& \left. - 2544n^2 - 1536n - 384) - \frac{32S_{-2}(n)}{3(n-1)n(n+1)(n+2)} - \frac{4S_1(n)}{3} - \frac{16\zeta_3}{3} \right]. \quad (C.2)
\end{aligned}$$

In view of applications at fixed n , we also summarize Z_{qA} and Z_{gA} for $n = 2, \dots, 10$,

$$\begin{aligned}
\frac{Z_{qA}|_{n=2}}{C_A} &= a_s^2 N_f \left(\frac{1}{6\epsilon^2} - \frac{2}{9\epsilon} \right) + a_s^3 N_f \left\{ \frac{1}{\epsilon^3} \left(\frac{4N_f}{27} + \frac{4C_F}{27} - \frac{7C_A}{12} \right) \right. \\
& + \frac{1}{\epsilon^2} \left[- \frac{5N_f}{27} + \frac{7C_F}{27} + \left(\frac{61}{72} - \frac{5(1-\xi)}{216} \right) C_A \right] \\
& + \frac{1}{\epsilon} \left[- \frac{67N_f}{486} + \left(\frac{4\zeta_3}{3} - \frac{895}{486} \right) C_F \right. \\
& \left. + C_A \left(\frac{7(1-\xi)}{432} - \zeta_3 + \frac{329}{432} \right) \right] \left. \right\}, \quad (C.3)
\end{aligned}$$

$$\begin{aligned}
\frac{Z_{qA}|_{n=4}}{C_A} &= a_s^2 N_f \left(\frac{11}{720\epsilon^2} - \frac{817}{43200\epsilon} \right) + a_s^3 N_f \left\{ \frac{1}{\epsilon^3} \left[\frac{11N_f}{810} + \frac{1727C_F}{64800} - \frac{3773C_A}{129600} \right] \right. \\
& + \frac{1}{\epsilon^2} \left[- \frac{601N_f}{48600} + \frac{209633C_F}{3888000} + \left(- \frac{11(1-\xi)}{28800} - \frac{14989}{2592000} \right) C_A \right] \\
& + \frac{1}{\epsilon} \left[- \frac{58907N_f}{2916000} + \left(\frac{11\zeta_3}{90} - \frac{7092241}{29160000} \right) C_F \right. \\
& \left. + C_A \left(- \frac{6179(1-\xi)}{1728000} - \frac{11\zeta_3}{120} + \frac{4233367}{29160000} \right) \right] \left. \right\}, \quad (C.4)
\end{aligned}$$

$$\begin{aligned}
 \frac{Z_{qA}|_{n=6}}{C_A} &= a_s^2 N_f \left\{ \frac{11}{2520\epsilon^2} - \frac{221}{42336\epsilon} \right\} + a_s^3 N_f \left\{ \frac{1}{\epsilon^3} \left[\frac{11N_f}{2835} + \frac{7799C_F}{793800} - \frac{2123C_A}{423360} \right] \right. \\
 &\quad + \frac{1}{\epsilon^2} \left[-\frac{3263N_f}{1190700} + \frac{1585219C_F}{66679200} + \left(\frac{209(1-\xi)}{2540160} - \frac{2683601}{177811200} \right) C_A \right] \\
 &\quad + \frac{1}{\epsilon} \left[-\frac{3611983N_f}{500094000} + \left(\frac{11\zeta_3}{315} - \frac{11494299613}{140026320000} \right) C_F \right. \\
 &\quad \left. \left. + C_A \left(-\frac{1533541(1-\xi)}{1066867200} - \frac{11\zeta_3}{420} + \frac{699651481}{14002632000} \right) \right] \right\}, \quad (C.5)
 \end{aligned}$$

$$\begin{aligned}
 \frac{Z_{qA}|_{n=8}}{C_A} &= a_s^2 N_f \left(\frac{37}{20160\epsilon^2} - \frac{15529}{7257600\epsilon} \right) + a_s^3 N_f \left\{ \right. \\
 &\quad \frac{1}{\epsilon^3} \left[\frac{37N_f}{22680} + \frac{365671C_F}{76204800} - \frac{35039C_A}{30481920} \right] \\
 &\quad + \frac{1}{\epsilon^2} \left[-\frac{10393N_f}{11430720} + \frac{119144021C_F}{9144576000} + \left(\frac{3737(1-\xi)}{43545600} - \frac{1352619283}{128024064000} \right) C_A \right] \\
 &\quad + \frac{1}{\epsilon} \left[-\frac{506429639N_f}{144027072000} + \left(\frac{37\zeta_3}{2520} - \frac{574815689173}{15122842560000} \right) C_F \right. \\
 &\quad \left. \left. + C_A \left(-\frac{26006693(1-\xi)}{36578304000} - \frac{37\zeta_3}{3360} + \frac{2769495158803}{120982740480000} \right) \right] \right\}, \quad (C.6)
 \end{aligned}$$

$$\begin{aligned}
 \frac{Z_{qA}|_{n=10}}{C_A} &= a_s^2 N_f \left(\frac{7}{7425\epsilon^2} - \frac{2641}{2450250\epsilon} \right) + a_s^3 N_f \left\{ \right. \\
 &\quad \frac{1}{\epsilon^3} \left[\frac{56N_f}{66825} + \frac{2411C_F}{882090} - \frac{37199C_A}{176418000} \right] \\
 &\quad + \frac{1}{\epsilon^2} \left[-\frac{24421N_f}{66156750} + \frac{820067201C_F}{101881395000} + \left(\frac{2491(1-\xi)}{39204000} - \frac{35096312513}{4890306960000} \right) C_A \right] \\
 &\quad + \frac{1}{\epsilon} \left[-\frac{3690429931N_f}{1833865110000} + \left(\frac{56\zeta_3}{7425} - \frac{352770630827569}{16944913616400000} \right) C_F \right. \\
 &\quad \left. \left. + C_A \left(-\frac{9807967(1-\xi)}{24149664000} - \frac{14\zeta_3}{2475} + \frac{23337173703212867}{1897830325036800000} \right) \right] \right\}, \quad (C.7)
 \end{aligned}$$

$$\begin{aligned}
 \frac{Z_{gA}|_{n=2}}{C_A} &= -\frac{a_s}{2\epsilon} + a_s^2 \left\{ \frac{1}{\epsilon^2} \left(\frac{19C_A}{24} - \frac{N_f}{3} \right) + \frac{1}{\epsilon} \left[\left(\frac{5(1-\xi)}{48} - \frac{35}{48} \right) C_A + \frac{7N_f}{18} \right] \right\} \\
 &\quad + a_s^3 \left\{ \frac{1}{\epsilon^3} \left[\frac{71C_A N_f}{54} - \frac{779C_A^2}{432} - \frac{4C_F N_f}{27} - \frac{2N_f^2}{9} \right] \right. \\
 &\quad + \frac{1}{\epsilon^2} \left[\left(\frac{5(1-\xi)}{108} - \frac{73}{36} \right) C_A N_f \right. \\
 &\quad \left. + \left(\frac{5}{288}(1-\xi)^2 - \frac{35(1-\xi)}{216} + \frac{2807}{864} \right) C_A^2 - \frac{16C_F N_f}{27} + \frac{7N_f^2}{27} \right] \\
 &\quad + \frac{1}{\epsilon} \left[C_A N_f \left(-\frac{7}{216}(1-\xi) + \frac{19\zeta_3}{9} - \frac{194}{243} \right) + C_A^2 \left(-\frac{65(1-\xi)^2}{1728} - \frac{16759}{7776} \right. \right. \\
 &\quad \left. \left. - \frac{11\zeta_3}{72} + (1-\xi) \left(\frac{5\zeta_3}{72} + \frac{377}{1728} \right) \right) + \left(\frac{3059}{972} - \frac{8\zeta_3}{3} \right) C_F N_f + \frac{11N_f^2}{54} \right] \left. \right\}, \quad (C.8)
 \end{aligned}$$

$$\begin{aligned}
 \frac{Z_{gA}|_{n=4}}{C_A} = & -\frac{a_s}{12\epsilon} + a_s^2 \left\{ \frac{1}{\epsilon^2} \left[-\frac{97C_A}{1440} - \frac{N_f}{18} \right] + \frac{1}{\epsilon} \left[\left(\frac{1-\xi}{320} - \frac{8641}{86400} \right) C_A + \frac{7N_f}{216} \right] \right\} \\
 & + a_s^3 \left\{ \frac{1}{\epsilon^3} \left[\frac{C_A N_f}{324} + \frac{9437C_A^2}{86400} - \frac{121C_F N_f}{32400} - \frac{N_f^2}{27} \right] \right. \\
 & + \frac{1}{\epsilon^2} \left[\left(\frac{1-\xi}{720} - \frac{277}{129600} \right) C_A N_f \right. \\
 & + \left. \left(-\frac{13(1-\xi)^2}{23040} + \frac{853(1-\xi)}{86400} - \frac{1520341}{15552000} \right) C_A^2 - \frac{224719C_F N_f}{1944000} + \frac{7N_f^2}{324} \right] \\
 & + \frac{1}{\epsilon} \left[C_A N_f \left(\frac{457(1-\xi)}{43200} + \frac{53\zeta_3}{150} - \frac{5748673}{23328000} \right) + C_A^2 \left(-\frac{2357(1-\xi)^2}{460800} \right. \right. \\
 & + (1-\xi) \left(\frac{37\zeta_3}{9600} + \frac{37199}{648000} \right) - \frac{\zeta_3}{2400} - \frac{166178237}{466560000} \\
 & \left. \left. + \left(\frac{11814181}{29160000} - \frac{4\zeta_3}{9} \right) C_F N_f + \frac{193N_f^2}{3888} \right] \right\}, \tag{C.9}
 \end{aligned}$$

$$\begin{aligned}
 \frac{Z_{gA}|_{n=6}}{C_A} = & -\frac{a_s}{30\epsilon} + a_s^2 \left\{ \frac{1}{\epsilon^2} \left(-\frac{653C_A}{10080} - \frac{N_f}{45} \right) \right. \\
 & + \frac{1}{\epsilon} \left[\left(-\frac{19(1-\xi)}{20160} - \frac{185093}{4233600} \right) C_A + \frac{13N_f}{2700} \right] \left. \right\} \\
 & + a_s^3 \left\{ \frac{1}{\epsilon^3} \left[-\frac{22C_A N_f}{567} + \left(\frac{7(1-\xi)}{57600} + \frac{600331}{12700800} \right) C_A^2 - \frac{121C_F N_f}{198450} - \frac{2N_f^2}{135} \right] \right. \\
 & + \frac{1}{\epsilon^2} \left[\left(\frac{11761}{381024} - \frac{19(1-\xi)}{45360} \right) C_A N_f - \frac{253187C_F N_f}{5556600} + \frac{13N_f^2}{4050} \right. \\
 & + \left. \left(-\frac{37(1-\xi)^2}{48384} + \frac{585157(1-\xi)}{169344000} - \frac{275019229}{1778112000} \right) C_A^2 \right] \\
 & + \frac{1}{\epsilon} \left[C_A N_f \left(\frac{88439(1-\xi)}{19051200} + \frac{103\zeta_3}{735} - \frac{203978813}{2000376000} \right) + C_A^2 \left(-\frac{195683(1-\xi)^2}{101606400} \right. \right. \\
 & + (1-\xi) \left(\frac{61\zeta_3}{35280} + \frac{4629063223}{213373440000} \right) + \frac{1219\zeta_3}{176400} - \frac{385478104231}{2240421120000} \\
 & \left. \left. + \left(\frac{4970689127}{35006580000} - \frac{8\zeta_3}{45} \right) C_F N_f + \frac{5231N_f^2}{243000} \right] \right\}, \tag{C.10}
 \end{aligned}$$

$$\begin{aligned}
 \frac{Z_{gA}|_{n=8}}{C_A} = & -\frac{a_s}{56\epsilon} + a_s^2 \left\{ \frac{1}{\epsilon^2} \left(-\frac{2749C_A}{56448} - \frac{N_f}{84} \right) \right. \\
 & + \frac{1}{\epsilon} \left[\left(-\frac{101(1-\xi)}{80640} - \frac{18855769}{711244800} \right) C_A - \frac{43N_f}{70560} \right] \left. \right\} + a_s^3 \left\{ \frac{1}{\epsilon^3} \left[\right. \right. \\
 & - \frac{11281C_A N_f}{317520} + \left(\frac{11(1-\xi)}{94080} + \frac{26350343}{2133734400} \right) C_A^2 - \frac{1369C_F N_f}{7620480} - \frac{N_f^2}{126} \\
 & \left. \left. + \frac{1}{\epsilon^2} \left[\left(\frac{68655941}{3200601600} - \frac{101(1-\xi)}{181440} \right) C_A N_f + C_A^2 \left(-\frac{1151(1-\xi)^2}{1935360} \right) \right] \right. \right.
 \end{aligned}$$

$$\begin{aligned}
 & + \left. \left(\frac{112333(1-\xi)}{213373440} - \frac{25862266399}{199148544000} \right) - \frac{22126259C_F N_f}{914457600} - \frac{43N_f^2}{105840} \right] \\
 & + \frac{1}{\epsilon} \left[C_A N_f \left(\frac{1022971(1-\xi)}{457228800} + \frac{169\zeta_3}{2268} - \frac{48152267873}{896168448000} \right) \right. \\
 & + C_A^2 \left(-\frac{4988429(1-\xi)^2}{4877107200} + (1-\xi) \left(\frac{295\zeta_3}{290304} + \frac{333783455249}{32262064128000} \right) \right. \\
 & + \left. \left. \frac{14029\zeta_3}{2540160} - \frac{730008394755263}{6775033466880000} \right) \right. \\
 & + \left. \left(\frac{835100678591}{12098274048000} - \frac{2\zeta_3}{21} \right) C_F N_f + \frac{1028201N_f^2}{88905600} \right] \Bigg\}, \tag{C.11}
 \end{aligned}$$

$$\begin{aligned}
 \frac{Z_{gA}|_{n=10}}{C_A} = & -\frac{1}{90} \frac{1}{\epsilon} a_s + a_s^2 \left\{ \frac{1}{\epsilon^2} \left(-\frac{N_f}{135} - \frac{369361C_A}{9979200} \right) \right. \\
 & + \frac{1}{\epsilon} \left[\left(-\frac{2491(1-\xi)}{2217600} - \frac{1030030931}{55324684800} \right) C_A - \frac{661N_f}{340200} \right] \Bigg\} \\
 & + a_s^3 \left\{ \frac{1}{\epsilon^3} \left[-\frac{784C_F N_f}{11026125} - \frac{327427C_A N_f}{11226600} - \frac{2}{405} N_f^2 \right. \right. \\
 & + \left. \left. \left(\frac{143(1-\xi)}{1451520} - \frac{2899927799}{829870272000} \right) C_A^2 \right] + \frac{1}{\epsilon^2} \left[-\frac{27254752C_F N_f}{1819310625} \right. \right. \\
 & + \left. \left. \left(\frac{2071551877}{155600676000} - \frac{2491(1-\xi)}{4989600} \right) C_A N_f - \frac{661N_f^2}{510300} \right. \right. \\
 & + \left. \left. \left(-\frac{12121(1-\xi)^2}{26611200} - \frac{100407731(1-\xi)}{147532492800} - \frac{15087045196453}{142000024320000} \right) C_A^2 \right] \right. \\
 & + \frac{1}{\epsilon} \left[\left(\frac{167914071891373}{4236228404100000} - \frac{8\zeta_3}{135} \right) C_F N_f \right. \\
 & + C_A N_f \left(\frac{158792849(1-\xi)}{138311712000} + \frac{251\zeta_3}{5445} - \frac{560285942701061}{17253002954880000} \right) \\
 & + C_A^2 \left(-\frac{79286117(1-\xi)^2}{122943744000} + (1-\xi) \left(\frac{29\zeta_3}{43560} + \frac{67751391198259}{12268802101248000} \right) \right. \\
 & + \left. \left. \frac{1333\zeta_3}{338800} - \frac{24201974825379990307}{318835494606182400000} \right) + \frac{8972717N_f^2}{1285956000} \right] \Bigg\}. \tag{C.12}
 \end{aligned}$$

D Standard QCD renormalization constants

In our computations, we need the QCD beta function to two-loop order [86, 87],

$$\beta_0 = \frac{11C_A}{3} - \frac{2N_f}{3}, \tag{D.1}$$

$$\beta_1 = -\frac{10C_A N_f}{3} + \frac{34C_A^2}{3} - 2C_F N_f, \tag{D.2}$$

where β_i is defined with the following convention,

$$\frac{da_s}{d \ln \mu} = \beta(a_s) = -2a_s \sum_{i=0}^{\infty} a_s^{i+1} \beta_i. \tag{D.3}$$

The computations of off-shell OMEs requires also wave function renormalizations. The quark and gluon field renormalization constants in the $\overline{\text{MS}}$ scheme are needed up to three-loop order [88, 89], and the ghost field renormalization constant is required up to two-loop order. We list all these ingredients in our convention (2.16):

$$Z_q^{(0)} = 1, \quad (\text{D.4})$$

$$Z_q^{(1)} = -C_F \frac{\xi}{\epsilon}, \quad (\text{D.5})$$

$$Z_q^{(2)} = \frac{C_F}{\epsilon} \left[\left(-\frac{\xi^2}{8} - \xi - \frac{25}{8} \right) C_A + \frac{3C_F}{4} + \frac{N_f}{2} \right] + \frac{C_F}{\epsilon^2} \left[\left(\frac{\xi^2}{4} + \frac{3\xi}{4} \right) C_A + \frac{\xi^2 C_F}{2} \right], \quad (\text{D.6})$$

$$\begin{aligned} Z_q^{(3)} = & \frac{C_F}{\epsilon^3} \left[C_A \left(\left(-\frac{\xi^3}{4} - \frac{3\xi^2}{4} \right) C_F + \frac{\xi N_f}{6} \right) + \left(-\frac{\xi^3}{12} - \frac{3\xi^2}{8} - \frac{31\xi}{24} \right) C_A^2 - \frac{1}{6} \xi^3 C_F^2 \right] \\ & + \frac{C_F}{\epsilon^2} \left[C_A \left(\left(\frac{\xi^3}{8} + \xi^2 + \frac{25\xi}{8} - \frac{11}{6} \right) C_F + \left(-\frac{\xi}{2} - \frac{47}{18} \right) N_f \right) \right. \\ & + \left. \left(\frac{\xi^3}{8} + \frac{3\xi^2}{4} + \frac{73\xi}{24} + \frac{275}{36} \right) C_A^2 + \left(\frac{1}{3} - \frac{\xi}{2} \right) C_F N_f - \frac{3\xi C_F^2}{4} + \frac{2N_f^2}{9} \right] \\ & + \frac{C_F}{\epsilon} \left[C_A \left(\left(\frac{143}{12} - 4\zeta_3 \right) C_F + \left(\frac{17\xi}{24} + \frac{287}{54} \right) N_f \right) - \frac{C_F N_f}{2} - \frac{C_F^2}{2} - \frac{5N_f^2}{27} \right. \\ & \left. + C_A^2 \left\{ -\frac{5\xi^3}{48} + \xi^2 \left(-\frac{\zeta_3}{8} - \frac{13}{32} \right) + \xi \left(-\frac{\zeta_3}{4} - \frac{263}{96} \right) + \frac{23\zeta_3}{8} - \frac{9155}{432} \right\} \right], \quad (\text{D.7}) \end{aligned}$$

$$Z_g^{(0)} = 1, \quad (\text{D.8})$$

$$Z_g^{(1)} = \frac{1}{\epsilon} \left[\left(\frac{13}{6} - \frac{\xi}{2} \right) C_A - \frac{2N_f}{3} \right], \quad (\text{D.9})$$

$$\begin{aligned} Z_g^{(2)} = & \frac{1}{\epsilon} \left[-\frac{5C_A N_f}{4} + \left(-\frac{\xi^2}{8} - \frac{11\xi}{16} + \frac{59}{16} \right) C_A^2 - C_F N_f \right] \\ & + \frac{1}{\epsilon^2} \left[\left(\frac{\xi}{3} + \frac{1}{2} \right) C_A N_f + \left(\frac{\xi^2}{4} - \frac{17\xi}{24} - \frac{13}{8} \right) C_A^2 \right], \quad (\text{D.10}) \end{aligned}$$

$$\begin{aligned} Z_g^{(3)} = & \frac{1}{\epsilon^2} \left[C_F \left\{ \left(\frac{\xi}{2} + \frac{31}{18} \right) C_A N_f - \frac{2N_f^2}{9} \right\} + \left(\frac{\xi^2}{12} + \frac{19\xi}{24} + \frac{481}{108} \right) C_A^2 N_f - \frac{25}{54} C_A N_f^2 \right. \\ & + \left. \left(\frac{7\xi^3}{48} + \frac{13\xi^2}{24} - \frac{143\xi}{96} - \frac{7957}{864} \right) C_A^3 \right] + \frac{1}{\epsilon} \left[C_F \left\{ \left(-4\zeta_3 - \frac{5}{108} \right) C_A N_f + \frac{11N_f^2}{27} \right\} \right. \\ & + C_A^2 N_f \left(\frac{\xi}{3} + 3\zeta_3 - \frac{911}{108} \right) + \frac{19}{27} C_A N_f^2 + C_A^3 \left\{ -\frac{7\xi^3}{96} + \xi^2 \left(-\frac{\zeta_3}{16} - \frac{11}{32} \right) \right. \\ & + \left. \xi \left(-\frac{\zeta_3}{4} - \frac{167}{96} \right) - \frac{3\zeta_3}{16} + \frac{9965}{864} \right\} + \frac{1}{3} C_F^2 N_f \left. \right] + \frac{1}{\epsilon^3} \left[\left(-\frac{\xi^2}{6} - \frac{5\xi}{12} - \frac{11}{9} \right) C_A^2 N_f \right. \\ & \left. + \frac{1}{9} C_A N_f^2 + \left(-\frac{\xi^3}{8} + \frac{\xi^2}{6} + \frac{47\xi}{48} + \frac{403}{144} \right) C_A^3 \right], \quad (\text{D.11}) \end{aligned}$$

$$Z_c^{(0)} = 1, \quad (\text{D.12})$$

$$Z_c^{(1)} = \frac{1}{\epsilon} \left(\frac{3}{4} - \frac{\xi}{4} \right) C_A, \quad (\text{D.13})$$

$$Z_c^{(2)} = \frac{1}{\epsilon^2} \left[\frac{C_A N_f}{4} + \left(\frac{3\xi^2}{32} - \frac{35}{32} \right) C_A^2 \right] + \frac{1}{\epsilon} \left[\left(\frac{\xi}{32} + \frac{95}{96} \right) C_A^2 - \frac{5C_A N_f}{24} \right]. \quad (\text{D.14})$$

Object	In supplementary material	Object	In supplementary material
\mathcal{F}_{ns}	Fns[q, q]	z_1	z1
\mathcal{F}_{qq}	Fs[q, q]	a_s	as
\mathcal{F}_{qg}	Fs[q, g]	a_s^b	asb
\mathcal{F}_{qc}	Fs[q, c]	ξ	xi
\mathcal{F}_{gq}	Fs[g, q]	ξ^b	xib
\mathcal{F}_{gg}	Fs[g, g]	ϵ	eps
\mathcal{F}_{gc}	Fs[g, c]	C_A	ca
$\mathcal{F}_{ABC,q}$	Fs[ABC, q]	C_F	cf
$\mathcal{F}_{ABC,g}$	Fs[ABC, g]	N_f	nf
$\mathcal{F}_{ABC,c}$	Fs[ABC, c]	$16d^{abc}d_{abc}$	d33c
$\mathcal{F}_{[ZO]_g^{\text{GV},(2)},q}$	Fs[zogGV2, q]	T_F	tf
$\mathcal{F}_{[ZO]_g^{\text{GV},(2)},g}$	Fs[zogGV2, g]	Harm. sum $S_{\dots}(n)$	S[... , n]
$\mathcal{F}_{[ZO]_g^{\text{GV},(2)},c}$	Fs[zogGV2, c]	G. H. sum $S_{\dots}(\dots; n)$	S[... , {...} , n]
		Riemann ζ_n	zeta[n]

Table 2. Objects appearing in this paper and notation used in the supplementary material.

E Instructions for supplementary material

The `Mathematica` files ‘`NSingletOMEs.m`’ and ‘`SingletOMEs.m`’ contain all results for two-parton OMEs in the non-singlet and singlet case, respectively. All OMEs are normalized according to (2.39), and expanded to order $a_s^j \epsilon^{3-j}$ ($j \leq 3$), where the highest order is relevant only at the four-loop level. A `Mathematica` notebook ‘`ExtractSpFromOMEs.nb`’ is used to combine all OMEs and to derive the physical anomalous dimensions. In the same notebook, we also compare our results with the literature results, which we assembled in ‘`RefNSsp.m`’ and ‘`RefSingletsp.m`’, finding perfect agreement for both, the non-singlet and the singlet physical anomalous dimensions. The file ‘`FRzogGV2.m`’ contains the result for the last line of equation (5.41). We also provide the results for the renormalization constants Z_{qA} and Z_{gA} shown in appendix C in the files ‘`zqA.m`’ and ‘`zgA.m`’, respectively.

The generic notation for two-parton OME with an insertion of a general twist-two operator is defined in (2.30). More explicitly, for OMEs with an operator insertion of O_q or O_g , we follow this notation closely, for example,

$$\begin{aligned}
 A_{qq} &= \langle g(p) | O_q | g(p) \rangle, \\
 A_{gc} &= \langle c(p) | O_g | c(p) \rangle,
 \end{aligned}
 \tag{E.1}$$

while for OMEs with the insertion of GV counterterms, we use slightly different notations, for example,

$$\begin{aligned}
 A_{ABC,g} &= \langle g(p) | O_{ABC} | g(p) \rangle, \\
 A_{[ZO]_g^{\text{GV},(2)},c} &= \langle c(p) | [ZO]_g^{\text{GV},(2)} | c(p) \rangle.
 \end{aligned}
 \tag{E.2}$$

Similarly, in (2.39) we use $\mathcal{F}_{ABC,q}$, $\mathcal{F}_{[ZO]_g^{\text{GV},(2)},q}$, and so on to represent form factors with the insertion of a GV counterterm. The above OMEs apply to the singlet case only. In the non-singlet case, we need a single OME,

$$\mathcal{F}_{ns} = \frac{1}{2N_c} \frac{\text{Tr} \left(\not{p} \langle q(p) | O_{q,k} | q(p) \rangle \right)}{(\Delta \cdot p)^n}. \quad (\text{E.3})$$

In table 2, we provide a list of the objects that appear in this paper along with the corresponding notation used in the supplementary material.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] K.G. Wilson, *Nonlagrangian models of current algebra*, *Phys. Rev.* **179** (1969) 1499 [[INSPIRE](#)].
- [2] Y. Frishman, *Light cone and short distances*, *Phys. Rept.* **13** (1974) 1 [[INSPIRE](#)].
- [3] D.J. Gross and F. Wilczek, *Asymptotically free gauge theories. 2*, *Phys. Rev. D* **9** (1974) 980 [[INSPIRE](#)].
- [4] J.D. Bjorken, *Asymptotic Sum Rules at Infinite Momentum*, *Phys. Rev.* **179** (1969) 1547 [[INSPIRE](#)].
- [5] J.D. Bjorken and E.A. Paschos, *Inelastic Electron Proton and γ -Proton Scattering, and the Structure of the Nucleon*, *Phys. Rev.* **185** (1969) 1975 [[INSPIRE](#)].
- [6] G. Altarelli and G. Parisi, *Asymptotic Freedom in Parton Language*, *Nucl. Phys. B* **126** (1977) 298 [[INSPIRE](#)].
- [7] Y.L. Dokshitzer, *Calculation of the Structure Functions for Deep Inelastic Scattering and e^+e^- Annihilation by Perturbation Theory in Quantum Chromodynamics*, *Sov. Phys. JETP* **46** (1977) 641 [[INSPIRE](#)].
- [8] V.N. Gribov and L.N. Lipatov, *Deep inelastic $e p$ scattering in perturbation theory*, *Sov. J. Nucl. Phys.* **15** (1972) 438 [[INSPIRE](#)].
- [9] G. Heinrich, *Collider Physics at the Precision Frontier*, *Phys. Rept.* **922** (2021) 1 [[arXiv:2009.00516](#)] [[INSPIRE](#)].
- [10] S. Moch, J.A.M. Vermaseren and A. Vogt, *The Three loop splitting functions in QCD: The Nonsinglet case*, *Nucl. Phys. B* **688** (2004) 101 [[hep-ph/0403192](#)] [[INSPIRE](#)].
- [11] A. Vogt, S. Moch and J.A.M. Vermaseren, *The Three-loop splitting functions in QCD: The Singlet case*, *Nucl. Phys. B* **691** (2004) 129 [[hep-ph/0404111](#)] [[INSPIRE](#)].
- [12] J.A. Dixon and J.C. Taylor, *Renormalization of wilson operators in gauge theories*, *Nucl. Phys. B* **78** (1974) 552 [[INSPIRE](#)].
- [13] H. Kluberg-Stern and J.B. Zuber, *Ward Identities and Some Clues to the Renormalization of Gauge Invariant Operators*, *Phys. Rev. D* **12** (1975) 467 [[INSPIRE](#)].
- [14] R. Hamberg and W.L. van Neerven, *The Correct renormalization of the gluon operator in a covariant gauge*, *Nucl. Phys. B* **379** (1992) 143 [[INSPIRE](#)].

- [15] J. Ablinger et al., *Massive 3-loop Ladder Diagrams for Quarkonic Local Operator Matrix Elements*, *Nucl. Phys. B* **864** (2012) 52 [[arXiv:1206.2252](#)] [[INSPIRE](#)].
- [16] E.G. Floratos, D.A. Ross and C.T. Sachrajda, *Higher Order Effects in Asymptotically Free Gauge Theories. II. Flavor Singlet Wilson Operators and Coefficient Functions*, *Nucl. Phys. B* **152** (1979) 493 [[INSPIRE](#)].
- [17] A. Gonzalez-Arroyo and C. Lopez, *Second Order Contributions to the Structure Functions in Deep Inelastic Scattering. III. The Singlet Case*, *Nucl. Phys. B* **166** (1980) 429 [[INSPIRE](#)].
- [18] S.D. Joglekar and B.W. Lee, *General Theory of Renormalization of Gauge Invariant Operators*, *Annals Phys.* **97** (1976) 160 [[INSPIRE](#)].
- [19] H. Kluberg-Stern and J.B. Zuber, *Renormalization of Nonabelian Gauge Theories in a Background Field Gauge. I. Green Functions*, *Phys. Rev. D* **12** (1975) 482 [[INSPIRE](#)].
- [20] H. Kluberg-Stern and J.B. Zuber, *Renormalization of Nonabelian Gauge Theories in a Background Field Gauge. II. Gauge Invariant Operators*, *Phys. Rev. D* **12** (1975) 3159 [[INSPIRE](#)].
- [21] J.C. Collins and R.J. Scalise, *The Renormalization of composite operators in Yang-Mills theories using general covariant gauge*, *Phys. Rev. D* **50** (1994) 4117 [[hep-ph/9403231](#)] [[INSPIRE](#)].
- [22] S. Moch, J.A.M. Vermaseren and A. Vogt, *The Three-Loop Splitting Functions in QCD: The Helicity-Dependent Case*, *Nucl. Phys. B* **889** (2014) 351 [[arXiv:1409.5131](#)] [[INSPIRE](#)].
- [23] B. Mistlberger, *Higgs boson production at hadron colliders at N^3 LO in QCD*, *JHEP* **05** (2018) 028 [[arXiv:1802.00833](#)] [[INSPIRE](#)].
- [24] C. Duhr, F. Dulat and B. Mistlberger, *Drell-Yan Cross Section to Third Order in the Strong Coupling Constant*, *Phys. Rev. Lett.* **125** (2020) 172001 [[arXiv:2001.07717](#)] [[INSPIRE](#)].
- [25] M.-X. Luo, T.-Z. Yang, H.X. Zhu and Y.J. Zhu, *Quark Transverse Parton Distribution at the Next-to-Next-to-Next-to-Leading Order*, *Phys. Rev. Lett.* **124** (2020) 092001 [[arXiv:1912.05778](#)] [[INSPIRE](#)].
- [26] M.A. Ebert, B. Mistlberger and G. Vita, *Transverse momentum dependent PDFs at N^3 LO*, *JHEP* **09** (2020) 146 [[arXiv:2006.05329](#)] [[INSPIRE](#)].
- [27] M.A. Ebert, B. Mistlberger and G. Vita, *N -jettiness beam functions at N^3 LO*, *JHEP* **09** (2020) 143 [[arXiv:2006.03056](#)] [[INSPIRE](#)].
- [28] M.-X. Luo, T.-Z. Yang, H.X. Zhu and Y.J. Zhu, *Unpolarized quark and gluon TMD PDFs and FFs at N^3 LO*, *JHEP* **06** (2021) 115 [[arXiv:2012.03256](#)] [[INSPIRE](#)].
- [29] D. Baranowski et al., *Beam functions for N -jettiness at N^3 LO in perturbative QCD*, *JHEP* **02** (2023) 073 [[arXiv:2211.05722](#)] [[INSPIRE](#)].
- [30] G. Falcioni and F. Herzog, *Renormalization of gluonic leading-twist operators in covariant gauges*, *JHEP* **05** (2022) 177 [[arXiv:2203.11181](#)] [[INSPIRE](#)].
- [31] J. Ablinger et al., *The 3-loop pure singlet heavy flavor contributions to the structure function $F_2(x, Q^2)$ and the anomalous dimension*, *Nucl. Phys. B* **890** (2014) 48 [[arXiv:1409.1135](#)] [[INSPIRE](#)].
- [32] J. Ablinger et al., *The 3-Loop Non-Singlet Heavy Flavor Contributions and Anomalous Dimensions for the Structure Function $F_2(x, Q^2)$ and Transversity*, *Nucl. Phys. B* **886** (2014) 733 [[arXiv:1406.4654](#)] [[INSPIRE](#)].

- [33] A. Behring et al., *The Polarized Three-Loop Anomalous Dimensions from On-Shell Massive Operator Matrix Elements*, *Nucl. Phys. B* **948** (2019) 114753 [[arXiv:1908.03779](#)] [[INSPIRE](#)].
- [34] J. Ablinger et al., *The unpolarized and polarized single-mass three-loop heavy flavor operator matrix elements $A_{gg,Q}$ and $\Delta A_{gg,Q}$* , *JHEP* **12** (2022) 134 [[arXiv:2211.05462](#)] [[INSPIRE](#)].
- [35] I. Bierenbaum et al., *$O(\alpha_s^2)$ polarized heavy flavor corrections to deep-inelastic scattering at $Q^2 \gg m^2$* , *Nucl. Phys. B* **988** (2023) 116114 [[arXiv:2211.15337](#)] [[INSPIRE](#)].
- [36] J. Ablinger et al., *The three-loop splitting functions $P_{qq}^{(2)}$ and $P_{gg}^{(2,N_F)}$* , *Nucl. Phys. B* **922** (2017) 1 [[arXiv:1705.01508](#)] [[INSPIRE](#)].
- [37] J. Blümlein, P. Marquard, C. Schneider and K. Schönwald, *The three-loop unpolarized and polarized non-singlet anomalous dimensions from off shell operator matrix elements*, *Nucl. Phys. B* **971** (2021) 115542 [[arXiv:2107.06267](#)] [[INSPIRE](#)].
- [38] R. Mertig and W.L. van Neerven, *The Calculation of the two loop spin splitting functions $P_{ij}^{(1)}(x)$* , *Z. Phys. C* **70** (1996) 637 [[hep-ph/9506451](#)] [[INSPIRE](#)].
- [39] W. Vogelsang, *A Rederivation of the spin dependent next-to-leading order splitting functions*, *Phys. Rev. D* **54** (1996) 2023 [[hep-ph/9512218](#)] [[INSPIRE](#)].
- [40] J. Blümlein, P. Marquard, C. Schneider and K. Schönwald, *The three-loop polarized singlet anomalous dimensions from off-shell operator matrix elements*, *JHEP* **01** (2022) 193 [[arXiv:2111.12401](#)] [[INSPIRE](#)].
- [41] G. Curci, W. Furmanski and R. Petronzio, *Evolution of Parton Densities Beyond Leading Order: The Nonsinglet Case*, *Nucl. Phys. B* **175** (1980) 27 [[INSPIRE](#)].
- [42] W. Furmanski and R. Petronzio, *Singlet Parton Densities Beyond Leading Order*, *Phys. Lett. B* **97** (1980) 437 [[INSPIRE](#)].
- [43] R.K. Ellis and W. Vogelsang, *The Evolution of parton distributions beyond leading order: The Singlet case*, [hep-ph/9602356](#) [[INSPIRE](#)].
- [44] A. Bassetto, G. Heinrich, Z. Kunszt and W. Vogelsang, *The Light cone gauge and the calculation of the two loop splitting functions*, *Phys. Rev. D* **58** (1998) 094020 [[hep-ph/9805283](#)] [[INSPIRE](#)].
- [45] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279 [[INSPIRE](#)].
- [46] S. Moch et al., *Low moments of the four-loop splitting functions in QCD*, *Phys. Lett. B* **825** (2022) 136853 [[arXiv:2111.15561](#)] [[INSPIRE](#)].
- [47] K.G. Chetyrkin and F.V. Tkachov, *Integration by Parts: The Algorithm to Calculate beta Functions in 4 Loops*, *Nucl. Phys. B* **192** (1981) 159 [[INSPIRE](#)].
- [48] J.A.M. Vermaseren, *New features of FORM*, [math-ph/0010025](#) [[INSPIRE](#)].
- [49] A. von Manteuffel and C. Studerus, *Reduze 2 — Distributed Feynman Integral Reduction*, [arXiv:1201.4330](#) [[INSPIRE](#)].
- [50] V. Shtabovenko, R. Mertig and F. Orellana, *New Developments in FeynCalc 9.0*, *Comput. Phys. Commun.* **207** (2016) 432 [[arXiv:1601.01167](#)] [[INSPIRE](#)].
- [51] V. Shtabovenko, *FeynCalc goes multiloop*, *J. Phys. Conf. Ser.* **2438** (2023) 012140 [[arXiv:2112.14132](#)] [[INSPIRE](#)].
- [52] F. Feng, *Apart: A Generalized Mathematica Apart Function*, *Comput. Phys. Commun.* **183** (2012) 2158 [[arXiv:1204.2314](#)] [[INSPIRE](#)].

- [53] R.N. Lee, *Presenting LiteRed: a tool for the Loop InTEgrals REDuction*, [arXiv:1212.2685](#) [[INSPIRE](#)].
- [54] A.V. Smirnov and F.S. Chuharev, *FIRE6: Feynman Integral REDuction with Modular Arithmetic*, *Comput. Phys. Commun.* **247** (2020) 106877 [[arXiv:1901.07808](#)] [[INSPIRE](#)].
- [55] J. Klappert, F. Lange, P. Maierhöfer and J. Usovitsch, *Integral reduction with Kira 2.0 and finite field methods*, *Comput. Phys. Commun.* **266** (2021) 108024 [[arXiv:2008.06494](#)] [[INSPIRE](#)].
- [56] S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, *Int. J. Mod. Phys. A* **15** (2000) 5087 [[hep-ph/0102033](#)] [[INSPIRE](#)].
- [57] T. Gehrmann and E. Remiddi, *Differential equations for two loop four point functions*, *Nucl. Phys. B* **580** (2000) 485 [[hep-ph/9912329](#)] [[INSPIRE](#)].
- [58] J.M. Henn, *Multiloop integrals in dimensional regularization made simple*, *Phys. Rev. Lett.* **110** (2013) 251601 [[arXiv:1304.1806](#)] [[INSPIRE](#)].
- [59] C. Meyer, *Algorithmic transformation of multi-loop master integrals to a canonical basis with CANONICA*, *Comput. Phys. Commun.* **222** (2018) 295 [[arXiv:1705.06252](#)] [[INSPIRE](#)].
- [60] C. Meyer, *Transforming differential equations of multi-loop Feynman integrals into canonical form*, *JHEP* **04** (2017) 006 [[arXiv:1611.01087](#)] [[INSPIRE](#)].
- [61] R.N. Lee, *Reducing differential equations for multiloop master integrals*, *JHEP* **04** (2015) 108 [[arXiv:1411.0911](#)] [[INSPIRE](#)].
- [62] R.N. Lee, *Libra: A package for transformation of differential systems for multiloop integrals*, *Comput. Phys. Commun.* **267** (2021) 108058 [[arXiv:2012.00279](#)] [[INSPIRE](#)].
- [63] P.A. Baikov and K.G. Chetyrkin, *Four Loop Massless Propagators: An Algebraic Evaluation of All Master Integrals*, *Nucl. Phys. B* **837** (2010) 186 [[arXiv:1004.1153](#)] [[INSPIRE](#)].
- [64] R.N. Lee, A.V. Smirnov and V.A. Smirnov, *Master Integrals for Four-Loop Massless Propagators up to Transcendentality Weight Twelve*, *Nucl. Phys. B* **856** (2012) 95 [[arXiv:1108.0732](#)] [[INSPIRE](#)].
- [65] E. Remiddi and J.A.M. Vermaseren, *Harmonic polylogarithms*, *Int. J. Mod. Phys. A* **15** (2000) 725 [[hep-ph/9905237](#)] [[INSPIRE](#)].
- [66] J.A.M. Vermaseren, *Harmonic sums, Mellin transforms and integrals*, *Int. J. Mod. Phys. A* **14** (1999) 2037 [[hep-ph/9806280](#)] [[INSPIRE](#)].
- [67] J. Blümlein and S. Kurth, *Harmonic sums and Mellin transforms up to two loop order*, *Phys. Rev. D* **60** (1999) 014018 [[hep-ph/9810241](#)] [[INSPIRE](#)].
- [68] J. Ablinger, *A Computer Algebra Toolbox for Harmonic Sums Related to Particle Physics*, MSc Thesis, Johannes Kepler University of Linz, AUSTRIA (2009) [[arXiv:1011.1176](#)] [[INSPIRE](#)].
- [69] J. Ablinger, *Computer Algebra Algorithms for Special Functions in Particle Physics*, Ph.D. Thesis, Johannes Kepler University of Linz, AUSTRIA (2012) [[arXiv:1305.0687](#)] [[INSPIRE](#)].
- [70] J. Ablinger, *The package HarmonicSums: Computer Algebra and Analytic aspects of Nested Sums*, *PoS* **LL2014** (2014) 019 [[arXiv:1407.6180](#)] [[INSPIRE](#)].
- [71] J. Ablinger, J. Blümlein and C. Schneider, *Harmonic Sums and Polylogarithms Generated by Cyclotomic Polynomials*, *J. Math. Phys.* **52** (2011) 102301 [[arXiv:1105.6063](#)] [[INSPIRE](#)].

- [72] J. Ablinger, J. Blümlein and C. Schneider, *Analytic and Algorithmic Aspects of Generalized Harmonic Sums and Polylogarithms*, *J. Math. Phys.* **54** (2013) 082301 [[arXiv:1302.0378](#)] [[INSPIRE](#)].
- [73] J. Ablinger, J. Blümlein, C.G. Raab and C. Schneider, *Iterated Binomial Sums and their Associated Iterated Integrals*, *J. Math. Phys.* **55** (2014) 112301 [[arXiv:1407.1822](#)] [[INSPIRE](#)].
- [74] A. von Manteuffel and R.M. Schabinger, *A novel approach to integration by parts reduction*, *Phys. Lett. B* **744** (2015) 101 [[arXiv:1406.4513](#)] [[INSPIRE](#)].
- [75] T. Peraro, *Scattering amplitudes over finite fields and multivariate functional reconstruction*, *JHEP* **12** (2016) 030 [[arXiv:1608.01902](#)] [[INSPIRE](#)].
- [76] M. Heller and A. von Manteuffel, *MultivariateApart: Generalized partial fractions*, *Comput. Phys. Commun.* **271** (2022) 108174 [[arXiv:2101.08283](#)] [[INSPIRE](#)].
- [77] J. Boehm et al., *IBP reduction coefficients made simple*, *JHEP* **12** (2020) 054 [[arXiv:2008.13194](#)] [[INSPIRE](#)].
- [78] J. Blümlein, P. Marquard, C. Schneider and K. Schönwald, *The two-loop massless off-shell QCD operator matrix elements to finite terms*, *Nucl. Phys. B* **980** (2022) 115794 [[arXiv:2202.03216](#)] [[INSPIRE](#)].
- [79] Y. Matiounine, J. Smith and W.L. van Neerven, *Two loop operator matrix elements calculated up to finite terms*, *Phys. Rev. D* **57** (1998) 6701 [[hep-ph/9801224](#)] [[INSPIRE](#)].
- [80] T.G. Birtwright, E.W.N. Glover and P. Marquard, *Master integrals for massless two-loop vertex diagrams with three offshell legs*, *JHEP* **09** (2004) 042 [[hep-ph/0407343](#)] [[INSPIRE](#)].
- [81] S. Moch, P. Uwer and S. Weinzierl, *Nested sums, expansion of transcendental functions and multiscale multiloop integrals*, *J. Math. Phys.* **43** (2002) 3363 [[hep-ph/0110083](#)] [[INSPIRE](#)].
- [82] T. Peraro, *FiniteFlow: multivariate functional reconstruction using finite fields and dataflow graphs*, *JHEP* **07** (2019) 031 [[arXiv:1905.08019](#)] [[INSPIRE](#)].
- [83] G. Falcioni and F. Herzog, private communication.
- [84] T. Gehrmann, A. von Manteuffel and T.-Z. Yang, *Renormalization of twist-two operators in QCD and its application to singlet splitting functions*, *PoS LL2022* (2022) 063 [[arXiv:2207.10108](#)] [[INSPIRE](#)].
- [85] S. Moch et al., *Four-Loop Non-Singlet Splitting Functions in the Planar Limit and Beyond*, *JHEP* **10** (2017) 041 [[arXiv:1707.08315](#)] [[INSPIRE](#)].
- [86] W.E. Caswell, *Asymptotic Behavior of Nonabelian Gauge Theories to Two Loop Order*, *Phys. Rev. Lett.* **33** (1974) 244 [[INSPIRE](#)].
- [87] D.R.T. Jones, *Two Loop Diagrams in Yang-Mills Theory*, *Nucl. Phys. B* **75** (1974) 531 [[INSPIRE](#)].
- [88] O.V. Tarasov, A.A. Vladimirov and A.Y. Zharkov, *The Gell-Mann-Low Function of QCD in the Three Loop Approximation*, *Phys. Lett. B* **93** (1980) 429 [[INSPIRE](#)].
- [89] S.A. Larin and J.A.M. Vermaseren, *The Three loop QCD Beta function and anomalous dimensions*, *Phys. Lett. B* **303** (1993) 334 [[hep-ph/9302208](#)] [[INSPIRE](#)].