



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

Transfusion in trauma: why and how should we change our current practice?

Theusinger, O M ; Spahn, D R ; Ganter, M T

DOI: <https://doi.org/10.1097/ACO.0b013e3283212c7c>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-24479>

Journal Article

Accepted Version

Originally published at:

Theusinger, O M; Spahn, D R; Ganter, M T (2009). Transfusion in trauma: why and how should we change our current practice? *Current Opinion in Anaesthesiology*, 22(2):305-312.

DOI: <https://doi.org/10.1097/ACO.0b013e3283212c7c>

Transfusion and trauma – Future strategies

Oliver M. Theusinger M.D.¹, Donat R. Spahn M.D., F.R.C.A.²

Michael T. Ganter M.D., D.E.A.A.³

¹ Resident, Institute of Anesthesiology, University Hospital and University of Zurich,
Switzerland

² Professor and Chairman, Institute of Anesthesiology, University Hospital and University
of Zurich, Switzerland

³ Senior staff member, Institute of Anesthesiology, University Hospital and University of
Zurich, Switzerland

Manuscript Number: ACO2202c

Address for correspondence:

Oliver M. Theusinger, MD
Institute of Anesthesiology
University Hospital and University of Zurich
Rämistrasse 100
CH-8091 Zurich, Switzerland
Tel: +41 44 255 26 96
Fax: +41 44 255 44 09
E-mail: oliver.theusinger@usz.ch

Abstract

Kommentar [OMT1]: Muss <200 sein, AKTUELL OK mit 199 Words

Kommentar [d2]: Mit diesen Modifikationen hat es nun 198 Wörter.

Purpose of review: Major trauma is often associated with hemorrhage and transfusion of blood and blood products, all associated with adverse clinical outcome. The aim of this review is to emphasize why bleeding and coagulation has to be monitored closely in trauma patients and to discuss the rationale behind modern and future transfusion strategies.

Recent findings: Hemorrhage is a major cause of early death after trauma. Besides the initial injuries, hemorrhage is significantly promoted by coagulopathy. Early identification of the underlying cause of hemorrhage with coagulation tests (routine and bedside) in conjunction with blood gas analysis allow early goal-directed treatment of coagulation disorders and anemia thereby stopping bleeding and reducing transfusion requirements. These treatment options have to be adapted to the civilian and non civilian sector. Transfusion of blood and its components is critical in the management of trauma hemorrhage, but is per se associated with adverse outcome. Decisions must weigh the potential benefits and harms.

Summary: Future transfusion strategies are based on early and continuous assessment of the bleeding and coagulation status of trauma patients. This allows specific and goal-directed treatment, thereby optimizing patient's coagulation status early, minimizing exposure to blood products, reducing costs and improving patient's outcome.

Keywords: Blood management, transfusion, hemorrhage

Introduction

Hemorrhage is known to be a major cause of early death after injury and has been shown to be responsible for 30% to 40% of trauma mortalities¹⁻³. Furthermore, hemorrhage with consecutive multiple transfusions has been shown to significantly worsen clinical outcomes.^{4,5}

Management priorities in trauma patients are to ensure adequate ventilation, oxygen delivery, hemorrhage control and to restore tissue perfusion to vital organs. Early and continuous re-assessment of the bleeding and coagulation status of trauma patients allows specific goal-directed treatment, thereby optimizing patient's coagulation status, minimizing exposure to blood products, reducing costs and improving patient's outcome.^{6,7}

Mechanism of hemorrhage in trauma patients

Hemorrhage reduces preload required to ensure adequate cardiac output and peripheral oxygen delivery. Inadequate tissue perfusion, not always associated with overt hypotension, can trigger a neuro-humoral cascade, leading to sequential organ failure. Mortality from established organ failure has not changed since it was first described almost 30 years ago⁸. Diagnosing and treating hemorrhage early remains imperative. The American College of Surgeons has developed the classification scheme stratifying blood loss from Stage 1 (<15% of total circulating blood volume) to Stage 4 (>40% of total circulating blood volume)⁹. Young people in good health may compensate well for large-volume blood loss, up to 50% of the total circulating blood volume. Then, they may

Kommentar [OMT3]: 50-100

AKTUELL OK mit 98 Words

Kommentar [OMT4]: 2500
Wörter

Gilt fuer gesamtes review exkl Refs

Kommentar [d5]: Den NEUEN Mechanismus (im Abstract angekündigt) sollte hier exakt beschrieben werden mit 1-2 Figuren um zu erklären warum es zu einer Hypokoagulation kommt und gleichzeitig zu einer Hyperfibrinolyse – Das ist einer der WICHTIGSTEN Kommentare

develop sudden cardiovascular compromise when compensatory mechanisms fail. Elderly people on the other side may tolerate much smaller blood losses only.

Anatomical bleeding

Active hemorrhage as a result of major injuries is life-threatening and leads to hemorrhagic shock and exsanguination if not treated readily. Bleeding may be stopped temporarily by external compression and tourniquets, however, surgical or interventional (e.g., arterial embolization) repair is required for final hemorrhage control.

Coagulopathic bleeding

An abnormal coagulation status is frequently present early after major trauma at admission to the emergency department and is associated with a 5-fold increase in mortality¹⁰⁻¹³. Traditionally, the acute traumatic coagulopathy has been thought to be due to consumption of coagulation factors, dilution from intravenous fluid therapy, hypothermia, and metabolic acidosis. However, it has recently been shown that none of these factors is initially responsible for the acute traumatic coagulopathy. These factors become significant only in the later phase of traumatic coagulopathy. Studies by Brohi et al¹¹⁻¹⁴ have described an early and previously unknown acute traumatic coagulopathy before any of the above mentioned traditional causes (lethal triade) of acute traumatic coagulopathy were present. These studies have shown that coagulopathy and mortality as well as transfusion requirements are to by linked with hypoperfusion of tissues and the protein c-pathway, this was also proven for traumatic brain injury. Low protein C levels are associated with prolongation of the partial thromboplastin and prothrombin times and hyperfibrinolysis with low levels of plasminogen activator inhibitor-1 and high D-dimer

levels. High thrombomodulin and low protein C levels are associated with increased mortality, blood transfusion requirements, acute renal injury, and reduced ventilator-free days.

The pathophysiology of fibrinolysis in association with injury is exacerbated by shock and this is mediated by deinhibition of tissue plasminogen activator through the consumption of plasminogen activator inhibitor-1. A low level plasminogen activator inhibitor-1, in combination with an increased release of plasminogen activators from the vessel wall will result in hyperfibrinolysis. Actually it seems that thrombin activatable fibrinolysis inhibitor is the main driver of fibrinolysis inhibition, and that reduction in thrombin activatable fibrinolysis inhibitor activation by the competitive binding of protein C to thrombin–thrombomodulin is the mechanism for derepression of fibrinolysis with activation of protein C. The consumption of plasminogen activator inhibitor-1 by activated protein C seems to be the most important clinical cause of hyperfibrinolysis.

Diagnosis and monitoring of bleeding

Assessment of hemorrhage and volume status. Blood pressure and heart rate are vital signs which are nonspecific to evaluate hemorrhage. Mixed and central venous oxygen saturation are more sensitive and reliable measurements of acute volume loss¹⁵⁻¹⁷. The degree of metabolic acidosis, as measured by the base deficit from an arterial blood gas sample, is helpful to evaluate the degree of shock. Base deficit has been shown to correlate with transfusion requirements, ICU stay, and ultimate outcome^{18,19}. During initial resuscitation, base deficit typically correlates with serum lactate level. Interestingly, the ability to clear lactate to normal is one of the most important predictors

of survival following hemorrhage and injury²⁰⁻²². Serial blood gas determinations (arterial and venous) may be helpful in determining whether blood loss is continuing or not.

Assessment of coagulation. Most commonly, routine laboratory-based coagulation tests (e.g., PT/INR, aPTT, fibrinogen), platelet numbers and hemoglobin concentration (Hb) are being used to assess the patients' current coagulation status²³. However, the value of these tests has been questioned in the acute bleeding situation because there are delays from blood sampling to obtaining results (45–60 min), coagulation tests are determined in plasma rather than whole blood, no information is available on platelet function and the assays are performed at a standard temperature of 37°C rather than the patient's temperature²⁴.

Point-of-care coagulation monitoring devices assessing the viscoelastic properties of the developing clot in whole blood, e.g., thrombelastography (TEG®) or rotation thrombelastometry (ROTEM®), may overcome several limitations of routine coagulation tests^{25,26}. In particular, these technologies allow to assess in vivo coagulation system interactions with platelets and red blood cells and provide useful information on platelet function²⁷. In addition, they may be performed at the bedside allowing faster turnaround times. Furthermore, the clot development can be visually displayed in real-time, the assay is fibrinolysis sensitive, and the coagulation analysis can be done at the patient's temperature²⁸. Nevertheless, results obtained from these in vitro tests must be carefully interpreted in the clinical context.

Present and future transfusion practice in trauma patients

Prevention of further bleeding

The initial phase after trauma - from injury to surgery/intervention – has to be minimized and further bleeding should be prevented. Hill et al²⁹ observed a significant decrease in mortality from shock by establishing a 60 minute emergency department time limit for patients in hemorrhagic shock. Additionally, Hoyt et al³⁰ showed that delayed transfer to theatre is an avoidable cause of death which can be minimized by reducing the time for diagnosis and resuscitation prior to surgery.

Without brain injury, a target systolic blood pressure of 80 to 100 mm Hg should be maintained until the major bleeding is stopped.⁶ Aggressive fluid therapy to preserve tissue oxygenation and to restore the circulating blood volume leads to dilution of coagulation factors, hypothermia, increased hydrostatic pressure and further bleeding. The concept of low-volume fluid resuscitation, called ‘permissive hypotension’, prior to surgical source control maintains tissue perfusion at a lower level but sufficient for a short period of time without the adverse effects mentioned above³¹. The low-volume approach is not to be applied in traumatic brain and spinal injuries as an adequate perfusion pressure is crucial to ensure tissue oxygenation of the injured central nervous system.

Transfusion of red blood cells (RBC)

Hemoglobin based transfusion triggers. According to current guidelines from the American Society of Anesthesiologist, RBC transfusion is recommended if the hemoglobin concentration drops below 6-10g/dl. Transfusions over 10 g/dl are rarely

indicated and, transfusions seem almost always indicated if hemoglobin falls below 6 g/dl.³² In Europe, a Hb target range of 7-9 g/dL is largely accepted in major trauma.⁶

Physiological transfusion triggers on the other hand are tachycardia, hypotension, oxygen extraction greater than 50%, mixed venous oxygen pressure of less than 32 mm Hg, increase of lactate and ECG changes^{6,33,34}. The depth of shock, hemodynamic response to resuscitation, and the rate of actual blood loss in the acutely bleeding and hemodynamically unstable patient may also be integrated into the indication for RBC transfusion³⁵. However, RBC transfusions should be used restrictively³⁶.

Alternatives to allogeneic red cell transfusion

In trauma, the only available alternative to allogeneic RBC transfusion in clinical practice is currently the intraoperative cell salvage. It has been shown efficacious for reducing allogeneic blood transfusions³⁷. The quality of salvaged blood seems to be better compared with stored packed RBCs, with less risk for the patient³⁷. However, cell salvage is only applicable when the operative site not contaminated.

Artificial oxygen carriers may represent a future alternative to RBC transfusions. There are two groups of artificial oxygen carrier currently under development: Synthetically manufactured perfluorocarbons, and hemoglobin-based oxygen carriers³⁸. However, none of these products has achieved market approval for Europe, US or Canada so far.

Transfusion of fresh frozen plasma (FFP) and platelets

Traditional indications for fresh frozen plasma are massive bleeding due to multiple factor deficiencies, emergency reversal of vitamin K antagonists, and treatment of

thrombotic thrombocytopenic purpura (TTP)³². However, the clinical efficacy of FFP is largely unproven^{6,39}. Large quantities of FFP (15 ml/kg body weight and more) are recommended to achieve an effect in massive bleeding^{6,40,41}. Interestingly, there are only expert-opinion but no evidence based transfusion thresholds published for FFP administration^{32,42}. However, there are significant adverse effects associated with FFP transfusions including volume overload (TACO), allergic reactions, transfusion-related acute lung injury (TRALI), transmission of infectious pathogens^{40,43,44} and a 3 fold increase of nosocomial infections⁴⁵.

In a bleeding patient, platelets should be kept $>50 \times 10^3 /\mu\text{l}$, and $>100 \times 10^3 /\mu\text{l}$ in patients with traumatic brain injury^{6,7}. Adverse effects of platelets are similar to FFP, except the much higher risk of bacterial contamination⁴⁶. Therefore, the indication to transfuse platelets is to be restrictive. As for FFP, there are only expert-opinions available for the indication of platelet transfusion, no studies on an evidence based level are available so far.

Replacement of specific coagulation factors

Fibrinogen. Fibrinogen is the substrate to form a clot. Several in-vitro and animal studies have shown that fibrinogen substitution is essential to reverse dilutional coagulopathy⁴⁷⁻⁴⁹. Furthermore, several human studies (civilian and non-civilian) confirmed these data, showing that early and aggressive replacement of fibrinogen in patients with severe hemorrhage and dilutional coagulopathy improves clot strength significantly and may lead to better survival⁵⁰⁻⁵³.

Factor XIII. Factor XIII is the key coagulation factor to stabilize the clot. Schroeder et al and Nielsen et al ⁵⁴⁻⁵⁶ have proven a relation between decreased factor XIII activity and reduced clot firmness (MCF) by computerized thrombelastography. Trauma and major hemorrhage is known to be a cause of acquired factor XIII deficiency ⁵⁷. It seems reasonable to substitute factor XIII early, thereby improving clot firmness and reducing bleeding ⁵⁸ as well as the use of blood products ^{59,60}.

Prothrombin complex concentrate. Prothrombin complex concentrates (PCC) provide a source of the four vitamin K dependent coagulation factors, and consequently, PCC is recommended both in Europe and the USA for emergent reversal of oral anticoagulants ⁶¹⁻⁶⁵. Furthermore, Bruce and Nokes ⁶⁶ recently demonstrated that the use of PCC in trauma patients leads to a considerable reduction in the use of blood products (FFPs, RBCs and cryoprecipitate) and that survival improved and bleeding stopped earlier. Therefore, PCC might have a place in control of trauma related bleeding, although this indication is currently off label.

rFVIIa. Recombinant activated factor VIII (rFVIIa) leads to a “thrombin burst” thereby transforming fibrinogen into fibrin. Before treating patients with rFVIIa, patients should fulfill certain criteria, for example thrombocytopenia and hypofibrinogenemia must be corrected ⁶⁷⁻⁶⁹. One study in trauma has shown a reduction in RBC transfusions in rFVIIa treated patients ⁷⁰. However, rFVIIa is not an alternative to surgical bleeding control, and its use in trauma is still an off label indication. Risks and benefits have to be carefully evaluated before usage and economic aspects taken into consideration ⁷¹.

Pharmacologic agents

Antifibrinolytics, such as aminocaproic acid (ϵ -aminocaproic acid) and tranexamic acid are used to inhibit overt fibrinolysis, which act by blocking the lysine-binding site on plasmin.⁶

Protamine sulfate reverses the anticoagulant effects of heparin by binding to it. It is a highly cationic peptide. It binds to heparin to form a stable ion pair which does not have anticoagulant activity.

Specific goal-directed transfusion, transfusion algorithms

Algorithms display decision-making treatment processes and problem-solving strategies by giving clearly defined and formalized guidelines. With the help of clinical algorithms, highly complex processes such as the management of the bleeding can be translated into a clearly structured, logical pathway.⁷² Specific goal-directed transfusion can be achieved by using such algorithms, leading to a reduction of blood component used and to a possible better outcome by stopping bleeding in trauma early.

At our institution, we recently implemented a transfusion algorithm for massively bleeding patients (Figure 1). This algorithm incorporates information obtained from patient's history, clinical presentation, routine coagulation laboratory and bed-site viscoelastic coagulation testings. Interestingly, our experience implementing and adhering to a transfusion algorithm is in accordance with previous studies showing significant reduction in number of transfusion administered, decrease of blood loss and

costs^{27,72,73}. For example, in the first 6 months after implementation of the algorithm, the use of FFP dropped by approximately 50% and RBC as well as platelet administration decreased by approximately 20% each. Despite a moderate increase in costs for point-of-care coagulation monitoring and more frequent administration of specific coagulation factor concentrates (fibrinogen, factor XIII, PCC, rFVIIa), we had a significant cost savings in this first 6 months after implementing our algorithm.

Adverse effects of transfusion

The administration of blood products is associated with adverse effects⁷⁴ including viral or bacterial⁷⁵ transmission. This is perceived to be under control in developed countries⁷⁶. However, there are other issues. RBC transfusion has been found to be a highly significant and independent factor for an increased mortality and morbidity in a variety of surgical situations⁷⁷. Also nosocomial infections are several fold more frequent in transfused vs. non-transfused patients.^{45,78,79} Transfusion-related acute lung injury (TRALI) is another risk^{80,81} occurring during or within 6 hours of transfusion⁸². The risk for TRALI is estimated to be 1 in 1000 to 1 in 4000 units transfused with a significant mortality⁸². Immune suppression or modulation potentially associated with multi organ failure (MOF) and an increased cancer recurrence is another adverse effect of allogeneic blood transfusions⁸³⁻⁸⁵, particularly with long storage times.⁸⁶

Differences between civilian and non civilian sector

In recent publications of the US military the use of fresh whole blood has been described for soldiers with life threatening injuries. Spinella found that surviving improved 48

hours and 30 days after whole blood transfusion compared to massive transfusion of stored red cells.^{87,88} The risk of transmission of infectious agents and other microorganisms remains higher for fresh whole blood than RBCs. In the civilian sector such protocols thus are not applicable.

Trauma exsanguination protocols like the one of Cotton⁸⁹ try to create whole blood conditions by transfusion of fixed amounts of RBCs, FFP and platelets. However, this does not seem to be an adequate strategy in civilian trauma management, the goal to achieve must be a specific management and replacement of blood components and factors according to on-line bed-side coagulation monitoring which may not be feasible in a military environment.

Conclusion

Although a lot of progress has been made in the field of trauma patients, treatment of massive bleeding still remains an interdisciplinary challenge for surgeons and anesthesiologists. Modern and future transfusion strategies are based on on-line bed-side coagulation monitoring with specific goal-directed administration of anti-fibrinolytics, coagulation factors, RBC, fresh frozen plasma and platelets to optimize coagulation early. This allows improving patient's outcome, minimizing exposure to blood products and reducing costs.

Kommentar [OMT6]: 50-100
Wörter, es sind 73

References

1. Sauaia A, Moore FA, Moore EE, Moser KS, Brennan R, Read RA, Pons PT: Epidemiology of trauma deaths: a reassessment. *J Trauma* 1995; 38: 185-93
2. Baker CC, Oppenheimer L, Stephens B, Lewis FR, Trunkey DD: Epidemiology of trauma deaths. *Am J Surg* 1980; 140: 144-50
3. Shackford SR, Mackersie RC, Holbrook TL, Davis JW, Hollingsworth-Fridlund P, Hoyt DB, Wolf PL: The epidemiology of traumatic death. A population-based analysis. *Arch Surg* 1993; 128: 571-5
4. Salim A, Hadjizacharia P, DuBose J, Brown C, Inaba K, Chan L, Margulies DR: Role of anemia in traumatic brain injury. *J Am Coll Surg* 2008; 207: 398-406
5. Malone DL, Dunne J, Tracy JK, Putnam AT, Scalea TM, Napolitano LM: Blood transfusion, independent of shock severity, is associated with worse outcome in trauma. *J Trauma* 2003; 54: 898-907
6. Spahn DR, Cerny V, Coats TJ, Duranteau J, Fernandez-Mondejar E, Gordini G, Stahel PF, Hunt BJ, Komadina R, Neugebauer E, Ozier Y, Riddez L, Schultz A, Vincent JL, Rossaint R: Management of bleeding following major trauma: a European guideline. *Crit Care* 2007; 11: R17
7. Rossaint R, Cerny V, Coats TJ, Duranteau J, Fernandez-Mondejar E, Gordini G, Stahel PF, Hunt BJ, Neugebauer E, Spahn DR: Key issues in advanced bleeding care in trauma. *Shock* 2006; 26: 322-31
8. Shoemaker WC, Appel PL, Bishop MH: Temporal patterns of blood volume, hemodynamics, and oxygen transport in pathogenesis and therapy of postoperative adult respiratory distress syndrome. *New Horiz* 1993; 1: 522-37
9. American College of Surgeons: Committee on Trauma: Advanced Trauma Life Support® for Doctors. 7th ed. 2004
10. Brohi K, Singh J, Heron M, Coats T: Acute traumatic coagulopathy. *J Trauma* 2003; 54: 1127-30
11. Brohi K, Cohen MJ, Ganter MT, Matthay MA, Mackersie RC, Pittet JF: Acute traumatic coagulopathy: initiated by hypoperfusion: modulated through the protein C pathway? *Ann Surg* 2007; 245: 812-8
12. Brohi K, Cohen MJ, Ganter MT, Schultz MJ, Levi M, Mackersie RC, Pittet JF: Acute coagulopathy of trauma: hypoperfusion induces systemic anticoagulation and hyperfibrinolysis. *J Trauma* 2008; 64: 1211-7
13. Brohi K, Cohen MJ, Davenport RA: Acute coagulopathy of trauma: mechanism, identification and effect. *Curr Opin Crit Care* 2007; 13: 680-5
14. Cohen MJ, Brohi K, Ganter MT, Manley GT, Mackersie RC, Pittet JF: Early coagulopathy after traumatic brain injury: the role of hypoperfusion and the protein C pathway. *J Trauma* 2007; 63: 1254-61
15. Scalea TM, Hartnett RW, Duncan AO, Atweh NA, Phillips TF, Sclafani SJ, Fuortes M, Shaftan GW: Central venous oxygen saturation: a useful clinical tool in trauma patients. *J Trauma* 1990; 30: 1539-43

16. Schmelzer TM, Perron AD, Thomason MH, Sing RF: A comparison of central venous and arterial base deficit as a predictor of survival in acute trauma. *Am J Emerg Med* 2008; 26: 119-23
17. Bannon MP, O'Neill CM, Martin M, Ilstrup DM, Fish NM, Barrett J: Central venous oxygen saturation, arterial base deficit, and lactate concentration in trauma patients. *Am Surg* 1995; 61: 738-45
18. Davis JW, Parks SN, Kaups KL, Gladen HE, O'Donnell-Nicol S: Admission base deficit predicts transfusion requirements and risk of complications. *J Trauma* 1996; 41: 769-74
19. Davis JW, Kaups KL, Parks SN: Base deficit is superior to pH in evaluating clearance of acidosis after traumatic shock. *J Trauma* 1998; 44: 114-8
20. Abramson D, Scalea TM, Hitchcock R, Trooskin SZ, Henry SM, Greenspan J: Lactate clearance and survival following injury. *J Trauma* 1993; 35: 584-8; discussion 588-9
21. Andel D, Kamolz LP, Roka J, Schramm W, Zimpfer M, Frey M, Andel H: Base deficit and lactate: early predictors of morbidity and mortality in patients with burns. *Burns* 2007; 33: 973-8
22. Kamolz LP, Andel H, Schramm W, Meissl G, Herndon DN, Frey M: Lactate: early predictor of morbidity and mortality in patients with severe burns. *Burns* 2005; 31: 986-90
23. Ciavarella D, Reed RL, Counts RB, Baron L, Pavlin E, Heimbach DM, Carrico CJ: Clotting factor levels and the risk of diffuse microvascular bleeding in the massively transfused patient. *Br J Haematol* 1987; 67: 365-8
24. Dzik WH: Predicting hemorrhage using preoperative coagulation screening assays. *Curr Hematol Rep* 2004; 3: 324-30
25. Ganter MT, Hofer CK: Coagulation monitoring: current techniques and clinical use of viscoelastic point-of-care coagulation devices. *Anesth Analg* 2008; 106: 1366-75
26. Kheirabadi BS, Crissey JM, Deguzman R, Holcomb JB: In vivo bleeding time and in vitro thrombelastography measurements are better indicators of dilutional hypothermic coagulopathy than prothrombin time. *J Trauma* 2007; 62: 1352-9; discussion 1359-61
27. Spalding GJ, Hartrumpf M, Sierig T, Oesberg N, Kirschke CG, Albes JM: Cost reduction of perioperative coagulation management in cardiac surgery: value of "bedside" thrombelastography (ROTEM). *Eur J Cardiothorac Surg* 2007; 31: 1052-7
28. Barrowcliffé TW, Cattaneo M, Podda GM, Bucciarelli P, Lussana F, Lecchi A, Toh CH, Hemker HC, Beguin S, Ingerslev J, Sorensen B: New approaches for measuring coagulation. *Haemophilia* 2006; 12 Suppl 3: 76-81
29. Hill DA, West RH, Roncal S: Outcome of patients with haemorrhagic shock: an indicator of performance in a trauma centre. *J R Coll Surg Edinb* 1995; 40: 221-4
30. Hoyt DB, Bulger EM, Knudson MM, Morris J, Ierardi R, Sugerman HJ, Shackford SR, Landercasper J, Winchell RJ, Jurkovich G, et al.: Death in the operating room: an analysis of a multi-center experience. *J Trauma* 1994; 37: 426-32
31. Stern SA: Low-volume fluid resuscitation for presumed hemorrhagic shock: helpful or harmful? *Curr Opin Crit Care* 2001; 7: 422-30

32. Practice guidelines for perioperative blood transfusion and adjuvant therapies: an updated report by the American Society of Anesthesiologists Task Force on Perioperative Blood Transfusion and Adjuvant Therapies. *Anesthesiology* 2006; 105: 198-208
33. Madjdpour C, Spahn DR: Allogeneic red blood cell transfusions: efficacy, risks, alternatives and indications. *Br J Anaesth* 2005; 95: 33-42
34. Spahn DR, Madjdpour C: Physiologic transfusion triggers: do we have to use (our) brain? *Anesthesiology* 2006; 104: 905-6
35. McIntyre L, Hebert PC, Wells G, Fergusson D, Marshall J, Yetisir E, Blajchman MJ: Is a restrictive transfusion strategy safe for resuscitated and critically ill trauma patients? *J Trauma* 2004; 57: 563-8
36. Earley AS, Gracias VH, Haut E, Sicoutris CP, Wiebe DJ, Reilly PM, Schwab CW: Anemia management program reduces transfusion volumes, incidence of ventilator-associated pneumonia, and cost in trauma patients. *J Trauma* 2006; 61: 1-5; discussion 5-7
37. Dai B, Wang L, Djaiani G, Mazer CD: Continuous and discontinuous cell-washing autotransfusion systems. *J Cardiothorac Vasc Anesth* 2004; 18: 210-7
38. Kocian R, Spahn DR: Haemoglobin, oxygen carriers and perioperative organ perfusion. *Best Pract Res Clin Anaesthesiol* 2008; 22: 63-80
39. Stanworth SJ, Brunskill SJ, Hyde CJ, McClelland DB, Murphy MF: Is fresh frozen plasma clinically effective? A systematic review of randomized controlled trials. *Br J Haematol* 2004; 126: 139-52
40. Dara SI, Rana R, Afessa B, Moore SB, Gajic O: Fresh frozen plasma transfusion in critically ill medical patients with coagulopathy. *Crit Care Med* 2005; 33: 2667-71
41. Ho AM, Karmakar MK, Dion PW: Are we giving enough coagulation factors during major trauma resuscitation? *Am J Surg* 2005; 190: 479-84
42. Gajic O, Dzik WH, Toy P: Fresh frozen plasma and platelet transfusion for nonbleeding patients in the intensive care unit: benefit or harm? *Crit Care Med* 2006; 34: S170-3
43. Toy P, Popovsky MA, Abraham E, Ambruso DR, Holness LG, Kopko PM, McFarland JG, Nathens AB, Silliman CC, Stronck D: Transfusion-related acute lung injury: definition and review. *Crit Care Med* 2005; 33: 721-6
44. Buddeberg F, Schimmer BB, Spahn DR: Transfusion-transmissible infections and transfusion-related immunomodulation. *Best Pract Res Clin Anaesthesiol* 2008; 22: 503-17
45. Sarani B, Dunkman WJ, Dean L, Sonnad S, Rohrbach JI, Gracias VH: Transfusion of fresh frozen plasma in critically ill surgical patients is associated with an increased risk of infection. *Crit Care Med* 2008; 36: 1114-8
46. Norda R, Tynell E, Akerblom O: Cumulative risks of early fresh frozen plasma, cryoprecipitate and platelet transfusion in Europe. *J Trauma* 2006; 60: S41-5
47. Ganter MT, MD, , Schmuck S, MD,, Hamiel CR, BS,, Wischmeyer PE, MD,, Heule D, MD, Zollinger A, MD, , Hofer CK, MD: Monitoring Recombinant Factor VIIa Treatment: Efficacy Depends on High Levels of Fibrinogen in a Model of Severe Dilutional Coagulopathy. *Journal of Cardiothoracic and Vascular Anesthesia* 2008; 22: 675-680

48. Monroe DM, Hoffman M: What does it take to make the perfect clot? *Arterioscler Thromb Vasc Biol* 2006; 26: 41-8
49. Innerhofer P, Fries D, Margreiter J, Klingler A, Kuhbacher G, Wachter B, Oswald E, Salner E, Frischhut B, Schobersberger W: The effects of perioperatively administered colloids and crystalloids on primary platelet-mediated hemostasis and clot formation. *Anesth Analg* 2002; 95: 858-65
50. Stinger HK, Spinella PC, Perkins JG, Grathwohl KW, Salinas J, Martini WZ, Hess JR, Dubick MA, Simon CD, Beekley AC, Wolf SE, Wade CE, Holcomb JB: The ratio of fibrinogen to red cells transfused affects survival in casualties receiving massive transfusions at an army combat support hospital. *J Trauma* 2008; 64: S79-85
51. Haas T, Fries D, Holz C, Innerhofer P, Streif W, Klingler A, Hanke A, Velik-Salchner C: Less impairment of hemostasis and reduced blood loss in pigs after resuscitation from hemorrhagic shock using the small-volume concept with hypertonic saline/hydroxyethyl starch as compared to administration of 4% gelatin or 6% hydroxyethyl starch solution. *Anesth Analg* 2008; 106: 1078-86
52. Haas T, Fries D, Velik-Salchner C, Reif C, Klingler A, Innerhofer P: The in vitro effects of fibrinogen concentrate, factor XIII and fresh frozen plasma on impaired clot formation after 60% dilution. *Anesth Analg* 2008; 106: 1360-5
53. Mittermayr M, Streif W, Haas T, Fries D, Velik-Salchner C, Klingler A, Oswald E, Bach C, Schnapka-Koepf M, Innerhofer P: Hemostatic changes after crystalloid or colloid fluid administration during major orthopedic surgery: the role of fibrinogen administration. *Anesth Analg* 2007; 105: 905-17
54. Schroeder V, Chatterjee T, Kohler HP: Influence of blood coagulation factor XIII and FXIII Val34Leu on plasma clot formation measured by thrombelastography. *Thromb Res* 2001; 104: 467-74
55. Nielsen VG, Kirklin JK, Hoogendoorn H, Ellis TC, Holman WL: Thrombelastographic method to quantify the contribution of factor XIII to coagulation kinetics. *Blood Coagul Fibrinolysis* 2007; 18: 145-50
56. Nielsen VG, Steenwyk BL, Gurley WQ: Contact activation prolongs clot lysis time in human plasma: role of thrombin-activatable fibrinolysis inhibitor and Factor XIII. *J Heart Lung Transplant* 2006; 25: 1247-52
57. Egbring R, Kroniger A, Seitz R: Factor XIII deficiency: pathogenic mechanisms and clinical significance. *Semin Thromb Hemost* 1996; 22: 419-25
58. Nielsen VG, Gurley WQ, Jr., Burch TM: The impact of factor XIII on coagulation kinetics and clot strength determined by thrombelastography. *Anesth Analg* 2004; 99: 120-3
59. Gerlach R, Raabe A, Zimmermann M, Siegemund A, Seifert V: Factor XIII deficiency and postoperative hemorrhage after neurosurgical procedures. *Surg Neurol* 2000; 54: 260-4
60. Godje O, Haushofer M, Lamm P, Reichart B: The effect of factor XIII on bleeding in coronary surgery. *Thorac Cardiovasc Surg* 1998; 46: 263-7
61. Ansell J, Hirsh J, Poller L, Bussey H, Jacobson A, Hylek E: The pharmacology and management of the vitamin K antagonists: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126: 204S-233S
62. Baglin TP, Keeling DM, Watson HG: Guidelines on oral anticoagulation (warfarin): third edition--2005 update. *Br J Haematol* 2006; 132: 277-85

63. Kohler M, Hellstern P, Lechler E, Uberfuhr P, Muller-Berghaus G: Thromboembolic complications associated with the use of prothrombin complex and factor IX concentrates. *Thromb Haemost* 1998; 80: 399-402
64. Evans G, Luddington R, Baglin T: Beriplex P/N reverses severe warfarin-induced overanticoagulation immediately and completely in patients presenting with major bleeding. *Br J Haematol* 2001; 115: 998-1001
65. Lorenz R, Kienast J, Otto U, Kiehl M, Schreiter D, Haertel S, Barthels M: Successful emergency reversal of phenprocoumon anticoagulation with prothrombin complex concentrate: a prospective clinical study. *Blood Coagul Fibrinolysis* 2007; 18: 565-70
66. Bruce D, Nokes TJ: Prothrombin complex concentrate (Beriplex P/N) in severe bleeding: experience in a large tertiary hospital. *Crit Care* 2008; 12: R105
67. Enomoto TM, Thorborg P: Emerging off-label uses for recombinant activated factor VII: grading the evidence. *Crit Care Clin* 2005; 21: 611-32
68. Ahonen J, Jokela R: Recombinant factor VIIa for life-threatening post-partum haemorrhage. *Br J Anaesth* 2005; 94: 592-5
69. Butwick AJ, Riley ET, Ahonen J, Jokela R: Recombinant factor VIIa for life-threatening post-partum haemorrhage. *Br J Anaesth* 2005; 95: 558; author reply 558
70. Boffard KD, Riou B, Warren B, Choong PI, Rizoli S, Rossaint R, Axelsen M, Kluger Y: Recombinant factor VIIa as adjunctive therapy for bleeding control in severely injured trauma patients: two parallel randomized, placebo-controlled, double-blind clinical trials. *J Trauma* 2005; 59: 8-15
71. Vincent JL, Rossaint R, Riou B, Ozier Y, Zideman D, Spahn DR: Recommendations on the use of recombinant activated factor VII as an adjunctive treatment for massive bleeding--a European perspective. *Crit Care* 2006; 10: R120
72. Steiner ME, Despotis GJ: Transfusion algorithms and how they apply to blood conservation: the high-risk cardiac surgical patient. *Hematol Oncol Clin North Am* 2007; 21: 177-84
73. O'Keefe T, Refaai M, Tchorz K, Forestner JE, Sarode R: A massive transfusion protocol to decrease blood component use and costs. *Arch Surg* 2008; 143: 686-90; discussion 690-1
74. Klein HG, Spahn DR, Carson JL: Red blood cell transfusion in clinical practice. *Lancet* 2007; 370: 415-26
75. Pomper GJ, Wu Y, Snyder EL: Risks of transfusion-transmitted infections: 2003. *Curr Opin Hematol* 2003; 10: 412-8
76. Goodnough LT: Risks of blood transfusion. *Anesthesiol Clin North America* 2005; 23: 241-52
77. Spahn DRM, H. Hofmann, A. Isbister, J.P.: Patient Blood Management – the pragmatic solution for the problems with blood transfusions. *Anesthesiology* 2008; 109
78. Claridge JA, Sawyer RG, Schulman AM, McLemore EC, Young JS: Blood transfusions correlate with infections in trauma patients in a dose-dependent manner. *Am Surg* 2002; 68: 566-72
79. Taylor RW, O'Brien J, Trottier SJ, Manganaro L, Cytron M, Lesko MF, Arnzen K, Cappadoro C, Fu M, Plisco MS, Sadaka FG, Veremakis C: Red blood cell

transfusions and nosocomial infections in critically ill patients. *Crit Care Med* 2006; 34: 2302-8; quiz 2309

80. Rana R, Fernandez-Perez ER, Khan SA, Rana S, Winters JL, Lesnick TG, Moore SB, Gajic O: Transfusion-related acute lung injury and pulmonary edema in critically ill patients: a retrospective study. *Transfusion* 2006; 46: 1478-83

81. Khan H, Belsher J, Yilmaz M, Afessa B, Winters JL, Moore SB, Hubmayr RD, Gajic O: Fresh-frozen plasma and platelet transfusions are associated with development of acute lung injury in critically ill medical patients. *Chest* 2007; 131: 1308-14

82. Toy P, Lowell C: TRALI--definition, mechanisms, incidence and clinical relevance. *Best Pract Res Clin Anaesthesiol* 2007; 21: 183-93

83. Amato A, Pescatori M: Perioperative blood transfusions for the recurrence of colorectal cancer. *Cochrane Database Syst Rev* 2006: CD005033

84. Moore FA, Moore EE, Sauaia A: Blood transfusion. An independent risk factor for postinjury multiple organ failure. *Arch Surg* 1997; 132: 620-5

85. Opelz G, Sengar DP, Mickey MR, Terasaki PI: Effect of blood transfusions on subsequent kidney transplants. *Transplant Proc* 1973; 5: 253-9

86. Koch CG, Li L, Sessler DI, Figueroa P, Hoeltge GA, Mihaljevic T, Blackstone EH: Duration of red-cell storage and complications after cardiac surgery. *N Engl J Med* 2008; 358: 1229-39

87. Spinella PC: Warm fresh whole blood transfusion for severe hemorrhage: U.S. military and potential civilian applications. *Crit Care Med* 2008; 36: S340-5

88. Spinella PC, Perkins JG, Grathwohl KW, Repine T, Beekley AC, Sebesta J, Jenkins D, Azarow K, Holcomb JB: Fresh whole blood transfusions in coalition military, foreign national, and enemy combatant patients during Operation Iraqi Freedom at a U.S. combat support hospital. *World J Surg* 2008; 32: 2-6

89. Cotton BA, Gunter OL, Isbell J, Au BK, Robertson AM, Morris JA, Jr., St Jacques P, Young PP: Damage control hematology: the impact of a trauma exsanguination protocol on survival and blood product utilization. *J Trauma* 2008; 64: 1177-83