



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

From clonal to sexual hybrids: Genetic recombination via triploids in all-hybrid populations of water frogs

Christiansen, D G; Reyer, H U

Abstract: Speciation via interspecific hybrids is very rare in animals, as compared to plants. Whereas most plants overcome the problem of meiosis between different chromosome sets by tetraploidization, animal hybrids often escape hybrid sterility by clonal reproduction. This comes at the expense of genetic diversity and the ability to purge deleterious mutations. However, here we show that all-hybrid populations of diploid (LR) and triploid (LLR and LRR) water frogs (*Pelophylax esculentus*) have secondarily acquired sexual reproduction. First, in a crossing experiment analyzed with microsatellite markers, triploid hybrids of both sexes and genotypes (LLR and LRR) recombined their homospecific genomes. Second, the great majority of natural populations investigated had low multilocus linkage disequilibrium, indicating a high recombination rate. As predicted from mating system models, the L genome had constant, low levels of linkage disequilibrium, whereas linkage disequilibrium in the R genome showed a significant reduction with increasing proportion of recombining triploids. This direct evidence of sexual reproduction in *P. esculentus* calls for a change of the conventional view of hybridogens as clonally reproducing diploids. Rather, hybridogens can be independent sexually reproducing units with an evolutionary potential.

DOI: <https://doi.org/10.1111/j.1558-5646.2009.00673.x>

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-24667>
Journal Article

Originally published at:

Christiansen, D G; Reyer, H U (2009). From clonal to sexual hybrids: Genetic recombination via triploids in all-hybrid populations of water frogs. *Evolution*, 63(7):1754-1768.
DOI: <https://doi.org/10.1111/j.1558-5646.2009.00673.x>

1 This article was published in *Evolution* 63-7: 1754–1768,
2 <http://www3.interscience.wiley.com/journal/122213330/abstract>
3 The definitive version is available at www.blackwell-synergy.com

4

5 **From clonal to sexual hybrids: genetic recombination via triploids**
6 **in all-hybrid populations of water frogs**

7

8

9 **Running title:** From clonal to sexual hybrid water frogs

10

11

12

13 **Ditte G. Christiansen^{1,2} and Heinz-Ulrich Reyer^{1,3}**

14

15 1) Ecology, Zoological Institute, University of Zurich, Winterthurerstrasse 190, CH-8057
16 Zurich, Switzerland.

17 2) E-mail: ditte.christiansen@zool.uzh.ch

18 3) E-mail: uli.reyer@zool.uzh.ch

19

20

21 **Speciation via interspecific hybrids is very rare in animals, as compared to plants.**

22 **Whereas most plants overcome the problem of meiosis between different chromosome**

23 **sets by tetraploidization, animal hybrids often escape hybrid sterility by clonal**

24 **reproduction. This comes at the expense of genetic diversity and the ability to purge**

25 **deleterious mutations. However, here we show that all-hybrid populations of diploid**

26 (LR) and triploid (LLR and LRR) water frogs (*P. esculentus*) have secondarily acquired
27 sexual reproduction. First, in a crossing experiment analyzed with microsatellite
28 markers, triploid hybrids of both sexes and genotypes (LLR and LRR) recombined their
29 homospecific genomes. Second, the great majority of natural populations investigated
30 had low multilocus linkage disequilibrium, indicating a high recombination rate. As
31 predicted from mating system models, the L genome had constant, low levels of linkage
32 disequilibrium, while linkage disequilibrium in the R genome showed a significant
33 reduction with increasing proportion of recombining triploids. This direct evidence of
34 sexual reproduction in *P. esculentus* calls for a change of the conventional view of
35 hybridogens as clonally reproducing diploids. Rather, hybridogens can be independent
36 sexually reproducing units with an evolutionary potential.

37

38 **Key words:** polyploidy, hybridogenesis, microsatellites, *Pelophylax esculentus*, *Rana*
39 *esculenta*

40

41 Hybridization instantly creates individuals with a new genetic composition and is therefore a
42 potentially powerful force in evolution (Wissemann 2007; Jiggins et al. 2008). Whether
43 hybridization leads to speciation depends on the hybrids' ability to survive and reproduce
44 (Arnold and Hodges 1995; Barton 2001; Chapman and Burke 2007). Two reproductive
45 challenges need to be overcome for establishment of new hybrid taxa: First, the hybrids must
46 be fertile, in spite of having two dissimilar chromosome sets which might interrupt meiosis
47 (Arnold and Hodges 1995; Chapman and Burke 2007). Second, the hybrids must be spatially
48 or reproductively isolated from the parental species (Wang et al. 2001; James and Abbott
49 2005; Chapman and Burke 2007).

50 Normal meiosis, as well as reproductive isolation, can instantly be restored by
51 tetraploidization. Via this process, hybridization has had a large impact on plant evolution

52 (Arnold 1997; Hegarty and Hiscock 2005; Wissemann 2007), while in animals, remarkably
53 few examples of tetraploid speciation are known (Orr 1990; Otto and Whitton 2000). In
54 animal hybrids, fertility and reproductive isolation are, however, often established by different
55 kinds of clonal reproduction which may or may not be accompanied by polyploidy. Among
56 clonal vertebrates, reptiles are parthenogenetic, while fishes and amphibians depend on sperm
57 from a sexual species for initiating embryogenesis (Vrijenhoek et al. 1989): In gynogenetic
58 taxa, the sperm activates, but usually does not fertilize the eggs. In hybridogenetic taxa,
59 fertilization takes place; yet, there is normally no recombination between the parental
60 genomes. This is because the paternal genome is usually excluded from the germ line prior to
61 meiosis while the remaining maternal genome is transmitted clonally (Dawley 1989).
62 Hybridity is restored each generation by matings with the paternal species. The hybrids' soma
63 is thus made up by both the sexual paternal and the clonal maternal genome, while the
64 hybrids' germ line contain only the latter.

65 For hybrid speciation to be of evolutionary importance, a third factor is crucial:
66 genetic recombination. Genetic recombination via sexual reproduction enhances genetic
67 diversity and is generally agreed to convey three important benefits: One, high genetic
68 diversity is required for defence against fast evolving parasites (Red Queen hypothesis,
69 Hamilton 1980). Two, the combination of beneficial mutations from different individuals
70 enhances the efficiency of selection (Fisher 1930; e.g. Colegrave 2002; Cooper 2007). Three,
71 and most importantly, the combination of deleterious mutations allows their purging from the
72 population (Muller 1932; e.g. Vrijenhoek 1994). Without recombination, clonal lines are
73 predicted to accumulate deleterious mutations via Muller's ratchet, which will eventually lead
74 to their extinction.

75 As a consequence of their clonal reproduction modes, parthenogenetic, gynogenetic
76 and hybridogenetic hybrid animal taxa lack the above mentioned advantages of genetic
77 diversity and the ability to purge mutations. Hence, they are generally considered to be

78 “evolutionary dead ends”, at least as far as individual lineages are concerned (e.g. Vrijenhoek
79 et al. 1989; Maynard Smith 1992). In agreement with this, strictly clonal taxa are, with very
80 few exceptions (Butlin 2002), distributed as short-lived tips on the tree of life mainly
81 comprised of sexual taxa (Simon et al. 2003).

82 However, at least genetic diversity seems to be higher in clonally reproducing taxa
83 than previously assumed, and various mechanisms have been described how this can be
84 achieved. First, clonal hybrids often arise recurrently from different progenitors. Hence, they
85 have a high genetic diversity possibly enabling them to fit different ecological niches (frozen
86 niche variation hypothesis, Vrijenhoek 1984). Recurrent origin of clonal and polyploid sexual
87 lineages is known from several plants (Soltis and Soltis 1999) and also from animals,
88 including ostracods (Little and Hebert 1997), fishes (Janko et al. 2003; Pala and Coelho
89 2005), reptiles (Moritz et al. 1989) and some anurans (Ptacek et al. 1994; Stöck et al. 2005).
90 Second, some allegedly asexual organisms are not strictly clonal but occasionally incorporate
91 new nuclear material from a sexual host (Hedges et al. 1992; Spolsky et al. 1992; Scharl et al.
92 1995). The most recent discovery of such a mechanism is “kleptogenesis” in unisexual
93 salamanders of the genus *Ambystoma* (Bogart et al. 2007): all-female lines can incorporate
94 (parts of) nuclear genomes from sperm from sympatric sexual species and presumably later
95 discard other parts of the genome. Third, in bisexual hybridogenetic species, like the edible
96 frog, *Pelophylax esculentus* (called *Rana esculenta* until Frost et al. 2006), and the Iberian
97 minnow, *Squalius alburnoides*, hybrid x hybrid matings lead to offspring with parental
98 genotype (cf. Fig. 1b). Although rarely viable, these offspring could recombine the otherwise
99 clonal genomes if they succeed in reproducing (Hotz et al. 1992; Alves et al. 1998; Vorburger
100 2001c). While the existence of these three mechanisms can not be denied, their potential for
101 lifting the doom of “evolutionary dead end” from the relevant hybrid taxa is subject to
102 discussion.

103 Here we investigate the potential for systematic and frequent sexual reproduction in
104 hybridogens through a mechanism called meiotic hybridogenesis. The term refers to the
105 possibility that in polyploid hybridogens of the general type AAB, the homospecific
106 chromosome sets from one parental species, A, recombine in a normal meiosis, whereas the
107 set from the other parental species, B, is discarded (Alves et al. 1998). Preferential pairing of
108 homologous chromosomes and elimination of the unmatched chromosomes has been shown
109 for a number of triploid fish and frog hybrids (see Morishima et al. 2008 and references
110 therein) but, so far, clear evidence for recombination through meiotic hybridogenesis comes
111 from one species only: the Iberian minnow, *S. alburnoides* (Crespo-Lopez et al. 2006).

112 It might be argued that meiotic hybridogenesis is a rare and special phenomenon
113 without much general relevance for the role of hybrids in animal evolution. However, meiotic
114 hybridogenesis is interesting as a newly discovered possibility for hybridogenetic hybrids to
115 obtain recombination in a regular, non-accidental way. Besides, the list of taxa with meiotic
116 hybridogenesis will surely grow: Firstly, with the increasing application of molecular tools to
117 organisms from different populations, the list of known hybridogens has grown recently and
118 is likely to grow further. Secondly, since hybridogenesis was originally discovered in the
119 diploid topminnow, *Poeciliopsis monacha-lucida* (Schulz 1969), polyploidy in hybridogens
120 increasingly appears to be the rule, rather than the exception. At present, polyploidy is known
121 from four of the six genera with hybridogenesis. Water frogs (*Pelophylax*, Berger 1967),
122 Iberian minnows (*Squalius*, Carmona et al. 1997), spined loaches (*Cobitis*, Saitoh et al. 2004)
123 and Oriental weatherloaches (*Misgurnus*, Morishima et al. 2008) exhibit polyploidy while
124 only hybridogenetic topminnows (*Poeciliopsis*) and stick insects (*Bacillus*, Bullini and
125 Nascetti 1990) are purely diploid. Polyploidy is also known from a hybridogenesis-related
126 mode of reproduction in the Batura toad (*Bufo viridis* complex, Stöck et al. 2002). We are
127 thus just at the beginning of discovering the diversity and implications of hybrid reproduction
128 modes.

129 Hence, investigating the extent of recombination during meiotic hybridogenesis and
130 thus the long-term evolutionary potential for intraspecific hybrids seems timely and
131 potentially relevant for more species than presently assumed. The edible frog, *Pelophylax*
132 *esculentus*, provides a particularly interesting system for such an investigation, because it is
133 the only hybrid yet known also to form self-sustaining, hybridogenetic, all-hybrid
134 populations. In the absence of the parental species, meiotic hybridogenesis is the sole
135 potential source of frequent recombination and could thus be of crucial evolutionary
136 importance for these populations. Moreover, *P. esculentus* comes in various mating systems
137 and, hence, offers an opportunity to study successive stages of incipient hybrid speciation.

138

139 **The *Pelophylax esculentus* systems**

140 *Pelophylax esculentus* (*Rana esculenta*) originated, and still originates, from interspecific
141 matings between the two sexual water frog species, *P. lessonae* (the pool frog, genotype LL)
142 and *P. ridibundus* (the marsh frog, genotype RR). The parental species, as well as the diploid
143 *P. esculentus* hybrid with the genomic composition LR, have wide distributions in Europe. In
144 the western part of this distribution area, LR excludes the L genome from the germ line prior
145 to meiosis and transmits the R genome to the gametes clonally. As a result, matings between
146 hybrids yield RR offspring, but these typically die due to homozygosity for deleterious
147 mutations in the clonal R genome (Vorburger 2001a and references therein; Guex et al. 2002).
148 In order to form a new generation of hybrid LR, *P. esculentus* is dependent on L gametes
149 obtained from mating with *P. lessonae* (LE system, Fig. 1). In parts of Eastern Europe the
150 pattern is reversed: hybrid LR excludes the R genome, produces L gametes and, therefore,
151 lives in sympatry and mates with *P. ridibundus* (RE system). In both of these diploid systems,
152 (reviewed by Graf and Polls Pelaz 1989) *P. esculentus* face disadvantages with respect to both
153 of its genomes: the one in the hybrid's germ line is clonal, while the other, sexual, genome
154 must for every generation be obtained by mating with the parental species. Various LE, RE

155 systems and *lessonae-esculentus-ridibundus* populations with both diploid and triploid *P.*
156 *esculentus* also exist (Günther 1991; Tunner and Heppich-Tunner 1992; Rybacki and Berger
157 2001), but unfortunately hardly anything is known about how these diverse and complicated
158 populations function.

159 The present study focuses on all-hybrid populations of *P. esculentus* (EE system) that,
160 by definition, live and reproduce without any of the parental species. Thus, the propagation of
161 both parental genomes, as well as any recombination within them, must be undertaken by
162 hybrids alone. All-hybrid populations are found in large areas of Denmark, southern Sweden
163 north-eastern Germany, and patchily in northern Poland and probably a few localities in south
164 eastern Europe (Mikulíček and Kotlík 2001; Rybacki and Berger 2001; Christiansen et al.
165 2005; Arioli 2007 cap. 5); (reviewed by Plötner 2005). These populations consist of diploid
166 (LR) and one or two types of triploid hybrids (LLR and LRR). LLR frogs of both sexes
167 provide L gametes while LRR make R gametes. Within the diploid LR, all males and some
168 females produce R gametes, while all females and a few males make unreduced LR gametes
169 yielding new triploids upon fusion with haploid gametes (Graf and Polls Pelaz 1989;
170 Christiansen et al. 2005; Arioli 2007 cap. 1; Jakob 2007 cap. 5). Sex determination is an xx-
171 xy system with a dominant male-determining y factor. The y factor is supposed to be present
172 in the L genome only (Graf and Polls Pelaz 1989; Berger and Günther 1991-1992), which
173 means that L genomes are either L_x or L_y while all R genomes are R_x . As a consequence, LLR
174 and LR come in both sexes, while the great majority of LRR are females (Jakob 2007 cap. 2
175 and the present study). In this way, the mix of di- and triploid hybrid frogs form self-
176 sustaining populations producing all gametes needed for a new generation of similar
177 composition (Fig. 1). Non-hybrid LL and RR offspring are also formed, but die off in natural
178 ponds during the tadpole stage (Arioli 2007 cap. 3).

179 In all three systems, clonally propagated *P. esculentus* genomes face the risk of
180 mutation accumulation. In the LE and RE systems, some accumulation can be tolerated, as the

181 clonal genome is constantly paired with a healthy parental genome in the hemiclinal hybrids
182 (confirmed in LE by Vorburger 2001b). Nevertheless, the lifespan of the clonal genomes in
183 diploid systems appears limited, as old clones are likely to become inviable or replaced by
184 new genomes that were more recently derived from primary hybridization between the
185 parental species. In the all-hybrid populations, the situation was, so far, unknown. It was often
186 assumed that the LLR recombine their two L chromosome sets after exclusion of the R, and
187 that, likewise, LRR recombine their two R sets after exclusion of the L genome (Günther et
188 al. 1979; Graf and Polls Pelaz 1989; Som and Reyer 2006a). Under this assumption, the all-
189 hybrid populations might be functionally sexual with a higher evolutionary potential than
190 diploid LE and RE system populations. However, experimental evidence for recombination in
191 triploids is scarce and controversial, due to low availability of polymorphic genetic markers.
192 Based on allozyme and sex data, Günther et al. (1979) probably found recombination in one
193 Polish LRR male (table 5, cross 25/26). Furthermore, Arioli (2007 cap. 1), using
194 microsatellite analysis on Swedish frogs, detected recombination in an LRR female, but not in
195 an LLR male. While these data demonstrate the capability of triploids to recombine, it
196 remains unclear whether recombination happens as a rule or as an exception and whether
197 there are sex- and/or genotype- (LLR vs. LRR) specific differences in the recombination rate.

198 Here we present the first crossing experiment with a sufficient number of frogs (30)
199 and polymorphic genetic markers (18) to conclude that intragenomic recombination takes
200 place in triploids of both sexes and genotypes (LLR and LRR). We also provide previously
201 unpublished microsatellite primers and new multiplex PCR protocols.

202 Confirming recombination in triploids does, however, not suffice to conclude that all-
203 hybrid populations are functionally sexual. Therefore, assessment of the impact of triploid-
204 mediated recombination on the genetic structure of the L and R genomes in wild populations
205 was needed. One might expect populations with many triploids to be highly recombined and
206 thus have low multilocus linkage disequilibrium. This, however, should be true only for the R

207 genome; not for the L genome. The reason for the difference is that R gametes can originate
208 from both recombining LRR and non-recombining LR frogs whereas L gametes come from
209 recombining LLR frogs alone (see Fig. 1). The monopoly on L gamete production guarantees
210 LLR frogs a large and constant reproductive contribution to the next generation and, hence,
211 should result in high recombination rates of L genomes, irrespective of the LLR/(LLR+LR)
212 ratio. This prediction was previously confirmed by a mathematical model (Som and Reyer
213 2006a), but empirical data are lacking. In contrast, recombination rates of R genomes should,
214 on average, be lower but increase with LRR/(LRR+LR) ratios. For this prediction, neither
215 theoretical nor empirical studies were available.

216 Here we show that linkage disequilibrium was low in a large sample of natural
217 populations from across the Danish and Swedish range, indicating that natural recombination
218 rates are sufficiently high for these all-hybrid populations to be functionally sexual. We also
219 provide evidence for the expected correlations between linkage disequilibrium and population
220 structure. Finally, we confirm that pond-specific influences and method-specific biases were
221 without importance for these results. In conclusion, the all-hybrid populations are an example
222 of a hybridogen that, in a unique way, has become an independent evolutionary unit with
223 sexual reproduction and thus a long-term evolutionary potential.

224

225 **Methods**

226 **Overview**

227 The study was carried out on Swedish and Danish all-hybrid populations, because these are
228 geographically isolated from populations with parental species (Christiansen et al. 2005;
229 Jakob 2007 cap. 2).

230 For direct evidence of whether triploids recombine, adult frogs were sampled,
231 genotyped and crossed and the offspring were reared and genotyped. Then, segregation and
232 linkage analyses were performed on the inheritance pattern of the microsatellite alleles

233 analyzed. Absence of linkage between the majority of loci, when compared pairwise, would
234 indicate recombination.

235 For investigating the level of recombination in natural populations, frogs were
236 sampled in ponds with different proportions of clone-propagating diploid (LR) and
237 recombining triploid frogs (LLR and LRR). All individuals were genotyped, and the
238 multilocus linkage disequilibria in the L and R genomes were calculated as r_d^- for each pond
239 separately. Low r_d^- values would indicate high levels of recombination. The effects of
240 genome, population structure, pond-specific effects and method-specific biases on r_d^- were
241 investigated to test the predictions outlined in the introduction and to test suspicions of
242 artifacts. Finally, F statistics were calculated, because non-random mating, resulting in high
243 F_{IS} values, would also affect r_d^- .

244

245 **Crosses**

246 Genetic variation in Swedish and Danish *P. esculentus* is very low (Christiansen et al. 2005;
247 Arioli 2007 cap. 4). To obtain genetic data at multiple heterozygous loci for linkage analysis,
248 it was therefore necessary to: 1) Screen both published and unpublished microsatellites for
249 polymorphism in Scandinavia and design multiplex PCRs with the final selection of 18
250 primer pairs. 2) Select the most heterozygous triploids of a large sample of frogs for crossing.
251 3) Raise the larvae to an age where offspring genotypes could be inferred reliably when the
252 heterozygous parents shared an allele. When parents share one allele, alleles or even whole
253 chromosomes missing in the offspring can lead to misinterpretation of the parental
254 contributions. This was of real concern, because many young larvae are aneuploid, i.e. they
255 have mixed, uninterpretable genotypes with extra or missing alleles (Christiansen et al. 2005
256 and unpublished data from the present study). Raising the larvae to metamorphosis ensured
257 that most aneuploid offspring died off and did not enter the analyses.

258 Crossing and rearing took place at Stensoffa Field Station, Scania, Sweden. Between
259 May 12 and 22, 2006, i.e. after their emergence from hibernation and before breeding, 269
260 frogs were caught at night using flashlight and dip net in one of the Danish (Alsønderup in
261 Christiansen et al. 2005) and 10 of the Swedish ponds included in the investigation of natural
262 populations described below. The frogs were marked individually with a transponder (Trovan
263 ID101, Euro I.D., DE), toe-clipped for DNA analysis, and kept at approximately 7° C while
264 the DNA samples were sent to the University of Zürich and analyzed for genome composition
265 (LLR, LR, LRR) and heterozygosity (at LL in LLR and RR in LRR). The triploids with most
266 heterozygous loci were preferred for the crossings, because recombination can only be
267 assessed from combinations of heterozygous loci. This preference made a balanced design of
268 source ponds impossible. Since males were more common than females among LLR frogs
269 and females were predominant among LRR frogs, LLR males and LRR females were picked
270 from a larger sample and were therefore more heterozygous than LLR females and LRR
271 males. Most L genome data therefore derived from males and most R genome data from
272 females.

273 Six crossing tables were designed, each having 3-4 females and 5-6 males including at
274 least one LLR, LR and LRR female and at least one LLR, two LR and one LRR male.
275 Substitute frogs were added if the sperm or egg quality looked suboptimal. All females were
276 crossed with all males within the same crossing table, so that all frogs were crossed to all
277 genotypes (half-sib design).

278 Offspring were produced on May 30, 2006 by artificial fertilization as described by
279 Berger *et al.* (1994). Sperm solutions from the testes of hormone-injected males were
280 distributed into 3-5 petri dishes per male. Eggs were then gently squeezed out of the hormone-
281 treated females and dropped directly into the individual sperm solutions of the 5-6 different
282 males, in small portions and in random order. The following day, the egg clumps were
283 transferred to 1 liter tubs with 1-2 cm of water and subdivided for better oxygen supply.

284 The water was changed every 2-4 days and the egg jelly was removed after hatching.
285 On June 12, when most tadpoles had just reached the feeding stage, 15 healthy-looking
286 tadpoles (or fewer, if 15 were not available) from each sibship were randomly selected for
287 rearing in 40 liters outdoors tubs covered with mesh lids allowing air and sunlight through,
288 but keeping predators out. Algae growing on the insides of the tubs, supplemented with rodent
289 pellets, ensured food *ad libitum*. Filamentous algae were regularly removed, fowling water
290 exchanged and *Daphnia sp.* added for good water quality. The tadpoles metamorphosed from
291 July 18 onwards. Slow-growing tadpoles were eventually moved indoors into smaller tubs,
292 where the last ones metamorphosed in mid October. Offspring that died early during rearing
293 disappeared, while offspring that died as metamorphs or nearly metamorphosing tadpoles
294 were attempted DNA-analyzed although they were sometimes rotten. In total, 1628 tadpoles
295 were selected for rearing, DNA samples were obtained from 1487 offspring (91%), and 1463
296 offspring (90%) were successfully genotyped.

297

298 **Natural populations**

299 Population structure was investigated in 54 Danish and 12 Swedish ponds from mid May to
300 mid August 2005. The Danish ponds were chosen as pairs of ecologically distinct ponds,
301 maximally 5 km apart, from across the area of distribution. At each location, approximately
302 30 frogs (predominately adults) were caught at night with flashlight and dip net, were
303 measured and had a toe tip cut for DNA analysis before being returned to their pond.

304 The Swedish ponds constituted 11 ecologically variable ponds in the center of the
305 small distribution area in Scania, Southern Sweden, and one from a satellite population near
306 Malmö, 18 km west of the others ("core ponds" in Jakob 2007 cap. 2). The Swedish ponds
307 were sampled as described above, but in both May and August, and the frogs were
308 additionally marked with a transponder for individual identification. The Swedish samples are
309 thus the sum of different individuals from the two catching rounds.

310 In total, 2296 Danish and Swedish frogs were caught and genotyped.

311

312 **Laboratory protocols**

313 DNA from the ethanol-stored toe-tips was extracted with Qiagen BioSprint 96 DNA Blood
314 Kit following Qiagen's protocol for tissue extraction. All samples were subjected to two
315 PCRs with nine primer pairs each. The reactions were of 5 µl and contained 0.8 µl DNA
316 extraction, 2.5 µl Qiagen Multiplex PCR Master mix and 1.7 µl primer mix. PCR 1 contained
317 primers Res16, Res20 (Zeisset et al. 2000), RICA5, RICA1b5 (Garner et al. 2000), Ca1b6,
318 Ga1a19, Re2CAGA3 (Arioli 2007 cap. 4), RICA2a34 and Rrid064A (table 1). PCR 2
319 contained Res22 (Zeisset et al. 2000), RICA18 (Garner et al. 2000), Rrid013A (Hotz et al.
320 2001), Rrid059A redesigned (Hotz et al. 2001 and table 1: forward primer redesigned to
321 extend the fragment amplified by 177 base pairs), Re1CAGA10 (Arioli 2007 cap. 4),
322 RICA1a27, ReGA1a23, Rrid169A and Rrid135A (table 1). Both forward and reverse primers
323 appeared in 0.1 µM (or rarer 0.2 µM) in the PCR. Of the forward primers, 8-40% were color
324 labeled with FAM, VIC, NED or PET. PCR 1 was given 15 min of initial denaturation at 95°
325 C, 30 cycles of 30s at 94° C, 90s at 57° C and 60s at 72° C and a final extension of 30 min at
326 60° C. PCR 2 was run similarly, but with 31 cycles with 60° C in stead of 57° C. 0.7µl of the
327 PCR products were run on an ABI 3730 Avant capillary sequencer with internal size standard
328 (GeneScan-500 LIZ) and the alleles were scored with the Genemapper software
329 (Applied_Biosystems 2004).

330

331 **Genotyping**

332 All samples were analyzed with 18 primer pairs amplifying loci in either the L genome, the R
333 genome or both. The 18 primers were scored at a total of 13 loci in each genome. With some
334 primers, genome specificity changed slightly with PCR conditions, i.e. typically
335 monomorphic L-specific alleles could arise or disappear beside the R allele(s) according to

336 annealing temperature or primer concentrations. However, monomorphic loci conveyed no
337 information of importance for the present study, and the choice of scoring or leaving out
338 particular loci for technical reasons would not bias the data on homozygosity/heterozygosity
339 which was the focus of this study.

340 All alleles scored were specific to either the L or the R genome. Allele specificity was
341 confirmed in *P. lessonae*, *P. esculentus* and *P. ridibundus* from Estonia, Latvia and Lithuania
342 (unpublished data), in non-hybrid LL and RR offspring from the crossings and through the
343 distribution of L and R specific alleles on LLR, LR and LRR frogs. Preliminary data from
344 German and Swiss samples indicated, however, that in these more southern populations with
345 higher genetic polymorphism, certain alleles were not genome-specific.

346 Four of the primer pairs (Res16, R1CA1b5, Ca1b6 and Ga1a19) amplifying both L and
347 R specific alleles were used to distinguish LLR, LR and LRR frogs by dosage effect, i.e. by
348 the relative intensities (peak heights) of the L and R alleles amplified (see Christiansen 2005).
349 L:R peak heights were evaluated separately per 96-well PCR, both per locus and per allele
350 combination within that locus. The great majority of the L:R peak height ratios clustered into
351 discrete groups corresponding to the LLR, LR and LRR genotypes. Samples producing
352 intermediate or extreme L:R ratios were subjected to repeated PCR analyses until each of the
353 four dosage effect loci clearly signaled LLR, LR or LRR. Assignment to LLR, LR or LRR
354 was thus determined independently at four loci. In non-hybrid offspring (LLL, LL, RR, RRR)
355 the peak height ratios of heterozygous L or R loci were used to determine ploidy in the same
356 way as just described. Not all loci and allele combinations proved diagnostic, but most did.

357 Samples that repeatedly gave conflicting results on genotype, i.e. had extra or missing
358 alleles at particular loci, were classified as mixed genotypes. Mixed genotypes, constituted
359 3.6% of the crossing experiment offspring and 2.1% (2.7% inclusive null alleles, see below)
360 of the natural pond samples and were excluded from data sets where the relevant loci could
361 not be scored unambiguously.

362 Null alleles, i.e. alleles missing according to the overall ploidy of the individual, can
363 be a nuisance in population genetics, because in high frequencies they bias estimates of allele
364 frequencies and heterozygosity. However, in this study, they were generally not a problem, as
365 they were often directly detectable and occurred in low frequencies only. The adults used for
366 crossings carried no problematic null alleles, as the analyses were made on the loci where
367 they were heterozygous for real alleles. Spontaneously missing alleles in mutant crossing
368 experiment offspring, as well as null alleles in the frogs from the natural populations, were all
369 directly detectable at the four dosage effect loci, and on average half of them were unmasked
370 and detectable in a hemizygous state at the remaining loci. For example, a null allele at an L
371 locus without dosage effect would be masked in LLR frogs but unmasked in LR and LRR
372 frogs. Individuals with detected null alleles were handled as mixed genotypes described
373 above. Only in two ponds was the same locus found missing in more than two frogs (i.e. six
374 and eight frogs respectively), indicating that undetected null alleles could occur at potentially
375 problematic frequencies in these ponds. In one of the two ponds, the entire locus was
376 therefore recoded as missing data. In the second pond, all individuals were hemizygous at that
377 locus, so that the null allele could always be detected. It was therefore coded as a real allele.
378 For determining LLR and LRR proportions in the ponds, mixed genotypes were assigned to
379 the most similar euploid genotype.

380

381 **Statistics: crossings**

382 The crossings yielded data from on 30 triploid frogs for segregation and linkage analyses. For
383 males, the analyses were based on 19-58 (mean 41) offspring and for females on 30-86 (mean
384 66) offspring, as females were on average mated to more partners than males.

385 Non-random segregation would indicate selection during the experiment or
386 unexpected genetic mechanisms. To check for random segregation at the heterozygous loci in
387 the parents, offspring allele frequencies were tested with Chi-square tests for homogeneity

388 with Yate's correction for continuity (Fowler and Cohen 1992). To correct for multiple tests
389 ($n = 55$ L and 57 R loci), sequential Bonferroni correction of the P values was calculated
390 according to Holm (1979) in the program MacBonferroni (Watkins 2002).

391 Linkage analysis involves analysis of the inheritance pattern at two loci that are
392 heterozygous in a parent (e.g. $Aa+Bb$). *Without* recombination, all pairs of loci should show
393 complete linkage, i.e. only two of the parent's allele combinations should be observed in the
394 offspring (e.g. $A+B$ and $a+b$). In contrast, *with* recombination all four possible combinations
395 should be found in the offspring ($A+B$, $a+b$, $A+b$ and $a+B$) in approximately equal
396 proportions of 0.25. Intermediate results, where the recombinant allele combinations ($A+b$
397 and $a+B$) are significantly less frequent than the parental ones ($A+B$ and $a+b$), would indicate
398 reduced recombination and would be hard to explain if deriving from the majority the locus
399 pairs. However, a few locus pairs must, by chance, be expected to have reduced or no
400 recombination, due to physical linkage. Linkage was investigated with 2×2 Chi-square tests
401 with Yate's correction for continuity (Fowler and Cohen 1992) for every pairwise
402 combination of loci that were heterozygous in the parent.

403

404 **Statistics: natural populations**

405 The rate of recombination is not easily measured directly. Instead, linkage disequilibrium
406 between multiple genetic markers was used for an indirect measure, as recombination and
407 linkage disequilibrium should be negatively related (see the discussion). Pairwise and
408 multilocus linkage disequilibria in natural populations were calculated as r_d^- , as
409 recommended by Halkett *et al.* (2005). r_d^- is an index of association adjusted for unequal
410 sample size, calculated by the program Multilocus (Agapow and Burt 2001). First, L and R
411 loci were divided into separate datasets. Then, the two homospecific allele sets in triploids
412 were split up into haploid data by recoding all but one randomly chosen heterozygous locus
413 into missing data. Recoding heterozygous loci into missing data is also how Multilocus

414 handles diploid data, according to the documentation file. Calculations were based on 20-71
415 (mean 37) haplotypes in Danish ponds and 56-110 (mean 78) in Swedish ponds. One pond
416 was excluded from the L and another from the R data set because less than our predefined
417 minimum of 20 haploid genotypes had been sampled. Two further ponds were excluded from
418 the L data and eight from the R data because no or only one locus was polymorphic. After
419 that, the genomes had 2-11 variable loci (mean 3.8 for the L and 5.2 for the R), i.e. loci with
420 at least 5 undeleted copies of an alternative allele.

421 Pairwise r^2_d was calculated in order check for locus pairs producing r^2_d values
422 differing significantly from the mean r^2_d of the remaining pairs, when tested pairwise (locus
423 pair in question vs. mean of remaining locus pairs) over all ponds. This pairwise within-pond
424 approach was necessary because overall linkage was expected to differ between ponds.
425 Significantly elevated linkage disequilibria could suggest physical linkage between the loci in
426 question, whereas linkage disequilibria lower than the mean would be difficult to explain.

427 Multilocus r^2_d were calculated for each genome in each pond to test the predicted
428 correlations between recombination and population structure outlined in the introduction. All
429 linear regressions, correlations and t-tests were performed in SPSS (2004). The L and R
430 slopes from the linear regressions were subjected to a test for difference between two
431 regression lines (Fowler and Cohen 1992).

432 The expected relationships between linkage disequilibrium and population structure
433 could be obscured by strong between-pond variation in the forces responsible for linkage
434 disequilibrium, i.e. founder effect, drift, migration and ecological selection on linked loci. If
435 these forces affect the L and R genomes to a similar extent, the magnitude of this problem
436 might be revealed by the degree of correlation between linkage disequilibrium in the L and R
437 genomes in the ponds. To test for such pond-specific effects, we correlated r^2_d values for the
438 L and R genome.

439 Genetic diversity varied between ponds and was generally lower in the L-specific than
440 the R-specific markers. To investigate whether the estimates of multilocus linkage
441 disequilibrium were affected by this variation in genetic diversity, we tested for correlation
442 between r_d^- and genetic diversity measured as expected heterozygosity summed over all loci
443 per genome. Expected heterozygosity was for each locus calculated as $H_E = 1 - (a_1^2 + a_2^2 +$
444 $a_3^2 \dots)$ from allele frequencies (a_1, a_2, a_3 etc) computed by the software, SPAGeDi (see
445 below).

446 As mentioned above, all but one of the heterozygous loci in triploid frogs had to be
447 excluded for the constructing haplotypes for calculating r_d^- . This affected the R genome the
448 most, as its higher genetic diversity resulted in many R-heterozygous LRR frogs. Ponds rich
449 in LRR frogs could thus theoretically have lower r_d^- values as a result of the lower resolution
450 after the exclusion of the many heterozygous loci. To investigate whether r_d^- was affected by
451 the resolution, it was tested whether the r_d^- values for the R genome were correlation with the
452 number of hemizygotes (LLR and LR which had no loci excluded) in the sample they were
453 calculated from.

454 To investigate inbreeding and population structuring, F_{IS} , F_{ST} and F_{IT} were calculated
455 in the program SPAGeDi (Hardy and Vekemans 2002), which accepts a mixture of different
456 ploidy levels. These F statistics were calculated for each genome separately so that with
457 respect to the L genome, LLR provided diploid data while LR and LRR provided haploid
458 data. Similarly, LLR and LR gave haploid R data while LRR gave diploid R data. Excluding
459 all haploid data from the analyses had very little effect on the results, though.

460

461 **Results**

462 **Crosses**

463 Recombination data was obtained from 7 LLR females, 10 LLR males, 7 LRR females and 6
464 LRR males. Due to multiple heterozygosity, most individuals provided data for several

465 pairwise locus combinations. The LLR frogs provided recombination data for a total of 18 out
466 of 21 possible pairwise combinations of 7 polymorphic L loci, and the LRR frogs for 47 of 66
467 possible combinations of 12 polymorphic R loci. All heterozygous loci in these triploids
468 demonstrated random segregation, i.e. none of the allele proportions differed significantly
469 from 0.5 at the 0.05 significance level after sequential Bonferroni correction performed within
470 each genome separately. All triploids produced three or four gamete types per locus pair,
471 corresponding to the two parental types and one or both recombinant types. All triploids thus
472 recombined all their loci, and only for one locus pair were not all four gamete types present.

473 The uncorrected P values for the Chi-square tested frequency distributions of the four
474 possible gamete types per locus pair are shown in Fig. 2. As parental and recombinant
475 gametes were indistinguishable because the genotypes of the parents of the frogs crossed were
476 unknown, insignificant P value deviations from zero do not necessarily imply reduced
477 recombination. Insignificant P value would also have resulted from randomly derived excess
478 of recombinant gametes and from uneven allele frequencies within the expected numbers of
479 parental and recombinant gametes. When considered individually, the $-\log(p)$ values
480 exceeding 1.30 were significant at the 0.05 level. After within-genome sequential Bonferroni
481 correction for the 65 tests in the L genome and the 91 tests in the R genome, however, only
482 four P values were significant. This indicates that the great majority of locus pairs were
483 unlinked and freely recombined.

484 The four locus pairs showing significant linkage occurred in four different frogs
485 (represented by four filled symbol types in Fig. 2) that all produced equilibrium offspring
486 frequencies at their remaining locus pairs. The linkage was therefore rather a property of the
487 loci than of the frogs involved. Unfortunately, replicate data was not obtained for the three
488 locus pairs giving the most significant P values in this study, but the pair with strongest
489 linkage, Re1CAGA10 + R1CA18 (L genome), was the same pair for which Arioli (2007 cap.
490 1) found no recombination. From the 40, 0, 0, 38 gamete frequency distribution in that and the

491 20, 0, 3, 23 gamete frequency distribution in the present study, it can be inferred that that
492 Re1CAGA10 and RICA18 are linked, i.e. situated closely together on the same chromosome.
493 Ca1b6 + Ga1a19 (R genome) had the offspring type distribution 10, 33, 18, 8 and Rrid169 +
494 Rrid059A (R genome) had 36, 16, 7, 27. These locus pairs thus appear weakly linked, but
495 replicate crossings would be needed to confirm linkage. Re2GAGA3 + Rrid135A appeared
496 significantly linked in one female with gamete frequency distribution 12, 13, 20, 17, but
497 unlinked in three other females. Overall, therefore, these two loci appear unlinked. Actually, a
498 mutation happened in the germ line of this female so that some of her offspring had a new
499 allele at locus Re2CAGA3. A rare allele at another locus confirmed that these offspring were
500 indeed hers. The offspring with the new allele were excluded from the analyses involving
501 Re2CAGA3, but when included by pooling the new and the lowest-frequency maternal allele,
502 from which it most probably mutated, all four P values for locus pairs including Re2CAGA3
503 dropped substantially and the significant value became clearly non-significant.

504 Nearly significant P values appeared for several other locus pairs, but also here
505 replicates raised the average for these loci to well above the 0.05 level, rendering no overall
506 indication of linkage. Males and females did not have significantly different mean P values
507 (male mean = 0.356, female mean = 0.285, t-test, $t_{154} = 1.606$, $P = 0.110$). Many species,
508 probably including *P. esculentus* (Burt et al. 1991), have lower crossing-over rates in males
509 than in females, but the present data set can neither confirm or disprove this for *P. esculentus*.

510

511 **Natural populations**

512 Triploids were found in all 55 ponds investigated, and both kinds (LLR and LRR) were found
513 in 82% of the ponds. The proportion of LLR varied from 0-100% while that of LRR varied
514 from 0-86% in the pond samples (Fig. 3). Of the 2296 frogs genotyped, only 0.2% were non-
515 hybrid. These were 5 LL from two Swedish ponds. Multilocus linkage disequilibrium,
516 measured as r^2 on a scale from zero to one, averaged 0.01 in the L genome and 0.11 in the R

517 genome, indicating that both genomes were well recombined in the majority of the natural
518 populations. Mean r_d^- in the R genome was, however, significantly higher than in the L
519 genome (t-test, $t_{111} = -3.819$, $P < 0.001$).

520 Multilocus disequilibrium in the L genome showed no relation with the proportion of
521 LLR individuals (linear regression: $F_{1,61} = 2.269$, $P = 0.137$, $r^2 = 0.036$). In contrast,
522 multilocus disequilibrium in the R genome was negatively associated with the proportion of
523 recombining LRR frogs among the R gamete-producing LR and LRR frogs (linear regression:
524 $F_{1,54} = 9.034$, $P = 0.004$, $r^2 = 0.143$, slope = -0.214). These results were thus fully in
525 accordance with the expectations. The slopes of the L and the R regressions were, however,
526 not significantly different ($t_{115} = 1.440$, $P = 0.153$).

527 The multilocus linkage disequilibria (r_d^-) in the L and the R genomes were not
528 positively correlated within ponds (Fig. 4). In fact, they were significantly negatively
529 correlated (Pearson correlation: $r_{55} = -0.374$, $P = 0.005$); even excluding the L outliers far left
530 and far right in Fig. 4. This indicates an absence of strong pond-specific effects affecting r_d^-
531 in the L and R genome simultaneously.

532 There was no correlation between r_d^- and genetic diversity, measured as the expected
533 heterozygosity summed over all loci (Pearson correlation for L and R data pooled: $r_{119} =$
534 0.013 , $P = 0.889$). The significant difference in mean multilocus disequilibrium between the L
535 and the R genome can therefore not be explained by lower polymorphism in the L specific
536 microsatellite loci, but only by differences in recombination rates. The r_d^- values for the R
537 genome showed also no correlation with the number of hemizygotes in the sample they were
538 calculated from (Pearson correlation: $r_{55} = 0.016$, $P = 0.904$). The significant relation between
539 LRR/(LR+LRR) and r_d^- in the R genome in Fig. 3b can therefore not be explained by
540 exclusion of heterozygous loci in LRR frogs, but must be attributed to differences in
541 recombination rates.

542 An analysis of pairwise r_d^- values showed that only two locus pair had r_d^- values
543 differing significantly from the mean pairwise r_d^- of the remaining locus pairs in the same
544 ponds (28 L and 63 R, paired t-tests with sequential Bonferroni correction within each
545 genome separately). These two locus pairs (L loci Res20 + Re1CAGA10 and the R loci
546 RICA1b5 + Rrid064A) both had significantly lower r_d^- than the remaining loci. Thus, most
547 locus pairs gave similar results within ponds and none gave elevated values suggesting
548 linkage. In spite of the tight linkage in the crossing experiment, pairwise r_d^- for Re1CAGA10
549 + RICA18 was not significantly different from the mean, even without Bonferroni correction
550 (paired t-test: $t_9 = 1.311$, $P = 0.222$). The same applies to the two potentially linked locus pairs
551 (Ca1b6 + Ga1a19: $t_{10} = -0.654$, $P = 0.528$; Rrid169 + Rrid059A: $t_{22} = 1.261$, $P = 0.221$).
552 Therefore, these three (potentially) linked locus pairs were not excluded from the analyses of
553 natural populations.

554 Global F_{IS} was very low in both the L and R genome, i.e. -0.007 and -0.008,
555 respectively, indicating random mating within ponds. Global F_{ST} values were rather high, i.e.
556 0.4561 and 0.6156 in the L and R genome, respectively, indicating much genetic structure
557 among ponds, which is in accordance with the expectations for a low-mobility animal. As a
558 consequence of the low F_{IS} , F_{IT} was very similar to F_{ST} for both genomes.

559

560 **Discussion**

561 Recombination was demonstrated in all 30 frogs tested in the crossing experiment including
562 both males and females of both LLR and LRR. As a consequence of such triploid-mediated
563 recombination, natural populations were found to have low multilocus linkage disequilibria.
564 In agreement with predictions from the asymmetrical propagation of L and R genomes in the
565 all-hybrid populations, L genomes were generally fully recombined while R genomes were
566 recombined according to the proportion of LRR triploids. The unique all-hybrid populations
567 of *P. esculentus* are thus functionally sexual; actually, they represent an obligate symbiosis of

568 two independent, functionally sexual genomes: the L and the R genome. Below, we will first
569 describe the genetic mechanisms underlying these results and then outline the evolutionary,
570 conceptual and conservation-political implications for all-hybrid populations and
571 hybridogenetic taxa.

572

573 **Recombination in all-hybrid populations**

574 In normal meiosis, the combined effects of random segregation of chromosomes and
575 chromosomal crossing-over assure equal proportions of parental and recombinant gametes for
576 most locus pairs. Reduced recombination rates due to physical linkage are, however, observed
577 between loci situated so closely together on the same chromosome that there is small
578 probability of crossing-over between them. A random sample of genetic markers for any kind
579 of organism might thus include a small proportion of linked loci. *P. esculentus* has 13
580 chromosomes per L or R set (e.g. Koref-Santibanez and Günther 1980). The physical
581 locations of our microsatellite loci on these chromosomes are unknown, but the results from
582 the crossing experiment suggested linkage between three of the 65 locus pairs investigated.
583 Loci Re1CAGA10 and R1CA18 showed strong linkage in a male crossed by us as well as in
584 one crossed by Arioli (2007 cap. 1); thus it can be inferred that these two loci are situated
585 close together. The apparent linkage of the two remaining locus pairs in this study was weaker
586 and assessed in only one frog each, so that linkage should not be concluded without further
587 verification. This discovery of one to three linked loci does not suggest variation in
588 recombination rates among individuals, as the three frogs with apparently linked loci had full
589 recombination at their remaining locus pairs investigated.

590 Selection took place in the crossing experiment, as dead and sick-looking tadpoles
591 were not reared, and 10% of the offspring chosen for rearing eluded genotyping – mainly by
592 dying. Only selection on the interaction of non-neutral loci linked to our markers could,
593 however, have affected the recombination results. Any such interaction effects were reduced

594 by crossing every parent to several mates of different genotypes. As no significant bias in the
595 segregation at any single locus was detected, bias of locus combinations by selection is
596 unlikely. Furthermore, selection on the interaction of linked non-neutral loci would most
597 likely bias the results towards less recombination, so it would not undermine the conclusion
598 of recombination.

599 Unlike linkage, linkage disequilibrium can arise between loci without physical
600 associations. Linkage disequilibrium, measured as r_d^- in the natural populations, is the net
601 result of generating and deteriorating forces. Linkage disequilibrium-generating forces
602 include founder effect, migration, drift, inbreeding and selection on linked genes, called
603 hitchhiking (Hedrick 2005). In clonal organisms, the entire genome hitchhikes with positively
604 selected genes. The deteriorating force is recombination. Linkage disequilibrium is a negative
605 linear function of recombination rate per generation, with half of the disequilibrium
606 disappearing per generation at 100% recombination (Hedrick 2005). Provided that r_d^- is a
607 good measure of linkage disequilibrium, that $LRR/(LR+LRR)$ was a fair substitute for
608 recombination frequency in the ponds, and that disequilibrium-generating forces did not
609 depend on population structure (e.g. on $LRR/(LR+LRR)$), linear relationships were therefore
610 expected in Fig. 3.

611 With low r_d^- irrespective of population structure in the L genome (Fig. 3a) and a
612 negative relationship between r_d^- and recombining triploids ($LRR/LR+LRR$) in the R genome
613 (Fig. 3b), the expectations outlined in the introduction were met. According to the model by
614 Som and Reyer (2006a), L genomes spend 2/3 of their generations in LLR frogs and 1/3 in
615 LR frogs, which means that they are recombined two out of three generations. The empirical
616 data from the present study shows that this recombination rate of 2/3, whatever the type and
617 strength of linkage disequilibrium-generating forces in the natural populations, is sufficient to
618 reduce r_d^- values to around zero (mean $r_d^- = 0.01$ on the scale from zero to one). For the R
619 genome, no theoretical model is available. Before a reliable model can be made, more

620 empirical data on the ratio of R and LR gametes produced by LR females and LR males is
621 needed, as this ratio is important for population dynamics and has been shown to vary
622 strongly between individuals and locations (Tunner and Heppich-Tunner 1991; Polls Pelaz
623 1994; Mikulíček and Kotlík 2001; Rybacki and Berger 2001; Christiansen et al. 2005; Arioli
624 2007 cap. 1; Jakob 2007 cap. 5). Central to such a model is also the question of why
625 populations vary in structure. Although it is commonly accepted that population structure of
626 *P. esculentus*, *P. lessonae* and/or *P. ridibundus* depend on ecological components (Pagano et
627 al. 2001; Holenweg Peter et al. 2002; Plötner 2005), attempts to identify the ecological
628 components determining population structure in Swedish all-hybrid populations were so far
629 rather inconclusive (Jakob 2007 cap. 3). In the absence of theoretical models, it was not
630 known what level of linkage disequilibrium to expect in the R genome of natural populations,
631 but the present empirical data show that it is generally low (mean $r_d^- = 0.11$), although the
632 genetic signature of clonal reproduction was visible in certain populations with few LRR
633 frogs. In clonal populations of other organisms, r_d^- values have been found to be considerably
634 higher than in the present study (e.g. Goyeau et al. 2007; Grundmann et al. 2008).
635 Unfortunately, no thorough studies on multilocus disequilibrium in the *R. esculenta* LE or RE
636 system have been conducted yet.

637 The variation not explained by the linear relations in Fig. 3 is expected to derive from
638 three main sources. 1) Error on the estimate of r_d^- from a random sample of 17-86 (mean 35)
639 individuals. 2) Error on the estimate of population structure, e.g. LRR/(LR+LRR), from the
640 same random sample and between-pond-variation in the ratio of R gametes from LR frogs. If
641 the proportion of R gametes made by LR frogs varies between ponds, this will add further
642 noise. 3) Between-pond variation in the strength of the various disequilibrium-generating
643 forces listed above. The combined effects of these three sources explain the rather large
644 variation for the R genome in Fig. 3b. For the L genome, population structure (source 2)
645 should have no relevance, however. Furthermore, if the recombination rate is so high that it

646 always overpowers the local disequilibrium-generating forces (source 3), as seems to be the
647 case in the L genome, variation comes only from the error on the estimate of r_d^- (source 1).
648 This explains the relative low variation in Fig. 3a. Unfortunately, disequilibrium-generating
649 forces are difficult to measure. The only disequilibrium-generating force, we could measure in
650 this study was inbreeding. The low F_{IS} values obtained indicated random mating, so that
651 inbreeding would have little effect on r_d^- . We did, however, test for those pond-specific
652 effects that affect the L and R genome similarly. The lack of a positive correlation between
653 r_d^- in the L and R genome across ponds (Fig. 4) indicates that such forces were absent. In
654 conclusion, the forces generating multilocus linkage disequilibrium in the natural populations
655 could not be indentified, but between-pond variation in their strength and composition did not
656 pose a problem in this study. On the contrary: the good match of observed with expected
657 relations in Fig. 3a and b shows that r_d^- can be a useful tool in studies of recombination.

658 The extreme positive outlier in Fig. 3a calls for a different explanation than those
659 given for residual variation. This explanation has to apply to the L genome only, as the high L
660 r_d^- value was not matched by a high R value (Fig. 4). Notably, in this pond, a null allele was
661 scored as a real L allele, because it did not pose a technical problem. As pairwise r_d^- values
662 were elevated for all locus pairs in this pond, the null allele cannot account for its outlier
663 status, however. Exclusion of the locus with the null allele reduced r_d^- to 0.29, i.e. the point
664 remained an outlier although less extreme. A better explanation for the high r_d^- value can be
665 derived from the pond's extreme left position in the Figure. Although necessary for
666 reproduction, LLR frogs were absent from our sample of 23 adults. Also notable, although not
667 exceptional for this pond, was that the population appeared small with few males, which are
668 more often LLR than females. We could therefore speculate that the L genomes in the
669 sampled frogs derived from very few LLR ancestors. A linkage disequilibrium in the L
670 genome caused by such a bottleneck in LLR frogs would persist for several generations of
671 recombination.

672

673 **Evolutionary consequences**

674 Triploids are not restricted to all-hybrid populations, but have been found in various
675 population types in Germany (Günther 1975), Poland (Rybacki and Berger 2001) and France
676 (Regnier and Neveu 1986). The ability to make diploid eggs giving rise to triploid individuals
677 provides all these *P. esculentus* populations with genetic recombination and potential
678 reproductive independence - two important steps in the direction of speciation. Where hybrids
679 live sympatrically with parental species, they do not reproduce independently, however, but
680 interbreed with the parental species. Here, recombination by triploids might be of little
681 genetic importance to the hybrids, because they can be supplied with recombined genomes
682 from the parental species. In contrast, the all-hybrid populations of Denmark and southern
683 Sweden must rely on recombination in triploids only, as they are isolated from the nearest
684 parental populations by sea or large stretches of uninhabited land, and non-hybrid LL and RR
685 offspring only very rarely survive to sexual maturity (Christiansen et al. 2005; Jakob 2007
686 cap. 2 and the present study). Here, *P. esculentus* has truly accomplished the transition from a
687 clonal, gamete-dependent hybrid to an independent, sexually reproducing evolutionary unit.

688 Although the all-hybrid populations have a combination of clonal and sexual
689 reproduction, the low multilocus linkage disequilibrium values indicate that the loci of natural
690 populations were well mixed. Selection should thus have the whole range of genetic
691 combinations to work on, enabling beneficial, as well as harmful, mutations to be combined
692 for fast adaption to changing environments (Fisher 1930) and for purging of deleterious
693 mutations (Muller 1932). This hybridogenetic reproduction mode also ensures continuous
694 genetic variation as a defense against fast evolving parasites (Red Queen hypothesis,
695 Hamilton 1980), since the combination of recombined and clonal gametes result in unique
696 individuals. The all-hybrid populations thus seem to have all the advantages of sexual
697 reproduction, including a long-term evolutionary potential. The ability of fast adaption to

698 changing environments might, however, be of more importance for the survival of *P.*
699 *esculentus*, given that habitat loss and climate change increasingly threaten amphibians
700 worldwide (Stuart et al. 2004).

701 It remains to be analyzed to what extent all-hybrid *P. esculentus* populations can also
702 benefit from the clonal reproduction of diploids. In general, potential benefits of clonal
703 reproduction include the possibility to save the costs of producing males and the ability to
704 propagate favorable gene combinations (Otto and Gerstein 2006). In all-hybrid *P. esculentus*
705 populations, the theoretical offspring sex ratio is only slightly female biased which is in
706 agreement with the mean observed adult sex ratio in large surveys (Som and Reyer 2006b;
707 Jakob 2007 cap. 2 and the present study). Thus, only a few percent of the cost of males might
708 be saved. Recombination takes place after maximum one generation in the L genome (Som
709 and Reyer 2006a) and after one to a few generations in the R genome, suggesting that
710 favorable gene combinations are not be preserved for long, unless physically linked.
711 Therefore, the benefit that all-hybrid populations of *P. esculentus* can potentially derive from
712 the clonal component in their reproduction appears small - in contrast to cyclical
713 parthenogens, such as aphids, rotifers, water fleas that have successfully combined the
714 advantages of sexual and clonal reproduction (Innes and Singleton 2000).

715 With sexual reproduction, the death of newly formed non-hybrid LL and RR in the all-
716 hybrid populations is intriguing, because it cannot be attributed to clonal propagation of the
717 genomes, as in the LE system. In the LE system, RR die because recessive deleterious
718 mutations have become fixed in the clonally propagated R genome of the diploid LR hybrids
719 (Vorburger 2001a; Guex et al. 2002). These deleterious mutations were either acquired
720 through Muller's ratchet or were already present at hemiclone formation (Vorburger 2001a).
721 In all-hybrid EE populations, both genomes are regularly recombined in triploid individuals,
722 the L when in LLR and the R when in LRR. Hence, fixation of deleterious mutations by
723 Muller's ratchet is unlikely, yet fixation may still have occurred by other mechanisms, for

724 example founder effect. Fixation and low genetic diversity is certainly observed at
725 microsatellite loci (Christiansen et al. 2005; Arioli 2007 cap. 4 and the present study).
726 Explanations for how genetic diversity became and remained this low in spite of the presence
727 of parental species just south of the German and north of the Swedish all-hybrid populations
728 are, however, lacking.

729 *P. esculentus* most closely resembles the Iberian minnow, *Squalius alburnoides* (also
730 called *Leuciscus*, *Rutilus* and *Tropidophoxinellus*, reviewed by Alves et al. 2001) of other
731 hybridogenetic taxa known: both hybrids often form mixed populations of di- and polyploid
732 hybrids and one or both parental genotypes. All-hybrid di- and triploid populations are,
733 however, not known from *S. alburnoides*. In stead, tetraploids occur in many *S. alburnoides*
734 populations and, in special habitats, tetraploids can constitute 73% of the mixed populations.
735 These tetraploids have an even sex ratio, have normal meiosis, produce tetraploid offspring
736 when mating with each other and appear to be reproductively isolated from other ploidy levels
737 (Cunha et al. 2008). The discovery of these mainly tetraploid populations strongly suggests
738 that meiotic hybridogenesis can act as a stepping stone to tetraploidization and ultimately to
739 speciation. In *P. esculentus*, tetraploidy has so far only been found in very low frequencies in
740 Swedish populations (Jakob 2007 cap. 2).

741 Given that recombination appears to be the rule in polyploid hybridogens and that
742 polyploidy in hybridogenetic taxa appears to be more common than previously assumed, the
743 prevailing view of hybridogens as clonally reproducing diploids may have to be changed.
744 Should the discoveries of hybridogenetic breeding systems continue to increase, which is
745 likely as more and more supposedly normal species are being genetically analyzed, this will
746 also affect our perception of the importance of hybridization for speciation in animals.

747 Studies on hybrids are also relevant from a conservation point of view. Modern man-
748 agement concepts stress the importance of conserving “evolutionary significant units”
749 (ESUs), i.e. populations representing significant adaptive variation; but how these units are to

750 be identified, is strongly debated (reviewed by Crandall et al. 2000). Hybrids, for instance, are
751 exempt from protection, because they do not seem to constitute independent evolutionary
752 lineages (Kraus 1995). While this may be true for F₁ progeny from many interspecific
753 matings, it is not true for parthenogenetic, gynogenetic and hybridogenetic taxa of hybrid
754 origin, which are capable of self propagation (Ranker and Arft 1994; Kraus 1995). This, plus
755 the finding that hybridogens like *P. esculentus* and *S. alburnoides* can form independent and
756 sexually reproducing populations, makes these organisms evolutionary significant units and
757 worthy of protection.

758

759 **Acknowledgements**

760 This study would not have been possible without the invaluable help from Lars Iversen,
761 Ursina Tobler and Eline Embrechts with field work, and from Sandra Röthlisberger with
762 laboratory work. Unpublished primers were provided by Trenton T. Garner and Hansjürg
763 Hotz, whom we warmly thank for letting us publish them. Ecogenics (Zurich) helped testing
764 primers in Danish and Swedish samples. Our acknowledgements also go to Josh Van Buskirk,
765 Peter Wandeler, Lukas Keller, Christian Mayer and anonymous reviewers for valuable inputs
766 to the methods and the manuscript. The study was supported by the Swiss National Science
767 Foundation (grant no. 31-64004.00 to H.-U. Reyer). Permits for catching frogs, toe-clipping
768 frogs and rearing tadpoles were obtained from the Danish and Swedish authorities (Skov- og
769 Naturstyrelsen SN 2001-441-0252, SNS-441-00047, Länsstyrelsen I Skaane Län 522-10481-
770 05, Djurskyddsmyndigheten M62-05, M71-06).

771

772 **References**

773 Agapow, P. M., and A. Burt. 2001. Indices of multilocus linkage disequilibrium. *Molecular*
774 *Ecology Notes* 1:101-102.

775 Alves, M. J., M. M. Coelho, and M. J. Collares-Pereira. 1998. Diversity in the reproductive
776 modes of females of the *Rutilus alburnoides* complex (Teleostei, Cyprinidae): A way
777 to avoid the genetic constraints of uniparentalism. *Molecular Biology and Evolution*
778 15:1233-1242.

779 Alves, M. J., M. M. Coelho, and M. J. Collares-Pereira. 2001. Evolution in action through
780 hybridisation and polyploidy in an Iberian freshwater fish: a genetic review. *Genetica*
781 111:375-385.

782 Applied Biosystems. 2004. Genemapper vers. 3.7.

783 Arioli, M. 2007. Reproductive patterns and population genetics in pure hybridogenetic water
784 frog populations of *Rana esculenta*. PhD thesis. University of Zurich, Ecology
785 department. Available at www.dissertationen.unizh.ch

786 Arnold, M. L. 1997. *Natural hybridization and evolution*. Oxford University Press, New
787 York.

788 Arnold, M. L., and S. A. Hodges. 1995. Are natural hybrids fit or unfit relative to their
789 parents? *Trends in Ecology & Evolution* 10:67-71.

790 Barton, N. H. 2001. The role of hybridization in evolution. *Molecular Ecology* 10:551-568.

791 Berger, L. 1967. Embryonal and larval development of F1 generation of green frogs different
792 combinations. *Acta Zoologica Cracoviensia*:123-162.

793 Berger, L., and R. Günther. 1991-1992. Inheritance patterns of water frog males from the
794 environments of nature reserve Steckby, Germany. *Zoologica Poloniae* 37:87-100.

795 Berger, L., M. Rybacki, and H. Hotz. 1994. Artificial fertilization of water frogs. *Amphibia-*
796 *Reptilia* 15:408-413.

797 Bogart, J. P., K. Bi, J. Z. Fu, D. W. A. Noble, and J. Niedzwiecki. 2007. Unisexual
798 salamanders (genus *Ambystoma*) present a new reproductive mode for eukaryotes.
799 *Genome* 50:119-136.

800 Bullini, L., and G. Nascetti. 1990. Speciation by hybridization in phasmids and other insects.
801 Canadian Journal of Zoology 68:1747-1760.

802 Burt, A., G. Bell, and P. H. Harvey. 1991. Sex-differences in recombination. Journal of
803 Evolutionary Biology 4:259-277.

804 Butlin, R. 2002. The costs and benefits of sex: new insights from old asexual lineages. Nature
805 Reviews Genetics 3:311-317.

806 Carmona, J. A., O. I. Sanjur, I. Doadrio, A. Machordom, and R. C. Vrijenhoek. 1997.
807 Hybridogenetic reproduction and maternal ancestry of polyploid Iberian fish: The
808 *Tropidophoxinellus alburnoides* complex. Genetics 146:983-993.

809 Chapman, M. A., and J. M. Burke. 2007. Genetic divergence and hybrid speciation. Evolution
810 61:1773-1780.

811 Christiansen, D. G. 2005. A microsatellite-based method for genotyping diploid and triploid
812 water frogs of the *Rana esculenta* hybrid complex. Molecular Ecology Notes 5:190-
813 193.

814 Christiansen, D. G., K. Fog, B. V. Pedersen, and J. J. Boomsma. 2005. Reproduction and
815 hybrid load in all-hybrid populations of *Rana esculenta* water frogs in Denmark.
816 Evolution 59:1348-1361.

817 Colegrave, N. 2002. Sex releases the speed limit on evolution. Nature 420:664-666.

818 Cooper, T. F. 2007. Recombination speeds adaptation by reducing competition between
819 beneficial mutations in populations of *Escherichia coli*. Plos Biology 5:1899-1905.

820 Crandall, K. A., O. R. P. Bininda-Emonds, G. M. Mace, and R. K. Wayne. 2000. Considering
821 evolutionary processes in conservation biology. Trends in Ecology & Evolution
822 15:290-295.

823 Crespo-Lopez, M. E., T. Duarte, T. Dowling, and M. M. Coelho. 2006. Modes of
824 reproduction of the hybridogenetic fish *Squalius alburnoides* in the Tejo and Guadiana
825 rivers: An approach with microsatellites. Zoology 109:277-286.

- 826 Cunha, C., I. Doadrio, and M. M. Coelho. 2008. Speciation towards tetraploidization after
827 intermediate processes of non-sexual reproduction. *Philosophical Transactions of the*
828 *Royal Society B* 363:2921-2929.
- 829 Dawley, R. M. 1989. An introduction to unisexual vertebrates. Pages 19-23 *in* R. M. Dawley
830 and J. P. Bogart, eds. *Evolution and ecology of unisexual vertebrates*. New York State
831 Museum Bulletin 466, New York State Museum, Albany, NY.
- 832 Fisher, R. A. 1930. *The genetic theory of natural selection*. Oxford University Press, Oxford
833 U.K.
- 834 Fowler, J., and L. Cohen. 1992. *Practical statistics for field biology*. John Wiley & Sons Ltd,
835 Chichester, West Sussex, UK.
- 836 Frost, D. R., T. Grant, J. Faivovich, R. H. Bain, A. Haas, C. F. B. Haddad, R. O. De Sa, A.
837 Channing, M. Wilkinson, S. C. Donnellan, C. J. Raxworthy, J. A. Campbell, B. L.
838 Blotto, P. Moler, R. C. Drewes, R. A. Nussbaum, J. D. Lynch, D. M. Green, and W. C.
839 Wheeler. 2006. The amphibian tree of life. *Bulletin of the American Museum of*
840 *Natural History* 297:8-370.
- 841 Garner, T. W. J., B. Gautschi, S. Rothlisberger, and H. U. Reyer. 2000. A set of CA repeat
842 microsatellite markers derived from the pool frog, *Rana lessonae*. *Molecular Ecology*
843 9:2173-2175.
- 844 Goyeau, H., F. Halkett, M. F. Zapater, J. Carlier, and C. Lannou. 2007. Clonality and host
845 selection in the wheat pathogenic fungus *Puccinia triticina*. *Fungal Genetics and*
846 *Biology* 44:474-483.
- 847 Graf, J. D., and M. Polls Pelaz. 1989. Evolutionary genetics of the *Rana esculenta* complex.
848 Pp. 289–302 *in* R. M. Dawley and J. P. Bogart, eds. *Evolution and ecology of*
849 *unisexual vertebrates*. New York State Museum Bulletin 466, New York State
850 Museum, Albany, NY.

- 851 Grundmann, M., S. W. Ansell, S. J. Russell, M. A. Koch, and J. C. Vogel. 2008. Hotspots of
852 diversity in a clonal world - the Mediterranean moss *Pleurochaete squarrosa* in
853 Central Europe. *Molecular Ecology* 17:825-838.
- 854 Guex, G. D., H. Hotz, and R. D. Semlitsch. 2002. Deleterious alleles and differential viability
855 in progeny of natural hemiclinal frogs. *Evolution* 56:1036-1044.
- 856 Günther, R. 1975. Zum natürlichen Vorkommen und zur Morphologie triploider Teichfrösche
857 "*Rana esculenta*", L., in der DDR (Anura, Ranidae). *Mitteilungen aus dem*
858 *Zoologischen Museum in Berlin* 51:145-158.
- 859 Günther, R. 1991. European waterfrogs (Anura, Ranidae) and the biospecies concept.
860 *Mitteilungen aus dem Museum fuer Naturkunde in Berlin Zoologische Reihe* 67:39-
861 53.
- 862 Günther, R., T. Uzzell, and L. Berger. 1979. Inheritance patterns in triploid *Rana*
863 "*esculenta*" (Amphibia, Salientia). *Mitteilungen aus dem Zoologischen Museum in*
864 *Berlin* 55:35-57.
- 865 Halkett, F., J. C. Simon, and F. Balloux. 2005. Tackling the population genetics of clonal and
866 partially clonal organisms. *Trends in Ecology & Evolution* 20:194-201.
- 867 Hamilton, W. D. 1980. Sex versus non-sex versus parasite. *Oikos* 35:282-290.
- 868 Hardy, O. J., and X. Vekemans. 2002. SPAGeDi: a versatile computer program to analyse
869 spatial genetic structure at the individual or population levels. *Molecular Ecology*
870 *Notes* 2:618-620.
- 871 Hedges, S. B., J. P. Bogart, and L. R. Maxson. 1992. Ancestry of unisexual salamanders.
872 *Nature* 356:708-710.
- 873 Hedrick, P. W. 2005. *Genetics of populations*, 3rd edition. Jones and Bartlett, Boston.
- 874 Hegarty, M. J., and S. J. Hiscock. 2005. Hybrid speciation in plants: new insights from
875 molecular studies. *New Phytologist* 165:411-423.

- 876 Holenweg Peter, A. K., H. U. Reyer, and G. A. Tietje. 2002. Species and sex ratio differences
877 in mixed populations of hybridogenetic water frogs: The influence of pond features.
878 *Ecoscience* 9:1-11.
- 879 Holm, S. 1979. A simple sequentially rejective multiple test procedure. *Scandinavian Journal*
880 *of Statistics* 6:65-70.
- 881 Hotz, H., P. Beerli, and C. Spolsky. 1992. Mitochondrial DNA reveals formation of
882 nonhybrid frogs by natural matings between hemiclinal hybrids. *Molecular Biology*
883 *and Evolution* 9:610-620.
- 884 Hotz, H., T. Uzzell, G. D. Guex, D. Alpers, R. D. Semlitsch, and P. Beerli. 2001.
885 Microsatellites: a tool for evolutionary genetic studies of western Palearctic water
886 frogs. *Mitteilungen aus dem Museum für Naturkunde Berlin, Zoologische Reihe*
887 *77:43–50*.
- 888 Innes, D. J., and D. R. Singleton. 2000. Variation in allocation to sexual and asexual
889 reproduction among clones of cyclically parthenogenetic *Daphnia pulex* (Crustacea:
890 Cladocera). *Biological Journal of the Linnean Society* 71:771-787.
- 891 Jakob, C. 2007. Structure and dynamics of pure hybridogenetic water frog populations of
892 *Rana esculenta* in Southern Sweden. PhD thesis. University of Zurich, Ecology
893 department. Available at www.dissertationen.unizh.ch
- 894 James, J. K., and R. J. Abbott. 2005. Recent, allopatric, homoploid hybrid speciation: The
895 origin of *Senecio squalidus* (Asteraceae) in the British Isles from a hybrid zone on
896 Mount Etna, Sicily. *Evolution* 59:2533-2547.
- 897 Janko, K., P. Kotlik, and P. Rab. 2003. Evolutionary history of asexual hybrid loaches
898 (*Cobitis*: Teleostei) inferred from phylogenetic analysis of mitochondrial DNA
899 variation. *Journal of Evolutionary Biology* 16:1280-1287.

900 Jiggins, C. D., C. Salazar, M. Linares, and J. Mavarez. 2008. Hybrid trait speciation and
901 Heliconius butterflies. Philosophical Transactions of the Royal Society B-Biological
902 Sciences 363:3047-3054.

903 Koref-Santibanez, S., and R. Günther. 1980. Karyological and serological studies in *Rana*
904 *lessonae*, *R. ridibunda* and in their hybrid *R. esculenta* (Amphibia, Anura). Genetica
905 52/53:195-207.

906 Kraus, F. 1995. The conservation of unisexual vertebrate populations. Conservation Biology
907 9:956-959.

908 Little, T. J., and P. D. N. Hebert. 1997. Clonal diversity in high arctic ostracodes. Journal of
909 Evolutionary Biology 10:233-252.

910 Maynard Smith, J. 1992. Clonal histories - age and the unisexual lineage. Nature 356:661-
911 662.

912 Mikulíček, P., and P. Kotlík. 2001. Two water frog populations from western Slovakia
913 consisting of diploid females and diploid and triploid males of the hybridogenetic
914 hybrid *Rana esculenta* (Anura, Ranidae). Mitteilungen aus dem Museum fuer
915 Naturkunde in Berlin Zoologische Reihe 77:59-64.

916 Morishima, K., H. Yoshikawa, and K. Arai. 2008. Meiotic hybridogenesis in triploid
917 *Misgurnus* loach derived from a clonal lineage. Heredity 100:581-586.

918 Moritz, C., W. M. Brown, L. D. Densmore, J. W. Wright, D. Vyas, S. Donellan, M. Adams,
919 and P. Baverstock. 1989. Genetic diversity and the dynamics of hybrid
920 parthenogenesis in *Cnemidophorus* (Teiidae) and *Heteronotia* (Gekkonidae). Pages
921 87-112 in R. M. Dawley and J. P. Bogart, eds. Evolution and Ecology of Unisexual
922 Vertebrates. New York State Museum Bulletin 466, New York State Museum,
923 Albany, NY

924 Muller, H. J. 1932. Some genetic aspects of sex. American Naturalist 66:118-138.

- 925 Orr, H. A. 1990. Why polyploidy is rarer in animals than in plants revisited. American
926 Naturalist 136:759-770.
- 927 Otto, S. P., and A. C. Gerstein. 2006. Why have sex? The population genetics of sex and
928 recombination. Biochemical Society Transactions 34:519-522.
- 929 Otto, S. P., and J. Whitton. 2000. Polyploid incidence and evolution. Annual Review of
930 Genetics 34:401-437.
- 931 Pagano, A., P. Joly, S. Plenet, A. Lehman, and O. Grolet. 2001. Breeding habitat partitioning
932 in the *Rana esculenta* complex: The intermediate niche hypothesis supported.
933 Ecoscience 8:294-300.
- 934 Pala, I., and M. M. Coelho. 2005. Contrasting views over a hybrid complex: Between
935 speciation and evolutionary "dead-end". Gene 347:283-294.
- 936 Plötner, J. 2005. Die westpaläarktischen Wasserfrösche. Laurenti-Verlag, Bielefeld.
- 937 Polls Pelaz, M. 1994. Modes of gametogenesis among kleptons of the hybridogenetic water
938 frog complex: an evolutionary synthesis. Zoologica Poloniae 39:123-138.
- 939 Ptacek, M. B., H. C. Gerhardt, and R. Sage. 1994. Speciation by polyploidy in treefrogs:
940 multiple origins of the tetraploid, *Hyla versicolor*. Evolution 48:898-908.
- 941 Ranker, T. A., and A. M. Arft. 1994. Allopolyploid species and the United States Endangered
942 Species Act. Conservation Biology 8:895-897.
- 943 Regnier, V., and A. Neveu. 1986. Specific structures in population of *Rana esculenta*
944 complex from different areas of Western France. Acta Oecologica - Oecologia
945 Applicata 7:3-26.
- 946 Rybacki, M., and L. Berger. 2001. Types of water frog populations (*Rana esculenta* complex)
947 in Poland. Mitteilungen aus dem Museum fuer Naturkunde in Berlin Zoologische
948 Reihe 77:51-57.
- 949 Saitoh, K., I. S. Kim, and E. H. Lee. 2004. Mitochondrial gene introgression between spined
950 Loaches via hybridogenesis. Zoological Science 21:795-798.

- 951 Schartl, M., I. Nanda, I. Schlupp, B. Wilde, J. T. Epplen, M. Schmid, and J. Parzefall. 1995.
952 Incorporation of subgenomic amounts of DNA as a compensation for mutational load
953 in a gynogenetic fish. *Nature* 373:68-71.
- 954 Simon, J. C., F. Delmotte, C. Rispe, and T. Crease. 2003. Phylogenetic relationships between
955 parthenogens and their sexual relatives: the possible routes to parthenogenesis in
956 animals. *Biological Journal of the Linnean Society* 79:151-163.
- 957 Soltis, D. E., and P. S. Soltis. 1999. Polyploidy: recurrent formation and genome evolution.
958 *Trends in Ecology and Evolution* 14:348-352.
- 959 Som, C., and H. U. Reyer. 2006a. Demography and evolution of pure hybridogenetic frog
960 (*Rana esculenta*) populations. *Evolutionary Ecology Research* 8:1235-1248.
- 961 Som, C., and H. U. Reyer. 2006b. Variation in sex ratio and evolutionary rate of hemiclonal
962 *Rana esculenta* populations. *Evolutionary Ecology* 20:159-172.
- 963 Spolsky, C. M., C. A. Phillips, and T. Uzzell. 1992. Antiquity of clonal salamander lineages
964 revealed by Mitochondrial DNA. *Nature* 356:706-708.
- 965 SPSS. 2004. vers. 13.0 for Windows.
- 966 Stöck, M., D. K. Lamatsch, C. Steinlein, J. T. Epplen, W. R. Grosse, R. Hock, T.
967 Klapperstuck, K. P. Lampert, U. Scheer, M. Schmid, and M. Schartl. 2002. A
968 bisexually reproducing all-triploid vertebrate. *Nature Genetics* 30:325-328.
- 969 Stöck, M., C. Steinlein, D. K. Lamatsch, M. Schartl, and M. Schmid. 2005. Multiple origins
970 of tetraploid taxa in the Eurasian *Bufo viridis* subgroup. *Genetica* 124:255-272.
- 971 Stuart, S. N., J. S. Chanson, N. A. Cox, B. E. Young, A. S. L. Rodrigues, D. L. Fischman, and
972 R. W. Waller. 2004. Status and trends of amphibian declines and extinctions
973 worldwide. *Science* 306:1783-1786.
- 974 Tunner, H. G., and S. Heppich-Tunner. 1991. Genome exclusion and two strategies of
975 chromosome duplication in oogenesis of a hybrid frog. *Naturwissenschaften* 78:32-34.

- 976 Tunner, H. G., and S. Heppich-Tunner. 1992. A new population system of water frogs
977 discovered in Hungary. Proceedings of the Sixth Ordinary General Meeting of the
978 Societas Europaea Herpetologica, Budapest 453-460.
- 979 Vorburger, C. 2001a. Fixation of deleterious mutations in clonal lineages: Evidence from
980 hybridogenetic frogs. *Evolution* 55:2319-2332.
- 981 Vorburger, C. 2001b. Heterozygous fitness effects of clonally transmitted genomes in
982 waterfrogs. *Journal of Evolutionary Biology* 14:602-610.
- 983 Vorburger, C. 2001c. Non-hybrid offspring from matings between hemiclonal hybrid
984 waterfrogs suggest occasional recombination between clonal genomes. *Ecology*
985 *Letters* 4:628-636.
- 986 Vrijenhoek, R. C. 1984. Ecological differentiation among clones: The frozen niche variation
987 model. Pages 217-231 in K. Wohrman and V. Loeshcke, eds. *Population Biology and*
988 *Evolution*. Springer-Verlag, Berlin-Heidelberg-New York.
- 989 Vrijenhoek, R. C. 1994. Unisexual fish: model systems for studying ecology and evolution.
990 *Annual Review of Ecology and Systematics* 25:71-96.
- 991 Vrijenhoek, R. C., R. M. Dawley, C. J. Cole, and J. P. Bogart. 1989. A list of the known
992 unisexual vertebrates. Pp. 19-23 in R. M. Dawley and J. P. Bogart, eds. *Evolution and*
993 *ecology of unisexual vertebrates*. New York State Museum Bulletin 466, New York
994 State Museum, Albany, NY.
- 995 Wang, X. R., A. E. Szmidt, and O. Savolainen. 2001. Genetic composition and diploid hybrid
996 speciation of a high mountain pine, *Pinus densata*, native to the Tibetan plateau.
997 *Genetics* 159:337-346.
- 998 Watkins, M. W. 2002. MacBonferroni vers. 1.00. Available at
999 <http://www.softpedia.com/get/Others/Home-Education/MacBonferroni.shtml>
- 1000 Wissemann, V. 2007. Plant evolution by means of hybridization. *Systematics and*
1001 *Biodiversity* 5:243-253.

1002 Zeisset, I., G. Rowe, and T. J. C. Beebee. 2000. Polymerase chain reaction primers for
 1003 microsatellite loci in the north European water frogs *Rana ridibunda* and *R. lessonae*.
 1004 Molecular Ecology 9:1173-1174.

1005

1006 **Tables**

1007 **Table 1.** Primer sequences not previously published.

Locus	Sequence 5' - 3'	Repeat	Genome specificity	Genbank ass. no.	Sequenced by
ReGA1a23	F: ATT GCT TTG GCA GTG AAG G R: TGA CAT CAC AGT GGG AGG AG	GA _n	L	EU445523	Garner <i>et al.</i> , Arioli and Jakob
RICA1a27	F: CAA ATG GGT CAT CCA CAC C R: GTT CAA GGG GGT CGA AAT AC	CA _n	L	EU445522	Garner <i>et al.</i>
RICA2a34	F: GCT CCA TGC CAA AAG TCT TC R: TTG GGT ATG ATA CTA CAA GCT ATG C	GT _n	L +R ¹	EU445521	Garner <i>et al.</i>
Rrid059A redesigned	F: TTG GAG ACA GAC TTC CGT AGG	CA _n	L ¹ +R	FJ024048	Hotz <i>et al.</i>
Rrid064A	F: TGT ACG GGC CTT TAG ACT GG R: AAC TTT TTG AAG GCC CCT TG	GT _nTA _n GT _n	R	EU445524	Hotz <i>et al.</i>
Rrid135A	F: TCT TTT GTT TTA GCG CAC CT R: CTG CCC GTC TAA GCA AGT GT	CA _n TA _n	R	EU445526	Hotz <i>et al.</i>
Rrid169A	F: CGG AAC TCC GCT TTA ATC AC R: CCC ATG TTG TCG TTG AGC TA	TA _n ...CA _n	R	EU445525	Hotz <i>et al.</i>

1008 ¹ monomorphic in this genome

1009

1010 **Figure legends**

1011 **Figure 1.** Adults, gametes and offspring of a) the LE system with *P. lessonae* and *P.*
 1012 *esculentus* and b) all-hybrid populations of *P. esculentus*. * denotes gametes that could be
 1013 recombined. Non-hybrid offspring from intraspecific *P. esculentus* matings are in parenthesis

1014 because they typically die before reproductive maturity. Note that in the LE system, the R
1015 genome is never recombined and the L genome is provided anew in every generation by *P.*
1016 *lessonae*. In the all-hybrid populations, both L and R genomes are supplied by hybrids and
1017 would regularly undergo recombination, if triploids have meiotic hybridogenesis.
1018

1019 **Figure 2.** Linkage analysis of various locus combinations in crossing experiment with triploid
1020 *P. esculentus*. The symbols represent $-\log_{10} P$ values from Chi-square tests of the frequency
1021 distributions of the four potential (two parental and two recombinant) gamete types produced.
1022 Circles = males, triangles = females. Most individuals were heterozygous at several loci and
1023 therefore contributed data for several locus pairs. Each point left of the dashed line indicate a
1024 freely recombined locus pair in a frog. Points right of the dashed line indicate significant
1025 linkage at the 0.05 level after sequential Bonferroni-correction within each genome
1026 separately. Filled symbols (circles and triangles pointing right, left and down) identify all P
1027 values derived from the four individuals that each gave a significant P value. The female
1028 identified by grey triangles pointing down had a mutation at Re2GACA3 in her germ line
1029 which she passed on to some of her offspring.


1030

1031 **Figure 3.** Multilocus linkage disequilibrium, r_d^- , as a function of the proportion of frogs
1032 producing recombined gametes in 66 *P. esculentus* populations from Denmark and Sweden.
1033 a) r_d^- in the L genome vs. recombining LLR frogs of the total number of frogs propagating L
1034 genomes (LLR+LR). Linear regression line dashed because non-significant. b) r_d^- in the R
1035 genome vs. recombining LRR/total R-propagating frogs; regression significant. r_d^- is an
1036 index of association adjusted for unequal sample size.

1037

1038 **Figure 4.** Multilocus linkage disequilibrium (r_d^-) in the L vs. the R genome in 56 ponds.

a) LE system



b) All-hybrid populations

