



University of
Zurich^{UZH}

Zurich Open Repository and
Archive

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2023

Observation of the $B_s c 1(3872) + \text{decay}$

LHCb Collaboration ; Bernet, R ; Müller, K ; Owen, P ; Serra, N ; Steinkamp, O ; et al

DOI: [https://doi.org/10.1007/JHEP07\(2023\)084](https://doi.org/10.1007/JHEP07(2023)084)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-253416>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

LHCb Collaboration; Bernet, R; Müller, K; Owen, P; Serra, N; Steinkamp, O; et al (2023). Observation of the $B_s c 1(3872) + \text{decay}$. Journal of High Energy Physics, 07:084.

DOI: [https://doi.org/10.1007/JHEP07\(2023\)084](https://doi.org/10.1007/JHEP07(2023)084)

Observation of the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decay



The LHCb collaboration

E-mail: Ivan.Belyaev@cern.ch

ABSTRACT: The first observation of the $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)\pi^+\pi^-$ decay is reported using proton-proton collision data, corresponding to integrated luminosities of 1, 2 and 6 fb^{-1} , collected by the LHCb experiment at centre-of-mass energies of 7, 8 and 13 TeV, respectively. The ratio of branching fractions relative to the $B_s^0 \rightarrow (\psi(2S) \rightarrow J/\psi\pi^+\pi^-)\pi^+\pi^-$ decay is measured to be

$$\frac{\mathcal{B}(B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-) \times \mathcal{B}(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)}{\mathcal{B}(B_s^0 \rightarrow \psi(2S)\pi^+\pi^-) \times \mathcal{B}(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)} = (6.8 \pm 1.1 \pm 0.2) \times 10^{-2},$$

where the first uncertainty is statistical and the second systematic. The mass spectrum of the $\pi^+\pi^-$ system recoiling against the $\chi_{c1}(3872)$ meson exhibits a large contribution from $B_s^0 \rightarrow \chi_{c1}(3872)(f_0(980) \rightarrow \pi^+\pi^-)$ decays.

KEYWORDS: B Physics, Hadron-Hadron Scattering, Quarkonium, Spectroscopy

ARXIV EPRINT: [2302.10629](https://arxiv.org/abs/2302.10629)

Contents

1	Introduction	1
2	Detector and simulation	2
3	Event selection	3
4	$B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays	4
5	Dipion mass spectrum	7
6	Systematic uncertainties	8
7	Summary	11
	The LHCb collaboration	18

1 Introduction

Decays of beauty hadrons to final states with charmonia provide a unique laboratory to study the properties of charmonia and charmonium-like states. A plethora of new states has been observed in such decays, including the $\chi_{c1}(3872)$ particle [1], pentaquark [2–6] and tetraquark [7–18] candidates as well as conventional charmonium states, such as the tensor D-wave $\psi_2(3823)$ meson [19–21]. The nature of many exotic charmonium-like candidates remains unclear. A comparison of their production rates with respect to those of conventional charmonium states in decays of beauty hadrons can shed light on their production mechanisms [22]. For example, the $D^*\bar{D}$ rescattering mechanism [23, 24] would give a large contribution to $\chi_{c1}(3872)$ production and affect the pattern of decay rates of beauty hadrons. There is a puzzling difference between the branching fractions for the $B^+ \rightarrow \chi_{c1}(3872)K^+$ and $B^0 \rightarrow \chi_{c1}(3872)K^0$ decays [25, 26]. It may be explained by a compact-tetraquark interpretation of the $\chi_{c1}(3872)$ state [27], which simultaneously accounts for the similarity of the branching fractions for the $B^0 \rightarrow \chi_{c1}(3872)K^0$ and $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ decays [16, 28]. Additional measurements on the $\chi_{c1}(3872)$ production in the decays of beauty hadrons, and in particular, decays of B_s^0 mesons, will be helpful for a better understanding of the nature of the $\chi_{c1}(3872)$ state.

In this paper, the first observation of the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decay is reported. This analysis is based on proton-proton (pp) collision data, corresponding to integrated luminosities of 1, 2 and 6 fb^{-1} , collected by the LHCb experiment at centre-of-mass energies of 7, 8 and 13 TeV, respectively. The $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decay is used as normalisation channel to

measure the ratio \mathcal{R} of the branching fractions of the $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-) \pi^+ \pi^-$ and $B_s^0 \rightarrow (\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) \pi^+ \pi^-$ decays

$$\mathcal{R} \equiv \frac{\mathcal{B}(B_s^0 \rightarrow \chi_{c1}(3872) \pi^+ \pi^-) \times \mathcal{B}(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-)}{\mathcal{B}(B_s^0 \rightarrow \psi(2S) \pi^+ \pi^-) \times \mathcal{B}(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)}. \quad (1.1)$$

The $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-) \pi^+ \pi^-$ and $B_s^0 \rightarrow (\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) \pi^+ \pi^-$ decays share the same final state, allowing for a large cancellation of systematic uncertainties.

2 Detector and simulation

The LHCb detector [29, 30] is a single-arm forward spectrometer covering the pseudo-rapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [31], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [32, 33] placed downstream of the magnet. The tracking system provides a measurement of the momentum of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/ c . The momentum scale is calibrated using samples of $J/\psi \rightarrow \mu^+ \mu^-$ and $B^+ \rightarrow J/\psi K^+$ decays collected concurrently with the data sample used for this analysis [34, 35]. The relative accuracy of this procedure is estimated to be 3×10^{-4} using samples of other fully reconstructed b hadrons, Υ and K_S^0 mesons. The minimum distance of a track to a primary pp -collision vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV/ c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors (RICH) [36]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter [37]. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [38].

The online event selection is performed by a trigger [39], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. The hardware trigger selects muon candidates with large transverse momentum or dimuon candidates with a large value of the product of the p_T of the muons. In the software trigger, two oppositely charged muons are required to form a good-quality vertex that is significantly displaced from every PV, with a dimuon mass exceeding 2.7 GeV/ c^2 .

Simulated events are used to describe signal shapes and to compute the efficiencies needed to determine the branching fraction ratio. In the simulation, pp collisions are generated using PYTHIA [40] with a specific LHCb configuration [41]. Decays of unstable particles are described by the EVTGEN package [42], in which final-state radiation is generated using PHOTOS [43]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [44, 45] as described in ref. [46]. The decays $B_s^0 \rightarrow \chi_{c1}(3872) \pi^+ \pi^-$ and $B_s^0 \rightarrow \psi(2S) \pi^+ \pi^+$ are simulated using

a phase-space decay model that is adjusted to match the mass distributions of the two-pion systems recoiling against the $\chi_{c1}(3872)$ and $\psi(2S)$ mesons in data. In the simulation $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ decays proceed via an S-wave $J/\psi\rho^0$ intermediate state [47–49]. The model described in refs. [50–55] is used for the $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ decays. The simulation is corrected to reproduce the transverse momentum and rapidity distributions of the B_s^0 mesons observed in data. To account for imperfections in the simulation of charged-particle reconstruction, the track reconstruction efficiency determined from simulation is corrected using data-driven techniques [56].

3 Event selection

Candidate $B_s^0 \rightarrow J/\psi\pi^+\pi^+\pi^-\pi^-$ decays are reconstructed using the $J/\psi \rightarrow \mu^+\mu^-$ decay mode. As explained in detail below, an initial selection criteria similar to those used in refs. [16, 57, 58] are used to reduce the background. Subsequently, a multivariate estimator, in the following referred as the MLP classifier, is applied. It is based on an artificial neural network algorithm [59, 60] configured with a cross-entropy cost estimator [61].

Muon and hadron candidates are identified using combined information from the RICH, calorimeter and muon detectors [62]. The candidates are required to have a transverse momentum greater than 550 MeV/c and 200 MeV/c for muons and pions, respectively. To ensure that the particles can be efficiently separated by the RICH detectors, pions are required to have a momentum between 3.2 and 150 GeV/c. To reduce the combinatorial background due to particles produced promptly in the pp interaction, only tracks that are inconsistent with originating from a PV are used. Pairs of oppositely charged muons consistent with originating from a common vertex are combined to form J/ψ candidates. The mass of the dimuon candidate is required to be between 3.05 and 3.15 GeV/c², corresponding to a range of approximately three times the $\mu^+\mu^-$ mass resolution, around the known mass of the J/ψ meson [25]. Selected J/ψ meson candidates are combined with two pairs of oppositely charged pions to form the $B_s^0 \rightarrow J/\psi\pi^+\pi^+\pi^-\pi^-$ candidates and a requirement on the quality of the common six-prong vertex is imposed. To improve the mass and decay time resolution, a kinematic fit [63] is used in which the momentum direction of the B_s^0 candidate is constrained to be collinear to the direction from the PV to the B_s^0 decay vertex, and a mass constraint on the J/ψ state is applied. A requirement on the χ^2 of this fit, χ_{fit}^2 , is imposed to reduce the background. The mass of selected $J/\psi\pi^+\pi^+\pi^-\pi^-$ combinations, $m_{J/\psi\pi^+\pi^+\pi^-\pi^-}$, is required to be between 5.30 and 5.48 GeV/c². The proper decay time of the B_s^0 candidates is required to be between 0.2 and 2.0 mm/c. The lower limit is used to reduce background from particles coming from the PV while the upper limit suppresses poorly reconstructed candidates. A possible feed down from $\Lambda_b^0 \rightarrow J/\psi p\pi^+\pi^-\pi^-$ decays, with the proton misidentified as a pion, is suppressed by rejecting the B_s^0 candidates whose mass, recalculated using the proton hypothesis for one of the pion candidates, is consistent with the known mass of the Λ_b^0 baryon [25].

The final selection of candidates using the MLP classifier is based on the p_T and pseudorapidity of J/ψ candidate, the χ^2 of the six-prong vertex, the value of χ_{fit}^2 , the proper decay time of the B_s^0 candidates, the transverse momenta of the pions,

the dipion mass from the $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ decays and the angle between the momenta of the J/ψ and B_s^0 mesons in the $\chi_{c1}(3872)$ rest frame. The MLP classifier is trained on a sample of simulated $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)\pi^+\pi^-$ decays and a background sample of $J/\psi\pi^+\pi^+\pi^-\pi^-$ combinations from the high mass sideband of the B_s^0 signal peak, $5.42 < m_{J/\psi\pi^+\pi^+\pi^-\pi^-} < 5.50 \text{ GeV}/c^2$. The low-mass sideband of the B_s^0 signal peak contains a large contribution from partially reconstructed b -hadron decays and therefore is not representative of the background under the B_s^0 signal peak. For the background sample, $J/\psi\pi^+\pi^-$ combinations with the $J/\psi\pi^+\pi^-$ mass consistent with the known masses of the $\chi_{c1}(3872)$ state [20, 64], $3.86 < m_{J/\psi\pi^+\pi^-} < 3.88 \text{ GeV}/c^2$, or $\psi(2S)$ meson [25], $3.68 < m_{J/\psi\pi^+\pi^-} < 3.69 \text{ GeV}/c^2$, are excluded. To avoid introducing a bias in the MLP evaluation, a k -fold cross-validation technique [65] with $k = 7$ is used. The requirement on the MLP classifier is chosen to maximize the Punzi figure of merit $\frac{\varepsilon}{\alpha/2 + \sqrt{B}}$ [66], where ε is the efficiency for the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ signal, $\alpha = 5$ is the target signal significance and B is the expected background yield. The efficiency ε is estimated using simulation. The expected background yield within the narrow mass window centred around the known masses of the B_s^0 and $\chi_{c1}(3872)$ mesons [25] is determined from fits to data.

The selected B_s^0 candidates with $J/\psi\pi^+\pi^-$ mass within the range $3.85 < m_{J/\psi\pi^+\pi^-} < 3.90 \text{ GeV}/c^2$ are considered as the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ candidates. Similarly, the B_s^0 candidates with the $J/\psi\pi^+\pi^-$ mass within the range $3.67 < m_{J/\psi\pi^+\pi^-} < 3.70 \text{ GeV}/c^2$ are considered as the $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ candidates. The same MLP classifier is used both for the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ candidates.

4 $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays

The signal yields for the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays are determined using a two-dimensional extended unbinned maximum-likelihood fit to the $J/\psi\pi^+\pi^+\pi^-\pi^-$ and $J/\psi\pi^+\pi^-$ mass distributions. Following refs. [16, 20, 67], to improve the resolution on the $J/\psi\pi^+\pi^-$ mass and to eliminate a small correlation between the $J/\psi\pi^+\pi^-$ and $J/\psi\pi^+\pi^+\pi^-\pi^-$ masses, the $J/\psi\pi^+\pi^-$ mass is computed constraining the mass of the B_s^0 candidate [63] to its known mass [25]. The fit is performed simultaneously in two separate regions of the $J/\psi\pi^+\pi^-$ mass, defined as the $\chi_{c1}(3872)$ region with $3.85 < m_{J/\psi\pi^+\pi^-} < 3.90 \text{ GeV}/c^2$, and the $\psi(2S)$ region with $3.67 < m_{J/\psi\pi^+\pi^-} < 3.70 \text{ GeV}/c^2$, which correspond to the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays, respectively.

In the $\chi_{c1}(3872)$ region, the two-dimensional fit model is defined as the sum of four components:

1. A signal component, corresponding to the $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)\pi^+\pi^-$ decay, described by the product of the B_s^0 and $\chi_{c1}(3872)$ signal templates, discussed in the next paragraph.

2. A component corresponding to $B_s^0 \rightarrow J/\psi \pi^+ \pi^+ \pi^- \pi^-$ decay, where the $J/\psi \pi^+ \pi^-$ combination does not originate from a $\chi_{c1}(3872)$ meson, parameterised by the product of the B_s^0 signal template and the phase-space function describing three-body combinations from five-body decays¹ $\Phi_{3,5}(m_{J/\psi \pi^+ \pi^-})$ [68], modified by a positive linear polynomial function.
3. A component corresponding to random combinations of the $\chi_{c1}(3872)$ state with a $\pi^+ \pi^-$ pair, parameterised as a product of the signal $\chi_{c1}(3872)$ template and a second-order positive-definite polynomial function [69].
4. A component corresponding to random $J/\psi \pi^+ \pi^+ \pi^- \pi^-$ combinations, parameterized as a two-dimensional non-factorizable positive polynomial function P_{bkg} , that is linear in each variable for a fixed value of the other variable.

In the $\psi(2S)$ region, the two-dimensional fit model is defined in an equivalent way with replacement of the signal $\chi_{c1}(3872)$ template with the signal $\psi(2S)$ template. In this approach we neglect possible interference effects between the $B_s^0 \rightarrow \chi_{c1}(3872) \pi^+ \pi^-$ and non-resonant $B_s^0 \rightarrow J/\psi \pi^+ \pi^+ \pi^- \pi^-$ contributions.

The B_s^0 signal shape is modelled with a modified Gaussian function with power-law tails on both sides of the distribution [70, 71]. The tail parameters are fixed from simulation, while the mass parameter of the B_s^0 meson is allowed to vary. The detector resolution taken from simulation is corrected by a scale factor, $s_{B_s^0}$, that accounts for a small discrepancy between data and simulation [16, 20] and is allowed to vary in the fit.

The $\chi_{c1}(3872)$ and $\psi(2S)$ signal shapes are modelled as relativistic S-wave Breit-Wigner functions convolved with the detector resolution. Due to the proximity of the $\chi_{c1}(3872)$ state to the $D^0 \bar{D}^{*0}$ threshold, modelling this component with a Breit-Wigner function may not be adequate [23, 72–75]. However, the analyses in refs. [20, 64] demonstrate that a good description of data is obtained with a Breit-Wigner line shape when the detector resolution is included. Mass parameters for the $\chi_{c1}(3872)$ and $\psi(2S)$ signals are allowed to vary in the fit, while the mass difference is Gaussian constrained to the known value [20]. The width parameters of the $\chi_{c1}(3872)$ and $\psi(2S)$ states are fixed to known values [20, 25] using a Gaussian constraint. The detector resolution functions are described by a symmetric modified Gaussian function with power-law tails on both sides of the distribution [70, 71], with all parameters determined from simulation. The resolutions are further corrected by a common scale factor, $s_{J/\psi \pi^+ \pi^-}$, that accounts for a small discrepancy between data and simulation and is allowed to vary in the fit.

The fit is performed using the B_s^0 mass and the resolution scale factors, $s_{B_s^0}$ and $s_{J/\psi \pi^+ \pi^-}$, as shared parameters between the $\chi_{c1}(3872) \pi^+ \pi^-$ and $\psi(2S) \pi^+ \pi^-$ final states. The $J/\psi \pi^+ \pi^+ \pi^- \pi^-$ and $J/\psi \pi^+ \pi^-$ mass distributions together with projections of the fit are shown in figure 1. The parameters of interest, namely the B_s^0 yields, masses of the B_s^0 meson, $\chi_{c1}(3872)$ and $\psi(2S)$ states, and the resolution scale factors are listed in table 1.

¹The phase-space mass distribution of a k -body combination of particles from an n -body decay is approximated by $\Phi_{k,n}(x) \propto x_*^{(3k-5)/2} (1-x_*)^{3(n-k)/2-1}$, where $x_* \equiv (x - x_{\min}) / (x_{\max} - x_{\min})$, and x_{\min} , x_{\max} denote the minimal and maximal values of x , respectively [68].




Figure 1. Distributions of the (left) $J/\psi\pi^+\pi^+\pi^-\pi^-$ and (right) $J/\psi\pi^+\pi^-$ mass of selected (top) $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ and (bottom) $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ candidates. Projections from the fit, described in the text, are overlaid. The $J/\psi\pi^+\pi^-$ mass distributions are shown for the $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ and $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ candidates within narrow $J/\psi\pi^+\pi^-$ mass ranges, $3.679 < m_{J/\psi\pi^+\pi^-} < 3.693 \text{ GeV}/c^2$ and $3.864 < m_{J/\psi\pi^+\pi^-} < 3.880 \text{ GeV}/c^2$, respectively. The $J/\psi\pi^+\pi^-$ mass distributions are shown for the B_s^0 candidates within a narrow $J/\psi\pi^+\pi^+\pi^-\pi^-$ mass range, $5.35 < m_{J/\psi\pi^+\pi^+\pi^-\pi^-} < 5.38 \text{ GeV}/c^2$.

Parameter	$B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$	$B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$
N	155 ± 23	1301 ± 47
$m_{\chi_{c1}(3872)}$ [MeV/ c^2]	3871.57 ± 0.09	—
$m_{\psi(2S)}$ [MeV/ c^2]	—	3686.08 ± 0.07
$m_{B_s^0}$ [MeV/ c^2]	5366.97 ± 0.23	
$s_{B_s^0}$	1.06 ± 0.03	
$s_{J/\psi\pi^+\pi^-}$	1.12 ± 0.03	

Table 1. Signal yields, hadron masses, and detector resolution scale factors from the simultaneous fit described in the text. The parameters $m_{B_s^0}$, $s_{B_s^0}$ and $s_{J/\psi\pi^+\pi^-}$ are shared in the fit among the two mass ranges. The uncertainties are statistical only.

The statistical significance for the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ signal is estimated using Wilks' theorem [76] to be 7.3 standard deviations.

The ratio of branching fractions \mathcal{R} , defined in eq. (1.1) is calculated as

$$\mathcal{R} = \frac{N_{B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-}}{N_{B_s^0 \rightarrow \psi(2S)\pi^+\pi^-}} \times \frac{\varepsilon_{B_s^0 \rightarrow \psi(2S)\pi^+\pi^-}}{\varepsilon_{B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-}}, \quad (4.1)$$

where the signal yields, $N_{B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-}$ and $N_{B_s^0 \rightarrow \psi(2S)\pi^+\pi^-}$, are taken from table 1 and $\varepsilon_{B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-}$ and $\varepsilon_{B_s^0 \rightarrow \psi(2S)\pi^+\pi^-}$ are the efficiencies for the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays, respectively. The efficiencies are defined as the product of the detector geometric acceptance and the reconstruction, selection, particle identification and trigger efficiencies. All of the efficiency contributions, except the particle identification efficiency, are determined using simulated samples. The efficiencies of the hadron identification are obtained as a function of particle momentum, pseudorapidity and number of charged tracks in the event using dedicated calibration samples of $D^{*+} \rightarrow (D^0 \rightarrow K^-\pi^+)\pi^+$ and $K_S^0 \rightarrow \pi^+\pi^-$ decays selected in data [36, 77]. The efficiency ratio is found to be

$$\frac{\varepsilon_{B_s^0 \rightarrow \psi(2S)\pi^+\pi^-}}{\varepsilon_{B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-}} = 0.57 \pm 0.01,$$

where the uncertainty is due to the limited size of the simulated samples. The efficiency ratio differs from unity due to the different p_T spectra of pions in the $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ and the $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ decays. The resulting value of \mathcal{R} is

$$\mathcal{R} = (6.8 \pm 1.1) \times 10^{-2}, \quad (4.2)$$

where the uncertainty is statistical.

5 Dipion mass spectrum

The mass spectra for the dipion system recoiling against the $\chi_{c1}(3872)$ and $\psi(2S)$ states are obtained using the sPlot technique [78], based on results of the two-dimensional fit described in previous section. The spectra shown in figure 2 exhibit significant deviations from the phase-space distribution and are similar to the $\pi^+\pi^-$ mass spectrum observed in the $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decay [79–81] and the S-wave $\pi^+\pi^-$ component in $D_s^+ \rightarrow \pi^+\pi^+\pi^-$ decays [82, 83].

In ref. [79] it has been found that the $\pi^+\pi^-$ mass spectrum from $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays can be described by a coherent sum of the $f_0(980)$ amplitude and a contribution from a high-mass scalar meson, initially identified as the $f_0(1370)$ state. References [84, 85] suggest a better interpretation of the high-mass state as the $f_0(1500)$ resonance. A dedicated amplitude analysis from ref. [81] followed this suggestion and confirmed the dominant $f_0(980)$ and $f_0(1500)$ contributions in $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays. Assuming a similarity with the $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays, the mass spectra of the dipion system recoiling against the $\chi_{c1}(3872)$ and $\psi(2S)$ states are described with a function consisting of two components:

- A component corresponding to a coherent sum of the scalar $f_0(980)$ and $f_0(1500)$ amplitudes and parameterised as

$$F(m) \propto mqp^3 \left| f\mathcal{A}_{f_0(980)}(m) + e^{i\varphi}\mathcal{A}_{f_0(1500)}(m) \right|^2, \quad (5.1)$$

where m is the $\pi^+\pi^-$ mass, q is the momentum of the π^+ meson in the $\pi^+\pi^-$ rest frame, p is the momentum of the $\pi^+\pi^-$ system in the B_s^0 rest frame, $\mathcal{A}_{f_0(980)}$ and $\mathcal{A}_{f_0(1500)}$ are the $f_0(980)$ and $f_0(1500)$ amplitudes, φ is a relative phase and the real coefficient f characterises the relative contributions of the $f_0(980)$ and $f_0(1500)$ components. The amplitude $\mathcal{A}_{f_0(1500)}$ is parameterised as a relativistic S-wave Breit-Wigner function, while the modified Flatté-Bugg amplitude [86, 87] (see eq. (18) in ref. [81]) is used for the $f_0(980)$ state.

- A component corresponding to incoherent nonresonant contribution and parameterised by the $\Phi_{2,3}(m)$ phase-space function [68].

The fit is performed simultaneously for the dipion mass spectra from the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^+$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^+$ decays. The shape parameters of the $f_0(980)$ and $f_0(1500)$ states are shared in the fit. To stabilise the fit, Gaussian constraints are applied to the parameters of the $f_0(980)$ state and the mass and width of the $f_0(1500)$ state, according to Solution I from ref. [81].

The fit results are shown in figure 2. This simplistic model qualitatively describes the major contributions to the dipion mass spectrum from $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays and supports the hypothesis of the dominant contribution of two S-wave resonances. The fit indicates the necessity of a dedicated analysis to properly account for the sub-leading contributions. The same model, consisting of two coherent contributions from the $f_0(980)$ and $f_0(1500)$ states describes well the dipion mass spectrum from the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decay. The statistical significance for the $B_s^0 \rightarrow \chi_{c1}(3872)f_0(980)$ decay is estimated using Wilks' theorem and found to be 9.1 standard deviations.

6 Systematic uncertainties

Due to the similar decay topologies, systematic uncertainties largely cancel in the ratio \mathcal{R} . The remaining contributions to systematic uncertainties are summarized in table 2 and discussed below.

The systematic uncertainties arising from imperfect knowledge of the signal and background shapes used for determination of the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ signal yields are estimated with alternative models. For the B_s^0 signal shape the bifurcated Student's t -distribution [88], Apollonios and Hypatia distributions [89] are tested as alternative models. The $J/\psi\pi^+\pi^-$ mass resolution functions are also modelled with Student's t -distribution and symmetric Apollonios functions. The degree of the polynomial functions used for the parameterisation of the background and the non-resonant $J/\psi\pi^+\pi^-$ components is varied by one unit. Exponential functions are used as components in alternative background models. In addition, the detector resolution scale factors




Figure 2. The background-subtracted mass spectra for the dipion system recoiling against the $\chi_{c1}(3872)$ or $\psi(2S)$ states for (left) $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ and (right) $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decays. The results of the fit, described in the text, are overlaid.

Source	$\sigma_{\mathcal{R}}$ [%]
Fit model	2.5
B_s^0 decay model	0.9
Efficiency corrections	0.1
Trigger efficiency	1.1
Data-simulation difference	2.0
Simulated sample size	0.5
Sum in quadrature	3.5

Table 2. Relative systematic uncertainties (in %) for the ratio of branching fractions. The sources are described in the text.

$s_{B_s^0}$ and $s_{J/\psi\pi^+\pi^-}$ are constrained to values of $s_{B_s^0} = 1.04 \pm 0.02$, $s_{J/\psi\pi^+\pi^-} = 1.06 \pm 0.02$ from ref. [16] and $s_{B_s^0} = 1.052 \pm 0.003$, $s_{J/\psi\pi^+\pi^-} = 1.048 \pm 0.004$ from ref. [20] using Gaussian constraints. For each alternative model the ratio of the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ signal yields is recalculated. The maximum relative deviation with respect to the baseline value is found to be 2.5% which is assigned as a relative systematic uncertainty for the ratio \mathcal{R} . The fit procedure itself is tested using a large sample of pseudoexperiments, generated using the default model with parameters extracted from data and the results are found to be unbiased.

The $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays are simulated as phase-space decays and corrected to reproduce the dipion mass spectra observed in data. A weighting procedure, based on a gradient boosted tree algorithm [90], is used for corrections of simulated samples. The systematic uncertainty related to the correction method is estimated by varying the hyper-parameters of the regression trees ensemble. The maximum deviation from the baseline value, 0.9%, is taken as the uncertainty associated with the unknown B_s^0 decay models.

An additional systematic uncertainty on the ratio \mathcal{R} arises due to differences between data and simulation. In particular, there are differences in the reconstruction efficiency of charged-particle tracks that do not cancel completely in the ratio due to the different kinematic distributions of the final-state particles. The track-finding efficiencies obtained from simulated samples are corrected using data-driven techniques [56]. The uncertainties related to the efficiency correction factors, together with the uncertainty on the hadron-identification efficiency due to the finite size of the calibration samples [36, 77], are propagated to the ratio of the total efficiencies using pseudoexperiments and amount to 0.1%. A systematic uncertainty on the ratio related to the knowledge of the trigger efficiencies is estimated by comparing the ratios of trigger efficiencies in data and simulation for large samples of $B^+ \rightarrow J/\psi K^+$ and $B^+ \rightarrow \psi(2S)K^+$ decays [91] and is taken to be 1.1%.

Remaining data-simulation differences, that are not previously discussed, are investigated by varying the selection criteria. The resulting variations in the ratios of the efficiency-corrected yields do not exceed 2%, which is taken as the corresponding systematic uncertainty. The final systematic uncertainty considered on the ratios of branching fractions is due to the knowledge of the ratios of efficiencies in eq. (4.1), limited by the size of simulated samples. It is determined to be 0.5%.

No systematic uncertainty is included for the admixture of the CP -odd and CP -even B_s^0 eigenstates in the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays [92], which is assumed to be the same for both channels. Analysis of the dipion spectrum from the $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decays in section 5 indicates that the final state is predominantly CP -odd, with the effective lifetime corresponding to the heavy-mass long-lifetime B_s^0 eigenstate [93–95]. The dipion spectrum from the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decays is also consistent with being predominantly CP -odd. In the extreme case that the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decay is an equal mixture of the short-lifetime and long-lifetime eigenstates, the corresponding ratio of branching fractions would change by 2.4%.

The statistical significance for the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decay is recalculated using Wilks' theorem for each alternative fit model, and the smallest value of 7.3 standard deviations is taken as the significance that includes the systematic uncertainty.

Alternative models are used also for the fit to the dipion mass spectra from the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ channels. In particular, Solution II from ref. [81] is investigated as an alternative external constraint for the $f_0(980)$ and $f_0(1500)$ parameters. The model with the $f_0(1370)$ resonance instead of the $f_0(1500)$ state is used as alternative model with a constraint to the known parameters of the $f_0(1370)$ state [25]. As an alternative model for the noncoherent nonresonant component, a product of the $\Phi_{2,3}$ phase-space function and the positive second-order polynomial function [69] has been probed. Also, a coherent nonresonant contribution, parameterised with a complex-valued constant function, is added to the function from eq. (5.1). The smallest significance for the $B_s^0 \rightarrow \chi_{c1}(3872)f_0(980)$ decay is found to be 7.4 standard deviations, which is taken as the significance including systematic uncertainty.

7 Summary

An observation of the $B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-$ decays with significance exceeding 7 standard deviations is reported using proton-proton collision data, corresponding to integrated luminosities of 1, 2 and 6 fb⁻¹, collected by the LHCb experiment at centre-of-mass energies of 7, 8 and 13 TeV, respectively. Using the $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ decay as a normalization channel, and neglecting the possible interference effects, the ratio of branching fractions is measured to be

$$\frac{\mathcal{B}(B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-) \times \mathcal{B}(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)}{\mathcal{B}(B_s^0 \rightarrow \psi(2S)\pi^+\pi^-) \times \mathcal{B}(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)} = (6.8 \pm 1.1 \pm 0.2) \times 10^{-2},$$

where the first uncertainty is statistical and the second systematic. The measured ratio of the branching fractions exceeds by a factor of three the analogous ratio for the $B_s^0 \rightarrow \chi_{c1}(3872)\phi$ and $B_s^0 \rightarrow \psi(2S)\phi$ decays reported in refs. [16, 28]. Using the known branching fractions for the $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$, and $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ decays [25], the branching fraction for the $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)\pi^+\pi^-$ is calculated to be

$$\mathcal{B}(B_s^0 \rightarrow \chi_{c1}(3872)\pi^+\pi^-) \times \mathcal{B}(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) = (1.6 \pm 0.3 \pm 0.1 \pm 0.3) \times 10^{-6},$$

where the third uncertainty is due to imprecise knowledge of the branching fractions for the $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ and $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ decays [25]. The obtained value is more than twice smaller than the branching fractions of the $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)\phi$ and the $B_s^0 \rightarrow (\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)(K^+K^-)_{\text{non-}\phi}$ decays [16, 25, 28].

The mass spectrum of the dipion system recoiling against the $\chi_{c1}(3872)$ state shows a similarity with those from the $B_s^0 \rightarrow \psi(2S)\pi^+\pi^-$ and $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays [81], compatible with a dominant S-wave contribution, and exhibits a large $B_s^0 \rightarrow \chi_{c1}(3872)f_0(980)$ component with significance exceeding 7 standard deviations. With a larger data sample and better understanding of the S-wave $\pi^+\pi^-$ scattering [96], a precise determination of the $f_0(980)$ component in this decay will be possible.

Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (U.S.A.). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), Polish WLCG (Poland) and NERSC (U.S.A.). We are indebted to the communities behind the multiple open-source software packages on which we depend. Indi-

vidual groups or members have received support from ARC and ARDC (Australia); Min-ciencias (Colombia); AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Ac-tions and ERC (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); GVA, XuntaGal, GENCAT and Prog. Atrac-ción Talento, CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] BELLE collaboration, *Observation of a narrow charmonium-like state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays*, *Phys. Rev. Lett.* **91** (2003) 262001 [[hep-ex/0309032](#)] [[INSPIRE](#)].
- [2] LHCb collaboration, *Observation of $J/\psi p$ resonances consistent with pentaquark states in $\Lambda_b^0 \rightarrow J/\psi K^- p$ decays*, *Phys. Rev. Lett.* **115** (2015) 072001 [[arXiv:1507.03414](#)] [[INSPIRE](#)].
- [3] LHCb collaboration, *Model-independent evidence for $J/\psi p$ contributions to $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays*, *Phys. Rev. Lett.* **117** (2016) 082002 [[arXiv:1604.05708](#)] [[INSPIRE](#)].
- [4] LHCb collaboration, *Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$* , *Phys. Rev. Lett.* **122** (2019) 222001 [[arXiv:1904.03947](#)] [[INSPIRE](#)].
- [5] LHCb collaboration, *Evidence for exotic hadron contributions to $\Lambda_b^0 \rightarrow J/\psi p \pi^-$ decays*, *Phys. Rev. Lett.* **117** (2016) 082003 [*Addendum ibid.* **117** (2016) 109902] [[arXiv:1606.06999](#)] [[INSPIRE](#)].
- [6] LHCb collaboration, *Evidence of a $J/\psi \Lambda$ structure and observation of excited Ξ^- states in the $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decay*, *Sci. Bull.* **66** (2021) 1278 [[arXiv:2012.10380](#)] [[INSPIRE](#)].
- [7] BELLE collaboration, *Observation of a resonance-like structure in the $\pi i^\pm \psi'$ mass distribution in exclusive $B \rightarrow K \pi^\pm \psi'$ decays*, *Phys. Rev. Lett.* **100** (2008) 142001 [[arXiv:0708.1790](#)] [[INSPIRE](#)].
- [8] BELLE collaboration, *Dalitz analysis of $B \rightarrow K \pi^+ \psi'$ decays and the $Z(4430)^+$* , *Phys. Rev. D* **80** (2009) 031104 [[arXiv:0905.2869](#)] [[INSPIRE](#)].
- [9] BELLE collaboration, *Experimental constraints on the spin and parity of the $Z(4430)^+$* , *Phys. Rev. D* **88** (2013) 074026 [[arXiv:1306.4894](#)] [[INSPIRE](#)].
- [10] LHCb collaboration, *Observation of the resonant character of the $Z(4430)^-$ state*, *Phys. Rev. Lett.* **112** (2014) 222002 [[arXiv:1404.1903](#)] [[INSPIRE](#)].
- [11] LHCb collaboration, *Model-independent confirmation of the $Z(4430)^-$ state*, *Phys. Rev. D* **92** (2015) 112009 [[arXiv:1510.01951](#)] [[INSPIRE](#)].
- [12] LHCb collaboration, *Observation of $J/\psi \phi$ structures consistent with exotic states from amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ decays*, *Phys. Rev. Lett.* **118** (2017) 022003 [[arXiv:1606.07895](#)] [[INSPIRE](#)].

- [13] LHCb collaboration, *Amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ decays*, *Phys. Rev. D* **95** (2017) 012002 [[arXiv:1606.07898](#)] [[INSPIRE](#)].
- [14] LHCb collaboration, *Evidence for an $\eta_c(1S)\pi^-$ resonance in $B^0 \rightarrow \eta_c(1S)K^+\pi^-$ decays*, *Eur. Phys. J. C* **78** (2018) 1019 [[arXiv:1809.07416](#)] [[INSPIRE](#)].
- [15] LHCb collaboration, *Model-independent observation of exotic contributions to $B^0 \rightarrow J/\psi K^+\pi^-$ decays*, *Phys. Rev. Lett.* **122** (2019) 152002 [[arXiv:1901.05745](#)] [[INSPIRE](#)].
- [16] LHCb collaboration, *Study of $B_s^0 \rightarrow J\psi\pi^+\pi^-K^+K^-$ decays*, *JHEP* **02** (2021) 024 [Erratum *ibid.* **04** (2021) 170] [[arXiv:2011.01867](#)] [[INSPIRE](#)].
- [17] LHCb collaboration, *Observation of new resonances decaying to $J/\psi K^{++}$ and $J/\psi\phi$* , *Phys. Rev. Lett.* **127** (2021) 082001 [[arXiv:2103.01803](#)] [[INSPIRE](#)].
- [18] LHCb collaboration, *Evidence for a new structure in the $J/\psi p$ and $J/\psi\bar{p}$ systems in $B_s^0 \rightarrow J/\psi p\bar{p}$ decays*, *Phys. Rev. Lett.* **128** (2022) 062001 [[arXiv:2108.04720](#)] [[INSPIRE](#)].
- [19] BELLE collaboration, *Evidence of a new narrow resonance decaying to $\chi_{c1}\gamma$ in $B \rightarrow \chi_{c1}\gamma K$* , *Phys. Rev. Lett.* **111** (2013) 032001 [[arXiv:1304.3975](#)] [[INSPIRE](#)].
- [20] LHCb collaboration, *Study of the $\psi_2(3823)$ and $\chi_{c1}(3872)$ states in $B^+ \rightarrow (J\psi\pi^+\pi^-)K^+$ decays*, *JHEP* **08** (2020) 123 [[arXiv:2005.13422](#)] [[INSPIRE](#)].
- [21] LHCb collaboration, *Study of charmonium and charmonium-like contributions in $B^+ \rightarrow J/\psi\eta K^+$ decays*, *JHEP* **04** (2022) 046 [[arXiv:2202.04045](#)] [[INSPIRE](#)].
- [22] L. Maiani, A.D. Polosa and V. Riquer, *A theory of X and Z multiquark resonances*, *Phys. Lett. B* **778** (2018) 247 [[arXiv:1712.05296](#)] [[INSPIRE](#)].
- [23] P. Artoisenet, E. Braaten and D. Kang, *Using line shapes to discriminate between binding mechanisms for the X(3872)*, *Phys. Rev. D* **82** (2010) 014013 [[arXiv:1005.2167](#)] [[INSPIRE](#)].
- [24] E. Braaten, L.-P. He and K. Ingles, *Production of X(3872) accompanied by a pion in B meson decay*, *Phys. Rev. D* **100** (2019) 074028 [[arXiv:1902.03259](#)] [[INSPIRE](#)].
- [25] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *PTEP* **2022** (2022) 083C01 [[INSPIRE](#)].
- [26] BELLE collaboration, *Study of the lineshape of X(3872) using B decays to $D^0\bar{D}^{*0}K$* , [[arXiv:2302.02127](#)] [[INSPIRE](#)].
- [27] L. Maiani, A.D. Polosa and V. Riquer, *X(3872) tetraquarks in B and B_s decays*, *Phys. Rev. D* **102** (2020) 034017 [[arXiv:2005.08764](#)] [[INSPIRE](#)].
- [28] CMS collaboration, *Observation of the $B_s^0 \rightarrow X(3872)\phi$ decay*, *Phys. Rev. Lett.* **125** (2020) 152001 [[arXiv:2005.04764](#)] [[INSPIRE](#)].
- [29] LHCb collaboration, *The LHCb detector at the LHC, 2008 JINST* **3** S08005 [[INSPIRE](#)].
- [30] LHCb collaboration, *LHCb detector performance*, *Int. J. Mod. Phys. A* **30** (2015) 1530022 [[arXiv:1412.6352](#)] [[INSPIRE](#)].
- [31] R. Aaij et al., *Performance of the LHCb vertex locator, 2014 JINST* **9** P09007 [[arXiv:1405.7808](#)] [[INSPIRE](#)].
- [32] LHCb Outer Tracker Group collaboration, *Performance of the LHCb outer tracker, 2014 JINST* **9** P01002 [[arXiv:1311.3893](#)] [[INSPIRE](#)].
- [33] LHCb Outer Tracker Group collaboration, *Improved performance of the LHCb outer tracker in LHC run 2, 2017 JINST* **12** P11016 [[arXiv:1708.00819](#)] [[INSPIRE](#)].

- [34] LHCb collaboration, *Measurement of the Λ_b^0 , Ξ_b^- and Ω_b^- baryon masses*, *Phys. Rev. Lett.* **110** (2013) 182001 [[arXiv:1302.1072](#)] [[INSPIRE](#)].
- [35] LHCb collaboration, *Precision measurement of D meson mass differences*, *JHEP* **06** (2013) 065 [[arXiv:1304.6865](#)] [[INSPIRE](#)].
- [36] LHCb RICH Group collaboration, *Performance of the LHCb RICH detector at the LHC*, *Eur. Phys. J. C* **73** (2013) 2431 [[arXiv:1211.6759](#)] [[INSPIRE](#)].
- [37] C. Abellán Beteta et al., *Calibration and performance of the LHCb calorimeters in run 1 and 2 at the LHC*, [arXiv:2008.11556](#) [[INSPIRE](#)].
- [38] A.A. Alves Jr. et al., *Performance of the LHCb muon system*, *2013 JINST* **8** P02022 [[arXiv:1211.1346](#)] [[INSPIRE](#)].
- [39] R. Aaij et al., *The LHCb trigger and its performance in 2011*, *2013 JINST* **8** P04022 [[arXiv:1211.3055](#)] [[INSPIRE](#)].
- [40] T. Sjöstrand, S. Mrenna and P.Z. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852 [[arXiv:0710.3820](#)] [[INSPIRE](#)].
- [41] LHCb collaboration, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, *J. Phys. Conf. Ser.* **331** (2011) 032047 [[INSPIRE](#)].
- [42] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [43] P. Golonka and Z. Was, *PHOTOS Monte Carlo: a precision tool for QED corrections in Z and W decays*, *Eur. Phys. J. C* **45** (2006) 97 [[hep-ph/0506026](#)] [[INSPIRE](#)].
- [44] J. Allison et al., *GEANT4 developments and applications*, *IEEE Trans. Nucl. Sci.* **53** (2006) 270 [[INSPIRE](#)].
- [45] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [46] LHCb collaboration, *The LHCb simulation application, Gauss: design, evolution and experience*, *J. Phys. Conf. Ser.* **331** (2011) 032023 [[INSPIRE](#)].
- [47] LHCb collaboration, *Determination of the $X(3872)$ meson quantum numbers*, *Phys. Rev. Lett.* **110** (2013) 222001 [[arXiv:1302.6269](#)] [[INSPIRE](#)].
- [48] LHCb collaboration, *Quantum numbers of the $X(3872)$ state and orbital angular momentum in its $\rho^0 J\psi$ decay*, *Phys. Rev. D* **92** (2015) 011102 [[arXiv:1504.06339](#)] [[INSPIRE](#)].
- [49] LHCb collaboration, *Observation of sizeable ω contribution to $\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi$ decays*, [arXiv:2204.12597](#) [[INSPIRE](#)].
- [50] K. Gottfried, *Hadronic transitions between quark-anti-quark bound states*, *Phys. Rev. Lett.* **40** (1978) 598 [[INSPIRE](#)].
- [51] M.B. Voloshin, *On dynamics of heavy quarks in nonperturbative QCD vacuum*, *Nucl. Phys. B* **154** (1979) 365 [[INSPIRE](#)].
- [52] M.E. Peskin, *Short distance analysis for heavy quark systems. 1. Diagrammatics*, *Nucl. Phys. B* **156** (1979) 365 [[INSPIRE](#)].
- [53] G. Bhanot and M.E. Peskin, *Short distance analysis for heavy quark systems. 2. Applications*, *Nucl. Phys. B* **156** (1979) 391 [[INSPIRE](#)].

- [54] M.B. Voloshin and V.I. Zakharov, *Measuring QCD anomalies in hadronic transitions between onium states*, *Phys. Rev. Lett.* **45** (1980) 688 [INSPIRE].
- [55] V.A. Novikov and M.A. Shifman, *Comment on the $\psi' \rightarrow J/\psi\pi\pi$ decay*, *Z. Phys. C* **8** (1981) 43 [INSPIRE].
- [56] LHCb collaboration, *Measurement of the track reconstruction efficiency at LHCb*, 2015 *JINST* **10** P02007 [arXiv:1408.1251] [INSPIRE].
- [57] LHCb collaboration, *Observation of $\Lambda_b^0 \rightarrow \psi(2S)pK^-$ and $\Lambda_b^0 \rightarrow J/\psi\pi^+\pi^-pK^-$ decays and a measurement of the Λ_b^0 baryon mass*, *JHEP* **05** (2016) 132 [arXiv:1603.06961] [INSPIRE].
- [58] LHCb collaboration, *Observation of the $\Lambda_b^0 \rightarrow \chi_{c1}(3872)pK^-$ decay*, *JHEP* **09** (2019) 028 [arXiv:1907.00954] [INSPIRE].
- [59] W.S. McCulloch and W. Pitts, *A logical calculus of the ideas immanent in nervous activity*, *Bull. Math. Biophys.* **5** (1943) 115.
- [60] F. Rosenblatt, *The perceptron: a probabilistic model for information storage and organization in the brain*, *Psychological Rev.* **65** (1958) 386.
- [61] J.-H. Zhong et al., *A program for the Bayesian neural network in the ROOT framework*, *Comput. Phys. Commun.* **182** (2011) 2655 [arXiv:1103.2854] [INSPIRE].
- [62] A. Powell et al., *Particle identification at LHCb*, *PoS ICHEP2010* (2010) 020.
- [63] W.D. Hulsbergen, *Decay chain fitting with a Kalman filter*, *Nucl. Instrum. Meth. A* **552** (2005) 566 [physics/0503191] [INSPIRE].
- [64] LHCb collaboration, *Study of the lineshape of the $\chi_{c1}(3872)$ state*, *Phys. Rev. D* **102** (2020) 092005 [arXiv:2005.13419] [INSPIRE].
- [65] S. Geisser, *Predictive inference: an introduction*, Chapman and Hall/CRC, New York, NY, U.S.A. (1993) [DOI:10.1201/9780203742310].
- [66] G. Punzi, *Sensitivity of searches for new signals and its optimization*, *eConf C030908* (2003) MODT002 [physics/0308063] [INSPIRE].
- [67] LHCb collaboration, *Study of B_c^+ meson decays to charmonia plus multihadron final states*, arXiv:2208.08660 [INSPIRE].
- [68] E. Byckling and K. Kajantie, *Particle kinematics*, John Wiley & Sons Inc., New York, NY, U.S.A. (1973).
- [69] S. Karlin and L.S. Shapley, *Geometry of moment spaces*, American Mathematical Society, Providence, RI, U.S.A. (1953).
- [70] LHCb collaboration, *Observation of J/ψ pair production in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Lett. B* **707** (2012) 52 [arXiv:1109.0963] [INSPIRE].
- [71] T. Skwarnicki, *A study of the radiative cascade transitions between the Υ' and Υ resonances*, Ph.D. thesis, Institute of Nuclear Physics, Krakow, Poland (1986) [INSPIRE].
- [72] C. Hanhart, Y.S. Kalashnikova, A.E. Kudryavtsev and A.V. Nefediev, *Reconciling the $X(3872)$ with the near-threshold enhancement in the $D^0\bar{D}^0$ final state*, *Phys. Rev. D* **76** (2007) 034007 [arXiv:0704.0605] [INSPIRE].
- [73] E. Braaten and J. Stapleton, *Analysis of $J/\psi\pi^+\pi^-$ and $D^0\bar{D}^0\pi^0$ decays of the $X(3872)$* , *Phys. Rev. D* **81** (2010) 014019 [arXiv:0907.3167] [INSPIRE].

- [74] Y.S. Kalashnikova and A.V. Nefediev, *Nature of $X(3872)$ from data*, *Phys. Rev. D* **80** (2009) 074004 [[arXiv:0907.4901](#)] [[INSPIRE](#)].
- [75] C. Hanhart, Y.S. Kalashnikova and A.V. Nefediev, *Interplay of quark and meson degrees of freedom in a near-threshold resonance: multi-channel case*, *Eur. Phys. J. A* **47** (2011) 101 [[arXiv:1106.1185](#)] [[INSPIRE](#)].
- [76] S.S. Wilks, *The large-sample distribution of the likelihood ratio for testing composite hypotheses*, *Annals Math. Statist.* **9** (1938) 60 [[INSPIRE](#)].
- [77] R. Aaij et al., *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in run 2*, *EPJ Tech. Instrum.* **6** (2019) 1 [[arXiv:1803.00824](#)] [[INSPIRE](#)].
- [78] M. Pivk and F.R. Le Diberder, *SPlot: a statistical tool to unfold data distributions*, *Nucl. Instrum. Meth. A* **555** (2005) 356 [[physics/0402083](#)] [[INSPIRE](#)].
- [79] LHCb collaboration, *First observation of $B_s^0 \rightarrow J/\psi f_0(980)$ decays*, *Phys. Lett. B* **698** (2011) 115 [[arXiv:1102.0206](#)] [[INSPIRE](#)].
- [80] LHCb collaboration, *Measurement of the CP violating phase ϕ_s in $\bar{B}_s^0 \rightarrow J/\psi f_0(980)$* , *Phys. Lett. B* **707** (2012) 497 [[arXiv:1112.3056](#)] [[INSPIRE](#)].
- [81] LHCb collaboration, *Measurement of resonant and CP components in $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays*, *Phys. Rev. D* **89** (2014) 092006 [[arXiv:1402.6248](#)] [[INSPIRE](#)].
- [82] BESIII collaboration, *Amplitude analysis of $D_s^+ \rightarrow \pi^+ \pi^- \pi^+$* , *Phys. Rev. D* **106** (2022) 112006 [[arXiv:2108.10050](#)] [[INSPIRE](#)].
- [83] LHCb collaboration, *Amplitude analysis of the $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$ decay*, [arXiv:2209.09840](#) [[INSPIRE](#)].
- [84] W. Ochs, *Spectroscopy with glueballs and the role of $f_0(1370)$* , *Acta Phys. Polon. Supp.* **6** (2013) 839 [[arXiv:1304.7634](#)] [[INSPIRE](#)].
- [85] W. Ochs, *The status of glueballs*, *J. Phys. G* **40** (2013) 043001 [[arXiv:1301.5183](#)] [[INSPIRE](#)].
- [86] S.M. Flatté, *Coupled-channel analysis of the $\pi\eta$ and $K\bar{K}$ systems near $K\bar{K}$ threshold*, *Phys. Lett. B* **63** (1976) 224 [[INSPIRE](#)].
- [87] D.V. Bugg, *Re-analysis of data on $a_0(1450)$ and $a_0(980)$* , *Phys. Rev. D* **78** (2008) 074023 [[arXiv:0808.2706](#)] [[INSPIRE](#)].
- [88] Student, *The probable error of a mean*, *Biometrika* **6** (1908) 1.
- [89] D. Martínez Santos and F. Dupertuis, *Mass distributions marginalized over per-event errors*, *Nucl. Instrum. Meth. A* **764** (2014) 150 [[arXiv:1312.5000](#)] [[INSPIRE](#)].
- [90] A. Rogozhnikov, *Reweighting with boosted decision trees*, *J. Phys. Conf. Ser.* **762** (2016) 012036 [[arXiv:1608.05806](#)] [[INSPIRE](#)].
- [91] LHCb collaboration, *Measurement of relative branching fractions of B decays to $\psi(2S)$ and J/ψ mesons*, *Eur. Phys. J. C* **72** (2012) 2118 [[arXiv:1205.0918](#)] [[INSPIRE](#)].
- [92] K. De Bruyn et al., *Branching ratio measurements of B_s decays*, *Phys. Rev. D* **86** (2012) 014027 [[arXiv:1204.1735](#)] [[INSPIRE](#)].
- [93] LHCb collaboration, *Measurement of the B_s effective lifetime in the $J/\psi f_0(980)$ final state*, *Phys. Rev. Lett.* **109** (2012) 152002 [[arXiv:1207.0878](#)] [[INSPIRE](#)].

- [94] D0 collaboration, B_s^0 lifetime measurement in the CP-odd decay channel $B_s^0 \rightarrow J/\psi f_0(980)$, *Phys. Rev. D* **94** (2016) 012001 [[arXiv:1603.01302](#)] [[INSPIRE](#)].
- [95] CMS collaboration, Measurement of b hadron lifetimes in pp collisions at $\sqrt{s} = 8$ TeV, *Eur. Phys. J. C* **78** (2018) 457 [Erratum *ibid.* **78** (2018) 561] [[arXiv:1710.08949](#)] [[INSPIRE](#)].
- [96] S. Ropertz, C. Hanhart and B. Kubis, A new parametrization for the scalar pion form factors, *Eur. Phys. J. C* **78** (2018) 1000 [[arXiv:1809.06867](#)] [[INSPIRE](#)].

The LHCb collaboration

R. Aaij [ID](#)³², A.S.W. Abdelmotteleb [ID](#)⁵⁰, C. Abellan Beteta⁴⁴, F. Abudinén [ID](#)⁵⁰, T. Ackernley [ID](#)⁵⁴, B. Adeva [ID](#)⁴⁰, M. Adinolfi [ID](#)⁴⁸, P. Adlarson [ID](#)⁷⁷, H. Afsharnia⁹, C. Agapopoulou [ID](#)¹³, C.A. Aidala [ID](#)⁷⁸, Z. Ajaltouni⁹, S. Akar [ID](#)⁵⁹, K. Akiba [ID](#)³², P. Albicocco [ID](#)²³, J. Albrecht [ID](#)¹⁵, F. Alessio [ID](#)⁴², M. Alexander [ID](#)⁵³, A. Alfonso Alberó [ID](#)³⁹, Z. Aliouche [ID](#)⁵⁶, P. Alvarez Cartelle [ID](#)⁴⁹, R. Amalric [ID](#)¹³, S. Amato [ID](#)², J.L. Amey [ID](#)⁴⁸, Y. Amhis [ID](#)^{11,42}, L. An [ID](#)⁴², L. Anderlini [ID](#)²², M. Andersson [ID](#)⁴⁴, A. Andreianov [ID](#)³⁸, M. Andreotti [ID](#)²¹, D. Andreou [ID](#)⁶², D. Ao [ID](#)⁶, F. Archilli [ID](#)^{31,t}, A. Artamonov [ID](#)³⁸, M. Artuso [ID](#)⁶², E. Aslanides [ID](#)¹⁰, M. Atzeni [ID](#)⁴⁴, B. Audurier [ID](#)⁷⁹, I.B. Bachiller Perea [ID](#)⁸, S. Bachmann [ID](#)¹⁷, M. Bachmayer [ID](#)⁴³, J.J. Back [ID](#)⁵⁰, A. Bailly-reyre¹³, P. Baladrón Rodríguez [ID](#)⁴⁰, V. Balagura [ID](#)¹², W. Baldini [ID](#)^{21,42}, J. Baptista de Souza Leite [ID](#)¹, M. Barbetti [ID](#)^{22,k}, R.J. Barlow [ID](#)⁵⁶, S. Barsuk [ID](#)¹¹, W. Barter [ID](#)⁵², M. Bartolini [ID](#)⁴⁹, F. Baryshnikov [ID](#)³⁸, J.M. Basels [ID](#)¹⁴, G. Bassi [ID](#)^{29,q}, B. Batsukh [ID](#)⁴, A. Battig [ID](#)¹⁵, A. Bay [ID](#)⁴³, A. Beck [ID](#)⁵⁰, M. Becker [ID](#)¹⁵, F. Bedeschi [ID](#)²⁹, I.B. Bediaga [ID](#)¹, A. Beiter⁶², S. Belin [ID](#)⁴⁰, V. Bellee [ID](#)⁴⁴, K. Belous [ID](#)³⁸, I. Belov [ID](#)³⁸, I. Belyaev [ID](#)³⁸, G. Benane [ID](#)¹⁰, G. Bencivenni [ID](#)²³, E. Ben-Haim [ID](#)¹³, A. Berezhnóy [ID](#)³⁸, R. Bernet [ID](#)⁴⁴, S. Bernet Andres [ID](#)⁷⁶, D. Berninghoff¹⁷, H.C. Bernstein⁶², C. Bertella [ID](#)⁵⁶, A. Bertolin [ID](#)²⁸, C. Betancourt [ID](#)⁴⁴, F. Betti [ID](#)⁴², I.a. Bezshyiko [ID](#)⁴⁴, J. Bhom [ID](#)³⁵, L. Bian [ID](#)⁶⁸, M.S. Bieker [ID](#)¹⁵, N.V. Biesuz [ID](#)²¹, P. Billoir [ID](#)¹³, A. Biolchini [ID](#)³², M. Birch [ID](#)⁵⁵, F.C.R. Bishop [ID](#)⁴⁹, A. Bitadze [ID](#)⁵⁶, A. Bizzeti [ID](#), M.P. Blago [ID](#)⁴⁹, T. Blake [ID](#)⁵⁰, F. Blanc [ID](#)⁴³, J.E. Blank [ID](#)¹⁵, S. Blusk [ID](#)⁶², D. Bobulska [ID](#)⁵³, V.B. Bocharnikov [ID](#)³⁸, J.A. Boelhave [ID](#)¹⁵, O. Boente Garcia [ID](#)¹², T. Boettcher [ID](#)⁵⁹, A. Boldyrev [ID](#)³⁸, C.S. Bolognani [ID](#)⁷⁴, R. Bolzonella [ID](#)^{21,j}, N. Bondar [ID](#)^{38,42}, F. Borgato [ID](#)²⁸, S. Borghi [ID](#)⁵⁶, M. Borsato [ID](#)¹⁷, J.T. Borsuk [ID](#)³⁵, S.A. Bouchiba [ID](#)⁴³, T.J.V. Bowcock [ID](#)⁵⁴, A. Boyer [ID](#)⁴², C. Bozzi [ID](#)²¹, M.J. Bradley⁵⁵, S. Braun [ID](#)⁶⁰, A. Brea Rodríguez [ID](#)⁴⁰, J. Brodzicka [ID](#)³⁵, A. Brossa Gonzalo [ID](#)⁴⁰, J. Brown [ID](#)⁵⁴, D. Brundu [ID](#)²⁷, A. Buonaura [ID](#)⁴⁴, L. Buonincontri [ID](#)²⁸, A.T. Burke [ID](#)⁵⁶, C. Burr [ID](#)⁴², A. Bursche⁶⁶, A. Butkevich [ID](#)³⁸, J.S. Butter [ID](#)³², J. Buytaert [ID](#)⁴², W. Byczynski [ID](#)⁴², S. Cadeddu [ID](#)²⁷, H. Cai⁶⁸, R. Calabrese [ID](#)^{21,j}, L. Calefice [ID](#)¹⁵, S. Cali [ID](#)²³, M. Calvi [ID](#)^{26,n}, M. Calvo Gomez [ID](#)⁷⁶, P. Campana [ID](#)²³, D.H. Campora Perez [ID](#)⁷⁴, A.F. Campoverde Quezada [ID](#)⁶, S. Capelli [ID](#)^{26,n}, L. Capriotti [ID](#)²⁰, A. Carbone [ID](#)^{20,h}, R. Cardinale [ID](#)^{24,l}, A. Cardini [ID](#)²⁷, P. Carniti [ID](#)^{26,n}, L. Carus¹⁴, A. Casais Vidal [ID](#)⁴⁰, R. Caspary [ID](#)¹⁷, G. Casse [ID](#)⁵⁴, M. Cattaneo [ID](#)⁴², G. Cavallero [ID](#)^{55,42}, V. Cavallini [ID](#)^{21,j}, S. Celani [ID](#)⁴³, J. Cerasoli [ID](#)¹⁰, D. Cervenkov [ID](#)⁵⁷, A.J. Chadwick [ID](#)⁵⁴, I. Chahrour [ID](#)⁷⁸, M.G. Chapman⁴⁸, M. Charles [ID](#)¹³, Ph. Charpentier [ID](#)⁴², C.A. Chavez Barajas [ID](#)⁵⁴, M. Chefdeville [ID](#)⁸, C. Chen [ID](#)¹⁰, S. Chen [ID](#)⁴, A. Chernov [ID](#)³⁵, S. Chernyshenko [ID](#)⁴⁶, V. Chobanova [ID](#)⁴⁰, S. Cholak [ID](#)⁴³, M. Chrzaszcz [ID](#)³⁵, A. Chubykin [ID](#)³⁸, V. Chulikov [ID](#)³⁸, P. Ciambriano [ID](#)²³, M.F. Cicala [ID](#)⁵⁰, X. Cid Vidal [ID](#)⁴⁰, G. Ciezarek [ID](#)⁴², P. Cifra [ID](#)⁴², G. Ciullo [ID](#)^{j,21}, P.E.L. Clarke [ID](#)⁵², M. Clemencic [ID](#)⁴², H.V. Cliff [ID](#)⁴⁹, J. Closier [ID](#)⁴², J.L. Cobbedick [ID](#)⁵⁶, V. Coco [ID](#)⁴², J. Cogan [ID](#)¹⁰, E. Cogneras [ID](#)⁹, L. Cojocariu [ID](#)³⁷, P. Collins [ID](#)⁴², T. Colombo [ID](#)⁴², L. Congedo [ID](#)¹⁹, A. Contu [ID](#)²⁷, N. Cooke [ID](#)⁴⁷, I. Corredoira [ID](#)⁴⁰, G. Corti [ID](#)⁴², B. Couturier [ID](#)⁴², D.C. Craik [ID](#)⁴⁴, M. Cruz Torres [ID](#)^{1,f}, R. Currie [ID](#)⁵², C.L. Da Silva [ID](#)⁶¹, S. Dadabaev [ID](#)³⁸, L. Dai [ID](#)⁶⁵, X. Dai [ID](#)⁵, E. Dall’Oco [ID](#)¹⁵, J. Dalseno [ID](#)⁴⁰, C. D’Ambrosio [ID](#)⁴², J. Daniel [ID](#)⁹, A. Danilina [ID](#)³⁸, P. d’Argent [ID](#)¹⁹, J.E. Davies [ID](#)⁵⁶, A. Davis [ID](#)⁵⁶, O. De Aguiar Francisco [ID](#)⁵⁶, J. de Boer [ID](#)⁴², K. De Bruyn [ID](#)⁷³, S. De Capua [ID](#)⁵⁶, M. De Cian [ID](#)⁴³,

U. De Freitas Carneiro Da Graca [ID](#)¹, E. De Lucia [ID](#)²³, J.M. De Miranda [ID](#)¹, L. De Paula [ID](#)²,
M. De Serio [ID](#)^{19,g}, D. De Simone [ID](#)⁴⁴, P. De Simone [ID](#)²³, F. De Vellis [ID](#)¹⁵, J.A. de Vries [ID](#)⁷⁴,
C.T. Dean [ID](#)⁶¹, F. Debernardis [ID](#)^{19,g}, D. Decamp [ID](#)⁸, V. Dedu [ID](#)¹⁰, L. Del Buono [ID](#)¹³,
B. Delaney [ID](#)⁵⁸, H.-P. Dembinski [ID](#)¹⁵, V. Denysenko [ID](#)⁴⁴, O. Deschamps [ID](#)⁹, F. Dettori [ID](#)^{27,i},
B. Dey [ID](#)⁷¹, P. Di Nezza [ID](#)²³, I. Diachkov [ID](#)³⁸, S. Didenko [ID](#)³⁸, L. Dieste Maronas⁴⁰, S. Ding [ID](#)⁶²,
V. Dobishuk [ID](#)⁴⁶, A. Dolmatov³⁸, C. Dong [ID](#)³, A.M. Donohoe [ID](#)¹⁸, F. Dordei [ID](#)²⁷,
A.C. dos Reis [ID](#)¹, L. Douglas⁵³, A.G. Downes [ID](#)⁸, P. Duda [ID](#)⁷⁵, M.W. Dudek [ID](#)³⁵, L. Dufour [ID](#)⁴²,
V. Duk [ID](#)⁷², P. Durante [ID](#)⁴², M. M. Duras [ID](#)⁷⁵, J.M. Durham [ID](#)⁶¹, D. Dutta [ID](#)⁵⁶, A. Dziurda [ID](#)³⁵,
A. Dzyuba [ID](#)³⁸, S. Easo [ID](#)⁵¹, U. Egede [ID](#)⁶³, V. Egorychev [ID](#)³⁸, C. Eirea Orro⁴⁰, S. Eisenhardt [ID](#)⁵²,
E. Ejopu [ID](#)⁵⁶, S. Ek-In [ID](#)⁴³, L. Eklund [ID](#)⁷⁷, M.E. Elashri [ID](#)⁵⁹, J. Ellbracht [ID](#)¹⁵, S. Ely [ID](#)⁵⁵,
A. Ene [ID](#)³⁷, E. Eppele [ID](#)⁵⁹, S. Escher [ID](#)¹⁴, J. Eschle [ID](#)⁴⁴, S. Esen [ID](#)⁴⁴, T. Evans [ID](#)⁵⁶,
F. Fabiano [ID](#)^{27,i}, L.N. Falcao [ID](#)¹, Y. Fan [ID](#)⁶, B. Fang [ID](#)^{11,68}, L. Fantini [ID](#)^{72,p}, M. Faria [ID](#)⁴³,
S. Farry [ID](#)⁵⁴, D. Fazzini [ID](#)^{26,n}, L.F. Felkowski [ID](#)⁷⁵, M. Feo [ID](#)⁴², M. Fernandez Gomez [ID](#)⁴⁰,
A.D. Fernez [ID](#)⁶⁰, F. Ferrari [ID](#)²⁰, L. Ferreira Lopes [ID](#)⁴³, F. Ferreira Rodrigues [ID](#)²,
S. Ferreres Sole [ID](#)³², M. Ferrillo [ID](#)⁴⁴, M. Ferro-Luzzi [ID](#)⁴², S. Filippov [ID](#)³⁸, R.A. Fini [ID](#)¹⁹,
M. Fiorini [ID](#)^{21,j}, M. Firlej [ID](#)³⁴, K.M. Fischer [ID](#)⁵⁷, D.S. Fitzgerald [ID](#)⁷⁸, C. Fitzpatrick [ID](#)⁵⁶,
T. Fiutowski [ID](#)³⁴, F. Fleuret [ID](#)¹², M. Fontana [ID](#)¹³, F. Fontanelli [ID](#)^{24,l}, R. Forty [ID](#)⁴²,
D. Foulds-Holt [ID](#)⁴⁹, V. Franco Lima [ID](#)⁵⁴, M. Franco Sevilla [ID](#)⁶⁰, M. Frank [ID](#)⁴², E. Franzoso [ID](#)^{21,j},
G. Frau [ID](#)¹⁷, C. Frei [ID](#)⁴², D.A. Friday [ID](#)⁵⁶, L. Frontini [ID](#)^{25,m}, J. Fu [ID](#)⁶, Q. Fuehring [ID](#)¹⁵,
T. Fulghesu [ID](#)¹³, E. Gabriel [ID](#)³², G. Galati [ID](#)^{19,g}, M.D. Galati [ID](#)³², A. Gallas Torreira [ID](#)⁴⁰,
D. Galli [ID](#)^{20,h}, S. Gambetta [ID](#)^{52,42}, M. Gandelman [ID](#)², P. Gandini [ID](#)²⁵, H.G. Gao [ID](#)⁶, Y. Gao [ID](#)⁷,
Y. Gao [ID](#)⁵, M. Garau [ID](#)^{27,i}, L.M. Garcia Martin [ID](#)⁵⁰, P. Garcia Moreno [ID](#)³⁹,
J. García Pardiñas [ID](#)⁴², B. Garcia Plana⁴⁰, F.A. Garcia Rosales [ID](#)¹², L. Garrido [ID](#)³⁹,
C. Gaspar [ID](#)⁴², R.E. Geertsema [ID](#)³², D. Gerick¹⁷, L.L. Gerken [ID](#)¹⁵, E. Gersabeck [ID](#)⁵⁶,
M. Gersabeck [ID](#)⁵⁶, T. Gershon [ID](#)⁵⁰, L. Giambastiani [ID](#)²⁸, V. Gibson [ID](#)⁴⁹, H.K. Gienza [ID](#)³⁶,
A.L. Gilman [ID](#)⁵⁷, M. Giovannetti [ID](#)²³, A. Gioventù [ID](#)⁴⁰, P. Gironella Gironell [ID](#)³⁹,
C. Giugliano [ID](#)^{21,j}, M.A. Giza [ID](#)³⁵, K. Gizdov [ID](#)⁵², E.L. Gkougkousis [ID](#)⁴², V.V. Gligorov [ID](#)^{13,42},
C. Göbel [ID](#)⁶⁴, E. Golobardes [ID](#)⁷⁶, D. Golubkov [ID](#)³⁸, A. Golutvin [ID](#)^{55,38}, A. Gomes [ID](#)^{1,2,b,a,†},
S. Gomez Fernandez [ID](#)³⁹, F. Goncalves Abrantes [ID](#)⁵⁷, M. Goncerz [ID](#)³⁵, G. Gong [ID](#)³,
I.V. Gorelov [ID](#)³⁸, C. Gotti [ID](#)²⁶, J.P. Grabowski [ID](#)⁷⁰, T. Grammatico [ID](#)¹³,
L.A. Granado Cardoso [ID](#)⁴², E. Graugés [ID](#)³⁹, E. Graverini [ID](#)⁴³, G. Graziani [ID](#), A. T. Grecu [ID](#)³⁷,
L.M. Greeven [ID](#)³², N.A. Grieser [ID](#)⁵⁹, L. Grillo [ID](#)⁵³, S. Gromov [ID](#)³⁸, B.R. Gruber Cazon [ID](#)⁵⁷, C.
Gu [ID](#)³, M. Guarise [ID](#)^{21,j}, M. Guittiere [ID](#)¹¹, P. A. Günther [ID](#)¹⁷, E. Gushchin [ID](#)³⁸, A. Guth¹⁴,
Y. Guz [ID](#)^{5,38,42}, T. Gys [ID](#)⁴², T. Hadavizadeh [ID](#)⁶³, C. Hadjivasiliou [ID](#)⁶⁰, G. Haefeli [ID](#)⁴³,
C. Haen [ID](#)⁴², J. Haimberger [ID](#)⁴², S.C. Haines [ID](#)⁴⁹, T. Halewood-leagas [ID](#)⁵⁴, M.M. Halvorsen [ID](#)⁴²,
P.M. Hamilton [ID](#)⁶⁰, J. Hammerich [ID](#)⁵⁴, Q. Han [ID](#)⁷, X. Han [ID](#)¹⁷, S. Hansmann-Menzemer [ID](#)¹⁷,
L. Hao [ID](#)⁶, N. Harnew [ID](#)⁵⁷, T. Harrison [ID](#)⁵⁴, C. Hasse [ID](#)⁴², M. Hatch [ID](#)⁴², J. He [ID](#)^{6,d},
K. Heijhoff [ID](#)³², F.H. Hemmer [ID](#)⁴², C. Henderson [ID](#)⁵⁹, R.D.L. Henderson [ID](#)^{63,50},
A.M. Hennequin [ID](#)⁵⁸, K. Hennessy [ID](#)⁵⁴, L. Henry [ID](#)⁴², J. Herd [ID](#)⁵⁵, J. Heuel [ID](#)¹⁴, A. Hicheur [ID](#)²,
D. Hill [ID](#)⁴³, M. Hilton [ID](#)⁵⁶, S.E. Hollitt [ID](#)¹⁵, J. Horswill [ID](#)⁵⁶, R. Hou [ID](#)⁷, Y. Hou [ID](#)⁸, J. Hu¹⁷,
J. Hu [ID](#)⁶⁶, W. Hu [ID](#)⁵, X. Hu [ID](#)³, W. Huang [ID](#)⁶, X. Huang⁶⁸, W. Hulsbergen [ID](#)³², R.J. Hunter [ID](#)⁵⁰,
M. Hushchyn [ID](#)³⁸, D. Hutchcroft [ID](#)⁵⁴, P. Ibis [ID](#)¹⁵, M. Idzik [ID](#)³⁴, D. Ilin [ID](#)³⁸, P. Ilten [ID](#)⁵⁹,
A. Inglessi [ID](#)³⁸, A. Iniukhin [ID](#)³⁸, A. Ishteev [ID](#)³⁸, K. Ivshin [ID](#)³⁸, R. Jacobsson [ID](#)⁴², H. Jage [ID](#)¹⁴,

S.J. Jaimes Elles [ID](#)⁴¹, S. Jakobsen [ID](#)⁴², E. Jans [ID](#)³², B.K. Jashal [ID](#)⁴¹, A. Jawahery [ID](#)⁶⁰,
 V. Jevtic [ID](#)¹⁵, E. Jiang [ID](#)⁶⁰, X. Jiang [ID](#)^{4,6}, Y. Jiang [ID](#)⁶, M. John [ID](#)⁵⁷, D. Johnson [ID](#)⁵⁸,
 C.R. Jones [ID](#)⁴⁹, T.P. Jones [ID](#)⁵⁰, S.J. Joshi [ID](#)³⁶, B. Jost [ID](#)⁴², N. Jurik [ID](#)⁴², I. Juszczak [ID](#)³⁵,
 S. Kandybei [ID](#)⁴⁵, Y. Kang [ID](#)³, M. Karacson [ID](#)⁴², D. Karpenkov [ID](#)³⁸, M. Karpov [ID](#)³⁸,
 J.W. Kautz [ID](#)⁵⁹, F. Keizer [ID](#)⁴², D.M. Keller [ID](#)⁶², M. Kenzie [ID](#)⁵⁰, T. Ketel [ID](#)³², B. Khanji [ID](#)¹⁵,
 A. Kharisova [ID](#)³⁸, S. Kholodenko [ID](#)³⁸, G. Khreich [ID](#)¹¹, T. Kirn [ID](#)¹⁴, V.S. Kirsebom [ID](#)⁴³,
 O. Kitouni [ID](#)⁵⁸, S. Klaver [ID](#)³³, N. Kleijne [ID](#)^{29,q}, K. Klimaszewski [ID](#)³⁶, M.R. Kmiec [ID](#)³⁶,
 S. Koliiev [ID](#)⁴⁶, L. Kolk [ID](#)¹⁵, A. Kondybayeva [ID](#)³⁸, A. Konoplyannikov [ID](#)³⁸, P. Kopciwicz [ID](#)³⁴,
 R. Kopecna¹⁷, P. Koppenburg [ID](#)³², M. Korolev [ID](#)³⁸, I. Kostiuk [ID](#)³², O. Kot⁴⁶, S. Kotriakhova [ID](#),
 A. Kozachuk [ID](#)³⁸, P. Kravchenko [ID](#)³⁸, L. Kravchuk [ID](#)³⁸, M. Kreps [ID](#)⁵⁰, S. Kretzschmar [ID](#)¹⁴,
 P. Krokovny [ID](#)³⁸, W. Krupa [ID](#)³⁴, W. Krzemien [ID](#)³⁶, J. Kubat¹⁷, S. Kubis [ID](#)⁷⁵, W. Kucewicz [ID](#)³⁵,
 M. Kucharczyk [ID](#)³⁵, V. Kudryavtsev [ID](#)³⁸, E.K. Kulikova [ID](#)³⁸, A. Kupsc [ID](#)⁷⁷, D. Lacarrere [ID](#)⁴²,
 G. Lafferty [ID](#)⁵⁶, A. Lai [ID](#)²⁷, A. Lampis [ID](#)^{27,i}, D. Lancierini [ID](#)⁴⁴, C. Landesa Gomez [ID](#)⁴⁰,
 J.J. Lane [ID](#)⁵⁶, R. Lane [ID](#)⁴⁸, C. Langenbruch [ID](#)¹⁴, J. Langer [ID](#)¹⁵, O. Lantwin [ID](#)³⁸, T. Latham [ID](#)⁵⁰,
 F. Lazzari [ID](#)^{29,r}, C. Lazzeroni [ID](#)⁴⁷, R. Le Gac [ID](#)¹⁰, S.H. Lee [ID](#)⁷⁸, R. Lefèvre [ID](#)⁹, A. Leflat [ID](#)³⁸,
 S. Legotin [ID](#)³⁸, P. Lenisa [ID](#)^{j,21}, O. Leroy [ID](#)¹⁰, T. Lesiak [ID](#)³⁵, B. Leverington [ID](#)¹⁷, A. Li [ID](#)³,
 H. Li [ID](#)⁶⁶, K. Li [ID](#)⁷, P. Li [ID](#)⁴², P.-R. Li [ID](#)⁶⁷, S. Li [ID](#)⁷, T. Li [ID](#)⁴, T. Li [ID](#)⁶⁶, Y. Li [ID](#)⁴, Z. Li [ID](#)⁶²,
 X. Liang [ID](#)⁶², C. Lin [ID](#)⁶, T. Lin [ID](#)⁵¹, R. Lindner [ID](#)⁴², V. Lisovskyi [ID](#)¹⁵, R. Litvinov [ID](#)^{27,i},
 G. Liu [ID](#)⁶⁶, H. Liu [ID](#)⁶, K. Liu [ID](#)⁶⁷, Q. Liu [ID](#)⁶, S. Liu [ID](#)^{4,6}, A. Lobo Salvia [ID](#)³⁹, A. Loi [ID](#)²⁷,
 R. Lollini [ID](#)⁷², J. Lomba Castro [ID](#)⁴⁰, I. Longstaff⁵³, J.H. Lopes [ID](#)², A. Lopez Huertas [ID](#)³⁹,
 S. López Soliño [ID](#)⁴⁰, G.H. Lovell [ID](#)⁴⁹, Y. Lu [ID](#)^{4,c}, C. Lucarelli [ID](#)^{22,k}, D. Lucchesi [ID](#)^{28,o},
 S. Luchuk [ID](#)³⁸, M. Lucio Martinez [ID](#)⁷⁴, V. Lukashenko [ID](#)^{32,46}, Y. Luo [ID](#)³, A. Lupato [ID](#)⁵⁶,
 E. Luppi [ID](#)^{21,j}, A. Lusiani [ID](#)^{29,q}, K. Lynch [ID](#)¹⁸, X.-R. Lyu [ID](#)⁶, R. Ma [ID](#)⁶, S. Maccolini [ID](#)¹⁵,
 F. Machefert [ID](#)¹¹, F. Maciuc [ID](#)³⁷, I. Mackay [ID](#)⁵⁷, V. Macko [ID](#)⁴³, L.R. Madhan Mohan [ID](#)⁴⁹,
 A. Maevskiy [ID](#)³⁸, D. Maisuzenko [ID](#)³⁸, M.W. Majewski³⁴, J.J. Malczewski [ID](#)³⁵, S. Malde [ID](#)⁵⁷,
 B. Malecki [ID](#)^{35,42}, A. Malinin [ID](#)³⁸, T. Maltsev [ID](#)³⁸, G. Manca [ID](#)^{27,i}, G. Mancinelli [ID](#)¹⁰,
 C. Mancuso [ID](#)^{11,25,m}, R. Manera Escalero³⁹, D. Manuzzi [ID](#)²⁰, C.A. Manzari [ID](#)⁴⁴,
 D. Marangotto [ID](#)^{25,m}, J.F. Marchand [ID](#)⁸, U. Marconi [ID](#)²⁰, S. Mariani [ID](#)⁴², C. Marin Benito [ID](#)³⁹,
 J. Marks [ID](#)¹⁷, A.M. Marshall [ID](#)⁴⁸, P.J. Marshall⁵⁴, G. Martelli [ID](#)^{72,p}, G. Martellotti [ID](#)³⁰,
 L. Martinazzoli [ID](#)^{42,n}, M. Martinelli [ID](#)^{26,n}, D. Martinez Santos [ID](#)⁴⁰, F. Martinez Vidal [ID](#)⁴¹,
 A. Massafferri [ID](#)¹, M. Materok [ID](#)¹⁴, R. Matev [ID](#)⁴², A. Mathad [ID](#)⁴⁴, V. Matiunin [ID](#)³⁸,
 C. Matteuzzi [ID](#)²⁶, K.R. Mattioli [ID](#)¹², A. Mauri [ID](#)⁵⁵, E. Maurice [ID](#)¹², J. Mauricio [ID](#)³⁹,
 M. Mazurek [ID](#)⁴², M. McCann [ID](#)⁵⁵, L. McConnell [ID](#)¹⁸, T.H. McGrath [ID](#)⁵⁶, N.T. McHugh [ID](#)⁵³,
 A. McNab [ID](#)⁵⁶, R. McNulty [ID](#)¹⁸, B. Meadows [ID](#)⁵⁹, G. Meier [ID](#)¹⁵, D. Melnychuk [ID](#)³⁶,
 S. Meloni [ID](#)^{26,n}, M. Merk [ID](#)^{32,74}, A. Merli [ID](#)^{25,m}, L. Meyer Garcia [ID](#)², D. Miao [ID](#)^{4,6}, H. Miao [ID](#)⁶,
 M. Mikhasenko [ID](#)^{70,e}, D.A. Milanes [ID](#)⁶⁹, E. Millard⁵⁰, M. Milovanovic [ID](#)⁴², M.-N. Minard^{8,†},
 A. Minotti [ID](#)^{26,n}, E. Minucci [ID](#)⁶², T. Miralles [ID](#)⁹, S.E. Mitchell [ID](#)⁵², B. Mitreska [ID](#)¹⁵,
 D.S. Mitzel [ID](#)¹⁵, A. Modak [ID](#)⁵¹, A. Mödden [ID](#)¹⁵, R.A. Mohammed [ID](#)⁵⁷, R.D. Moise [ID](#)¹⁴,
 S. Mokhnenko [ID](#)³⁸, T. Mombächer [ID](#)⁴⁰, M. Monk [ID](#)^{50,63}, I.A. Monroy [ID](#)⁶⁹, S. Monteil [ID](#)⁹,
 G. Morello [ID](#)²³, M.J. Morello [ID](#)^{29,q}, M.P. Morgenthaler [ID](#)¹⁷, J. Moron [ID](#)³⁴, A.B. Morris [ID](#)⁴²,
 A.G. Morris [ID](#)¹⁰, R. Mountain [ID](#)⁶², H. Mu [ID](#)³, E. Muhammad [ID](#)⁵⁰, F. Muheim [ID](#)⁵²,
 M. Mulder [ID](#)⁷³, K. Müller [ID](#)⁴⁴, C.H. Murphy [ID](#)⁵⁷, D. Murray [ID](#)⁵⁶, R. Murta [ID](#)⁵⁵,
 P. Muzzetto [ID](#)^{27,i}, P. Naik [ID](#)⁴⁸, T. Nakada [ID](#)⁴³, R. Nandakumar [ID](#)⁵¹, T. Nanut [ID](#)⁴², I. Nasteva [ID](#)²,

M. Needham [ID](#)⁵², N. Neri [ID](#)^{25,m}, S. Neubert [ID](#)⁷⁰, N. Neufeld [ID](#)⁴², P. Neustroev³⁸, R. Newcombe⁵⁵,
 J. Nicolini [ID](#)^{15,11}, D. Nicotra [ID](#)⁷⁴, E.M. Niel [ID](#)⁴³, S. Nieswand¹⁴, N. Nikitin [ID](#)³⁸, N.S. Nolte [ID](#)⁵⁸,
 C. Normand [ID](#)^{8,i,27}, J. Novoa Fernandez [ID](#)⁴⁰, G.N Nowak [ID](#)⁵⁹, C. Nunez [ID](#)⁷⁸,
 A. Oblakowska-Mucha [ID](#)³⁴, V. Obraztsov [ID](#)³⁸, T. Oeser [ID](#)¹⁴, S. Okamura [ID](#)^{21,j}, R. Oldeman [ID](#)^{27,i},
 F. Oliva [ID](#)⁵², C.J.G. Onderwater [ID](#)⁷³, R.H. O’Neil [ID](#)⁵², J.M. Otalora Goicochea [ID](#)²,
 T. Ovsianikova [ID](#)³⁸, P. Owen [ID](#)⁴⁴, A. Oyanguren [ID](#)⁴¹, O. Ozcelik [ID](#)⁵², K.O. Padeken [ID](#)⁷⁰,
 B. Pagare [ID](#)⁵⁰, P.R. Pais [ID](#)⁴², T. Pajero [ID](#)⁵⁷, A. Palano [ID](#)¹⁹, M. Palutan [ID](#)²³, G. Panshin [ID](#)³⁸,
 L. Paolucci [ID](#)⁵⁰, A. Papanestis [ID](#)⁵¹, M. Pappagallo [ID](#)^{19,g}, L.L. Pappalardo [ID](#)^{21,j},
 C. Pappenheimer [ID](#)⁵⁹, W. Parker [ID](#)⁶⁰, C. Parkes [ID](#)^{56,42}, B. Passalacqua [ID](#)^{21,j}, G. Passaleva [ID](#)²²,
 A. Pastore [ID](#)¹⁹, M. Patel [ID](#)⁵⁵, C. Patrignani [ID](#)^{20,h}, C.J. Pawley [ID](#)⁷⁴, A. Pellegrino [ID](#)³²,
 M. Pepe Altarelli [ID](#)⁴², S. Perazzini [ID](#)²⁰, D. Pereima [ID](#)³⁸, A. Pereiro Castro [ID](#)⁴⁰, P. Perret [ID](#)⁹,
 K. Petridis [ID](#)⁴⁸, A. Petrolini [ID](#)^{24,l}, S. Petrucci [ID](#)⁵², M. Petruzzo [ID](#)²⁵, H. Pham [ID](#)⁶²,
 A. Philippov [ID](#)³⁸, R. Piandani [ID](#)⁶, L. Pica [ID](#)^{29,q}, M. Piccini [ID](#)⁷², B. Pietrzyk [ID](#)⁸, G. Pietrzyk [ID](#)¹¹,
 M. Pili [ID](#)⁵⁷, D. Pinci [ID](#)³⁰, F. Pisani [ID](#)⁴², M. Pizzichemi [ID](#)^{26,n,42}, V. Placinta [ID](#)³⁷, J. Plews [ID](#)⁴⁷,
 M. Plo Casasus [ID](#)⁴⁰, F. Polci [ID](#)^{13,42}, M. Poli Lener [ID](#)²³, A. Poluektov [ID](#)¹⁰, N. Polukhina [ID](#)³⁸,
 I. Polyakov [ID](#)⁴², E. Polycarpo [ID](#)², S. Ponce [ID](#)⁴², D. Popov [ID](#)^{6,42}, S. Poslavskii [ID](#)³⁸,
 K. Prasanth [ID](#)³⁵, L. Promberger [ID](#)¹⁷, C. Prouve [ID](#)⁴⁰, V. Pugatch [ID](#)⁴⁶, V. Puill [ID](#)¹¹, G. Punzi [ID](#)^{29,r},
 H.R. Qi [ID](#)³, W. Qian [ID](#)⁶, N. Qin [ID](#)³, S. Qu [ID](#)³, R. Quagliani [ID](#)⁴³, N.V. Raab [ID](#)¹⁸, B. Rachwal [ID](#)³⁴,
 J.H. Rademacker [ID](#)⁴⁸, R. Rajagopalan⁶², M. Rama [ID](#)²⁹, M. Ramos Pernas [ID](#)⁵⁰, M.S. Rangel [ID](#)²,
 F. Ratnikov [ID](#)³⁸, G. Raven [ID](#)³³, M. Rebollo De Miguel [ID](#)⁴¹, F. Redi [ID](#)⁴², J. Reich [ID](#)⁴⁸,
 F. Reiss [ID](#)⁵⁶, C. Remon Alepuz⁴¹, Z. Ren [ID](#)³, P.K. Resmi [ID](#)⁵⁷, R. Ribatti [ID](#)^{29,q}, A.M. Ricci [ID](#)²⁷,
 S. Ricciardi [ID](#)⁵¹, K. Richardson [ID](#)⁵⁸, M. Richardson-Slipper [ID](#)⁵², K. Rinnert [ID](#)⁵⁴, P. Robbe [ID](#)¹¹,
 G. Robertson [ID](#)⁵², E. Rodrigues [ID](#)^{54,42}, E. Rodriguez Fernandez [ID](#)⁴⁰, J.A. Rodriguez Lopez [ID](#)⁶⁹,
 E. Rodriguez Rodriguez [ID](#)⁴⁰, D.L. Rolf [ID](#)⁴², A. Rollings [ID](#)⁵⁷, P. Roloff [ID](#)⁴², V. Romanovskiy [ID](#)³⁸,
 M. Romero Lamas [ID](#)⁴⁰, A. Romero Vidal [ID](#)⁴⁰, J.D. Roth^{78,†}, M. Rotondo [ID](#)²³, M.S. Rudolph [ID](#)⁶²,
 T. Ruf [ID](#)⁴², R.A. Ruiz Fernandez [ID](#)⁴⁰, J. Ruiz Vidal [ID](#)⁴¹, A. Ryzhikov [ID](#)³⁸, J. Ryzka [ID](#)³⁴,
 J.J. Saborido Silva [ID](#)⁴⁰, N. Sagidova [ID](#)³⁸, N. Sahoo [ID](#)⁴⁷, B. Saitta [ID](#)^{27,i}, M. Salomoni [ID](#)⁴²,
 C. Sanchez Gras [ID](#)³², I. Sanderswood [ID](#)⁴¹, R. Santacesaria [ID](#)³⁰, C. Santamarina Rios [ID](#)⁴⁰,
 M. Santimaria [ID](#)²³, E. Santovetti [ID](#)^{31,t}, D. Saranin [ID](#)³⁸, G. Sarpis [ID](#)¹⁴, M. Sarpis [ID](#)⁷⁰, A. Sarti [ID](#)³⁰,
 C. Satriano [ID](#)^{30,s}, A. Satta [ID](#)³¹, M. Saur [ID](#)¹⁵, D. Savrina [ID](#)³⁸, H. Sazak [ID](#)⁹,
 L.G. Scantlebury Smead [ID](#)⁵⁷, A. Scarabotto [ID](#)¹³, S. Schael [ID](#)¹⁴, S. Scherl [ID](#)⁵⁴, A. M. Schertz [ID](#)⁷¹,
 M. Schiller [ID](#)⁵³, H. Schindler [ID](#)⁴², M. Schmelling [ID](#)¹⁶, B. Schmidt [ID](#)⁴², S. Schmitt [ID](#)¹⁴,
 O. Schneider [ID](#)⁴³, A. Schopper [ID](#)⁴², M. Schubiger [ID](#)³², N. Schulte [ID](#)¹⁵, S. Schulte [ID](#)⁴³,
 M.H. Schune [ID](#)¹¹, R. Schwemmer [ID](#)⁴², B. Sciascia [ID](#)²³, A. Sciuccati [ID](#)⁴², S. Sellam [ID](#)⁴⁰,
 A. Semennikov [ID](#)³⁸, M. Senghi Soares [ID](#)³³, A. Sergi [ID](#)^{24,l}, N. Serra [ID](#)⁴⁴, L. Sestini [ID](#)²⁸,
 A. Seuthe [ID](#)¹⁵, Y. Shang [ID](#)⁵, D.M. Shangase [ID](#)⁷⁸, M. Shapkin [ID](#)³⁸, I. Shchernerov [ID](#)³⁸,
 L. Shchutska [ID](#)⁴³, T. Shears [ID](#)⁵⁴, L. Shekhtman [ID](#)³⁸, Z. Shen [ID](#)⁵, S. Sheng [ID](#)^{4,6},
 V. Shevchenko [ID](#)³⁸, B. Shi [ID](#)⁶, E.B. Shields [ID](#)^{26,n}, Y. Shimizu [ID](#)¹¹, E. Shmanin [ID](#)³⁸,
 R. Shorkin [ID](#)³⁸, J.D. Shupperd [ID](#)⁶², B.G. Siddi [ID](#)^{21,j}, R. Silva Coutinho [ID](#)⁶², G. Simi [ID](#)²⁸,
 S. Simone [ID](#)^{19,g}, M. Singla [ID](#)⁶³, N. Skidmore [ID](#)⁵⁶, R. Skuza [ID](#)¹⁷, T. Skwarnicki [ID](#)⁶²,
 M.W. Slater [ID](#)⁴⁷, J.C. Smallwood [ID](#)⁵⁷, J.G. Smeaton [ID](#)⁴⁹, E. Smith [ID](#)⁴⁴, K. Smith [ID](#)⁶¹,
 M. Smith [ID](#)⁵⁵, A. Snoch [ID](#)³², L. Soares Lavra [ID](#)⁹, M.D. Sokoloff [ID](#)⁵⁹, F.J.P. Soler [ID](#)⁵³,
 A. Solomin [ID](#)^{38,48}, A. Solovev [ID](#)³⁸, I. Solovyev [ID](#)³⁸, R. Song [ID](#)⁶³, F.L. Souza De Almeida [ID](#)²,

B. Souza De Paula ², B. Spaan ^{15,†}, E. Spadaro Norella ^{25,m}, E. Spedicato ²⁰, J.G. Speer ¹⁵, E. Spiridenkov ³⁸, P. Spradlin ⁵³, V. Sriskaran ⁴², F. Stagni ⁴², M. Stahl ⁴², S. Stahl ⁴², S. Stanislaus ⁵⁷, E.N. Stein ⁴², O. Steinkamp ⁴⁴, O. Stenyakin ³⁸, H. Stevens ¹⁵, D. Strelalina ³⁸, Y.S. Su ⁶, F. Suljik ⁵⁷, J. Sun ²⁷, L. Sun ⁶⁸, Y. Sun ⁶⁰, P.N. Swallow ⁴⁷, K. Swientek ³⁴, A. Szabelski ³⁶, T. Szumlak ³⁴, M. Szymanski ⁴², Y. Tan ³, S. Taneja ⁵⁶, M.D. Tat ⁵⁷, A. Terentev ⁴⁴, F. Teubert ⁴², E. Thomas ⁴², D.J.D. Thompson ⁴⁷, H. Tilquin ⁵⁵, V. Tisserand ⁹, S. T'Jampens ⁸, M. Tobin ⁴, L. Tomassetti ^{21,j}, G. Tonani ^{25,m}, X. Tong ⁵, D. Torres Machado ¹, D.Y. Tou ³, C. Tripl ⁴³, G. Tuci ⁶, N. Tuning ³², A. Ukleja ³⁶, D.J. Unverzagt ¹⁷, A. Usachov ³³, A. Ustyuzhanin ³⁸, U. Uwer ¹⁷, V. Vagnoni ²⁰, A. Valassi ⁴², G. Valenti ²⁰, N. Valls Canudas ⁷⁶, M. Van Dijk ⁴³, H. Van Hecke ⁶¹, E. van Herwijnen ⁵⁵, C.B. Van Hulse ^{40,v}, M. van Veghel ³², R. Vazquez Gomez ³⁹, P. Vazquez Regueiro ⁴⁰, C. Vázquez Sierra ⁴², S. Vecchi ²¹, J.J. Velthuis ⁴⁸, M. Veltri ^{22,u}, A. Venkateswaran ⁴³, M. Veronesi ³², M. Vesterinen ⁵⁰, D. Vieira ⁵⁹, M. Vieites Diaz ⁴³, X. Vilasis-Cardona ⁷⁶, E. Vilella Figueras ⁵⁴, A. Villa ²⁰, P. Vincent ¹³, F.C. Volle ¹¹, D. vom Bruch ¹⁰, V. Vorobyev ³⁸, N. Voropaev ³⁸, K. Vos ⁷⁴, C. Vrahas ⁵², J. Walsh ²⁹, E.J. Walton ⁶³, G. Wan ⁵, C. Wang ¹⁷, G. Wang ⁷, J. Wang ⁵, J. Wang ⁴, J. Wang ³, J. Wang ⁶⁸, M. Wang ²⁵, R. Wang ⁴⁸, X. Wang ⁶⁶, Y. Wang ⁷, Z. Wang ⁴⁴, Z. Wang ³, Z. Wang ⁶, J.A. Ward ^{50,63}, N.K. Watson ⁴⁷, D. Websdale ⁵⁵, Y. Wei ⁵, B.D.C. Westhenry ⁴⁸, D.J. White ⁵⁶, M. Whitehead ⁵³, A.R. Wiederhold ⁵⁰, D. Wiedner ¹⁵, G. Wilkinson ⁵⁷, M.K. Wilkinson ⁵⁹, I. Williams ⁴⁹, M. Williams ⁵⁸, M.R.J. Williams ⁵², R. Williams ⁴⁹, F.F. Wilson ⁵¹, W. Wislicki ³⁶, M. Witek ³⁵, L. Witola ¹⁷, C.P. Wong ⁶¹, G. Wormser ¹¹, S.A. Wotton ⁴⁹, H. Wu ⁶², J. Wu ⁷, K. Wyllie ⁴², Z. Xiang ⁶, Y. Xie ⁷, A. Xu ⁵, J. Xu ⁶, L. Xu ³, L. Xu ³, M. Xu ⁵⁰, Q. Xu ⁶, Z. Xu ⁹, Z. Xu ⁶, D. Yang ³, S. Yang ⁶, X. Yang ⁵, Y. Yang ⁶, Z. Yang ⁵, Z. Yang ⁶⁰, L.E. Yeomans ⁵⁴, V. Yeroshenko ¹¹, H. Yeung ⁵⁶, H. Yin ⁷, J. Yu ⁶⁵, X. Yuan ⁶², E. Zaffaroni ⁴³, M. Zavertyaev ¹⁶, M. Zdybal ³⁵, M. Zeng ³, C. Zhang ⁵, D. Zhang ⁷, L. Zhang ³, S. Zhang ⁶⁵, S. Zhang ⁵, Y. Zhang ⁵, Y. Zhang ⁵⁷, Y. Zhao ¹⁷, A. Zharkova ³⁸, A. Zhelezov ¹⁷, Y. Zheng ⁶, T. Zhou ⁵, X. Zhou ⁷, Y. Zhou ⁶, V. Zhovkovska ¹¹, X. Zhu ³, X. Zhu ⁷, Z. Zhu ⁶, V. Zhukov ^{14,38}, Q. Zou ^{4,6}, S. Zucchelli ^{20,h}, D. Zuliani ²⁸, G. Zunica ⁵⁶

¹ Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

² Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³ Center for High Energy Physics, Tsinghua University, Beijing, China

⁴ Institute Of High Energy Physics (IHEP), Beijing, China

⁵ School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁶ University of Chinese Academy of Sciences, Beijing, China

⁷ Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁸ Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁹ Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

¹⁰ Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

¹¹ Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

¹² Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

¹³ LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

- ¹⁴ *I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ¹⁵ *Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ¹⁶ *Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ¹⁷ *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ¹⁸ *School of Physics, University College Dublin, Dublin, Ireland*
- ¹⁹ *INFN Sezione di Bari, Bari, Italy*
- ²⁰ *INFN Sezione di Bologna, Bologna, Italy*
- ²¹ *INFN Sezione di Ferrara, Ferrara, Italy*
- ²² *INFN Sezione di Firenze, Firenze, Italy*
- ²³ *INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ²⁴ *INFN Sezione di Genova, Genova, Italy*
- ²⁵ *INFN Sezione di Milano, Milano, Italy*
- ²⁶ *INFN Sezione di Milano-Bicocca, Milano, Italy*
- ²⁷ *INFN Sezione di Cagliari, Monserrato, Italy*
- ²⁸ *Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy*
- ²⁹ *INFN Sezione di Pisa, Pisa, Italy*
- ³⁰ *INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³¹ *INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³² *Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³³ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ³⁴ *AGH — University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ³⁵ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ³⁶ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³⁷ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³⁸ *Affiliated with an institute covered by a cooperation agreement with CERN*
- ³⁹ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁰ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴¹ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia — CSIC, Valencia, Spain*
- ⁴² *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴³ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁴ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁵ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁶ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁷ *University of Birmingham, Birmingham, United Kingdom*
- ⁴⁸ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁹ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁰ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵¹ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵² *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵³ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁴ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁵ *Imperial College London, London, United Kingdom*
- ⁵⁶ *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁷ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁸ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁹ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁰ *University of Maryland, College Park, MD, United States*
- ⁶¹ *Los Alamos National Laboratory (LANL), Los Alamos, NM, United States*
- ⁶² *Syracuse University, Syracuse, NY, United States*

- ⁶³ School of Physics and Astronomy, Monash University, Melbourne, Australia, associated to ⁵⁰
- ⁶⁴ Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²
- ⁶⁵ Physics and Micro Electronic College, Hunan University, Changsha City, China, associated to ⁷
- ⁶⁶ Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to ³
- ⁶⁷ Lanzhou University, Lanzhou, China, associated to ⁴
- ⁶⁸ School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³
- ⁶⁹ Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ¹³
- ⁷⁰ Universität Bonn — Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany, associated to ¹⁷
- ⁷¹ Eotvos Lorand University, Budapest, Hungary, associated to ⁴²
- ⁷² INFN Sezione di Perugia, Perugia, Italy, associated to ²¹
- ⁷³ Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ³²
- ⁷⁴ Universiteit Maastricht, Maastricht, Netherlands, associated to ³²
- ⁷⁵ Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland, associated to ³⁵
- ⁷⁶ DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain, associated to ³⁹
- ⁷⁷ Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, associated to ⁵³
- ⁷⁸ University of Michigan, Ann Arbor, MI, United States, associated to ⁶²
- ⁷⁹ Departement de Physique Nucleaire (SPhN), Gif-Sur-Yvette, France

^a Universidade de Brasília, Brasília, Brazil

^b Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil

^c Central South U., Changsha, China

^d Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China

^e Excellence Cluster ORIGINS, Munich, Germany

^f Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras

^g Università di Bari, Bari, Italy

^h Università di Bologna, Bologna, Italy

ⁱ Università di Cagliari, Cagliari, Italy

^j Università di Ferrara, Ferrara, Italy

^k Università di Firenze, Firenze, Italy

^l Università di Genova, Genova, Italy

^m Università degli Studi di Milano, Milano, Italy

ⁿ Università di Milano Bicocca, Milano, Italy

^o Università di Padova, Padova, Italy

^p Università di Perugia, Perugia, Italy

^q Scuola Normale Superiore, Pisa, Italy

^r Università di Pisa, Pisa, Italy

^s Università della Basilicata, Potenza, Italy

^t Università di Roma Tor Vergata, Roma, Italy

^u Università di Urbino, Urbino, Italy

^v Universidad de Alcalá, Alcalá de Henares, Spain

[†] Deceased