



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2010

Astrophysical limitations to the identification of dark matter: Indirect neutrino signals vis-à-vis direct detection recoil rates

Serpico, P D ; Bertone, G

Abstract: A convincing identification of dark matter (DM) particles can probably be achieved only through a combined analysis of different detections strategies, which provides an effective way of removing degeneracies in the parameter space of DM models. In practice, however, this program is made complicated by the fact that different strategies depend on different physical quantities, or on the same quantities but in a different way, making the treatment of systematic errors rather tricky. We discuss here the uncertainties on the recoil rate in direct-detection experiments and on the muon rate induced by neutrinos from dark matter annihilations in the Sun, and we show that, contrarily to the local DM density or overall cross section scale, irreducible astrophysical uncertainties affect the two rates in a different fashion, therefore limiting our ability to reconstruct the parameters of the dark matter particles. By varying within their respective errors astrophysical parameters such as the escape velocity and the velocity dispersion of dark matter particles, we show that the uncertainty on the relative strength of the neutrino and direct-detection signal is as large as a factor of 2 for typical values of the parameters, but can be even larger in some circumstances.

DOI: <https://doi.org/10.1103/PhysRevD.82.063505>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-41579>

Journal Article

Accepted Version

Originally published at:

Serpico, P D; Bertone, G (2010). Astrophysical limitations to the identification of dark matter: Indirect neutrino signals vis-à-vis direct detection recoil rates. *Physical Review D*, 82(6):063505.

DOI: <https://doi.org/10.1103/PhysRevD.82.063505>

Astrophysical limitations to the identification of dark matter: indirect neutrino signals vis-à-vis direct detection recoil rates

Pasquale D. Serpico

*Physics Department, Theory Group, CERN, CH-1211 Geneva 23, Switzerland and
LAPTH, UMR 5108, 9 chemin de Bellevue - BP 110, 74941 Annecy-Le-Vieux, France*

Gianfranco Bertone

*Institut d'Astrophysique de Paris, France. UMR7095-CNRS UPMC,
98bis Boulevard Arago, 75014 Paris, France and
Institute for Theoretical Physics, University of Zurich, 8057 Zurich, Switzerland*

(Dated: June 17, 2010)

A convincing identification of dark matter (DM) particles can probably be achieved only through a combined analysis of different detections strategies, which provides an effective way of removing degeneracies in the parameter space of DM models. In practice, however, this program is made complicated by the fact that different strategies depend on different physical quantities, or on the same quantities but in a different way, making the treatment of systematic errors rather tricky. We discuss here the uncertainties on the recoil rate in direct detection experiments and on the muon rate induced by neutrinos from dark matter annihilations in the Sun, and we show that, contrarily to the local DM density or overall cross section scale, irreducible astrophysical uncertainties affect the two rates in a different fashion, therefore limiting our ability to reconstruct the parameters of the dark matter particle. By varying within their respective errors astrophysical parameters such as the escape velocity and the velocity dispersion of dark matter particles, we show that the uncertainty on the relative strength of the neutrino and capture signal is as large as a factor of two for typical values of the parameters, but can be even larger in some circumstances.

PACS numbers: 95.35.+d, 14.60.Lm

CERN-PH-TH/2010-139, LAPTH-020/2010

I. INTRODUCTION

The detection of Dark Matter (DM) particles in accelerators or with direct and indirect searches would be of paramount importance for particle physics and cosmology. A convincing identification of DM, however, will require a combined analysis of different experimental strategies (see Refs. [1–4] for recent reviews). From LHC data alone, for instance, it will be difficult to reconstruct the relic density of the newly discovered particles (if any), even if one assumes to know the underlying theory (see e.g. Ref. [5] and references therein). However, combining LHC data e.g. with direct detection data significantly improves the reconstruction [6]. One could cite several other examples of complementarity of different detection techniques, such as the combination of different targets in direct detection experiments [7, 8], or multi-wavelength analyses in indirect detection experiments [9].

Although the combination of different detection strategies provides an effective way of removing degeneracies in the parameter space of DM model, a combined analysis is made complicated by the fact that one needs to put together results that depend on different physical quantities, or on the same quantities but in a different way, a circumstance that makes the treatment of systematic errors (i.e. depending on the underlying model assumptions) rather problematic.

The statistical and systematic uncertainties affecting the reconstruction of DM particle from various experi-

ments have been discussed widely in literature. One recent example is the the uncertainty on the local DM density, affecting both the rate of nuclear recoils in direct detection experiments, and the flux of upgoing muons induce by DM annihilations in the Sun. The statistical error on this quantity, based on dynamical constraints of various tracers in the Galaxy, have been discussed in Refs.[12, 13] (see also Ref. [14]). The systematic error due to the poor knowledge of the DM density profile in the Galaxy, and in particular to its possibly triaxiality has been discussed in Ref. [15]. Another recent example is the solar model dependence of the DM-induced neutrino signal [16].

Here we want to assess a minimal level of uncertainty in the normalization of the neutrino flux from the Sun (and, a fortiori, the Earth) due to the ignorance on crucial astrophysical parameters, in particular concerning the DM phase space distribution. While the impact of these uncertainties for direct dark matter searches has been long acknowledged (see e.g. [17]) and it is still actively investigated (see for example [18]) the impact for indirect neutrino detection has been investigated less extensively. Occasionally, it has been noted that some effects (e.g. clumpiness and dark disk, see Sec. III.5 below) may enhance the “baseline” flux, but little attention has been paid to the robustness of the baseline flux from an ordinary, smooth halo. Also, it is of interest to address the question of what would be the uncertainty on the prediction for the indirect neutrino signal, should a direct detection be used for normalization. Finally, since this

uncertainty turns out to be the dominant one, its value also sets the accuracy needed for several nuclear and particle physics input parameters, derived either empirically or via theoretical computations.

Here, we study this issue also in light of recent results from numerical simulations. The structure of the article is the following: in Sec. II we review the dependence of the recoil and muon rates on key astrophysical parameters. In Sec. III we discuss the systematic uncertainties inevitably associated with these parameters, including time-dependent effects. In Sec. IV we discuss the impact of these uncertainties on theoretical predictions and in Sec. V we discuss the results and present our conclusions.

II. THEORETICAL SETUP

The theory of DM experiments has been widely discussed in literature. We summarize in this section the dependence of the predicted detection rates on some key parameters.

A. Recoil rate in direct detection experiments

We start from direct detection experiments, for which, unless otherwise stated, we follow the notation of [19]. Apart for numerical constants of order one, the differential number of recoil events per unit time and target mass writes

$$\frac{dR}{dE} \sim \frac{\sigma_{0,A}}{\mu_{\chi,A}^2} \frac{\rho_{\odot}}{m_{\chi}} F_A^2(E) \int dv \frac{f_1(v)}{v} \Theta(v - v_{\min}) \quad (1)$$

where $\mu_{\chi,A} = m_{\chi} M_A / (m_{\chi} + M_A)$, $v_{\min} = \sqrt{\frac{EM_A}{2\mu_{\chi}^2}}$ and

$$f_1(v) \equiv \int d\Omega v^2 f(\mathbf{v}) \Rightarrow \int_0^{\infty} dv f_1(v) = 1, \quad (2)$$

where m_{χ} is the DM mass, M_A the target atom mass, $\sigma_{0,A}$ the relevant cross-section, F_A the form factor, ρ_{\odot} the local density of DM, and $f(\mathbf{v})$ is the 3D velocity distribution. Note that $f_1(v)$ denotes the velocity distribution of DM particles *at the Earth*. If one has the distribution of velocity in the “halo rest frame”, in general one has to correct for: i) time-dependent velocity of the detector with respect to this halo, including Earth motion with respect to the Sun and Solar motion with respect to the halo; ii) Gravitational effects, including in general both the gravitational focusing of unbound particles (see e.g. [20] and refs therein) and the population of gravitationally bound DM particles (see [21, 22] and refs. therein).

We shall not enter the details of the form factor dependence to avoid introducing too many independent parameters: for our purposes, we shall use the parameterization in terms of the nuclear mass from [1]

$$F_A^2(E) = \exp(-m_A R_A^2 E/3) \quad (3)$$

with $R_A = 1 \text{ fm}[0.3 + 0.91 \sqrt[3]{m_{A,\text{GeV}}}]$. The overall rate R of interest here is obtained by integrating Eq. (1) over energy, starting from the threshold E_T (experiment-dependent). So, the input parameters to specify in this case are: $\{m_A, E_T, m_{\chi}, \sigma_{0,A}, \mathbf{y}\}$, where \mathbf{y} denotes collectively the astrophysical parameters.

B. Muon rate induced by neutrinos from DM annihilations in the Sun

As mentioned above, the detection of a high-energy neutrino flux from the center of the Sun or the Earth would provide a very convincing evidence for DM annihilations. The rate of events in neutrino telescopes depends on the number of particles that are captured, and on their fate inside the star or planet.

The number of DM particles captured in the Sun/Earth obeys the following time evolution equation

$$\dot{N} = C - C_A N^2 \quad (4)$$

where C is the capture rate and C_A (see below) regulates DM annihilations. If both coefficient are constant, solving for $N(t)$ one can derive

$$\Gamma_A(t) = \frac{C}{2} \tanh^2\left(\frac{t}{\tau_{\text{eq}}}\right), \quad \tau_{\text{eq}} = (C C_A)^{-1/2}. \quad (5)$$

In the limit where steady state is reached within timescales much shorter than the lifetime of the Sun, $t_{\odot} \simeq 4.6 \times 10^9 \text{ yr}$, one has $\dot{N} = 0$, $N = \sqrt{C/C_A}$, and the annihilation rate writes

$$\Gamma_A = \frac{C_A}{2} N_{\text{eq}}^2 = \frac{C}{2}. \quad (6)$$

In this regime, the normalization of the signal only depends on C . More generally, the present value $\Gamma_A(t_{\odot})$ depends also on τ_{eq} , i.e. on C_A . C_A can be written in terms of effective volumes $V_{1,2}$, as

$$C_A = \langle \sigma_A v \rangle \frac{V_2}{V_1^2}, \quad V_j \simeq \left(\frac{3 m_{\text{Pl}}^2 T_{\odot}}{2 j m_{\chi} \rho_{\odot}} \right)^{3/2} \quad (7)$$

with T_{\odot} , ρ_{\odot} respectively the central temperature and density of the body under consideration [1]. Note that the above formula already has an uncertainty due to the only approximate assumption of thermalization, homogeneous conditions of the core, etc. In the other useful limit $\tau_{\text{eq}} \gg t_{\odot}$,

$$\Gamma_A(t_{\odot}) \approx 0.5 C^2 C_A t_{\odot}^2. \quad (8)$$

This has the following implications for the large τ_{eq} regime: i) the signal is quadratic in C ; ii) there is an additional dependence on C_A (i.e. linear in $\langle \sigma_A v \rangle (\frac{m_{\chi} \rho_{\odot}}{T_{\odot}})^{3/2}$) as well as a quadratic one on t_{\odot} . Even neglecting particle physics unknowns, this means that additional astrophysical uncertainties enter. For the Sun, different models (see

e.g. Ref. [23, 24]) yield different predictions for these parameters at the $\sim 1\%$ level. For the Earth, uncertainties are much larger. For example, according to [25], Earth's core temperature is estimated at around 5000 to 7000 degrees Celsius, therefore it carries an error of $\sim 15\%$. We note incidentally that despite these uncertainties, it is possible to obtain useful constraints on DM particles, by requiring that they do not over-heat the Earth [26].

Focusing on C , we shall first remind that for most particle physics models (including Kaluza-Klein and most neutralino DM models), the capture in the Sun is actually dominated by spin-dependent interactions on hydrogen [3]. This implies among others that we can consider the form factor ≈ 1 , which greatly simplifies the formulae. Although exceptions do exist, in the following we are interested in discussing the uncertainties induced by the astrophysical (rather than nuclear/particle physics) input, so we shall limit ourselves to this single contribution in the capture.

From rewriting Eqs. (2.8,2.13) of [27] in our notation we get

$$C = \sigma_{0,p} \frac{\rho_{\odot} \epsilon_p M_{\odot}}{m_{\chi} m_p} \times \int_0^1 d\mathcal{M} \nu^2(\mathcal{M}) \int_0^{u_{\max}} du \frac{f_1(u)}{u} \left[1 - \frac{u^2}{u_{\max}^2} \right], \quad (9)$$

where

$$u_{\max}(\mathcal{M}) \equiv \frac{\sqrt{4 m_{\chi} m_p}}{m_{\chi} - m_p} \nu(\mathcal{M}) \quad (10)$$

and ν is the escape velocity from the unit shell volume considered, in turn depending on the distance from the center r ; to a good approximation [10, 16],

$$\begin{aligned} \nu^2(r) &= v_{\odot}^2 - \mathcal{M}(r)(v_{\odot}^2 - v_s^2), \\ v_{\odot} &\simeq 1355 \text{ km/s}, \quad v_s \simeq 818 \text{ km/s} \end{aligned} \quad (11)$$

where $\mathcal{M}(r)$ is the mass within the radius r , in units of the solar mass. To derive the above equation for the capture rate, we replaced the Sun with a pure hydrogen sphere and introduced the factor $\epsilon_p < 1$ to account for the fact that only a fraction of the Sun is made of hydrogen. What is crucial here is the approximation that the density of hydrogen traces the total one of the Sun, which amounts to a small ($\sim 8\%$) overestimate of the capture [16]. Note that the consequences of our approximations are further softened by the fact that we are only concerned with *relative* response of the observables to variations in the astrophysical inputs, so that overall scale mismatches are of no importance here.

The main input parameters in this case are thus $\{m_{\chi}, \sigma_{0,p}, \mathbf{y}\}$. Further simplifications might be obtained by specifying the function f_1 , but we do not proceed further along this path, since we want to keep some generality concerning f_1 , whose key parameters are described in the following section.




FIG. 1: Timescale to reach equilibrium between capture and annihilation (Eq. (5)) as a function of the muon rate on a terrestrial detector for DM models in a pMSSM scenario, as implemented in DarkSUSY [28]. Solid circles correspond to models that fulfil the constraints on the cosmological relic abundance. The two horizontal lines correspond to the solar system age and 1/4 of the solar revolution time in the Galaxy. The vertical line shows the approximate sensitivity of IceCube plus DeepCore in 5 years of data taking [29].

Neutrino experiments such as IceCube are mostly sensitive to the flux of up-going muons induced by high-energy neutrinos. The rate of μ 's induced by a neutrino flux $\Phi_{\nu}(E_{\nu})$ is given by

$$\Gamma_{\mu} = V_{\phi}(\delta) \int_{E_{\mu}^{\text{thr}}}^{m_{\chi}} dE_{\nu} \int_0^{y_{\nu}} dy A(E_{\mu}) P_{\mu}(E_{\nu}, y) \Phi_{\nu}(E_{\nu}) \quad (12)$$

where $y_{\nu} = 1 - E_{\mu}^{\text{thr}}/E_{\nu}$ and E_{μ}^{thr} is the muon threshold energy of the experiment. $A(E_{\mu})$ is the effective area of the detector and $P_{\mu}(E_{\nu}, y)$ is the probability that a neutrino of energy E_{ν} interacts with a nucleon producing a muon of energy $E_{\mu} \equiv (1 - y)E_{\nu}$ above the detector threshold energy, and can be estimated as

$$P_{\mu}(E_{\nu}, y) \simeq \frac{\tilde{\rho}}{m_p} \tilde{R}(E_{\mu}, E_{\mu}^{\text{thr}}) \sigma(E_{\nu}, y) \quad (13)$$

where $\tilde{\rho}$ is the density of the medium (typically water/ice), $\tilde{R}(E_{\mu}, E_{\mu}^{\text{thr}})$ is the muon range in that medium, *i.e.* the distance traveled by muons before their energy drops below E_{μ}^{thr} , and $\sigma(E_{\nu}, y) \equiv d\sigma_{CC}^{\nu N}(E_{\nu}, y)/dy$ is the differential cross section for neutrino–nucleon charged–current scattering (for further details see e.g. Ref. [3] and references therein). In our case, the neutrino flux is simply given by

$$\phi_{\nu} = \frac{\Gamma_A}{4\pi D^2} \frac{dN_{\nu}}{dE} \quad (14)$$

where dN_{ν}/dE is the neutrino spectrum per annihilation and D is the distance of the Sun to the Earth (Astronomical Unit).

We show in Fig. 1 the timescale to reach equilibrium between capture and annihilation as a function of the muon rate on a terrestrial detector for DM models in a pMSSM (“Phenomenological Minimal Super-Symmetric Standard Model”) scenario, as implemented in DarkSUSY [28]. For reference, we also show with horizontal curves two characteristic timescales of the solar system: its age, and 1/4 of the revolution time around the Galactic center τ_{gal} , i.e. the time to span a quadrant. The vertical line shows the approximate sensitivity of IceCube plus DeepCore in 5 years of data taking (e.g. Ref. [29]). It is clear that most interesting model in this scenario have $\tau_{\text{eq}} \in (0.2 \div 1) \tau_{\text{gal}}$, which makes “galactic” effects potentially important. Also note that for the Earth equilibrium is rarely reached in interesting models.

III. ASTROPHYSICAL UNCERTAINTIES

A key ingredient to make predictions for, or interpret, DM experiments is $v_c(R_\odot)$, the velocity that a test-mass would have on a circular orbit in the Galactic Plane at the solar distance R_\odot from the GC. Note that, to a good approximation, the Milky Way mass distributions is axisymmetric: one can find a cylindrical coordinate system $\{R, \phi, z\}$ centered on the GC where the potential is independent of ϕ , with the $z = 0$ plane dubbed “Galactic Plane” (see below the discussion of deviations from axial symmetry). In such a system, a particle moving in the plane has no way to know if the potential is spherically symmetric, or merely cylindrically symmetric. On the top of that, if the particle has no radial component of the velocity vector, but only an azimuthal one ($v_c(R) = R d\phi/dt = v_\phi$), it will move effectively in a circular orbit at distance R , whose force is generated by the mass enclosed within (R), i.e.

$$\frac{GM(R)}{R^2} = \frac{v_c(R)^2}{R}. \quad (15)$$

Note, however, that realistic stars have orbits varying in all coordinates. For stars having initial velocities in the plane $z = 0$, one has exactly $\langle v_z \rangle = 0$, while the orbit has a “rosetta” shape in the $\{R, \phi\}$ plane, with R varying between a minimum and a maximum value. If the star moves off-plane, then another oscillation takes place above/below the plane (for more details, see e.g. Chap. 3 in [30]). So, in general to derive v_c requires identifying the circular motion of the “local standard of rest” (LSR), with respect to which actual stars in our neighborhood (including the Sun) possess relative motions. In [12], this was evaluated from a compilation of data to be $v_c = 245 \pm 10$ km/s, but note that other observables may lead to a lower values, closer to 220 ± 20 km/s recommended by the IAU [31]. For the purposes of our estimate, we assume a central value of $v_c = 235$ km/s, affected by a $\sim 10\%$ error. How reasonable this is can be also inferred from the collection of data shown e.g. in Ref. [32].

1. The halo typical velocity

The halo velocity distribution cannot be directly observed, but must be inferred from a combination of observations and theoretical considerations. In this inference, several sources of error enter. First of all, one assumes that the *rotation curve is constant* over a significant radius around the solar position, i.e. $v_c \simeq \text{const}$. As shown by visual inspection of Fig. 1 in [32], this approximation holds at not better than $\sim 10\%$ level. For a *spherically symmetric system* this implies that the gravitational potential Φ is logarithmic, since

$$v_c^2(r) = r \frac{d\Phi}{dr} \Rightarrow \Phi(r) = v_c^2 \log(r) + \text{const}. \quad (16)$$

Until now, we assumed nowhere that the DM dominates the overall potential. If we did so, we would find the isothermal profile $\rho \propto r^{-2}$ as unique solution satisfying the above (approximate) observational condition and symmetry condition, and the Maxwell-Boltzmann (MB) distribution $f(\mathbf{v}) \propto \exp\left(-\frac{v^2}{v_0^2}\right)$ with $v_0 = v_c$ as velocity distribution. However, fit models suggest that DM only contributes a fraction ζ of the v_c^2 at the solar system. For examples, the two models in table 2.3 of [30] yield $\zeta \approx \{0.38, 0.64\}$. Note also that this fraction is changing with radius, too. So, in order to derive the link between, say, the average dispersion in DM velocity and the DM profile one needs further input, either from fits to the data or simulations. For a density of DM which goes as power-law $\rho(r) \propto r^{-\beta}$ in the neighborhood of the Sun, the Eddington equation (see again [30]) yields a phase space distribution function for DM which has a dispersion

$$\sigma_v^2 \equiv \frac{1}{3} \langle |\mathbf{v}|^2 \rangle = \frac{v_c^2}{\beta}. \quad (17)$$

it is immediate to check that assuming that the only v -dependence enters as $f(\mathbf{v}) \propto \exp\left(-\frac{v^2}{v_0^2}\right)$, one gets

$$\sigma_v^2 \equiv \langle v^2 \rangle = \int d^3\mathbf{v} v^2 f(\mathbf{v}) = \frac{v_0^2}{2} \Rightarrow v_0 = v_c \sqrt{\frac{2}{\beta}}, \quad (18)$$

where the second equality is required for Eq. (17) to hold. Note that it has been often acritically assumed (perhaps following historical papers as [33] and [1]) that $v_0 = v_c$, but this strictly requires $\beta = 2$ at the solar radius. If one fits the DM profile with another popular model from numerical simulations, say NFW [34] with a scale radius $\simeq 20$ kpc, there is a departure from this value of about 20%. Alone, such uncertainty on the radial profile of DM at the solar radius implies an error of $\sim 10\%$ on the equality $v_c = v_0$. As described previously, a similar level of error appears inherent to the assumption of the constancy of v_c . Other errors, as the limitations of the spherical approximation (and thus isotropy of the space

and velocity dependence of the DM profile) are more difficult to assess, but likely present as well, with comparable or larger amplitude. We shall comment on that a bit further in the following.

In summary, an estimate of the typical velocity parameter v_0 is

$$v_0 \approx v_c, \quad (19)$$

with a $\sim 10\%$ value due to the uncertainty on the measured v_c (and thus correlated with its true value) and probably an uncertainty *at least twice as big* which is inherent to the theoretical assumptions. The latter errors are unfortunately of systematic nature and hardly reducible in the near future.

2. Escape velocity

From *model-dependent* considerations similar to the ones of the above section one can also derive that expressions for the escape velocity as a function of Galactic parameters, see e.g. [33]. Equivalently, based on milder assumptions and on surveys of high velocity stars, one can infer constraints such as [35]

$$498 \text{ km/s} < v_{\text{max}}(R_{\odot}) < 608 \text{ km/s}, \quad (20)$$

with a median value of 544 km/s [35] which we adopt in the following as benchmark. Note that since theoretically one expects that $v_{\text{max}}^2 \propto v_c^2$, it is not surprising that a similar statistical error of $\sim 10\%$ is found for both quantities. On the other hand, the local escape velocity depends on *global* properties of the dark matter halo. Hence, to be conservative we shall treat the uncertainty on v_{max} as independent from the uncertainty on v_0 as inferred from v_c .

3. Shape of the velocity distribution

In the above considerations, for simplicity we parametrized the uncertainty on the velocity distribution via the typical velocity in the halo, or equivalently v_0 . But the event rates are not only sensitive to the second moment of the distribution, but also to its shape. There is no guarantee that the velocity distribution function is truly MB, and actually several numerical simulations (see [36, 37]) suggest that noticeable deviations are present. To get a feeling for the sensitivity of the relevant observables to deviations from the MB shape, we parameterize the deviation from a MB. distribution as

$$f(\mathbf{v}, \alpha) \propto \left[1 + (\alpha - 1) \left(\frac{v}{\langle v \rangle} - 1 \right) \right] f_{\text{MB}}(v) \quad (21)$$

with $\alpha \in [0.7 \div 1.3]$, as suggested e.g. from the right panel in Fig. 5 of [36]. Note that if the capture equilibration is long, the capture may probe the average distribution,

while the direct detection probes the present one (see below) which is another effect linked to the possible shape deviation, not explored here.

What we shall employ is the Eq. (21), with a MB cut at the v_{max} , correctly normalized to one. To that distribution one has then to apply a Galilean transformation to account for the motion of the Sun with respect to the DM halo, see below.

4. Motion of the Sun with respect to the DM halo and time-dependent effects

Another effect on the recoil and capture rate is due to the fact that the Sun is in motion with respect to the DM halo “rest frame” with a velocity \mathbf{v}_{\odot} (since we are interested not in directional signals, only the modulus v_{\odot} is going to matter in what follows.) Naively, one would expect

$$v_{\odot} \approx v_c. \quad (22)$$

The situation is actually more complicated than that: the Sun is not rotating uniformly around the GC, i.e. it is not at rest with respect to the LSR frame. Then, one has a relative motion between sun and DM halo given by

$$\mathbf{v}_{\odot} = \mathbf{v}_{\text{LSR}} + \mathbf{V}_{\text{loc}}. \quad (23)$$

From the recent analysis in [38] one has the relative velocity determination $\mathbf{V}_{\text{loc}} = \{U, V, W\} = \{11.1 \pm 0.7 \pm 1, 12.2 \pm 5 \pm 2, 7.3 \pm 0.4 \pm 0.5\}$ km/s respectively radially inwards (U), in the direction of the Galactic rotation (V), and vertically upwards (W), with the former error statistical and the second one systematic. Taken at face value, this only amounts to an upward correction of about $\sim 6\%$ to the naive estimate $v_{\odot} \approx v_c$, with an uncertainty which is even smaller, thus providing a sub-leading error. However, the situation is different when time-dependent effects are taken into account. In fact, as we have seen, one has to contemplate the possibility that the equilibration time for capture in the Sun is comparable to (or longer than) a fraction of its orbital time $\sim 2 \times 10^8$ yr. This situation is verified in a significant fraction of SUSY models leading to a large muon flux, as shown in Fig. 1 (see also the discussion in Ref. [16]). For the signal of the Earth, this is true for practically all interesting models.

The fact that the capture signal depends on the long-term past history of the Sun (and Earth) immediately changes the importance of the solar motion effect mentioned previously: differently from the first term at the RHS of Eq. (23), the second term is time-dependent due to the variation of velocity of the Sun along its orbit in the Galaxy, with typical timescale of $\sim 2 \times 10^8$ yr.

To be more quantitative, one would need to know the actual orbit of the Sun in the Galaxy *and* the equilibration timescale for the case realized in nature. Both have significant sources of error: i) the error in the “initial”




FIG. 2: *Left Panel:* Effective density as a function of the position in a circle of radius R_{\odot} around the GC, for different choices of τ_{eq} . *Right Panel:* Maximum relative difference between the effective and instantaneous density, as a function of the time to reach equilibrium τ_{eq} , in units of a galactic year τ_{gal} .

(actually final, since the evolution is performed backwards!) conditions, in particular the velocity vector \mathbf{V}_{loc} ; ii) the only approximate knowledge of the Galactic potential; iii) the incomplete knowledge of particle physics and astrophysical parameters determining τ_{eq} . As a result, rather than “a refinement” to the naive estimate we can conservatively consider this kind of effects as an additional source of error. For illustration, we take the results of the numerical integration performed in [39] for the model 2 in [40] (note that compared with those results the local velocity component has been significantly revised in the analysis reported in [38]). As a result, that the distance from the GC varies between -2% and +7% from the present value, while the velocity results to anticorrelate almost exactly, with an equal (in modulus) and opposite sign variation (the motion is indeed quasi-keplerian). As a consequence, the two effects almost cancel out for the capture rate in the Sun, so this time-dependence is not the main source of error *within the simple model considered*.

The situation is actually different if we move beyond a smooth and isotropic halo approximation: dropping the isotropy means that the local density might differ from the averaged one (to which the Solar capture is sensitive for long equilibration times). This difference has been recently evaluated in [15] based on numerical simulations: in the ones including approximately the effects of baryons, this error is about 20% ÷ 30%. In those without baryons, depending on the ignorance of our orbit within the triaxial halo of DM, the error is as large as a factor ~ 2 . To account for triaxiality, a simple model with a dependence of the density along the Galactic Plane as $\rho(\phi) = \rho_{\odot}[1 + \varepsilon \sin(2\pi\phi)]$ can be taken as representative of results with baryons, for $\varepsilon = 0.3$. Of

course, since we do not know the phase at which we are at current time, one can define an effective density

$$\langle \rho \rangle_{\text{eff}}(\phi) = \frac{\tau_{\text{gal}}}{\tau_{\text{eq}}} \int_{-\frac{\tau_{\text{eq}}}{\tau_{\text{gal}}}}^0 \rho(\phi + \varphi) d\varphi \quad (24)$$

which is probed by the capture rate and is a function of ϕ . We show in figure 2 the effective density as a function of position in a circle of size R_{\odot} around the GC, for different choices of τ_{eq} in units of τ_{gal} . We also show in the same figure (right panel) the maximum relative difference between the effective and instantaneous density, as a function of the time to reach equilibrium τ_{eq} , in units of a galactic year τ_{gal} . This is obtained at each point by scanning over the unknown ϕ . In both cases, we fix $\varepsilon = 0.3$. Two comments are in order: the maximal error obtained when $\tau_{\text{eq}}/\tau_{\text{gal}}$ is very large reflects the simple fact that the capture probes the real average value ρ_{\odot} , while the recoil the instantaneous value which can differ as much as $\varepsilon = 0.3$ from the average (black, solid curve in the left panel of Fig. 2). However, the situation is even more interesting when $\tau_{\text{eq}}/\tau_{\text{gal}} \lesssim 0.6$ (which Fig. 1 hints to be a physically interesting range): in this case the relative error can range from almost negligible to above 40%, within only a factor 3 variation of τ_{eq} . This suggests that, even if we knew that the ellipticity of the Milky Way halo is at the 30% level, and even by assuming that an eventual neutrino detection hints at $\tau_{\text{eq}}/\tau_{\text{gal}} \lesssim 1$, the lack of a detailed knowledge of the particle physics and astrophysics limit the estimate of this error conservatively at the 40% level. Needless to say, should the baryon disk effect be less prominent than what suggested by simulations, this effect could be a *factor of several* larger. The actual impact of baryons on the DM structure is in fact still subject to debate.

5. Additional halo (sub)structures

Concerning the non-smoothness of the halo, an attempt to address its impact on detection rates was performed in Ref. [41], where the effects of the enhanced density while crossing a sub-halo were considered. For our purposes of assessing reasonable errors, rather than exploring what can happen in principle if such an encounter takes place, it is more interesting to assess the probability that such an encounter takes place at all. If we focus on subhalos of mass $M_{\text{sh}} = 10^{-6}M_{\odot}$, we can work out the characteristic size, number density therefore average separation for these objects. The Sun encounters a subhalo of mass M_{sh} and size r_{sh} with a frequency $\Gamma_{\text{sh}} = 4\pi v_{\text{tot}} n_{\text{sh}} r_{\text{sh}}^2$. Assuming that a fraction f of the DM is in the form of clumps, and inserting typical values, we find

$$\Gamma_{\text{sh}} \sim f 10^{-6} \text{yr}^{-1} \frac{v_{\text{sh}}}{220 \text{km s}^{-1}} \frac{10^{-6} M_{\odot}}{M_{\text{sh}}} \left(\frac{r_{\text{sh}}}{0.01 \text{pc}} \right)^2 \quad (25)$$

To perform this estimate n_{sh} we have followed the usual strategy of extrapolating the mass function of subhalos to very small masses, and assumed that the spatial distribution of subhalos traces the smooth DM distribution. The second assumption is most likely a very crude approximation, as the most recent numerical simulation have shown that the distribution of subhalos is *anti-biased*, such that if $\rho(r)$ is the total (smooth plus subhalos) DM density profile, $n_{\text{sh}}(r) \propto r\rho(r)$. The fraction f of DM in subhalos is therefore a function of r and it tends to deplete the number of clumps at the solar radius. The results of Ref. [42] point towards a more realistic value¹ of $f \sim 10^{-2}$. This estimate suggests that the timescale for an encounter is comparable to (a relevant fraction of) the orbital time of the Sun in the Galaxy. On the other hand, the crossing time t_{sh} for such an encounter is very short, of the order of a century. Hence, already on this basis we can conclude that significant alterations of the annihilation yield, while possible in principle, are very unlikely. We also note that this estimate is not very dependent on the exact mass of the sub-halo considered: since one has roughly $r_{\text{sh}} \sim M_{\text{sh}}^{1/3}$, and $t_{\text{sh}} \propto r_{\text{sh}}$, it follows the weak dependence $\Gamma_{\text{sh}} \sim M_{\text{sh}}^{-1/3}$, while $\Gamma_{\text{sh}} t_{\text{sh}}$ which determines the probability of “living during a crossing” stays virtually constant and very small.

It is worth mentioning that it is not even sufficient that an encounter takes place in order for a significantly enhanced capture to happen. In fact, as a first approximation we can think of each subhalo as orbiting the galaxy with a *typical* velocity w sampled from the $f(\mathbf{v})$ distribution. Neglecting the v_{max} cutoff and assuming a MB

distribution, the 1D distribution at the Sun is readily written as

$$f_1^{\text{sh}}(w) \approx \frac{2w}{\sqrt{\pi} v_0 v_{\odot}} \sinh \left[\frac{2w v_{\odot}}{v_0^2} \right] \exp \left(-\frac{w^2 + v_{\odot}^2}{v_0^2} \right). \quad (26)$$

The particles in each sub-halo have a dispersion σ around w . However, as long as $\sigma \ll v_0$ (as expected for the smallest, hence more abundant, sub-halos) this only brings in a small correction. Roughly speaking, as long as long $w < u_{\text{max}}$ the DM particles will be captured, when instead in an encounter one has $w > u_{\text{max}}$ only a very small fraction of the DM population could be captured (see Eq. (10)). For fiducial parameters in Eq. (26) and taking into account that for the Sun $u_{\text{max}} \simeq 200 \text{km/s}$ for a 100 GeV particle, one finds that only $\sim 15\%$ of the encounters will lead to a(n efficient) capture. For the Earth, this fraction is negligible.

The scenario could be further enriched if one consider the possible existence of the “dark disk” [43, 44] due to the dragging effect of the baryonic disk on the DM halo. The crucial aspect is that the (typically subleading by number) population of DM disk particles has a relatively low lag velocity with respect to the stars, $v_{\text{lag}} \sim 0 \div 150 \text{km/s}$, and a comparably low velocity dispersion σ_d . As a consequence, the Sun and the Earth effectively see a colder DM gas, a fact that eases captures. For the Sun, the scaling relation for the capture reported in [44] suggests that, depending on parameters, the annihilation signal can obtain a correction which ranges from marginal (a few percent) up to more than one order of magnitude. For the Earth the dependence is by far more dramatic, usually amounting to orders of magnitude. This confirms once again that the normalization of the signal from the Earth is extremely dependent on the details of the (very) cold tail of DM phase space distribution and, while still potentially useful for a serendipitous discovery of DM, can tell us very little on the particle physics.

IV. HALO UNCERTAINTIES ON DM DETECTION RATES

We are interested in estimating the variations in the direct experiment recoil rate R as well as capture rate C when astrophysical parameters vary within their uncertainties and/or they vary over time. The two functions depend in general from non-astrophysical parameters \mathbf{x} (DM mass, mass of the target nucleus, etc.) as well as some astrophysical parameters \mathbf{y} .

For a first glimpse at the overall dependence of the observables on parameters, we show in Fig. 3 a series of “scatter plots” in the R vs. C plane as a function of the \mathbf{x} , scanning over the min-med-max values of the \mathbf{y} parameters. In particular, we consider as x variable the DM mass (top panel), different values of the energy threshold for direct detection (center) and different values of the target mass A (bottom). The latter are chosen as follows: fiducial value $A = 73$ for an “intermediate

¹ It is perhaps curious to note that if an unrealistically large value of f is used, the actual annihilation signal could be *lowered* with respect to its naive equilibrium value in the limit of smooth halo [41].




FIG. 3: Impact of varying astrophysical parameters (see text for range of variation), in the $R-C$ plane, for different values of the DM mass (top) different values of the energy threshold (center) and different values of target mass A (bottom).

mass” target, such as Germanium; $A = 131$ for a heavy target material as Xenon; $A = 28$ for a relatively light target, such as Silicon (these identifications should not be taken too seriously but to investigate the dependence on the target mass, since the form-factor functions are only schematic). The \mathbf{y} 's are varied as follows: v_0 at fiducial value as well as varied by $\pm 10\%$ (which, as discussed, is likely a lower limit to the error range), v_\odot at fiducial value and with $\pm 10\%$ variations, α at the values 0.7, 1, 1.3 and v_{\max} is set at 498, 544, and 608 km/s. The observables are normalized to the values assumed for fiducial parameters. It is obvious that, in all cases, the uncertainty on the capture can have excursions of $\sim 40\%$, while the range span by R is typically within $\pm 20\%$ of its fiducial value, although it can increase for light targets, high thresholds, or heavy particles. What is more important, there are (anti)correlations between variations in R and C , whose magnitude and even *sign* depend on the variable considered. This is illustrated by diagonal “trends” in the dispersion plots, whose slope can change both quantitatively and in sign. It is not unusual that the relative uncertainty of the ratio C/R , say, is larger than the error on each quantity C , R due to this effect. Just for a qualitative illustration of this effect, we computed the correlation coefficient of the R and C for the scanned points. The dataset cross-correlation when varying $m_\chi=20, 100, 500$ GeV varies from -0.12 to -0.15 and to $+0.55$; when varying $E_{\text{th}}=5, 10, 20$ keV rises from 0.003 to -0.15 and to -0.42 ; when varying A from 28 to 73 and 131 rises respectively from -0.50 to -0.15 and to $+0.37$. Of course, this has no truly statistical meaning (we assumed “flat” priors for the variables in their range and did not account for the existing correlations among them), but it clearly illustrates the trend of qualitative change of the correlation when underlying parameters are changed.

For a more compact and quantitative exploration of the parameter dependence, it turns useful to define the logarithmic derivatives:

$$\gamma_i(\mathbf{x}, \mathbf{y}_0) \equiv \left. \frac{\partial \ln C(\mathbf{x}, \mathbf{y})}{\partial \ln y_i} \right|_{\mathbf{y}=\mathbf{y}_0}, \quad (27)$$

$$\varrho_i(\mathbf{x}, \mathbf{y}_0) \equiv \left. \frac{\partial \ln R(\mathbf{x}, \mathbf{y})}{\partial \ln y_i} \right|_{\mathbf{y}=\mathbf{y}_0}. \quad (28)$$

The functions γ_i and ϱ_i are sufficient (to leading order) in describing the sensitivities to the quantities of interest. Note also that the change in C/R as a consequence of a change $\delta_i \equiv \delta y_i/y_0$ can be written, in the linear approximation, as (no sum over repeated indices)

$$\delta_i \left(\frac{C}{R} \right) \approx \left. \frac{\partial}{\partial y_i} \frac{C}{R} \right|_{\mathbf{y}_0} \delta y_i = \frac{C}{R} \left[\gamma_i - \varrho_i \right]_{\mathbf{y}_0} \delta_i. \quad (29)$$

The values of the γ_i and ϱ_i coefficients for fiducial values of \mathbf{y} and different values of the \mathbf{x} are reported in Tab. I and Tab. II.

i	$\gamma_i(20)$	$\varrho_i(20)$	$\gamma_i(100)$	$\varrho_i(100)$	$\gamma_i(500)$	$\varrho_i(500)$
v_0	-0.80	1.04	-0.98	-0.04	-0.90	-0.22
v_\odot	-0.94	1.00	-1.64	0.15	-1.92	-0.05
v_{\max}	-0.12	3.21	-0.13	0.77	-0.13	0.50
α	-0.15	0.22	-0.16	0.004	-0.12	-0.04

TABLE I: The response functions for capture and direct recoil rate with respect to different halo parameters for three values of the mass $m_\chi = 20, 100, 500$ GeV, and for the fiducial values of the parameters $\alpha = 1$, $A = 73$, $E_{\text{th}} = 10$ keV, $v_c = 235$ km/s, $v_0 = 235$ km/s, $v_{\max} = 544$ km/s. Note that the product $\sigma\rho$ enters only as a scaling, and similarly for both observables (i.e. response is 1).

From these tables it is clear that the dependence on the parameters is actually different in the two cases. Take the second row of Tab. I: it means that an uncertainty in the velocity of the sun in the halo is responsible for a factor $|-1.64 - 0.15| \simeq 1.8$ times larger uncertainty in the *relative* normalization of the two rates. From that table, one immediately notices the general trend that, when the recoil rate only probes the high-energy tail of the distribution (light DM particles, high thresholds, light targets) the dependencies on virtually all parameters are *opposite* for R and C : the capture in fact always prefers the low-energy tail of the distribution, where particles are slower and thus easier to capture. For higher target masses, a competing effect arises due to the form-factor suppression, so that mixed trends arise. On the other hand, in the limit $E_{\text{th}} \rightarrow 0$, the dependence on the halo parameters goes to zero since the whole phase space is actually probed via direct direction. Note that the effect of the shape parameter α is rather modest, and the relative uncertainties due to the non-MB nature of the distribution can be estimated to be $< 10\%$. A posteriori, this justifies a simple parametric approach to explore this variable.

i	$\varrho_i(A=28)$	$\varrho_i(A=131)$	$\varrho_i(5\text{ keV})$	$\varrho_i(20\text{ keV})$
v_0	0.14	-0.19	-0.10	0.11
v_\odot	0.32	0.01	0.07	0.31
v_{\max}	1.12	0.52	0.69	0.95
α	0.05	-0.02	-0.01	0.04

TABLE II: As in Tab. I, for the direct detection rate, keeping $m_\chi = 100$ GeV and varying the target mass and the threshold.

On the top of the different sensitivity to the variables, another source of uncertainty is the fact that the two signals sample different *time averages* of the physical quantities, as already argued in Sec. III.4 (see Fig. 2). Accounting for a $\sim 40\%$ maximal error on the ratio of observables estimated on the basis of simulation results, we can conservatively conclude that there is likely a relative uncertainty on C/R which reaches a factor of ~ 2 , just due to “halo astrophysics”.

V. CONCLUSIONS

A conservative estimate of the error on the capture rate suggests that the normalization of the neutrino signal from DM annihilation in the Sun/Earth brings not only the uncertainty given by the local DM density ρ_\odot , but often more importantly uncertainties connected with the velocity distribution function, as well as our motion in the halo. We showed that the error on the capture C can easily reach $\approx 40\%$, which translates into a comparable or larger error on the annihilation signal. Even more important, the relative uncertainty on the normalization of the annihilation to recoil signal is typically larger, easily up to a factor two in presence of typical equilibration times of $\sim 10^7 \div 10^8$ yr. For cases where equilibration times are billions of years or larger (as typical for the Earth) the uncertainties are likely at least few times larger. The same happens if the effects of baryons in reducing Galactic halo triaxiality are smaller than what estimated at present from numerical simulations.

We also studied the sensitivity of the observables to different input parameters, which was shown to be often opposite for the direct and indirect signals, due to the preferential probe of the high tail part of the velocity distribution for the recoil rate. Interestingly, at least for some variables this anticorrelation can be reversed depending on masses of DM and instrumental parameters.

It is worth noting that these “halo” uncertainties appear larger than uncertainties coming from solar composition and/or nuclear/particle physics (for a recent estimate see [16]) and thus provide in most cases the dominant limiting factor in the extracting particle physics information from a future detection of a DM neutrino signal, even when normalizing the rate to some direct detection event rates. This is true in particular if no or little spectral information is available. Enhancements in the annihilation rate are a generic expectation of the presence of a “dark disk” created by interaction of the halo particles with the baryonic disk, but its effect can range from few percent to orders of magnitude. The role of substructures appears instead irrelevant, barring for highly unlikely circumstances. In principle, the velocity distribution of WIMP DM can be reconstructed from direct dark matter detection data in the range of velocities probed (provided the DM mass is sufficiently constrained), but $\mathcal{O}(100)$ events are needed to start improving the knowledge of $f_1(v)$ to better than what inferred indirectly via considerations as those reported in Sec. III (see e.g. [45]).

The considerations developed in this article have several implications, whose exploration is left for future works. One obvious consequence is that great care must be paid when comparing the constraints on the spin-dependent scattering cross section inferred from direct detection with those inferred by neutrino experiments (see e.g. [46–48]), especially when light DM particles are considered, as well as combinations of experiments with different target masses. This is notably the case of the

comparison of the DAMA and CoGent results with the exclusion plots obtained by other experiments (see e.g. Ref. [49] and references therein).

Another improvement over the first estimate provided in this article would consist in providing a more statistically sound assessment of the error range in the parameters governing the velocity structure of the halo (and the motion of the Sun through it), along the lines of what done in [12] for the local halo density. Here we implicitly assumed flat priors in the parameter space and uncorrelated variations of the different parameters. Needless to

say, such refinements would become compelling for extracting particle physics parameters if the present generation of WIMP DM detectors were to show any evidence for these particles.

Acknowledgments

We would like to thank F. Iocco and G. Lake for discussions during the initial stages of this work.

-
- [1] G. Jungman, M. Kamionkowski and K. Griest, *Phys. Rept.* **267** (1996) 195.
- [2] L. Bergstrom, *Rept. Prog. Phys.* **63** (2000) 793.
- [3] G. Bertone, D. Hooper and J. Silk, *Phys. Rept.* **405** (2005) 279.
- [4] *Particle Dark Matter: Observations, Models and Searches*, ed. G. Bertone, 2010, Cambridge University Press
- [5] P. Nath *et al.*, *Nucl. Phys. Proc. Suppl.* **200-202** (2010) 185.
- [6] G. Bertone, D. G. Cerdeno, M. Fornasa, R. R. de Austri and R. Trotta, arXiv:1005.4280 [hep-ph].
- [7] M. Drees and C. L. Shan, *JCAP* **0806** (2008) 012.
- [8] G. Bertone, D. G. Cerdeño, J. I. Collar and B. C. Odom, *Phys. Rev. Lett.* **99** (2007) 151301.
- [9] S. Profumo and P. Ullio, arXiv:1001.4086 [astro-ph.HE].
- [10] A. Gould, *Astrophys. J.* **388**, 338 (1992).
- [11] G. Bertone, arXiv:0710.5603 [astro-ph].
- [12] R. Catena and P. Ullio, arXiv:0907.0018 [astro-ph.CO].
- [13] L. E. Strigari and R. Trotta, *JCAP* **0911** (2009) 019.
- [14] P. Salucci, F. Nesti, G. Gentile and C. F. Martins, arXiv:1003.3101 [astro-ph.GA].
- [15] M. Pato, O. Agertz, G. Bertone, B. Moore and R. Teyssier, arXiv:1006.1322 [astro-ph.HE].
- [16] J. Ellis, K. A. Olive, C. Savage and V. C. Spanos, *Phys. Rev. D* **81**, 085004 (2010).
- [17] P. Belli, R. Cerulli, N. Fornengo and S. Scopel, *Phys. Rev. D* **66**, 043503 (2002).
- [18] M. Kuhlen *et al.*, *JCAP* **1002**, 030 (2010).
- [19] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, *Comput. Phys. Commun.* **180**, 747 (2009).
- [20] M. S. Alenazi and P. Gondolo, *Phys. Rev. D* **74**, 083518 (2006).
- [21] J. Lundberg and J. Edsjo, *Phys. Rev. D* **69**, 123505 (2004).
- [22] A. H. G. Peter, *Phys. Rev. D* **79**, 103531 (2009).
- [23] http://www.mpa-garching.mpg.de/~aldos/solar_main.htm
- [24] G. Dogan, A. Bonanno and J. Christensen-Dalsgaard, arXiv:1004.2215 [astro-ph.SR].
- [25] Kubala, Bizy & Mahan Rao “Earth’s Core Temperature” *Byrdand Black*, 1996; D. Alfe, D. Alfe, M. J. Gillan, and G. D. Price (2003) *Mineralogical Magazine* **67**, 113-123.
- [26] G. D. Mack, J. F. Beacom and G. Bertone, *Phys. Rev. D* **76** (2007) 043523.
- [27] A. Gould, *Astrophys. J.* **321**, 571 (1987).
- [28] P. Gondolo, J. Edsjo, P. Ullio, L. Bergstrom, M. Schelke and E. A. Baltz, *JCAP* **0407** (2004) 008.
- [29] F. Halzen and D. Hooper, *New J. Phys.* **11**, 105019 (2009).
- [30] J. Binney and S. Tremaine, “Galactic Dynamics”, Princeton Series in Astrophysics, 2nd edition (2008).
- [31] F. J. Kerr and D. Lynden-Bell, *Mon. Not. Roy. Astron. Soc.* **221**, 1023 (1986).
- [32] Y. Sofue, M. Honma and T. Omodaka, arXiv:0811.0859 [astro-ph].
- [33] A. K. Drukier, K. Freese and D. N. Spergel, *Phys. Rev. D* **33**, 3495 (1986).
- [34] J. F. Navarro *et al.*, *Astrophys. J.* 462 563 (1996).
- [35] M. C. Smith *et al.*, *Mon. Not. Roy. Astron. Soc.* **379**, 755 (2007).
- [36] M. Vogelsberger *et al.*, arXiv:0812.0362 [astro-ph].
- [37] M. Zemp *et al.*, arXiv:0812.2033 [astro-ph].
- [38] R. Schoenrich, J. Binney and W. Dehnen, arXiv:0912.3693 [astro-ph.GA].
- [39] D. R. Gies and J. W. Helsel, *Astrophys. J.* **626**, 844 (2005).
- [40] W. Dehnen and J. Binney, *Mon. Not. Roy. Astron. Soc.* **298**, 387 (1998).
- [41] S. M. Koushiappas and M. Kamionkowski, *Phys. Rev. Lett.* **103**, 121301 (2009).
- [42] J. Diemand, M. Kuhlen, P. Madau, M. Zemp, B. Moore, D. Potter and J. Stadel, *Nature* **454** (2008) 735.
- [43] J. I. Read, L. Mayer, A. M. Brooks, F. Governato and G. Lake, arXiv:0902.0009 [astro-ph.GA].
- [44] T. Bruch, A. H. G. Peter, J. Read, L. Baudis and G. Lake, *Phys. Lett. B* **674**, 250 (2009).
- [45] M. Drees and C. L. Shan, *JCAP* **0706**, 011 (2007).
- [46] T. Flacke, A. Menon, D. Hooper and K. Freese, arXiv:0908.0899 [hep-ph].
- [47] M. Blennow, H. Melbeus and T. Ohlsson, *JCAP* **1001**, 018 (2010).
- [48] P. Agrawal, Z. Chacko, C. Kilic and R. K. Mishra, arXiv:1003.5905 [hep-ph].
- [49] C. Savage, G. Gelmini, P. Gondolo and K. Freese, arXiv:1006.0972 [astro-ph.CO].