



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

Simulations of recoiling massive black holes

Guedes, J ; Madau, P ; Mayer, L ; Kuhlen, M ; Diemand, J ; Zemp, M

Abstract: The coalescence of black hole binaries is a significant source of gravitational wave radiation. The typically asymmetric nature of this emission, which carries linear momentum, can result in the recoil of the black hole remnant with velocities in the range $100 < V_{\text{recoil}} < 3750$ km s⁻¹. The detectability of recoiling massive black holes (MBH) as off-nuclear QSOs is tightly connected with the properties of the host galaxy, which determine the MBH's orbit and fuel reservoir. We present the results of N-body simulations of recoiling MBHs in high-resolution, non-axisymmetric potentials. We find that if the recoil velocities are high enough to reach regions of the galaxy dominated by the generally triaxial dark matter distribution, the return time is significantly extended when compared to a spherical distribution. We also perform simulations of recoiling MBHs traveling in gas merger remnants, where large amounts of gas have been funneled to the central regions. In this case, the MBHs remain within $R < 1$ kpc from the center of the host even for high recoil velocities ($V_{\text{recoil}} = 1200$ km s⁻¹) due to the compactness of the remnant galaxy's nuclear disk. We discuss the implications of both scenarios for detectability.

DOI: <https://doi.org/10.1017/S1743921310006393>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-41665>

Conference or Workshop Item

Published Version

Originally published at:

Guedes, J; Madau, P; Mayer, L; Kuhlen, M; Diemand, J; Zemp, M (2009). Simulations of recoiling massive black holes. In: IAU Symposium 267: Evolution of Galaxies and Central Black Holes: Feeding and Feedback, Rio de Janeiro, Brazil, 10 August 2009 - 14 August 2009. Cambridge University Press, 262.

DOI: <https://doi.org/10.1017/S1743921310006393>

Simulations of Recoiling Massive Black Holes

Javiera Guedes¹, Piero Madau¹, Lucio Mayer², Michael Kuhlen³,
Jürg Diemand², and Marcel Zemp⁴

¹Department of Astronomy & Astrophysics, University of California, Santa Cruz,
1156 High Street, Santa Cruz, CA 95062, USA
Email: javiera@ucolick.org

²Institute for Theoretical Physics, University of Zürich, Winterthurerstrasse 190, CH-9057
Zürich, Switzerland

³Theoretical Astrophysics Center, University of California Berkeley 601 Campbell Hall,
Berkeley, CA 94720, USA

⁴Department of Astronomy, University of Michigan, Ann Arbor, MI 48109, USA

Abstract. The coalescence of black hole binaries is a significant source of gravitational wave radiation. The typically asymmetric nature of this emission, which carries linear momentum, can result in the recoil of the black hole remnant with velocities in the range $100 < V_{\text{recoil}} < 3750 \text{ km s}^{-1}$. The detectability of recoiling massive black holes (MBH) as off-nuclear QSOs is tightly connected with the properties of the host galaxy, which determine the MBH's orbit and fuel reservoir. We present the results of N -body simulations of recoiling MBHs in high-resolution, non-axisymmetric potentials. We find that if the recoil velocities are high enough to reach regions of the galaxy dominated by the generally triaxial dark matter distribution, the return time is significantly extended when compared to a spherical distribution. We also perform simulations of recoiling MBHs traveling in gas merger remnants, where large amounts of gas have been funneled to the central regions. In this case, the MBHs remain within $R < 1 \text{ kpc}$ from the center of the host even for high recoil velocities ($V_{\text{recoil}} = 1200 \text{ km s}^{-1}$) due to the compactness of the remnant galaxy's nuclear disk. We discuss the implications of both scenarios for detectability.

Keywords. galaxies: nuclei, gravitational waves, methods: n-body simulations

Axisymmetry is typically broken in real galaxy mergers. To grasp the extent to which deviations from symmetry are important to the detectability of recoiling MBHs as off-nuclear QSOs, we first study the effect of the triaxiality of the dark matter distribution using full N -body simulations of MBHs embedded in the main halo of the Via Lactea I simulation (Diemand *et al.* 2007). We use MBH orbits in pure dark matter simulations to constrain a model of a composite galaxy (a stellar spheroid plus a triaxial halo) and find that ejected MBHs with initial recoil velocities $V_{\text{recoil}} > 500 \text{ km s}^{-1}$ do not return to the center of the host within a Hubble time (Guedes *et al.* 2008, 2009). To assess the effect of gas, we run simulations of recoiling MBHs in the high-resolution merger of Mayer *et al.* (2007). In this case, MBHs are not removed from the nuclear region. In both cases, detectability is challenging. In dry mergers, for instance, apocenter distances are high but the MBH fuel reservoir is scarce, while the opposite is true in major gas mergers.

References

- Diemand, J., Kuhlen, M., & Madau, P. 2007, *ApJ*, 657, 262
Guedes, J., Diemand, J., Zemp, M., Kuhlen, M., Madau, P., & Mayer, L. 2008, *AN*, 329, 1004
Guedes, J., Madau, P., Kuhlen, M., Diemand, J., & Zemp, M. 2009, *ApJ*, 702, 890
Mayer, L., Kazantzidis, S., Madau, P., Colpi, M., Quinn, T., & Wadsley, J. 2007, *Science*, 316, 1874