



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2010

---

## **A new crystal modification of diammonium hydrogen phosphate, (NH<sub>4</sub>)<sub>2</sub>(HPO<sub>4</sub>)**

Kunz, P C ; Wetzl, C ; Spingler, B

DOI: <https://doi.org/10.1107/S1600536810009839>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-44992>

Journal Article

Originally published at:

Kunz, P C ; Wetzl, C ; Spingler, B (2010). A new crystal modification of diammonium hydrogen phosphate, (NH<sub>4</sub>)<sub>2</sub>(HPO<sub>4</sub>). *Acta Crystallographica. Section E, Structure Reports Online*, E66(Pt. 4):i26-i27.

DOI: <https://doi.org/10.1107/S1600536810009839>

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W.T.A. Harrison, J. Simpson and  
M. Weil

## A new crystal modification of diammonium hydrogen phosphate, $(\text{NH}_4)_2(\text{HPO}_4)$

Peter C. Kunz, Corinna Wetzel and Bernhard Spingler

*Acta Cryst.* (2010). E66, i26–i27

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

# A new crystal modification of diammonium hydrogen phosphate, (NH<sub>4</sub>)<sub>2</sub>(HPO<sub>4</sub>)

Peter C. Kunz,<sup>a\*</sup> Corinna Wetzel<sup>a</sup> and Bernhard Spingler<sup>b</sup>

<sup>a</sup>Heinrich-Heine-Universität Düsseldorf, Institut für Anorganische Chemie und Strukturchemie I, Universitätsstrasse 1, D-40225 Düsseldorf, Germany, and

<sup>b</sup>Anorganisch-Chemisches Institut, Universität Zürich-Irchel, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

Correspondence e-mail: peter.kunz@uni-duesseldorf.de

Received 8 February 2010; accepted 16 March 2010

Key indicators: single-crystal X-ray study;  $T = 183$  K; mean  $\sigma(P-O) = 0.002$  Å;  $R$  factor = 0.048;  $wR$  factor = 0.158; data-to-parameter ratio = 29.7.

The addition of hexafluoridophosphate salts (ammonium, silver, thallium or potassium) is usually used to precipitate complex cations from aqueous solutions. It has long been known that  $PF_6^-$  is sensitive towards hydrolysis under acidic conditions [Gebala & Jones (1969). *J. Inorg. Nucl. Chem.* **31**, 771–776; Plakhotnyk *et al.* (2005). *J. Fluorine Chem.* **126**, 27–31]. During the course of our investigation into coinage metal complexes of diphosphine ligands, we used ammonium hexafluoridophosphate in order to crystallize  $[Ag(\text{diphosphine})_2]PF_6$  complexes. From these solutions we always obtained needle-like crystals which turned out to be the title compound,  $2NH_4^+ \cdot HPO_4^{2-}$ . It was received as the hydrolysis product of  $NH_4PF_6$ . The crystals are a new modification of diammonium hydrogen phosphate. In contrast to the previously published polymorph [Khan *et al.* (1972). *Acta Cryst.* **B28**, 2065–2069],  $Z'$  of the title compound is 2. In the new modification of the title compound, there are eight molecules of  $(NH_4)_2(HPO_4)$  in the unit cell. The structure consists of  $PO_3OH$  and  $NH_4$  tetrahedra, held together by  $O-H \cdots O$  and  $N-H \cdots O$  hydrogen bonds.

## Related literature

For the study of another crystal modification of the title compound, see: Khan *et al.* (1972). For the hydrolysis of hexafluoridophosphates, see: Akbayeva *et al.* (2006); Deifel *et al.* (2008); Fernandez-Galan *et al.* (1994); Gebala & Jones (1969); Nikitenko *et al.* (2007); Plakhotnyk *et al.* (2005).

## Experimental

### Crystal data

$2NH_4^+ \cdot HPO_4^{2-}$   
 $M_r = 132.06$   
 Monoclinic,  $P2_1/c$

$a = 11.2868$  (3) Å  
 $b = 15.3466$  (4) Å  
 $c = 6.41894$  (19) Å

$\beta = 90.795$  (3)°  
 $V = 1111.74$  (5) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation

$\mu = 0.42$  mm<sup>-1</sup>  
 $T = 183$  K  
 $0.44 \times 0.17 \times 0.11$  mm

### Data collection

Oxford Xcalibur Ruby CCD diffractometer  
 Absorption correction: multi-scan *CrysAlis PRO* (Oxford Diffraction, 2009)  
 $T_{\min} = 0.891$ ,  $T_{\max} = 0.955$

22537 measured reflections  
 5384 independent reflections  
 4400 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.158$   
 $S = 1.21$   
 5384 reflections  
 181 parameters  
 16 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.82$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.69$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1–H1 $\cdots$ O7 <sup>i</sup>	0.78 (4)	1.80 (4)	2.570 (2)	168 (4)
O5–H5 $\cdots$ O8 <sup>ii</sup>	0.85 (4)	1.79 (4)	2.632 (2)	170 (4)
N11–H11A $\cdots$ O4 <sup>iii</sup>	0.86 (2)	1.89 (2)	2.747 (2)	175 (4)
N11–H11B $\cdots$ O8 <sup>i</sup>	0.86 (2)	2.10 (2)	2.951 (2)	171 (3)
N11–H11C $\cdots$ O3 <sup>iv</sup>	0.88 (2)	1.99 (2)	2.852 (3)	169 (3)
N11–H11D $\cdots$ O4 <sup>i</sup>	0.88 (2)	2.01 (2)	2.870 (2)	167 (3)
N12–H12A $\cdots$ O6 <sup>v</sup>	0.87 (2)	1.91 (2)	2.755 (2)	165 (3)
N12–H12B $\cdots$ O5 <sup>vi</sup>	0.88 (2)	2.16 (2)	3.008 (3)	161 (3)
N12–H12C $\cdots$ O6 <sup>i</sup>	0.87 (2)	1.99 (2)	2.827 (2)	161 (3)
N12–H12D $\cdots$ O2	0.88 (2)	1.88 (2)	2.754 (2)	175 (3)
N13–H13A $\cdots$ O3	0.86 (2)	1.92 (2)	2.773 (2)	172 (3)
N13–H13B $\cdots$ O2 <sup>vii</sup>	0.86 (2)	1.96 (2)	2.822 (2)	175 (3)
N13–H13C $\cdots$ O6 <sup>v</sup>	0.89 (2)	1.95 (2)	2.830 (2)	168 (3)
N13–H13D $\cdots$ O2 <sup>ii</sup>	0.86 (2)	1.96 (2)	2.820 (2)	176 (3)
N14–H14A $\cdots$ O7	0.87 (2)	1.90 (2)	2.771 (2)	174 (3)
N14–H14B $\cdots$ O8 <sup>vii</sup>	0.86 (2)	2.01 (2)	2.859 (2)	171 (3)
N14–H14C $\cdots$ O3 <sup>iv</sup>	0.86 (2)	1.92 (2)	2.784 (2)	178 (3)
N14–H14D $\cdots$ O4	0.89 (2)	1.89 (2)	2.771 (2)	171 (3)

Symmetry codes: (i)  $-x, -y + 1, -z + 1$ ; (ii)  $x, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (iii)  $-x, y + \frac{1}{2}, -z + \frac{3}{2}$ ; (iv)  $-x, -y + 1, -z + 2$ ; (v)  $x + 1, y, z + 1$ ; (vi)  $x + 1, y, z$ ; (vii)  $x, y, z + 1$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

BS thanks the University of Zürich for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BR2137).

## References

- Akbayeva, D., Vaira, M., Costantini, S., Peruzzini, M. & Stoppioni, P. (2006). *Dalton Trans.* pp. 389–395.  
 Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.  
 Deifel, N. P., Holman, K. T. & Cahil, C. (2008). *Chem. Commun.* pp. 6037–6038.

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Fernandez-Galan, R., Manzano, B., Otero, A., Lanfranchi, M. & Pellinghelli, M. (1994). *Inorg. Chem.* **33**, 2309–2312.
- Gebala, A. & Jones, M. (1969). *J. Inorg. Nucl. Chem.* **31**, 771–776.
- Khan, A. A., Roux, J. P. & James, W. J. (1972). *Acta Cryst.* **B28**, 2065–2069.
- Nikitenko, S., Berthon, C. & Moisy, P. (2007). *C. R. Chim.* **10**, 1122–1127.
- Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction, Yarnton, England.
- Plakhotnyk, A., Ernst, L. & Schmutzler, R. (2005). *J. Fluorine Chem.* **126**, 27–31.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2010). E66, i26-i27 [ doi:10.1107/S1600536810009839 ]

## A new crystal modification of diammonium hydrogen phosphate, $(\text{NH}_4)_2(\text{HPO}_4)$

P. C. Kunz, C. Wetzel and B. Spingler

### Comment

The addition of hexafluorophosphate salts (ammonium, silver, thallium or potassium) is usually used to precipitate complex cations from aqueous solutions. It is long known, that  $\text{PF}_6^-$  is sensitive towards hydrolysis under acidic conditions (Gebala and Jones 1969; Plakhotnyk, Ernst *et al.* 2005). In organic solvents the spatial hydrolysis to intermediate species as HF,  $\text{POF}_3$  and  $\text{PO}_2\text{F}_2^-$  is observed (Fernandez-Galan, Manzano *et al.* 1994; Akbayeva, Vaira *et al.* 2006; Nikitenko, Berthon *et al.* 2007) and under hydrothermal conditions this reaction is used for the formation of phosphate materials (Deifel, Holman *et al.* 2008).

During the course of our investigation into coinage metal complexes of diphosphine ligands we used ammonium hexafluorophosphate in order to crystallise  $[\text{Ag}(\text{diphosphine})_2]\text{PF}_6$  complexes. From these solutions we always obtained needle-like crystals which turned out to be the title compound  $(\text{NH}_4)_2(\text{HPO}_4)$  as the product of hydrolysis of  $\text{NH}_4\text{PF}_6$ .

A modification of diammonium hydrogen phosphate is known with the cell parameters  $a = 11.043$  (6),  $b = 6.700$  (3),  $c = 8.031$  (4),  $\beta = 113.42$  (3) $^\circ$  and  $Z = 4$ . (Khan, Roux *et al.* 1972) In the new modification of the title compound reported here, there are eight molecules of  $(\text{NH}_4)_2(\text{HPO}_4)$  in the unit cell. The structure consists of  $\text{PO}_4$  and  $\text{NH}_4$  tetrahedra, held together by  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds. These hydrogen bonds are more or less linear ( $169^\circ < \angle(\text{DHA}) < 178^\circ$ ). Of the four P-O bonds in both  $\text{PO}_4$  tetrahedra, one is longer than the remaining three, typical of a  $\text{O}_3\text{P}(\text{OH})$  group. The two different  $\text{HPO}_4^-$  molecules ( $\text{P1O}_4$  and  $\text{P2O}_4$ ) are hydrogen bonded to ten and seven ammonium ions, respectively. In hydrogen phosphate P2, one  $\text{NH}_4^+$  molecule is bound to O5 and O7, two to O8 and three to O6. In the other hydrogen phosphate, the three non-protonated O-atoms O2, O3 and O4 are bound to three  $\text{NH}_4^+$  molecules each, whereas the protonated O1 is only bound to one  $\text{NH}_4^+$  molecule. In the hydrogen phosphate P1 the hydroxyl group O1H1 forms a hydrogen bridge to O7 ( $d = 1.80$  (4) Å) and in the hydrogen phosphate P2 the hydroxyl group O5H5 forms a hydrogen bridge to O8 ( $d = 1.79$  (4) Å). The N—O distances around the four-coordinated ammonium ions N13 and N14 are within the range of  $2.77 < d < 2.86$  Å; The N $\cdots$ O distances around N12 fall within this range with the exception of N12 $\cdots$ O5 which is significantly longer ( $d_{\text{N12O5}} = 3.007$  Å). A very different picture is found around N11. Here, five neighbouring O-atoms are found, three of which are in a shorter distance ( $2.74 < d < 2.87$  Å) and two in a longer distance ( $d_{\text{N11O5}} = 2.951$  Å and  $d_{\text{N11O1}} = 3.048$  Å). The fifth N $\cdots$ O contact may be a result of dynamic or static disorder of the ammonium ion or that each N atom in addition to three normal N—H $\cdots$ O bonds also formed one bifurcated bond. This is in contrast to the other modification, in which each ammonium ion has five N $\cdots$ O contacts smaller than 3.4 Å. Since Khan and Roux (Khan, Roux *et al.* 1972) only reported that they used "a commercially supplied crystalline sample", we cannot compare the crystallization conditions that lead to the two different crystal forms.

## Experimental

A suitable crystal was covered with oil (Infineum V8512, formerly known as Paratone N), mounted on top of a glass fibre and immediately transferred to the diffractometer. The final model was checked for higher symmetry with help of the program PLATON (Spek, 2009).

## Refinement

All hydrogen atoms were located by difference Fourier synthesis and refined with fixed individual displacement parameters [ $U(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ ] at an O-H distance of 0.87 Å.

## Figures





Fig. 1. H-bonded network in the solid-state of the title compound. Displacement ellipsoids are drawn at a 50 % level, H-atoms are represented as capped sticks.

## diammonium hydrogen phosphate

### Crystal data



$$M_r = 132.06$$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$$a = 11.2868 (3) \text{ \AA}$$

$$b = 15.3466 (4) \text{ \AA}$$

$$c = 6.41894 (19) \text{ \AA}$$

$$\beta = 90.795 (3)^\circ$$

$$V = 1111.74 (5) \text{ \AA}^3$$

$$Z = 8$$

$$F(000) = 560$$

$$D_x = 1.578 \text{ Mg m}^{-3}$$

Mo  $K\alpha$  radiation,  $\lambda = 0.7107 \text{ \AA}$

Cell parameters from 11185 reflections

$$\theta = 2.7\text{--}37.6^\circ$$

$$\mu = 0.42 \text{ mm}^{-1}$$

$$T = 183 \text{ K}$$

Needle, colourless

$$0.44 \times 0.17 \times 0.11 \text{ mm}$$

### Data collection

Oxford Xcalibur Ruby CCD diffractometer

Radiation source: Enhance (Mo) X-ray Source graphite

Detector resolution: 10.4498 pixels  $\text{mm}^{-1}$

$\omega$  oscillation scan

Absorption correction: multi-scan

CrysAlis PRO (Oxford Diffraction, 2009)

$$T_{\text{min}} = 0.891, T_{\text{max}} = 0.955$$

5384 independent reflections

4400 reflections with  $I > 2\sigma(I)$

$$R_{\text{int}} = 0.033$$

$$\theta_{\text{max}} = 36.3^\circ, \theta_{\text{min}} = 2.7^\circ$$

$$h = -18 \rightarrow 18$$

$$k = -25 \rightarrow 25$$

$$l = -9 \rightarrow 10$$

22537 measured reflections

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.048$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.158$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.21$	$w = 1/[\sigma^2(F_o^2) + (0.0508P)^2 + 1.9491P]$
5384 reflections	where $P = (F_o^2 + 2F_c^2)/3$
181 parameters	$(\Delta/\sigma)_{\max} < 0.001$
16 restraints	$\Delta\rho_{\max} = 0.82 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.69 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.12341 (16)	0.48227 (11)	0.6812 (3)	0.0260 (4)
H1	0.181 (4)	0.502 (3)	0.635 (6)	0.039*
O2	0.25104 (13)	0.34866 (10)	0.7264 (2)	0.0180 (3)
O3	0.13920 (15)	0.40992 (11)	1.0319 (2)	0.0198 (3)
O4	0.02645 (13)	0.34042 (10)	0.7334 (3)	0.0190 (3)
O5	-0.41866 (14)	0.29289 (11)	0.4731 (3)	0.0209 (3)
H5	-0.368 (3)	0.259 (3)	0.529 (6)	0.031*
O6	-0.47633 (13)	0.40326 (10)	0.2142 (2)	0.0174 (3)
O7	-0.30247 (15)	0.43252 (11)	0.4549 (2)	0.0209 (3)
O8	-0.28270 (13)	0.32102 (10)	0.1683 (2)	0.0182 (3)
P1	0.13662 (4)	0.39195 (3)	0.79806 (7)	0.01228 (10)
P2	-0.36645 (4)	0.36515 (3)	0.32105 (7)	0.01211 (10)
N11	0.03108 (16)	0.66864 (12)	0.7030 (3)	0.0185 (3)
H11A	0.009 (3)	0.7219 (13)	0.720 (5)	0.028*
H11B	0.1058 (17)	0.667 (2)	0.730 (5)	0.028*
H11C	-0.017 (3)	0.638 (2)	0.780 (5)	0.028*



## supplementary materials

---

H11D	0.018 (3)	0.658 (2)	0.570 (3)	0.028*
N12	0.47645 (17)	0.41255 (13)	0.7925 (3)	0.0188 (3)
H12A	0.500 (3)	0.402 (2)	0.920 (3)	0.028*
H12B	0.523 (3)	0.382 (2)	0.711 (5)	0.028*
H12C	0.479 (3)	0.4675 (13)	0.760 (5)	0.028*
H12D	0.4031 (19)	0.395 (2)	0.770 (5)	0.028*
N13	0.29995 (16)	0.32905 (12)	1.2991 (3)	0.0182 (3)
H13A	0.252 (3)	0.351 (2)	1.207 (4)	0.027*
H13B	0.285 (3)	0.339 (2)	1.428 (3)	0.027*
H13C	0.3748 (18)	0.345 (2)	1.278 (5)	0.027*
H13D	0.288 (3)	0.2742 (12)	1.282 (5)	0.027*
N14	-0.18493 (17)	0.42049 (12)	0.8356 (3)	0.0178 (3)
H14A	-0.227 (3)	0.425 (2)	0.720 (4)	0.027*
H14B	-0.221 (3)	0.394 (2)	0.933 (4)	0.027*
H14C	-0.169 (3)	0.4726 (14)	0.877 (5)	0.027*
H14D	-0.1132 (19)	0.398 (2)	0.814 (5)	0.027*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0185 (7)	0.0193 (7)	0.0403 (10)	0.0021 (6)	0.0007 (6)	0.0145 (7)
O2	0.0140 (6)	0.0182 (6)	0.0220 (7)	0.0022 (5)	0.0025 (5)	-0.0015 (5)
O3	0.0225 (7)	0.0215 (7)	0.0153 (6)	0.0000 (5)	-0.0005 (5)	-0.0035 (5)
O4	0.0142 (6)	0.0179 (6)	0.0249 (7)	-0.0037 (5)	-0.0022 (5)	-0.0034 (5)
O5	0.0183 (6)	0.0207 (7)	0.0237 (7)	0.0025 (5)	0.0055 (5)	0.0099 (5)
O6	0.0145 (6)	0.0189 (6)	0.0188 (6)	0.0015 (5)	-0.0032 (5)	0.0032 (5)
O7	0.0221 (7)	0.0242 (7)	0.0165 (6)	-0.0048 (6)	-0.0035 (5)	-0.0031 (5)
O8	0.0150 (6)	0.0206 (7)	0.0192 (6)	-0.0010 (5)	0.0046 (5)	-0.0033 (5)
P1	0.01141 (19)	0.01162 (19)	0.01380 (19)	-0.00003 (14)	-0.00053 (14)	-0.00007 (14)
P2	0.01113 (18)	0.0140 (2)	0.01122 (18)	-0.00061 (14)	0.00038 (13)	0.00073 (14)
N11	0.0167 (7)	0.0184 (7)	0.0205 (8)	0.0015 (6)	-0.0007 (6)	-0.0001 (6)
N12	0.0170 (7)	0.0226 (8)	0.0169 (7)	0.0013 (6)	-0.0008 (6)	-0.0018 (6)
N13	0.0160 (7)	0.0199 (7)	0.0187 (7)	-0.0015 (6)	-0.0001 (5)	0.0002 (6)
N14	0.0178 (7)	0.0197 (7)	0.0159 (7)	-0.0006 (6)	0.0001 (5)	-0.0016 (6)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—P1	1.5821 (17)	N11—H11D	0.877 (18)
O1—H1	0.78 (4)	N12—H12A	0.869 (18)
O2—P1	1.5287 (15)	N12—H12B	0.878 (18)
O3—P1	1.5257 (16)	N12—H12C	0.868 (18)
O4—P1	1.5262 (15)	N12—H12D	0.879 (18)
O5—P2	1.5955 (16)	N13—H13A	0.863 (18)
O5—H5	0.85 (4)	N13—H13B	0.862 (18)
O6—P2	1.5254 (15)	N13—H13C	0.892 (18)
O7—P2	1.5201 (16)	N13—H13D	0.859 (18)
O8—P2	1.5295 (15)	N14—H14A	0.874 (18)
N11—H11A	0.862 (18)	N14—H14B	0.856 (18)
N11—H11B	0.859 (18)	N14—H14C	0.861 (18)

N11—H11C	0.876 (18)	N14—H14D	0.890 (18)
P1—O1—H1	116 (3)	H11C—N11—H11D	110 (3)
P2—O5—H5	115 (3)	H12A—N12—H12B	107 (3)
O3—P1—O4	111.44 (9)	H12A—N12—H12C	113 (3)
O3—P1—O2	111.70 (9)	H12B—N12—H12C	111 (3)
O4—P1—O2	112.43 (9)	H12A—N12—H12D	112 (3)
O3—P1—O1	107.95 (10)	H12B—N12—H12D	108 (3)
O4—P1—O1	104.71 (10)	H12C—N12—H12D	106 (3)
O2—P1—O1	108.21 (9)	H13A—N13—H13B	118 (3)
O7—P2—O6	111.73 (9)	H13A—N13—H13C	112 (3)
O7—P2—O8	111.74 (9)	H13B—N13—H13C	107 (3)
O6—P2—O8	112.79 (9)	H13A—N13—H13D	101 (3)
O7—P2—O5	107.68 (9)	H13B—N13—H13D	105 (3)
O6—P2—O5	103.69 (9)	H13C—N13—H13D	113 (3)
O8—P2—O5	108.73 (9)	H14A—N14—H14B	114 (3)
H11A—N11—H11B	107 (3)	H14A—N14—H14C	107 (3)
H11A—N11—H11C	105 (3)	H14B—N14—H14C	109 (3)
H11B—N11—H11C	119 (3)	H14A—N14—H14D	112 (3)
H11A—N11—H11D	105 (3)	H14B—N14—H14D	112 (3)
H11B—N11—H11D	110 (3)	H14C—N14—H14D	102 (3)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 $\cdots$ O7 <sup>i</sup>	0.78 (4)	1.80 (4)	2.570 (2)	168 (4)
O5—H5 $\cdots$ O8 <sup>ii</sup>	0.85 (4)	1.79 (4)	2.632 (2)	170 (4)
N11—H11A $\cdots$ O4 <sup>iii</sup>	0.86 (2)	1.89 (2)	2.747 (2)	175 (4)
N11—H11B $\cdots$ O8 <sup>i</sup>	0.86 (2)	2.10 (2)	2.951 (2)	171 (3)
N11—H11C $\cdots$ O3 <sup>iv</sup>	0.88 (2)	1.99 (2)	2.852 (3)	169 (3)
N11—H11D $\cdots$ O4 <sup>i</sup>	0.88 (2)	2.01 (2)	2.870 (2)	167 (3)
N12—H12A $\cdots$ O6 <sup>v</sup>	0.87 (2)	1.91 (2)	2.755 (2)	165 (3)
N12—H12B $\cdots$ O5 <sup>vi</sup>	0.88 (2)	2.16 (2)	3.008 (3)	161 (3)
N12—H12C $\cdots$ O6 <sup>i</sup>	0.87 (2)	1.99 (2)	2.827 (2)	161 (3)
N12—H12D $\cdots$ O2	0.88 (2)	1.88 (2)	2.754 (2)	175 (3)
N13—H13A $\cdots$ O3	0.86 (2)	1.92 (2)	2.773 (2)	172 (3)
N13—H13B $\cdots$ O2 <sup>vii</sup>	0.86 (2)	1.96 (2)	2.822 (2)	175 (3)
N13—H13C $\cdots$ O6 <sup>v</sup>	0.89 (2)	1.95 (2)	2.830 (2)	168 (3)
N13—H13D $\cdots$ O2 <sup>ii</sup>	0.86 (2)	1.96 (2)	2.820 (2)	176 (3)
N14—H14A $\cdots$ O7	0.87 (2)	1.90 (2)	2.771 (2)	174 (3)
N14—H14B $\cdots$ O8 <sup>vii</sup>	0.86 (2)	2.01 (2)	2.859 (2)	171 (3)
N14—H14C $\cdots$ O3 <sup>iv</sup>	0.86 (2)	1.92 (2)	2.784 (2)	178 (3)
N14—H14D $\cdots$ O4	0.89 (2)	1.89 (2)	2.771 (2)	171 (3)

Symmetry codes: (i)  $-x, -y+1, -z+1$ ; (ii)  $x, -y+1/2, z+1/2$ ; (iii)  $-x, y+1/2, -z+3/2$ ; (iv)  $-x, -y+1, -z+2$ ; (v)  $x+1, y, z+1$ ; (vi)  $x+1, y, z$ ; (vii)  $x, y, z+1$ .

Fig. 1

