**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Possible hazardous effects of hydrofluoric acid and recommendations for treatment approach: a review

Özcan, M ; Allahbeickaraghi, A ; Dündar, M

Abstract: Hydrofluoric acid (HF) is commonly used for conditioning the glass ceramics either prior to cementation or for intraoral repair in prosthetic and restorative dentistry. The present study offers a review of chemical properties of HF used, highlight the possible hazardous effects of this agent, and to recommend the treatment approach for potential risks. Available published information documented in PubMed, Medline, and Picarta literature databases was reviewed. Additional information was derived from scientific reports, medical and chemical textbooks, handbooks, product information, manufacturers' instructions, Internet web sites of the HF manufacturers. No report was found on the incidence of the hazardous effects of HF in dentistry. Reports from other fields presented incidences of acute and chronic symptoms in exposure to HF. While acute symptoms include skin or nail burns, chronic ones involve systemic toxicity, eye injuries, inhalation and ingestion-related symptoms that can be even fatal. HF can be harmful and particularly aggressive to soft tissues, but symptoms may not be apparent immediately after exposure. The hazardous effects are not based on the pH value, but on the toxicity of HF. Potential hazards of HF known from other applications than dentistry should be considered also in dental applications. Especially the clinicians, who often deal with adhesive cementation or repair of glass ceramics, should take necessary precautions for possible hazards of HF.

DOI: <https://doi.org/10.1007/s00784-011-0636-6>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-62951>

Journal Article

Published Version

Originally published at:

Özcan, M; Allahbeickaraghi, A; Dündar, M (2012). Possible hazardous effects of hydrofluoric acid and recommendations for treatment approach: a review. *Clinical Oral Investigations*, 16(1):15-23.

DOI: <https://doi.org/10.1007/s00784-011-0636-6>

Possible hazardous effects of hydrofluoric acid and recommendations for treatment approach: a review

Mutlu Özcan · Arezo Allahbeickaraghi · Mine Dündar

Received: 2 September 2010 / Accepted: 26 October 2011 / Published online: 9 November 2011
© Springer-Verlag 2011

Abstract Hydrofluoric acid (HF) is commonly used for conditioning the glass ceramics either prior to cementation or for intraoral repair in prosthetic and restorative dentistry. The present study offers a review of chemical properties of HF used, highlight the possible hazardous effects of this agent, and to recommend the treatment approach for potential risks. Available published information documented in PubMed, Medline, and Picarta literature databases was reviewed. Additional information was derived from scientific reports, medical and chemical textbooks, handbooks, product information, manufacturers' instructions, Internet web sites of the HF manufacturers. No report was found on the incidence of the hazardous effects of HF in dentistry. Reports from other fields presented incidences of acute and chronic symptoms in exposure to HF. While acute symptoms include skin or nail burns, chronic ones involve systemic toxicity, eye injuries, inhalation and ingestion-related symptoms that can be even fatal. HF can be harmful and particularly aggressive to soft tissues, but symptoms may not be apparent immediately after exposure. The hazardous effects are not based on the pH

value, but on the toxicity of HF. Potential hazards of HF known from other applications than dentistry should be considered also in dental applications. Especially the clinicians, who often deal with adhesive cementation or repair of glass ceramics, should take necessary precautions for possible hazards of HF.

Keywords Dental ceramics · Dentistry · Hazardous compounds · Hydrofluoric acid

Introduction

Highly concentrated, pure aqueous solution of hydrofluoric acid (HF) was first prepared in 1809 and hydrogen fluoride was first obtained at the end of the eighteenth century [1, 2]. From chemistry perspective, HF is the inorganic acid of elemental fluorine. Hydrogen fluoride is produced by the reaction of calcium fluoride (CaF_2) and sulfuric acid in a reaction furnace at 200°C, which is then cooled and stored as a colorless liquid [1, 2]. The main source of CaF_2 to produce hydrogen fluoride is the mineral fluorspar. The volatile HF is condensed and purified by distillation [1, 2]. HF can be produced through processes such as aluminum melting, ceramic production, and phosphate fertilizers. It may also occur as an impurity in different chemicals or can be produced through the reaction of other chemicals [1]. It is the starting material for the production of most organic and inorganic chemicals containing the fluorine atom. Fluorine gas is the most important fluorine compound for industrial use.

HF solutions can be formed when fluoride salts are acidified with acidic solutions. As it is a weak acid, hydrogen fluoride in water for instance, is not completely ionized. Its conjugate base, the fluoride ion F^- , can re-associate to form HF in solutions with low pH [3]. In strong

M. Özcan (✉)
University of Zürich, Dental Materials Unit,
Center for Dental and Oral Medicine, Clinic for Fixed
and Removable Prosthodontics and Dental Materials Science,
Plattenstrasse 11,
CH-8032 Zurich, Switzerland
e-mail: mutluozcan@hotmail.com

A. Allahbeickaraghi
Dental School, University of Groningen,
Antonius Deusinglaan 1,
9713 AV Groningen, The Netherlands

M. Dündar
Department of Prosthodontics, School of Dentistry,
Ege University,
35100 Izmir, Turkey

acidic solutions, the pH of the fluoride-containing mixture will be low and the free F^- formed from the ionization of the fluoride salt in aqueous solution will exist as HF. However, in acidic aqueous suspensions, solubility increases by the presence of the acid, and with this route a significant quantity of HF can be produced [1].

Early versions of HF were aqueous solutions. Aqueous HF is commonly formulated for products with lower concentration. The major use of aqueous HF for occupational purposes are etching and polishing glass and ceramics, removing sand from metal castings, and cleaning processes for floors and walls [3]. Other important uses include fabric rust removal, etching silicon in semiconductors, cleaning stone and marble, and manufacturing inorganic fluoride compounds [4, 5].

Since the introduction of glass-based ceramics and discovery of the benefits of adhesive cementation in dentistry, HF started to be used to condition restorative and prosthetic devices [6]. Strong bonds are formed between resin-based materials and dental ceramics that are etched with HF [7]. This is based on the affinity of fluoride to silicon that is higher than to oxygen: $4HF + SiO_2 \rightarrow SiF_4 + 2H_2O$. By the selective removal of the glassy matrix, principally micro-mechanical retention is achieved between the resin and the etched ceramic. The greatest advantage of the use of HF for ceramics is that its chair-side application is very simple, and for this reason, sophisticated conditioning or cleaning methods may not be necessary [8].

In 1998, in fields other than dentistry, HF was acknowledged as a highly hazardous chemical due to its toxicity, corrosiveness, and reactivity [1]. It can be used in the anhydrous (100%) form or in aqueous solutions where the latter range in concentration from <1–70%. Severity and rapidity of onset of signs and symptoms depends on the concentration, duration of exposure, and penetration to the exposed tissue [6]. When accidentally released, HF can diffuse as a dense vapor and aerosol. HF itself is not flammable, explosive, or oxidizing, but it is a highly reactive chemical. Hence, its storage conditions and handling must be taken into account when using this acid. Also, contact with metals can lead to the formation of hydrogen gas, which forms explosive mixtures in air at concentrations between 4% and 75% [1, 2]. The reaction occurs at concentrations of approximately less than 65% of aqueous HF but over time at higher concentrations of anhydrous HF. In closed containers, hydrogen formation can cause dangerously high pressures [9, 10]. As the reaction of hydrogen fluoride or concentrated HF with water is exothermic, it may react violently on mixing. Concentrated aqueous solutions can fume in the air, and heat is produced upon reaction.

The objective of this review was to summarize the chemical properties, applications of HF used in medical or

other fields as well as in restorative and prosthetic dentistry, to investigate the incidence of reports on hazardous effects of HF in dentistry, to highlight the possible hazardous effects of this agent in general, and to recommend the treatment approach for potential risks, based on the published information available in the chemical and medical fields.

Materials and methods

Original scientific papers listed in PubMed, Medline, Picarta up to the end of 2010 on HF were included in this review using the following search terms: hydrofluoric acid, acid etching, surface conditioning, ceramics, fracture, repair, hydrofluoric acid burns, hydrofluoric acid treatment, hydrofluoric acid etching, workplace poisoning, poisoning, fatalities, cementation, ceramics, intra-oral repair, fluorohydric acid, fluoric acid, and hydrogen fluoride solution. Only articles in English were selected. Additional information was derived from scientific reports, medical and chemical textbooks, handbooks, product information, manufacturers' instructions, Internet web sites of the HF manufacturers.

Results

While no reports of incidence have been found on the hazardous effects due to exposure to HF in the dental literature, low incidence of reactions or even death presented as case reports have been reported in the case of significant or long exposure to HF in other fields. Potential hazards in case of significant exposure and the therapy approach to each of the symptom could be summarized as follows.

Acute hazardous effects

As with all inorganic acids, due to hydrogen ion degeneration and corrosion, HF can burn the skin [11–13]. After penetrating into the skin, it moves quickly to the deeper tissue layers and releases the freely dissociable fluoride ion. Toxicity of this ion is based on its strong reactivity. The mechanism of HF injury is thought to be binding of fluoride ions to tissue calcium and magnesium cations to form insoluble salt. These interfere with cellular metabolism, inducing cellular death and necrosis [4]. HF burns could be as a consequence of the dehydrating effect of this agent and the low pH. Skin contact with HF, even dilute solutions (0.1%) can cause painful second- and third-degree burns that heal very slowly [10]. Burns have a rather characteristic appearance and present as severely reddened, swollen areas with blanched, whitish regions that rapidly progress to blistering, and necrosis accompanied with pain [14]. Latent damages could also be observed since fluoride ion penetrates through all layers of

the epidermis, dermis, and even the deeper subcutaneous tissues, causing severe destruction, necrosis, and injuries to the underlying bone due to decalcification [15]. The effect can continue for several days, causing increased tissue damage at a later stage. The damage is directly related to the concentration of the acid and the duration of contact. HF burns are classified according to the acid concentrations of 0–20%, 21–50%, and greater than 50% [15–19]. Skin contact with hydrogen fluoride, or solutions containing more than 50% HF, produces immediate pain. Reactions to more dilute solutions (concentration 20–50%) may be delayed for minutes to several hours. The accompanying pain is excruciating and persistent, and healing is delayed [15–19]. Erythema and pain may be delayed up to 24 h in exposure to dilute HF solutions (<20%).

Different types of therapy options have been suggested for HF burns. The aim of all treatment methods is to chemically sequester the fluoride ion and to prevent extensive, deep tissue destruction [20]. The first treatment of chemical burns should always be thorough and copious lavage of the wound to clean the unreacted surface chemical and dilute that is already in contact with the skin [20]. The second phase of the treatment aims to inactivate the free fluoride ion by promoting the formation of an insoluble fluoride salt. It was reported that the calcium ion could decrease the toxic effects of sodium fluoride in tissue [20, 21]. For this reason, calcium gluconate gel (2.5%) is widely used for first aid and/or primary treatment of skin burns due to exposure to HF [22–30]. The gel is applied promptly and repeatedly onto the burned area, up to several hours. Using this gel may be more convenient for dilute acid burns that occur in exposure to commercial products such as rust removers, aluminum cleaners using etching solutions [28]. If good relief of pain is not obtained after 30–40 min, alternative methods of treatment such as calcium gluconate injections or benzalkonium chloride soaks should be considered. Calcium gluconate injection (5%) is indicated for large burns (over 160 cm²) or in cases where initial treatment has been delayed [31–34].

Burns of the fingers and nails permit penetration of fluoride ions but prevent gels from being effective. Therefore, it may be necessary to drill, split, or even remove nails to allow the topical treatment methods to be effective. However, it was reported that in less than 10% of the burns, removal of the nail was necessary [31–34]. If burned nail is immersed in benzalkonium chloride (Zephrian®) solution immediately, it may be possible to avoid removing the nail.

Early aggressive surgery with primary excision of necrotic skin areas has also been recommended as an emergency measure to prevent deep tissue destruction and systemic absorption of fluoride ions [35–41]. When painful indolent ulcer remains resistant to all conservative measures, surgery may be considered with the possibility of unacceptable cosmetic results.

Chronic hazardous effects

HF is not classified as a carcinogen agent since no data exists supporting this. However, increased rates of cancer have been observed in workers exposed to a mixture of chemicals that included fluoride, but no single chemical was identified and reported as the cause of the cancer [13]. Yet, possible effects on reproduction have been postulated [42]. Studies on women exposed to fluoride for occupational reasons showed increased rates of menstrual irregularities. Animal studies also showed that fluoride impairs reproduction and increases fetal bone and teeth malformation [43].

Systemic toxicity

In case of severe inhalation or ingestion of HF, systemic toxicity is of concern.

To produce HF, CaF₂ reacts with sulfuric acid:

This process releases energy and occurs very quickly. Although the pH value of HF is only 3.45, HF is soluble in water by a strong reaction at every concentration. Since it is very hygroscopic, it has the ability to dehydrate substances. Hence, it can quickly penetrate into skin and other soft tissues. When HF is exposed in concentrations of about 5% to human skin, initially, in most cases, no pain or other symptoms can be observed, but after several hours, a painful burn reaction may become apparent [44–48]. There is some evidence that when fluoride combines with calcium and phosphate, intracellular levels increase [2, 44]. Binding of fluoride to body calcium is one of the mechanisms of HF toxicity [47–52]. One serious consequence of exposure to HF is the decrease in serum calcium (hypocalcemia) and other metabolic changes, which may result in a fatal outcome if not recognized and treated [53–56]. Hypocalcemia should be considered as a potential risk in all instances of inhalation or ingestion, especially when skin burns exceed 160 cm². Fluoride also binds to potassium and magnesium ions leading to myocardial irritation and arrhythmia. Chronic exposure to HF may cause skeletal fluorosis that can be fatal [57–61]. Hemodialysis has been reported to be an effective therapy for cases of severe systemic intoxication [57–61].

Eye injuries

The eye is highly susceptible to HF liquid or vapor exposures. Contact of HF with eyes may result in permanent eye damage [58–60]. Ocular exposure may denude the cornea and conjunctivae epithelium and lead

to stromal corneal edema, conjunctivae ischemia, sloughing, and chemosis [58, 59]. Deeper penetration of the fluoride ion may cause corneal opacification and necrosis of the anterior chamber structures. Usually, the effects are noted within 3 days, although Hatai et al. described a case of delayed corneal damage, it was not apparent until 4 days after the exposure [59]. Latent complications included recurrent corneal epithelial erosions, keratoconjunctivitis sicca, progressive vascularization of the corneal stroma, and symblepharon [58]. As for cutaneous burns, the most important therapy is prompt and efficient lavage. This must be immediate and continued for at least 15–30 min using either water or normal saline [62–66].

Browne advised the use of a few drops of 10% calcium gluconate after irrigation [62]. Trevino et al. recommended 1% gluconate in saline as an irrigant followed by 1% calcium gluconate drops every 2–3 h for several days [63]. They have reported complete recovery using this treatment in 18 out of 19 patients, of whom 13 had completely opacified corneas prior to treatment [62]. Hatai et al. achieved quick, complete recovery of a large corneal erosion caused by 49% HF using a repeated instillation of 1% calcium gluconate drops combined with conventional treatment [59]. Although others also advised the use 1% calcium gluconate drops [41, 67], its use remains empirical. McCulley et al. compared irrigation with water, normal saline, magnesium chloride, lanthanum chloride, 0.2% Hyamine, 0.03% and 0.05% Zephiran as well as topical ointments of magnesium oxide and magnesium sulfate [68]. These solutions were also tried as a 1-ml subconjunctival injection. They showed that the only beneficial treatments were irrigation with water, isotonic saline, or isotonic magnesium chloride. Procedures involving repetitive irrigation with these agents caused increased rates of corneal ulceration. The same group has also studied the mechanism of eye injury by HF, demonstrating a clear dose-related response [68]. After irrigation, it is mandatory to request a prompt ophthalmological opinion. Instillation of cycloplegics to dilate the pupil may be required, and some advocate steroid drops to lessen fibroblast formation in the cornea [63, 69]. Rubinfeld et al. are the only group who reported the use of debridement of the sloughing necrotic conjunctiva in an attempt to reduce the fluoride level further and aid in healing [60]. However, speed of delivery of the irrigation remains the most important step among all trials [58].

Inhalation

Inhalation of low concentrations of HF can cause severe respiratory damage, irritation of the nose, sore throat, lungs, pulmonary edema, burning sensation, cough, dizziness, headache, nausea, shortness of breath, vomiting, and eye/skin irritation [70, 71]. Animal studies have also showed increased rates of kidney and liver damage as a result of HF

inhalation [42, 72–74]. Anhydrous HF is very volatile with a boiling point of only 19.5°C, and even the aqueous solutions (boiling points: 75% HF, 56°C; 60% HF, 89°C) produce a pungent odor. The recommended level of HF in the working atmosphere should not exceed 3 ppm which could be noticed by smelling [75].

A particular hazard, as noted by Mayer and Guelich, occurs when clothing is contaminated since this can produce high concentrations in the inspired air [70]. They reported the case of three workmen who died of pulmonary complications following splashes to their cotton clothing with 70% HF, all succumbing within 2 h of the incident. Dieffenbacher and Thompson also reported about a petroleum worker whose face and neck were in contact with 10% HF for only 30–35 s [76]. Despite intravenous and subcutaneous calcium gluconate application, he died 2 h later. Post-mortem findings showed thick mucus with blood in the bronchi, edematous lungs with hemorrhagic patches, and sub-serous hemorrhages in the pleurae. Green-dyke and Hodge described similar post-mortem appearances in two men who died following a serious airway exposure to HF of unspecified concentration [77]. High blood fluoride levels were also present in these patients. Chela and Huguet recorded the murder of a young woman who had thrown HF to her face [78]. She died from acute respiratory insufficiency 3 h after exposure.

Treatment of inhalational injuries begins with removing the patient immediately from the contaminated area and administering humidified 100% oxygen as soon as possible. In an attempt to neutralize the fluoride ion, some authorities recommend a nebulized solution of 2.5–3.0% calcium gluconate [41, 63]. If hemorrhage occurs, then bronchoscopy would be necessary to avoid lobe collapse, and transfusion may also be required [66]. Although corticosteroids have been tried, they should be carefully indicated [63, 79, 80].

Ingestion

The first recorded fatality from fluoride poisoning was in 1873 when King reported the case of a 35-year-old man who died within 35 min after drinking about 15 ml of HF [81]. This demonstrates the speed of absorption of the fluoride ion by the gastrointestinal tract. Since then, many cases of fluoride poisoning have been recorded due to ingestion [82–90]. Initial symptoms often include nausea, vomiting, and abdominal pain. Ingestion of even relatively dilute solutions of HF may be rapidly fatal.

Management of HF ingestion essentially remains as the dilution, removal, and neutralization of the acid. Stimulation to cause vomiting is not recommended since this may cause further oral, esophageal, or pulmonary injury. The patient should be encouraged to drink either water or preferably milk that has been shown to be effective in

precipitating fluoride ions in vitro [90]. At the hospital, gastric lavage can be considered and some advise the addition of 10% calcium gluconate to the lavage fluid [91].

Extreme exposure and death

HF at concentrations of 30 parts per million (ppm) is immediately dangerous to life [70, 71]. Fatalities have been reported from a skin exposure to as little as 2.5% of body surface area. Once it penetrates the skin, it slowly disassociates into the hydrogen and fluoride ion. The fluoride ion affects tissue integrity and metabolism by liquefaction necrosis, decalcification, destruction of bone, and production of insoluble salts. This will eventually result in loss of calcium from the bones. The development of hypocalcemia can be rapidly fatal because calcium is important for muscles, including cardiac muscle. Without calcium, many metabolic pathways are broken down.

Precautions to avoid HF exposure

Ventilation

Local exhaust ventilation is generally preferred because it can control the emissions of the contaminant at its source, preventing dispersion of it into the general working area [92]. For emergencies or instances where the exposure levels are not known, the use of a face-piece positive-pressure, air-supplied respirator would be useful. It was warned that air-purifying respirators do not protect workers in oxygen-deficient atmospheres [92].

Skin protection

Protective clothing with long sleeves, air-inflated suits with masks, goggles, including boots or safety shoes with polyvinyl chloride or neoprene should be worn [92].

Eye protection

Chemical safety goggles and/or full-face shields should be used where splashing is possible. Fountains and quick drench facilities should be maintained in the work area [92].

Summary of major acute and chronic symptoms observed after exposure to HF and the first aid measures are listed in Table 1.

HF use in dentistry and clinical manifestations

HF types and quantities

Concentrations of HF used in dentistry are usually small, ranging between 5% and 15%. They are available either in gel

or liquid form. It is indicated to be used both in vivo and ex vivo. While the majority of the HF manufacturers for dental use recommend its use with a rubber dam, others provide a gel product, the so-called dental dam to protect adjacent gingival and soft tissues from HF. In order to avoid the likelihood of HF exposure and to make it more visible, coloring dyes are added to make the product red, yellow, or green.

Metals

The application of HF (7.5% and 9.5%) for metals usually takes place in the dental laboratories primarily for stripping and cleaning metals. When a precious metal is cast with a ceramic insert to produce a cylindrical hole that will receive a pin, HF is used to dissolve the ceramic insert to the casting [93].

Ceramics

The most common use of HF in dentistry is for conditioning glassy-matrix ceramic surfaces to adhere resin-based materials either prior to cementation or during repairs [94–97]. HF selectively etches and dissolves the glass ceramic and cause physical alteration of the surface creating micromechanical retention. The ceramic is then rinsed with water spray. For cementation purposes, HF etched restoration is placed in neutralizing solution of CaCO_3 and/or NaOCl . HF gel is further removed in an ultrasonic bath. For intraoral repair applications, ultrasonic cleaning of HF is not possible since the restoration cannot be removed [97]. HF application is then followed by silane coupling agent. Acid concentrations and etching times vary depending on the specifications of the ceramics to receive optimized adhesion of the resin-based material.

Composites

Although some manufacturers claim that HF is effective for conditioning resin-based composites also for cementation and repair purposes, poor results were reported [98]. When composites are exposed to HF acid gel, water monolayer may penetrate via voids to fillers that, in turn, may disorganize the siloxane network formed from the condensation of intermolecular silanol groups, which is responsible for stabilizing the filler-resin interface. For this reason, the use of HF acid gel was not found to be a predictable option for conditioning composite materials for repair or relamination purposes.

Acidulated phosphate fluoride

The hazards, caustic effects to soft tissues and the danger for clinical use for hydrofluoric acids are well-known by

Table 1 Major symptoms observed after exposure to hydrofluoric acid and the first aid measures

Hazards	Possible symptoms	First aid ^a
Acute hazardous effects		
Skin burns	Redness, pain, blisters, swellings, necrosis accompanied with pain	Rinse skin with water, calcium gluconate gel (2.5%) application or calcium luconate injection, depending on the severity, remove affected skin
Nail injuries	Necrotic skin around the nail, indulent ulcer	Drilling, splitting, or removing nail, immersing the nail in benzalkonium chloride
Chronic hazardous effects		
Systemic toxicity	Painful burn, hypocalcaemia, myocardial problems, arrhythmia	Immediate transfer to a medical center, Haemodialysis
Eye injuries	Corneal eudema, conjunctivae ischemia, delayed corneal damage, keratoconjunctivitis sicca, vascularization of the corneal stroma	Efficient lavage (15–30 min) with water or isotonic saline followed by application of calcium gluconate drops, consulting an ophthalmologist
Inhalation	Irritation of the nose, sore throat, lungs, burning sensation, cough, dizziness, headache, nausea, shortness of breath, vomiting, eye, skin, pulmonary edema, possible kidney, liver damage, hypocalcemia, death	Immediate transfer to a medical center, removing the subject from the contaminated area, nebulized solution of calcium gluconate depending on the severity bronchoscopy, transfusion
Ingestion	Nausea, vomiting, and abdominal pain	Drink water, milk, gastric lavage

^a In all cases, a doctor should be consulted

the manufacturers and the researchers. For this reason, some studies questioned whether 1.23% acidulated phosphate fluoride gels might serve as a safe and effective substitute for etching ceramic surfaces to bond composite resin to silanized ceramics [99–101]. It was concluded that 1.23% acidulated phosphate fluoride gels can substitute 9.5% HF but prolonged etching times were required with the lower concentrations of HF [102]. However, the specimens etched with 4% acidulated phosphate fluoride gel, showed a wider statistical spread than the ones etched with 9.6% HF. This suggested that HF etching might well produce a more reliable and consistent result but this has not been confirmed, as the sample size of these studies was too small [100, 102].

Topical fluoride rinses and gels

Therapeutic fluoride products are also applied on teeth for preventive purposes especially in pediatric dentistry. However, HF is a minor ingredient in topical therapeutic fluoride. Necessary precautions should still be considered using a rubber dam during their application.

Concluding remarks

Since no incidence on hazardous effects of HF has been reported in the dental literature, this literature review led to the conclusion that risks of in vivo and ex vivo use of HF,

considering the concentrations used in dentistry appears to be unwarranted. Since repair actions are rather seldom, particularly the clinicians who indicate glass ceramic restorations would be more likely to be exposed to HF during cementation procedures. Concentrated solutions cause immediate pain and produce a surface burn similar to other common acids with erythematic responses, blistering, and necrosis [52]. HF concentrations of 14.5% produce symptoms immediately and concentrations of 12% may take up to an hour. Reaction to exposure of 7% HF or less, may take several hours before onset of symptoms, resulting in delayed symptoms with the deeper penetration of the HF. For occupational exposure, American Conference of Industrial Hygienists reported the lowest limits by giving a maximum value (a limit that should never be exceeded for even an “instantaneous” exposure period of 3 ppm; 2.3 mg/m³) [103]. This exposure limit (3 ppm) has also been given as 8 h time weighed average level by the Occupational Safety and Health Administration [103]. On the other hand, the National Institute of Occupational Safety and Health considered Dangerous To Life and Health value as 30 ppm [103]. Taking these values into account, for the dental professionals and patients, a risk analysis could be performed based on the frequency of the use of HF. Due to the continuous use of HF in dentistry, the patients should be monitored even after cementation or fixation of the restorations. Knowing potential hazards of HF from other

fields, media other than HF gel have certainly potential in the future for conditioning ceramic surfaces in dentistry.

Acknowledgments The authors thank to Assoc. Prof. Dr. H.G. Orhan, Ege University, School of Pharmacy, Department of Toxicology, Izmir, Turkey, for the helpful discussion and comments regarding to toxicity.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Litovitz TL, Klein-Schwartz W, Dyer KS, Shannon M, Lee S, Powers M (1998) 1997 Annual Report of the American Association of Poison Control Centers Toxic Exposure Surveillance System. *Am J Emerg Med* 16:443–497
- Litovitz TL, Klein-Schwartz W, Caravati EM, Youniss J, Crouch B, Lee S (1999) 1998 Annual Report of the American Association of Poison Control Centers Toxic Exposure Surveillance System. *Am J Emerg Med* 17:435–487
- McClure FJ (1933) Review of fluorine and its physiological effects. *Physiol Rev* 13:277–300
- Fact Sheet Environment: Health & Safety information for the Berkley Campus. Division of Industrial Hygiene, National Institute of Health (1943) Hydrofluoric acid burns. *Indust Med* 12:634–637
- Shewmake SW, Anderson BG (1979) Hydrofluoric acid burns. A report of a case and review of the literature. *Arch Dermatol* 115:593–596
- Calamia JR (1989) Clinical evaluation of etched porcelain veneers. *Am J Dent* 2:9–15
- Diaz-Arnold AM, Schneider RL, Aquilino SA (1989) Bond strengths of intraoral porcelain repair materials. *J Prosthet Dent* 61:305–309
- Özcan M, Pfeiffer P, Nergiz I (1998) A brief history and current status of metal/ceramic surface conditioning concepts for resin bonding in dentistry. *Quintessence Int* 29:713–724
- Caravati EM (1988) Acute hydrofluoric acid exposure. *Am J Emerg Med* 6:143–150
- Bertolini JC (1992) Hydrofluoric acid: a review of toxicity. *J Emerg Med* 10:163–168
- Harris JC, Rumack BH, Bregman DJ (1981) Comparative efficacy of injectable calcium and magnesium salts in the therapy of hydrofluoric acid burns. *Clin Toxicol* 18:1027–1032
- Dunn B, MacKinnon MA, Knowlden NF, Billmaier DJ, Derelanko MJ, Rusch GM (1996) Hydrofluoric acid dermal burns: an assessment of treatment efficacy using an experimental pig model. *J Occup Environ Med* 38:507–514
- Gallerani M, Bettoli V, Peron L, Manfredini R (1998) Systemic and topical effects of intradermal hydrofluoric acid. *Am J Emerg Med* 16:521–522
- Doyle I (1989) Treatment of hydrofluoric acid exposure. *J Occup Med* 30:52–63
- Klauder JV, Shelanski L, Gabriel K (1955) Industrial uses of compounds of fluorine and oxalic acids. *Arch Indust Health* 12:412–419
- Janzekovic Z (1970) A new concept in skin grafting of burns. *J Traumatol* 10:1103–1108
- Reynolds KE, Whitford GB, Pashley DH (1978) Acute fluoride toxicity: the influence of acid base status. *Toxicol App Pharmacol* 45:415–427
- McIvor ME (1990) Acute fluoride toxicity: pathophysiology and management. *Drug Saf* 5:79–85
- Whelanski RH (1990) Hydrofluoric acid burns of the skin. *J Hand Surg* 15:52–63
- Paterson JD (1956) Control of the fluorine hazard. *Brit J Indust Med* 16:301–303
- Henry JA, Hla KK (1992) Intravenous regional calcium gluconate perfusion for hydrofluoric acid burns. *J Toxicol Clin Toxicol* 30:203–207
- Jones AT (1939) The treatment of hydrofluoric acid burns. *J Ind Hyg Toxicol* 2:205–212
- Chick LR, Borah G (1990) Calcium carbonate gel therapy for hydrofluoric acid burns of the hand. *Plast Reconstr Surg* 86:935–940
- Henry JA, Hla KK (1992) Intravenous regional calcium gluconate perfusion for hydrofluoric acid burns. *Clin Toxicol* 30:203–207
- Burkhart KK, Brent J, Kirk MA, Baker DC, Kulig KW (1994) Hydrofluoric acid burns. *Ann Emerg Med* 23:110–121
- Cox RD, Osgood KA (1994) Evaluation of intravenous magnesium sulphate for the treatment of hydrofluoric acid burns. *J Toxicol Clin Toxicol* 32:123–136
- Graudins A, Burns MJ, Aaron CK (1997) Regional intravenous infusion of calcium gluconate for hydrofluoric acid burns of the upper extremity. *Ann Emerg Med* 30:604–607
- Yasuda H, Honda S, Yamamoto O, Asahi M (1999) Therapeutic effect of topical calcium gluconate for hydrofluoric acid burn-time limit for the start of treatment. *J Occup Med* 21:209–216
- Aguilera IM, Vaughan RS (2000) Calcium and the anaesthetist. *Anaesthesia* 55:779–790
- Yamashita M, Suzuki M, Hirai H, Kajigaya H (2001) Iontophoretic delivery of calcium for experimental hydrofluoric acid burns. *Crit Care Med* 29:1575–1578
- Vance MV, Curry SC, Kunkell DB, Ryan PJ, Ruggeri SB (1986) Digital hydrofluoric acid burns: treatment with intra-arterial calcium infusion. *Ann Emerg Med* 15:890–896
- Roberts JR, Merigian KS (1988) Acute hydrofluoric acid exposure. *Am J Emerg Med* 7:125–126
- Burkhart KK, Brent J, Kirk MA, Baker DC, Kulig KW (1995) Comparison of topical magnesium and calcium treatment for dermal hydrofluoric acid burns. *Ann Emerg Med* 21:9–13
- Aaron CK, Burns MJ, Graudins A (1997) Regional intravenous infusion of calcium gluconate for hydrofluoric acid burns of the upper extremity. *Ann Emerg Med* 30:604–607
- Dale RH (1951) Treatment of hydrofluoric acid burns. *Brit Med J* 1:728–732
- Craig RDP (1964) Hydrofluoric acid burns. *Brit J Plas Surg* 17:52–59
- Schwamweber HC (1969) Treatment of hydrofluoric acid burns of the skin by immediate surgical excision. *Indust Med* 38:31–32
- Tschantz P, Favre H (1975) Necrotizing pancreatitis after formic acid and hydrofluoric acid burns. *Schweiz Rundsch Med Prax* 64:487–490
- Kohnlein HE, Achinger R (1982) A new method of treatment of hydrofluoric acid burns of the extremities. *Chirurgie Plast* 6:297–305
- Velvart J (1983) Arterial perfusion for hydrofluoric acid burns. *Human Exposure Toxicol* 2:233–238
- MacKinnon MA (1986) Treatment of hydrofluoric acid burns. *J Occup Med* 28:804
- Wing JS, Brender JD, Sanderson LM, Perrotta DM, Beauchamp RA (1991) Acute health effects in a community after a release of hydrofluoric acid. *Arch Environ Health* 46:155–160
- Morris JB, Smith JA (1982) Regional deposition and absorption of inhaled hydrogen fluoride in the rat. *Toxicol App Pharmacol* 62:81–89
- Lewis GK (1959) Chemical burns. *Am J Surg* 98:928–937

45. Hamilton M (1975) Hydrofluoric acid burns. *Occup Health* 27:468–470
46. Edelman P (1986) Hydrofluoric acid burns. *Occup Med* 1:89–103
47. Chan KM, Svancarek WP, Creer M (1987) Fatality due to acute hydrofluoric acid exposure. *J Clin Toxicol* 25:333–339
48. Wilkes GJ (1993) Intravenous regional calcium gluconate for hydrofluoric acid burns of the digits. *Emer Med* 5:155–158
49. McIvor ME, Cummings CE, Mower MM (1987) Sudden cardiac death from acute fluoride intoxication: the role of potassium. *Ann Emer Med* 16:777–781
50. Mullett T, Zoeller T, Bingham H, Pepine CJ, Prida CX, Castenholz R, Kirby R (1987) Fatal hydrofluoric acid cutaneous exposure with refractory ventricular fibrillation. *J Burn Care Rehabil* 2:216–219
51. Muriale L, Lee E, Genovese J, Trend S (1996) Fatality due to acute fluoride poisoning following dermal contact with hydrofluoric acid in a palynology laboratory. *Ann Occup Hyg* 40:705–710
52. Mayer TG (1998) Hydrofluoric acid burns. *Am J Emer Med* 32:116–127
53. Waldbott GL (1963) Acute fluoride intoxication. *Acta Med Scand* 174:5–44
54. Tepperman PB (1980) Fatality due to acute systemic fluoride poisoning following a hydrofluoric acid skin burn. *J Occup Med* 22:691–692
55. Mayer TG, Gross PL (1985) Fatal systemic fluorosis due to hydrofluoric acid burns. *Am J Emer Med* 14:149–153
56. Chela A, Reig R, Sanz P, Huguet E, Corbella J (1989) Death due to hydrofluoric acid. *Am J For Med Pathol* 10:47–48
57. Watson AA, Oliver JS, Thorpe JA (1973) Accidental death due to inhalation of hydrofluoric acid. *Med Sci Law* 13:277–279
58. McCulley JP, Whiting DW, Pettitt MG, Lauber SE (1983) Hydrofluoric acid burns of the eye. *J Occup Med* 25:447–450
59. Hatai JK, Weber JN, Doizaki K (1986) Hydrofluoric acid burns of the eye: report of possible delayed toxicity. *J Toxicol Cut* 5:179–184
60. Rubinfeld RS, Silbert DI, Arentsen JJ, Laibson PR (1992) Ocular hydrofluoric acid burns. *Am J Ophthalmol* 114:420–423
61. Chan BS, Duggin GG (1997) Survival after massive hydrofluoric acid ingestion. *J Toxicol Clin Toxicol* 35:307–310
62. Browne TD (1974) The treatment of hydrofluoric acid burns. *J Occup Med* 24:80–89
63. Trevino MA, Herrmann GH, Sprout WI (1983) Treatment of severe hydrofluoric acid exposures. *J Occup Med* 25:861–863
64. MacKinnon MA (1988) Hydrofluoric acid burns. *Dermatol Clin* 6:67–74
65. Zoeller T, Bingham H, Pepine CJ (1989) Hydrofluoric acid burns. *J Burn Care Rehabil* 35:16–19
66. Upfal M, Doyle C (1990) Medical management of hydrofluoric acid exposure. *J Occup Med* 32:726–731
67. Pegg SP, Siu S, Gillett G (1985) Intra-arterial infusions in the treatment of hydrofluoric acid burns. *Burns* 2:440–443
68. McCulley JP (1990) Ocular hydrofluoric acid burns: animal model, mechanism of injury and therapy. *Trans Am Ophthalmol Soc* 88:649–684
69. Vaughn D, Asbury T (1983) General Ophthalmology. Los Altos, California: Lange 39.
70. Mayer L, Guelich J (1963) Hydrogen fluoride inhalation and burns. *Arch Environ Health* 7:445–447
71. Vance MV, Curry SC, Kunkel DB (1983) Treatment of hydrofluoric acid burns. *Emer Med* 17:15–56
72. Morris JB, Smith FA (1978) Regional deposition and absorption of inhaled hydrogen fluoride. *Toxicol App Pharmacol* 45:415–427
73. Braun J, Stoss H, Zober A (1984) Intoxication following inhalation of hydrogen fluoride. *Arch Toxicol* 56:50–54
74. Kono K, Yoshida Y, Watanabe M (1992) Serum fluoride as an indicator of occupational hydrofluoric acid exposure. *Inf Arch Occup Environ Health* 64:343–346
75. Sevvitt S (1957) Burns: pathology and therapeutic applications. Butterworth, London, pp 330–333
76. Dieffenbacher PF, Thompson JH (1996) Burns from exposure to anhydrous hydrofluoric acid. *J Occup Med* 4:325–326
77. Greendyke RM, Hodge HC (1964) Accidental death due to hydrofluoric acid. *J Foren Sci* 9:383–390
78. Chela A, Huguet E (1987) Hydrofluoric acid burns. *Am J Foren Med* 5:87–110
79. Iverson RE, Laub DR, Madison MS (1988) A radical approach to severe hydrofluoric acid burns. A case report. *J Occup Med* 30:873–874
80. Foster DE, Barone JA (1989) Rectal hydrofluoric acid exposure. *Clin Pharmacol* 8:516–518
81. King R (1973) Poisoning by hydrofluoric acid. *Trans Pathol Soc* 24:98–100
82. McNally WD (1923) Four deaths caused by sodium fluoride. *J Am Med Assoc* 23:811–813
83. Fullerton WW (1930) Two rather uncommon fatal cases of poisoning. *N Eng J Med* 203:423
84. Maletz L (1935) Report of a fatal case of fluoride poisoning. *N Eng J Med* 213:370–372
85. Greenwood DA (1940) Fluoride intoxication. *Physiol Rev* 20:582–616
86. Lidbeck WL, Hill IB, Beeman JA (1943) Acute sodium fluoride poisoning. *J Am Med Assoc* 121:826–827
87. Rabinowitch IM (1945) Acute fluoride poisoning. *Canad Med Assoc* 52:345–349
88. Waldbott GL (1959) Urticaria due to fluoride. *Acta Allerg* 13:456–468
89. Siedlecki JT (1965) Hydrofluoric acid burns. Questions and answers. *J Am Med Assoc* 191:353
90. Yolken R, Konecny P, McCarthy P (1976) Acute fluoride poisoning. *Paediatrics* 58:90–93
91. Satremski ES, Grande GA, Ling LJ (1992) Survival following hydrofluoric acid ingestion. *Am Emer Med* 21:1396–1399
92. Catton V (1983) Prevention and treatment of hydrofluoric acid burns. *Occup Health* 35:227–231
93. Wood DJ, Bubbs NL, Millar BJ, Dunne SM (1997) Preliminary investigation of a novel retentive system for hydrofluoric acid etch-resistant dental ceramics. *J Prosthet Dent* 78:275–280
94. Calamia JR (1985) Etched porcelain veneers: the current state of the art. *Quintessence Int* 1:5–12
95. Stangel I, Nathanson D, Hsu CS (1987) Shear strength of the composite bond to etched porcelain. *J Dent Res* 66:1460–1465
96. Thurmond JW, Barkmeier W, Wilwerding TM (1994) Effect of porcelain surface treatments on bond strengths of composite resin bonded to porcelain. *J Prosthet Dent* 72:355–359
97. Özcan M (2003) Evaluation of alternative intraoral repair techniques for fractured ceramic-fused-to-metal restorations. *J Oral Rehabil* 30:194–203
98. Özcan M, Alander P, Vallittu PK, Huysmans MC, Kalk W (2005) Effect of three surface conditioning methods to improve bond strength of particulate filler resin composites. *J Mater Sci Mater Med* 16:21–27

99. Sposetti V, Shen C, Levin A (1986) The effect of topical fluoride application on porcelain restorations. *J Prosthet Dent* 55:677–686
100. Lacy AM, Laluz J, Watanabe LG, Dellinges M (1988) Effect of porcelain treatments on the bond strength of composite. *J Prosthet Dent* 60:288–292
101. Della Bona A, Van Noort R, Sear VS (1995) Tensile bond strength of resin composite bonded to ceramic. *J Dent Res* 74:1591–1596
102. Brentel AS, Özcan M, Valandro LF, Alarça LG, Amaral R, Bottino MA (2007) Microtensile bond strength of a resin cement to a feldspathic ceramic after different etching and silanization regimens in dry and aged conditions. *Dent Mater* 23:1323–1331
103. Threshold Limit Values (TLVs) for Chemical Substances and Physical Agents and Biological Exposure Indices (BEIs®), American Conference of Governmental Industrial Hygienists, Inc., p 42. Cincinnati, Ohio, 2005. <http://www.osha.gov> 2005