



University of  
Zurich<sup>UZH</sup>

Zurich Open Repository and  
Archive

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **Inflammation marker and risk of pancreatic cancer: a nested case-control study within the EPIC cohort**

Grote, V A ; Kaaks, R ; Nieters, A ; Tjønneland, A ; Halkjær, J ; Overvad, K ; Skjelbo Nielsen, M R ;  
Boutron-Ruault, M C ; Clavel-Chapelon, F ; Racine, A ; Teucher, B ; Becker, S ; Pischon, T ; Boeing, H ;  
Trichopoulou, A ; Cassapa, C ; Stratigakou, V ; Palli, D ; Krogh, V ; Tumino, R ; Vineis, P ; Panico, S ; Rodríguez,  
L ; Duell, E J ; Sánchez, M J ; Dorronsoro, M ; Navarro, C ; Gurrea, A B ; Siersema, P D ; Peeters, P H M ; Ye, W ;  
Sund, M ; Lindkvist, B ; Johansen, D ; Khaw, K T ; Wareham, N ; Allen, N E ; Travis, R C ; Fedirko, V ; Jenab, M ;  
Michaud, D S ; Chuang, S C ; Romaguera, D ; Bueno-de-Mesquita, H B ; Rohrmann, S

DOI: <https://doi.org/10.1038/bjc.2012.172>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-69751>

Journal Article

Accepted Version

Originally published at:

Grote, V A ; Kaaks, R ; Nieters, A ; Tjønneland, A ; Halkjær, J ; Overvad, K ; Skjelbo Nielsen, M R ; Boutron-Ruault, M C ; Clavel-Chapelon, F ; Racine, A ; Teucher, B ; Becker, S ; Pischon, T ; Boeing, H ; Trichopoulou, A ; Cassapa, C ; Stratigakou, V ; Palli, D ; Krogh, V ; Tumino, R ; Vineis, P ; Panico, S ; Rodríguez, L ; Duell, E J ; Sánchez, M J ; Dorronsoro, M ; Navarro, C ; Gurrea, A B ; Siersema, P D ; Peeters, P H M ; Ye, W ; Sund, M ; Lindkvist, B ; Johansen, D ; Khaw, K T ; Wareham, N ; Allen, N E ; Travis, R C ; Fedirko, V ; Jenab, M ; Michaud, D S ; Chuang, S C ; Romaguera, D ; Bueno-de-Mesquita, H B ; Rohrmann, S (2012). Inflammation marker and risk of pancreatic cancer: a nested case-control study within the EPIC cohort. *British Journal of Cancer*, 106(11):1866-1874.

DOI: <https://doi.org/10.1038/bjc.2012.172>

1 **Inflammation marker and risk of pancreatic cancer: a nested case-control**  
2 **study within the EPIC cohort**

3

4 Running title: Inflammation and pancreatic cancer risk

5

6 Verena A Grote<sup>1</sup>, Rudolf Kaaks<sup>1\*</sup>, Alexandra Nieters<sup>2,1</sup>, Anne Tjønneland<sup>3</sup>, Jytte  
7 Halkjær<sup>3</sup>, Kim Overvad<sup>4</sup>, Michael R. Skjelbo Nielsen<sup>5</sup>, Marie C Boutron-Ruault<sup>6,7</sup>,  
8 Françoise Clavel-Chapelon<sup>6,7</sup>, Antoine Racine<sup>6,7</sup>, Birgit Teucher<sup>1</sup>, Susen Becker<sup>1,8</sup>,  
9 Tobias Pischon<sup>9</sup>, Heiner Boeing<sup>10</sup>, Antonia Trichopoulou<sup>11,12</sup>, Christina Cassapa<sup>12</sup>,  
10 Vasiliki Stratigakou<sup>12</sup>, Domenico Palli<sup>13</sup>, Vittorio Krogh<sup>14</sup>, Rosario Tumino<sup>15</sup>, Paolo  
11 Vineis<sup>16,17</sup>, Salvatore Panico<sup>18</sup>, Laudina Rodríguez<sup>19</sup>, Eric J Duell<sup>20,21</sup>, María-José

---

<sup>1</sup> Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Im Neuenheimer Feld 581, 69120 Heidelberg, Germany

<sup>2</sup> Centre of Chronic Immunodeficiency, CCI Molecular Epidemiology, University Medical Center Freiburg, Breisacher St. 117, 79106 Freiburg, Germany

<sup>3</sup> Institute of Cancer Epidemiology, Danish Cancer Society, Strandboulevarden 49, 2100 Copenhagen, Denmark

<sup>4</sup> Department of Epidemiology, School of Public Health, Aarhus University, Nordre Ringgade 1, 8000 Aarhus, Denmark

<sup>5</sup> Center for Cardiovascular Research, Aarhus University Hospital, Hobrovej 18, 9100 Aalborg, Denmark

<sup>6</sup> INSERM, Centre for Research in Epidemiology and Population Health, U1018, Institut Gustave Roussy, 114 rue Edouard Vaillant, 94805 Villejuif, France

<sup>7</sup> Paris South University, UMRS 1018, Villejuif, France

<sup>8</sup> Institute of Laboratory Medicine, Clinical Chemistry and Molecular Diagnostics, University Hospital Leipzig, Paul-List-Str. 13-15, 4103 Leipzig, Germany

<sup>9</sup> Molecular Epidemiology Group, Max Delbrück Center for Molecular Medicine (MDC), Robert-Rössle-Straße 10, 13125 Berlin-Buch, Germany

<sup>10</sup> Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Arthur-Scheunert-Allee 114/116, 14558 Nuthetal, Germany

<sup>11</sup> WHO Collaborating Center for Food and Nutrition Policies, Department of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Athens, Greece

<sup>12</sup> Hellenic Health Foundation, 10-12 Tetrapoleos Street, 115 27 Athens, Greece

<sup>13</sup> Molecular and Nutritional Epidemiology Unit, Cancer Research and Prevention Institute - ISPO, Ponte Nuovo, Via delle Oblate n.2, 50141 Florence, Italy

<sup>14</sup> Nutritional Epidemiology Unit, Fondazione IRCCS Istituto Nazionale Tumori, Via Venezian, 1, 20133 Milano, Italy

<sup>15</sup> Cancer Registry and Histopathology Unit, "Civile M.P.Arezzo" Hospital, Via Civile, 97100 Ragusa, Italy

<sup>16</sup> School of Public Health, Imperial College London, Norfolk Place, W2 1PG London, UK

<sup>17</sup> HuGeF Foundation, Torino, Italy

1 Sánchez<sup>21,22</sup>, Miren Dorronsoro<sup>21,23</sup>, Carmen Navarro<sup>21,24</sup>, Aurelio Barricarte  
2 Gurrea<sup>21,25</sup>, Peter D. Siersema<sup>26</sup>, Petra HM Peeters<sup>27,28</sup>, Weimin Ye<sup>29,30</sup>, Malin Sund<sup>31</sup>,  
3 Björn Lindkvist<sup>32</sup>, Dorthe Johansen<sup>33</sup>, Kay-Tee Khaw<sup>34</sup>, Nick Wareham<sup>35</sup>, Naomi E  
4 Allen<sup>36</sup>, Ruth C Travis<sup>36</sup>, Veronika Fedirko<sup>37</sup>, Mazda Jenab<sup>37</sup>, Dominique S

---

<sup>18</sup> Department of Clinical and Experimental Medicine, Federico II University, Corso Umberto I, 80138 Naples, Italy

<sup>19</sup> Public Health and Participation Directorate, Health and Health Care Services Council, Ciriaco Miguel Vigil 9, 33009 Asturias, Spain

<sup>20</sup> Unit of Nutrition, Environment and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology (ICO-IDIBELL), L'Hospitalet de Llobregat, Avda Gran Via 199-203, 8907 Barcelona, Spain

<sup>21</sup> CIBER Epidemiología y Salud Pública (CIBERESP), Spain

<sup>22</sup> Andalusian School of Public Health, Cuesta del Observatorio 4, 18011 Granada, Spain

<sup>23</sup> Public Health Division of Gipuzkoa, Basque Regional Health Department, Avda. Navarra, 4, 20013 San Sebastian, Spain

<sup>24</sup> Department of Epidemiology, Murcia Regional Health Authority, Ronda de Levante, 11, 30008 Murcia, Spain

<sup>25</sup> Navarre Public Health Institute, Polígono de Landaben C/ F, 31012 Pamplona, Spain

<sup>26</sup> Department of Gastroenterology and Hepatology, University Medical Centre Utrecht (UMCU), Heidelberglaan 100, 3584 CX Utrecht, The Netherlands

<sup>27</sup> Julius Center, University Medical Center Utrecht, Huispost Str. 6.131, 3508 GA Utrecht, The Netherlands

<sup>28</sup> Department of Epidemiology and Biostatistics, School of Public Health, Faculty of Medicine, Imperial College, London, UK

<sup>29</sup> Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Box 281, 171 77 Stockholm, Sweden

<sup>30</sup> The Medical Biobank at Umeå University, Umeå, Sweden

<sup>31</sup> Departments of Surgical and Perioperative Sciences, Surgery and Public Health and Clinical Medicine, Nutrition Research, Umeå University, 1A, 9 tr, Kirurgcentrum, 952, 901 85 Umeå, Sweden

<sup>32</sup> Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Box 100, 205 02 Gothenburg, Sweden

<sup>33</sup> Department of Surgery, Skåne University Hospital, Lund University, Box 117, 205 02 Malmö, Sweden

<sup>34</sup> Department of Public Health and Primary Care, University of Cambridge, Robinson Way, CB2 0SR Cambridge, UK

<sup>35</sup> MRC Epidemiology Unit, Hills Road, CB2 0QQ Cambridge, UK

<sup>36</sup> Cancer Epidemiology Unit, Nuffield Department of Clinical Medicine, University of Oxford, Old Road Campus, OX3 7LF Oxford, UK

<sup>37</sup> Nutritional Epidemiology Group, International Agency for Research on Cancer (IARC-WHO), 150 Cours Albert Thomas, 69372 Lyon, France

1 Michaud<sup>38,39</sup>, Shu-Chun Chuang<sup>38</sup>, Dora Romaguera<sup>38</sup>, H Bas Bueno-de-Mesquita<sup>26,40</sup>,

2 Sabine Rohrmann<sup>1,41</sup>

3

4 **\*Correspondence to:**

5 Rudolf Kaaks

6 Division of Cancer Epidemiology c020

7 German Cancer Research Center (DKFZ)

8 Im Neuenheimer Feld 581

9 69120 Heidelberg

10 Germany

11 Email: [r.kaaks@dkfz.de](mailto:r.kaaks@dkfz.de)

12

13

14 Word count abstract: 247

15 Word count main document: 3917

16

---

<sup>38</sup> School of Public Health, Imperial College, Norfolk Place, W2 1PG London, United Kingdom

<sup>39</sup> Division of Biology and Medicine, Brown University, Providence, RI, USA

<sup>40</sup> National Institute for Public Health and the Environment (RIVM), PO Box 1, 3720 BA Bilthoven, The Netherlands

<sup>41</sup> Division of Cancer Epidemiology and Prevention, Institute of Social and Preventive Medicine, Hirschengraben 84, 8001 Zurich, Switzerland

1 **Abstract (247 words)**

2

3 **Background:** Established risk factors for pancreatic cancer include smoking, long-  
4 standing diabetes, high body fatness, and chronic pancreatitis, all of which can be  
5 characterized by aspects of inflammatory processes. However, prospective studies  
6 investigating the relation between inflammatory markers and pancreatic cancer risk are  
7 scarce.

8 **Methods:** We conducted a nested case-control study within the European Prospective  
9 Investigation into Cancer and Nutrition (EPIC), measuring pre-diagnostic blood levels  
10 of C-reactive protein (CRP), interleukin 6 (IL-6) and soluble receptors of tumour  
11 necrosis factor  $\alpha$  (sTNF-R1, R2) in 455 pancreatic cancer cases and 455 matched  
12 controls. Odds ratios (OR) were estimated using conditional logistic regression models.

13 **Results:** None of the inflammatory markers were significantly associated with risk of  
14 pancreatic cancer overall, although a borderline significant association was observed  
15 for higher circulating sTNF-R2 (crude OR = 1.52 [95% CI 0.97 – 2.39], highest vs.  
16 lowest quartile). In women, however, higher sTNF-R1 levels were significantly  
17 associated with risk of pancreatic cancer (crude OR = 1.97 [95% CI 1.02-3.79]). For  
18 sTNF-R2, risk associations seemed to be stronger for diabetic individuals and those  
19 with a higher BMI.

20 **Conclusion:** Prospectively, CRP and IL-6 do not seem to play a role in our study with  
21 respect to risk of pancreatic cancer, whereas sTNF-R1 seemed to be a risk factor in  
22 women and sTNF-R2 might be a mediator in the risk relationship between overweight  
23 and diabetes with pancreatic cancer. Further large prospective studies are needed to  
24 clarify the role of pro-inflammatory proteins and cytokines in the pathogenesis of  
25 exocrine pancreatic cancer..

26

1 Key words: inflammation, pancreatic cancer, EPIC, CRP, IL-6, TNF receptor

2

## 1 **Introduction**

2 Evidence is accumulating that systemic low-grade chronic inflammation in  
3 addition to local inflammation in the pancreas is involved in the pathogenesis of  
4 pancreatic cancer (Farrow & Evers, 2002; McKay *et al*, 2008; Whitcomb, 2004).  
5 Research findings pointing to this direction include the documented relationship of  
6 pancreatic cancer risk with chronic pancreatitis (Raimondi *et al*, 2010), as well as with  
7 smoking (Lynch *et al*, 2009; Vrieling *et al*, 2010), pre-existing and long-standing  
8 diabetes mellitus (Huxley *et al*, 2005), and excess weight (Genkinger *et al*, 2010), all of  
9 which are known or suggestive determinants of low-grade inflammatory states  
10 (Goncalves *et al*, 2011; Hotamisligil, 2006; Kolb & Mandrup-Poulsen, 2005; Whitcomb,  
11 2004).

12 Even though the mechanisms by which chronic inflammation leads to  
13 carcinogenesis are not fully understood, it is generally accepted that inflammation  
14 results in repeated DNA damage and in the accumulation of genetic defects (McKay *et*  
15 *al*, 2008). However, pro-inflammatory cytokines and growth factors are also released in  
16 response to the tumour, making it difficult to distinguish between cause and effect in  
17 the inflammatory processes (McKay *et al*, 2008).

18 Circulating C-reactive protein (CRP) concentration, an acute-phase protein  
19 produced in the liver, is increased in pancreatic cancer patients (Barber *et al*, 1999;  
20 Moses *et al*, 2009; Mroczko *et al*, 2010), most likely as part of the systemic  
21 inflammatory response to the tumour. Interleukin-6 (IL-6) and tumour necrosis factor  
22 alpha (TNF- $\alpha$ ) are up-regulating factors of CRP and have also been shown to be  
23 increased in pancreatic cancer patients (Barber *et al*, 1999; Ebrahimi *et al*, 2004;  
24 Moses *et al*, 2009; Mroczko *et al*, 2010; Talar-Wojnarowska *et al*, 2009). Prospectively,  
25 increased levels of CRP have inconsistently been associated with pancreatic cancer  
26 risk. To our knowledge, prospective studies on the association of IL-6, TNF- $\alpha$ , or its  
27 receptors with risk of pancreatic cancer are lacking.

1 We measured prediagnostic concentrations of CRP, IL-6, and soluble TNF receptors  
2 (sTNF-R1 and R2) in blood samples of 455 primary exocrine pancreatic cancer cases  
3 and 455 individually matched controls within the Prospective Investigation into Cancer  
4 and Nutrition (EPIC) as possible reflections of either pancreatic cancer or a metabolic  
5 risk factor potentially increasing pancreatic cancer risk by aggravating pancreatic  
6 inflammatory disease.

7

## 8 **Materials and Methods**

### 9 *Study Population*

10 The European Prospective Investigation into Cancer and Nutrition (EPIC) is a  
11 large cohort study conducted in 23 centres in ten European countries (Denmark,  
12 France, Germany, Greece, Italy, the Netherlands, Spain, Sweden, and the United  
13 Kingdom). Detailed descriptions of study design, population, and baseline data  
14 collection of the cohort can be found elsewhere (Haftenberger *et al*, 2002; Riboli *et al*,  
15 2002). Briefly, about 370 000 women and 150 000 men were enrolled between 1992  
16 and 2000. Participants provided information on dietary habits and lifestyle factors, and  
17 in addition, weight, height, and waist and hip circumferences were measured at  
18 baseline. Each participant provided informed consent, and the local ethical review  
19 committees approved the EPIC cohort study as well as the current project.

20

### 21 *Blood Sample Collection and Storage*

22 In the seven EPIC core countries (France, Germany, Greece, Italy, the  
23 Netherlands, Spain, and the United Kingdom), blood samples were collected at  
24 baseline, based on a standardized protocol and aliquoted in plastic straws (plasma,  
25 serum, erythrocytes, and buffy coat for DNA). The aliquoted specimens were then  
26 stored in a central biorepository in liquid nitrogen (-196°C). In Sweden, all samples  
27 were stored locally in freezers at -70°C and in Denmark in nitrogen vapour (-150°C). In



1 the present study, Norway was excluded because blood samples were only recently  
2 collected and very few pancreatic cancer cases have been diagnosed after blood  
3 donation.

4

#### 5 *Follow-up for Cancer Incidence and Vital Status*

6 In six of the participating countries (Denmark, Italy, the Netherlands, Spain,  
7 Sweden, and the United Kingdom), follow-up of cancer cases was based on population  
8 registries. In the other three countries (France, Germany, and Greece), a combination  
9 of methods was used including health insurance records, cancer and pathology  
10 registries, and active follow-up through study subjects and their next-of-kin. In all EPIC  
11 centres, data on vital status is collected from mortality registries at the regional or  
12 national level, which is combined with health insurance data (France) or data collected  
13 by active follow-up (Greece). Cases reported in this study were all diagnosed up to the  
14 latest dates of complete follow-up, which was between December 2002 and 2005,  
15 depending on the study centre. For Germany, Greece, and France, the end of follow-up  
16 was the last known contact, date of diagnosis, or date of death, whichever came first.

17

#### 18 *Selection of Case and Control Subjects*

19 Up to December 2006, follow-up has led to the identification of 578 incident  
20 cases of non-endocrine pancreas cancer that were coded according to ICD-10 (C25.0-  
21 25.3, 25.7-25.9), and for 455 of these cases blood specimens were available.  
22 Exclusion criteria were occurrence of other malignant tumours preceding the diagnosis  
23 of pancreatic cancer, except for non-melanoma skin cancer. Of the 455 cases, 334  
24 (76%) were microscopically confirmed and the remaining 24% were diagnosed by  
25 imaging results, physical examination, or clinical symptoms. Most tumours occurred in  
26 the head of the pancreas (42%), followed by body (7%) and tail (5%), while the rest of  
27 the tumours were of unknown localization. For each case, one control subject was

1 selected, that was alive and free of cancer at the time the index case was diagnosed,  
2 using an incidence density sampling procedure. All identified cases were matched with  
3 one control by centre, sex, age at blood collection (+/- 3 years), date of blood donation  
4 (+/- 3 months), time of blood donation (+/- 2 hours), fasting status (<3 hours, 3-6 hours,  
5 >6 hours after last meal) and use of hormones (oral contraceptive pill, hormone or  
6 oestrogen replacement therapy).

7

#### 8 *Laboratory Assays*

9 Plasma (in Scandinavian samples) and serum concentrations of CRP were  
10 measured by multiplex immunoassays using the Fluorokine MAP Obesity Base Kit  
11 (R&D Systems, Inc., Minneapolis, MN, USA). IL-6 and sTNF receptors were measured  
12 by enzyme linked immune sorbent assays (ELISA) using the Quantikine kit (R&D  
13 Systems, Inc., Minneapolis, MN, USA). The total amount of free receptor plus the total  
14 amount of receptor bound to TNF is measured with this method. All measurements  
15 were performed in our specialized immunoassay laboratory of the Division of Cancer  
16 Epidemiology (German Cancer Research Center, Heidelberg, Germany). Samples of  
17 cases and matched controls were analyzed within the same analytical batch. Intra-  
18 batch and inter-batch coefficients of variation were 6.6 and 10.8% for IL-6, 3.6 and  
19 4.1% for sTNF-R1, 5.5 and 11.0% for sTNF-R2, and 10.3 and 11.6% for CRP. Units of  
20 IL-6 are expressed as pg/mL, of sTNF receptors as ng/mL, and of CRP as mg/L. One  
21 batch during the sTNF-R2 measurements did not perform well and, therefore, 70  
22 subjects were excluded due to technically invalid results (all from Malmo, Sweden).

23

#### 24 *Statistical Analysis*

25 Case and control differences across baseline characteristics were assessed by  
26 paired t-tests (continuous variables) or by generalized McNemar's Test (categorical  
27 variables). Spearman's partial rank correlation coefficients [r] adjusted for age, sex, and

1 EPIC recruitment centre were used to assess the strength of associations between  
2 waist circumference, waist-hip-ratio, BMI, glycated haemoglobin (HbA1c), and  
3 inflammatory markers, as well as for the correlation between the inflammatory markers.

4 Odds ratios (OR) and corresponding 95% confidence intervals (CI) for pancreatic  
5 cancer at different serum levels of IL-6, sTNF receptors, and CRP were calculated by  
6 conditional logistic regression models, using the exposure assessments of the matched  
7 case-control sets. Continuous measurements of the inflammatory markers were log<sub>2</sub>  
8 transformed to achieve approximate normality. In this scale, a unit increase  
9 corresponds to a doubling of concentration. Quartile cut-points were based on the  
10 distribution of biomarkers among controls. Sex-specific quartile cut-points had a  
11 negligible effect on risk estimates and were, therefore, not applied. Modelling the  
12 median within each quartile as a continuous variable was used to assess linear trends  
13 in ORs. Testing the model fit for categorical vs. continuous models resulted in very  
14 similar AICs, with a slightly better fit for the latter model.

15 Inflammatory markers may be downstream in the causal chain of excess body  
16 weight, smoking, or diabetes and pancreatic cancer. Alternatively, other pathways  
17 might explain associations of these conditions with risk of pancreatic cancer and,  
18 hence, inflammatory markers may be independently related to cancer or not at all. We  
19 tried to elucidate these rather complex and yet unknown relationships in our study by  
20 applying different adjustment models and by performing several subgroups analyses.  
21 All these models and methods are of exploratory nature in our study.

22 Potential confounding of factors other than those controlled for by matching were  
23 examined by assessing the association of these factors with pancreatic cancer risk  
24 using unconditional logistic regression models adjusted for matching factors, by  
25 correlation analyses, and by including these as additional factors in conditional logistic  
26 regression models. BMI, waist-hip-ratio, waist circumference, alcohol consumption,  
27 current and past tobacco smoking, and diabetes were considered as potential

1 confounders. Variables remained in the models, if they were associated with pancreatic  
2 cancer, correlated with the inflammatory markers, or changed the  $\beta$  estimate by more  
3 than 10%. Based on these conditions, BMI as a continuous variable and smoking as a  
4 categorical variable [never smoking, former smoking (quitting smoking < 10 years ago,  
5  $\geq 10$  years ago), current smoking (< 10, 10 - 20,  $\geq 10$  cigarettes a day), missing] were  
6 considered as confounding factors and remained in the multivariate adjusted model. To  
7 assess a possible confounding effect of diabetes on the risk associations, we controlled  
8 for diabetes in further exploratory analyses. Subjects were defined as diabetics if they  
9 self-reported the condition in the baseline questionnaire at recruitment (n=52) and/or  
10 had glycated haemoglobin (HbA1c) levels  $\geq 6.5\%$  in the current study (n=93). This  
11 percentage is used as a cut-off for diabetes diagnosis (ADA, 2009). HbA1c has been  
12 measured previously in the same study population (Grote *et al*, 2011). Physical activity  
13 and socio-economic status did not markedly change the risk estimates and were,  
14 therefore, not included in the final model.

15 Subgroup analyses were performed to assess possible effect modifications by  
16 sex, diabetes and smoking status, by median age (62 years), waist circumference  
17 (96cm for men, 80 for women), waist-hip-ratio (0.95 for men, 0.80 for women), and  
18 median BMI (26.2 kg/m<sup>2</sup> for men, 24.6 for women), or by lag-time (time between blood  
19 collection and diagnosis of pancreatic cancer,  $\leq$  vs.  $>$  5 years). Cross-product terms  
20 were added in logistic regression models and Wald tests were performed to examine  
21 whether any apparent heterogeneity of effect was significant. To limit reverse causation  
22 bias which could occur when the advanced tumour causes changes in inflammatory  
23 marker levels, we performed subgroup analyses with two years of follow-up as a cut-  
24 point ( $\leq$  vs.  $>$  2 years).

25 All statistical analyses were conducted using the Statistical Analysis System  
26 (SAS) software package, Version 9.2 (SAS Institute Inc., Cary, North Carolina, USA).  
27 All statistical tests were two-tailed and significant at the 5%-level.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

**Results**

Baseline characteristics of pancreatic cancer cases and matched control subjects are shown in **Table 1**. Mean age at recruitment into the initial cohort was 58 years and mean age of cases at pancreatic cancer diagnosis was 63 years, resulting in mean follow-up time of 5.3 years for cases (range 0-13). Female pancreatic cancer cases had a significantly higher BMI and waist-circumference than corresponding controls, but no difference in waist-hip-ratio was observed. For men, however, no significant difference for any of the anthropometric measures comparing cases and controls was seen. A higher percentage of cases currently smoked compared to controls (31 vs. 22%). At baseline, cases also reported more often to be diabetic and/or had HbA1c levels  $\geq 6.5\%$  compared with controls (14 vs. 8%). However, these results are not mutually adjusted and serve descriptive purposes only.

Among controls, sTNF-R1 and sTNF-R2 showed a high degree of correlation. The correlation of circulating CRP levels with IL-6, sTNF-R1, and sTNF-R2 concentrations was relatively high with Spearman's rank correlation coefficients up to 0.44. BMI, waist circumference, and waist-hip-ratio correlated moderately with CRP and IL-6 and to lesser extent with sTNF-R1 but not with sTNF-R2 concentrations (**Table 2**). Participants with diabetes (self-reported at baseline and/or HbA1c  $\geq 6.5\%$ ) and those who smoked had higher levels of CRP and IL-6 than non-diabetics (Table 2). Mutual adjustments for smoking categories and/or BMI resulted in unaltered (diabetes) or stronger associations (smoking, data not shown).

The potential confounders or effect modifiers overweight (OR = 1.05 [95% CI 1.01-1.08], per 5 BMI units), smoking (OR = 1.84 [95% CI 1.30-2.60], current vs. never), and diabetes (OR = 1.74 [95% CI 1.12-2.71]) were associated with risk of pancreatic cancer in our study.

1 Pancreatic cancer risk tended to be increased with higher levels of sTNF-R2  
2 (crude OR = 1.52 [95% CI 0.97 – 2.39] comparing highest with lowest quartiles, p-trend  
3 over quartiles = 0.07); but these associations were not significant at the 5%-level, and  
4 BMI and smoking adjustments attenuated the risks of pancreatic cancer (**Table 3**).  
5 Elevated CRP (crude OR = 1.36 [95% CI 0.92 – 2.01], p-trend = 0.26), IL-6 (OR = 1.30  
6 [95% CI 0.84 – 2.00], p-trend = 0.61), and sTNF-R1 levels (OR = 1.23 [95% CI 0.78-  
7 1.94], p-trend = 0.23) showed no significant association with risk of pancreatic cancer.  
8 Adjustments for HbA1c levels and mutually for the other inflammatory markers in  
9 addition to BMI and smoking categories attenuated risk estimates for elevated levels of  
10 inflammatory markers closer to 1.0 (data not shown). Exclusion of subjects with CRP  
11 levels above 10 mg/L (as this is more likely an indication for an acute rather than a  
12 chronic inflammatory state) had no effect on the association between CRP levels and  
13 pancreatic cancer risk (data not shown). Women tended to be at increased pancreatic  
14 cancer risks for higher CRP or sTNF receptor levels, and specifically so for sTNF-R1,  
15 although risk estimates were inconsistently significant between categorical and  
16 continuous analyses and between crude and BMI and smoking adjusted models (Table  
17 3).

18  
19 Tests for heterogeneity of continuous sTNF receptors, adjusted for matching  
20 factors, resulted in statistically significant differences in pancreatic cancer risk by  
21 median BMI, diabetes and smoking status, but not by median waist circumference,  
22 waist-hip-ratio or median age. Compared to never smokers, risks in former and current  
23 smokers were elevated albeit not statistical significant. Diabetics (p interaction = 0.001)  
24 and subjects with a BMI above the median (p interaction = 0.04) had a significantly  
25 higher risk of pancreatic cancer with elevated levels of sTNF-R2 than non-diabetics or  
26 subjects with lower than median BMI, respectively (**Figure 1b**). Adjusting subgroup  
27 analyses for BMI, smoking categories, HbA1c levels, and/or mutually for inflammatory

1 markers attenuated the risk estimates to non-significance (data not shown).  
2 Interestingly, higher circulating CRP and IL-6 levels tended to be related to increased  
3 pancreatic cancer risk in leaner subjects, although ORs and tests for interaction were  
4 not statistically significant (**Figures 1c/d**).

5

## 6 **Discussion**

7 In our nested case-control study of 455 pancreatic cancer subjects and 455  
8 individually matched controls higher circulating levels of sTNF-R2, but not of sTNF-R1,  
9 CRP, and IL-6 levels tended to be positively associated with the risk of pancreatic  
10 cancer. Stratification by sex revealed significantly increased pancreatic cancer risks in  
11 women for higher sTNF-R1 levels. Positive associations between sTNF-R2 and  
12 pancreatic cancer seemed to be likely for diabetic subjects, those with a higher BMI,  
13 and possibly also for smokers.

14

15 In the acute-phase response to tissues damage, infection, inflammation, or  
16 malignant neoplasia, CRP is increasingly produced by hepatocytes, predominantly  
17 under control by IL-6. CRP binds to damaged cell membranes or apoptotic cells,  
18 forming an aggregate that activates the complement pathway, resulting in the  
19 phagocytosis of the damaged cells and in increased pro-inflammatory  
20 pathophysiological effects. CRP, therefore, reflects ongoing inflammation and/or tissue  
21 damage and functions as a pro-inflammatory mediator. In this context, it may not only  
22 be a marker of a disease, but it may also contribute to pathogenesis (Pepys &  
23 Hirschfield, 2003). In several small hospital-based case-control studies, CRP levels  
24 were significantly higher in pancreatic cancer cases compared to chronic pancreatitis  
25 patients or controls (Barber *et al*, 1999; Moses *et al*, 2009; Mroczko *et al*, 2010). In  
26 addition, elevated levels of CRP were associated with a poor prognosis in pancreatic  
27 cancer patients (McKay *et al*, 2008). Prospectively, no association was observed in a

1 Greek study with 14 pancreatic cancer cases (Trichopoulos *et al*, 2006), whereas a  
2 weak decrease in pancreatic cancer risk with an OR of 0.94 [95% CI 0.89-0.99] was  
3 seen among 311 cases in the Alpha-Tocopherol, Beta-Carotene Cancer Prevention  
4 Study (ATBC) cohort of male Finish smokers (Douglas *et al*, 2010). The same authors  
5 did not find an association in the Ovarian Cancer Screening Trial (PLCO) or in  
6 combined analyses of both cohorts. Our results are in line with the prospective Greek  
7 and PLCO study showing no association of CRP with risk of pancreatic cancer.

8

9 No prospective study has been conducted so far to assess the association of  
10 circulating TNF- $\alpha$ , its soluble receptors, or IL-6 levels with risk of pancreatic cancer,  
11 both up-regulators of CRP. TNF- $\alpha$  is a pro-inflammatory cytokine produced by many  
12 cell types, including cancer cells, upon exogenous noxious stimuli. The effects of TNF-  
13  $\alpha$  are mediated mainly by two receptors, TNF-R1 and TNF-R2, which also circulate in  
14 soluble forms upon shedding. TNF receptor activation leads to induction of genes  
15 involved in inflammation and cell survival, resulting in the activation of nuclear factor- $\kappa$ B  
16 (NF- $\kappa$ B). However, if NF- $\kappa$ B activation is inadequate, apoptosis is mediated via  
17 accumulation of reactive oxygen species as a late response to TNF- $\alpha$ . This cytokine,  
18 thus, is not only involved in maintenance of the immune system, but also in  
19 pathological processes such as malignant diseases. The majority of cell types and  
20 tissues express both receptor types (Balkwill, 2006) and among colon cancer patients it  
21 has been shown that the concentrations of sTNF-Rs correlate with the stage of disease  
22 as tumour cells have a greater tendency than non-malignant cells to shed forms of their  
23 cell surface proteins (Aderka, 1996). Soluble TNF receptors can serve as TNF  
24 antagonists, carrier proteins of TNF, slow release reservoirs for TNF, and stabilizers of  
25 TNF-bioactivity. It is not known, however, whether the two soluble receptors have  
26 distinct or similar functions (Aderka, 1996), and based on this, we cannot explain, why  
27 we observed a potential increase in pancreatic cancer risk for elevated sTNF-R2 but



1 not for sTNF-R1. It might be, however, that sTNF-R2 plays a more prominent role in  
2 pancreatic cancer development. This aspect needs to be explored in functional studies.  
3 So far, TNF- $\alpha$  and/or the soluble receptors have been assessed in hospital-based  
4 case-control studies with pancreatic cancer patients, observing either higher levels of  
5 TNF- $\alpha$ /soluble TNF receptors among pancreatic cancer subjects than among controls  
6 (healthy volunteers or chronic pancreatitis patients (Barber *et al*, 1999; Talar-  
7 Wojnarowska *et al*, 2009)), or no difference in serum levels (Ebrahimi *et al*, 2004). To  
8 our knowledge, our nested-case control study within the prospective EPIC cohort study  
9 is the first to address the association of sTNF receptors with risk of pancreatic cancer,  
10 and we observed a non-significant increase in risk overall, which was more apparent  
11 for sTNF-R2 than sTNF-R1, and which was attenuated after adjustments for smoking  
12 status, BMI, and HbA1c levels or diabetes status. It is unclear why we found a  
13 difference in risk between men and women with elevated risks for increasing levels of  
14 sTNF-R1 in women only.

15  
16 As with TNF- $\alpha$ , pancreatic cancer patients' IL-6 concentrations have shown to be  
17 higher than in healthy controls in hospital-based case-control studies (Barber *et al*,  
18 1999; Ebrahimi *et al*, 2004; Moses *et al*, 2009; Mroczko *et al*, 2010). In contrast to  
19 these observations, in our prospective study we did not find elevated pre-diagnostic IL-  
20 6 concentrations in subjects who became pancreatic cancer cases later in time  
21 compared to non-cancer controls at baseline. IL-6 is synthesized by many cell types in  
22 response to stimulation from TNF- $\alpha$  and IL-1 and indirectly regulates cell proliferation  
23 and apoptosis through its activation of other factors. Therefore, IL-6 plays a role in  
24 chronic inflammation, which may enhance cancer development (Hodge *et al*, 2005).  
25 However, due to the small number of prospective studies so far investigating the  
26 relationship of IL-6 with cancer, a recent published review concluded that it is yet

1 impossible to determine whether IL-6 is causally related to cancer (Heikkilä *et al*,  
2 2008).

3  
4 It has been shown in a wide range of studies that CRP, IL-6, TNF- $\alpha$ , and TNF  
5 receptor levels vary by body weight, with higher levels among overweight or obese  
6 compared to normal weight subjects, and with decreasing levels during weight loss  
7 (Forsythe *et al*, 2008; Himmerich *et al*, 2006). Furthermore, compared with never  
8 smokers, cigarette smokers also have significantly higher levels of CRP and IL-6, and  
9 possibly also of TNF receptors (Fernandez-Real *et al*, 2003). Finally, subclinical  
10 systemic inflammation has been reported in type 2 diabetes (Kolb & Mandrup-Poulsen,  
11 2005), including elevated levels of the aforementioned and evaluated parameters in our  
12 study. In our study, elevated levels of CRP, IL-6 and sTNF-R1 correlated with excess  
13 weight and, in addition, higher CRP and IL-6 levels were associated with smoking and  
14 diabetes.

15 Furthermore, overweight, smoking, or diabetic participants at baseline were at  
16 increased pancreatic cancer risk. This risk was even stronger if overweight or diabetic  
17 participants had elevated levels of sTNF-R2, even though this marker was not  
18 correlated with BMI or associated with diabetes in controls. This can be interpreted as  
19 sTNF-R2 being a mediator of the relationship between overweight and/or diabetes and  
20 pancreatic cancer. A similar scenario is likely for sTNF-R1, but our results do not  
21 clearly support this hypothesis (**Figure 1a**). In contrast, stratification by median BMI,  
22 diabetes or smoking status resulted in similar weak risk estimates for elevated CRP  
23 and IL-6 concentrations. It seems as if, regardless of the presence of a putative  
24 pancreatic cancer risk factor (overweight, diabetes, and smoking), these inflammatory  
25 markers are not associated with pancreatic cancer risk themselves. In addition, they  
26 also do not appear to be in the causal chain between risk factor and cancer.

1           Some strengths and limitations of our study should be mentioned. Although a  
2 single measurement of a biomarker, as assessed in our study, could result in random  
3 misclassification, CRP, IL-6, and sTNF receptors have been shown to be reliably  
4 measured over time (Clendenen *et al*, 2010; Gu *et al*, 2009). A major strength of our  
5 study is that questionnaire data and blood samples were collected prospectively  
6 around the same time point, prior to pancreatic cancer diagnosis, which reduces the  
7 possibility of reverse causation bias to some extent. In addition, pancreatic cancer risk  
8 seemed to be stronger for elevated sTNF receptor levels among subjects with longer  
9 follow-up times. A limitation of our study is that information on pancreatic or liver  
10 disorders, on inflammatory diseases, or on use of anti-inflammatory drugs was not  
11 recorded for most of the EPIC centres; therefore, controlling for these potential  
12 confounders was not possible. Consequently, we cannot exclude the possibility that the  
13 observed suggestive increased pancreatic cancer risk among individuals with elevated  
14 sTNF-R2 levels may partly be due to chronic pancreatitis or impaired liver function, for  
15 example. Furthermore, number of subjects in specific subgroups were rather small,  
16 thus, we cannot rule out that results obtained from these analyses are chance findings.  
17 Further large prospective studies are needed to verify our results in the respective  
18 subgroups with sufficient power to detect significant risk associations.

19

## 20 **Conclusion**

21           Prospectively, CRP and IL-6 do so seem to play a role in our study with respect  
22 to risk of pancreatic cancer, whereas sTNF-R1 seemed to be a risk factor in women  
23 and sTNF-R2 might be a mediator in the risk relationship between overweight and  
24 diabetes with pancreatic cancer. In order to clarify the role of pro-inflammatory proteins  
25 and cytokines in the pathogenesis of exocrine pancreatic cancer, more prospective  
26 studies in large settings are needed, controlling for the potential bias of other conditions  
27 and stratifying by sex.

## **Acknowledgments**

We thank Mrs. Laure Dossus for greatly appreciated statistical support and Miss Britta Lederer and Miss Sigrid Henke for their excellent work in performing the immunoassays. We would also like to take the opportunity to thank the anonymous referees for greatly improving our manuscript.

## **Grant support / funding**

VAG was funded by a grant from the German Research Foundation, Graduiertenkolleg 793: Epidemiology of communicable and chronic noncommunicable diseases and their interrelationships. This work was supported by WCRF UK and WCRF International, grant no 2009/39. The coordination of EPIC is financially supported by the European Commission (DG-SANCO) and the International Agency for Research on Cancer. The national cohorts are supported by Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l'Éducation Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM) (France); Deutsche Krebshilfe, Deutsches Krebsforschungszentrum (DKFZ) and Federal Ministry of Education and Research (Germany); Ministry of Health and Social Solidarity, Stavros Niarchos Foundation and Hellenic Health Foundation (Greece); Italian Association for Research on Cancer (AIRC) and National Research Council (Italy); Dutch Ministry of Public Health, Welfare and Sports (VWS), Netherlands Cancer Registry (NKR), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF), Statistics Netherlands (The Netherlands); ERC-2009-AdG 232997 and Nordforsk, Nordic Centre of Excellence programme on Food, Nutrition and Health. (Norway); Health Research Fund (FIS), Regional Governments of Andalucía, Asturias, Basque Country, Murcia (No 6236) and Navarra, ISCIII RETIC (RD06/0020) (Spain); Swedish Cancer Society,

Swedish Scientific Council and Regional Government of Skåne and Västerbotten (Sweden); Cancer Research UK, Medical Research Council, Stroke Association, British Heart Foundation, Department of Health, Food Standards Agency, and Wellcome Trust (United Kingdom).

## References

- ADA (2009) International Expert Committee report on the role of the A1C assay in the diagnosis of diabetes. *Diabetes Care* **32**(7): 1327-34
- Aderka D (1996) The potential biological and clinical significance of the soluble tumor necrosis factor receptors. *Cytokine Growth Factor Rev* **7**(3): 231-40
- Balkwill F (2006) TNF-alpha in promotion and progression of cancer. *Cancer Metastasis Rev* **25**(3): 409-16
- Barber MD, Fearon KC, Ross JA (1999) Relationship of serum levels of interleukin-6, soluble interleukin-6 receptor and tumour necrosis factor receptors to the acute-phase protein response in advanced pancreatic cancer. *Clin Sci (Lond)* **96**(1): 83-7
- Clendenen TV, Arslan AA, Lokshin AE, Idahl A, Hallmans G, Koenig KL, Marrangoni AM, Nolen BM, Ohlson N, Zeleniuch-Jacquotte A, Lundin E (2010) Temporal reliability of cytokines and growth factors in EDTA plasma. *BMC Res Notes* **3**: 302
- Douglas JB, Silverman DT, Weinstein SJ, Graubard BI, Pollak MN, Tao Y, Virtamo J, Albanes D, Stolzenberg-Solomon RZ (2010) Serum C-reactive protein and risk of pancreatic cancer in two nested, case-control studies. *Cancer Epidemiol Biomarkers Prev* **20**(2): 359-69
- Ebrahimi B, Tucker SL, Li D, Abbruzzese JL, Kurzrock R (2004) Cytokines in pancreatic carcinoma: correlation with phenotypic characteristics and prognosis. *Cancer* **101**(12): 2727-36
- Farrow B, Evers BM (2002) Inflammation and the development of pancreatic cancer. *Surg Oncol* **10**(4): 153-69
- Fernandez-Real JM, Broch M, Vendrell J, Ricart W (2003) Smoking, fat mass and activation of the tumor necrosis factor-alpha pathway. *Int J Obes Relat Metab Disord* **27**(12): 1552-6
- Forsythe LK, Wallace JM, Livingstone MB (2008) Obesity and inflammation: the effects of weight loss. *Nutr Res Rev* **21**(2): 117-33
- Genkinger JM, Spiegelman D, Anderson KE, Bernstein L, van den Brandt PA, Calle EE, English DR, Folsom AR, Freudenheim JL, Fuchs CS, Giles GG, Giovannucci E, Horn-Ross PL, Larsson SC, Leitzmann M, Mannisto S, Marshall JR, Miller AB, Patel AV, Rohan TE, Stolzenberg-Solomon RZ, Verhage BA, Virtamo J, Willcox BJ, Wolk A, Ziegler RG, Smith-Warner SA (2010) A pooled analysis of 14 cohort studies of anthropometric factors and pancreatic cancer risk. *Int J Cancer*
- Goncalves RB, Coletta RD, Silv rio KG, Benevides L, Casati MZ, da Silva JS, Nociti FH, Jr. (2011) Impact of smoking on inflammation: overview of molecular mechanisms. *Inflamm Res* **60**(5): 409-24
- Grote VA, Rohrmann S, Nieters A, Dossus L, Tjonneland A, Halkjaer J, Overvad K, Fagherazzi G, Boutron-Ruault MC, Morois S, Teucher B, Becker S, Sluik D, Boeing H,

Trichopoulou A, Lagiou P, Trichopoulos D, Palli D, Pala V, Tumino R, Vineis P, Panico S, Rodriguez L, Duell EJ, Molina-Montes E, Dorransoro M, Huerta JM, Ardanaz E, Jeurnink SM, Beulens JW, Peeters PH, Sund M, Ye W, Lindkvist B, Johansen D, Khaw KT, Wareham N, Allen N, Crowe F, Jenab M, Romieu I, Michaud DS, Riboli E, Romaguera D, Bueno-de-Mesquita HB, Kaaks R (2011) Diabetes mellitus, glycated haemoglobin and C-peptide levels in relation to pancreatic cancer risk: a study within the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort. *Diabetologia*

Gu Y, Zeleniuch-Jacquotte A, Linkov F, Koenig KL, Liu M, Velikokhatnaya L, Shore RE, Marrangoni A, Toniolo P, Lokshin AE, Arslan AA (2009) Reproducibility of serum cytokines and growth factors. *Cytokine* **45**(1): 44-9

Haftenberger M, Lahmann PH, Panico S, Gonzalez CA, Seidell JC, Boeing H, Giurdanella MC, Krogh V, Bueno-de-Mesquita HB, Peeters PH, Skeie G, Hjartaker A, Rodriguez M, Quiros JR, Berglund G, Janlert U, Khaw KT, Spencer EA, Overvad K, Tjonneland A, Clavel-Chapelon F, Tehard B, Miller AB, Klipstein-Grobusch K, Benetou V, Kiriazi G, Riboli E, Slimani N (2002) Overweight, obesity and fat distribution in 50- to 64-year-old participants in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr* **5**(6B): 1147-62

Heikkilä K, Ebrahim S, Lawlor DA (2008) Systematic review of the association between circulating interleukin-6 (IL-6) and cancer. *Eur J Cancer* **44**(7): 937-45

Himmerich H, Fulda S, Linseisen J, Seiler H, Wolfram G, Himmerich S, Gedrich K, Pollmacher T (2006) TNF-alpha, soluble TNF receptor and interleukin-6 plasma levels in the general population. *Eur Cytokine Netw* **17**(3): 196-201

Hodge DR, Hurt EM, Farrar WL (2005) The role of IL-6 and STAT3 in inflammation and cancer. *Eur J Cancer* **41**(16): 2502-12

Hotamisligil GS (2006) Inflammation and metabolic disorders. *Nature* **444**(7121): 860-7

Huxley R, Ansary-Moghaddam A, Berrington de González A, Barzi F, Woodward M (2005) Type-II diabetes and pancreatic cancer: a meta-analysis of 36 studies. *Br J Cancer* **92**(11): 2076-83

Kolb H, Mandrup-Poulsen T (2005) An immune origin of type 2 diabetes? *Diabetologia* **48**(6): 1038-50

Lynch SM, Vrieling A, Lubin JH, Kraft P, Mendelsohn JB, Hartge P, Canzian F, Steplowski E, Arslan AA, Gross M, Helzlsouer K, Jacobs EJ, LaCroix A, Petersen G, Zheng W, Albanes D, Amundadottir L, Bingham SA, Boffetta P, Boutron-Ruault MC, Chanock SJ, Clipp S, Hoover RN, Jacobs K, Johnson KC, Kooperberg C, Luo J, Messina C, Palli D, Patel AV, Riboli E, Shu XO, Rodriguez Suarez L, Thomas G, Tjonneland A, Tobias GS, Tong E, Trichopoulos D, Virtamo J, Ye W, Yu K, Zeleniuch-Jacquette A, Bueno-de-Mesquita HB, Stolzenberg-Solomon RZ (2009) Cigarette smoking and pancreatic cancer: a pooled analysis from the pancreatic cancer cohort consortium. *Am J Epidemiol* **170**(4): 403-13

McKay CJ, Glen P, McMillan DC (2008) Chronic inflammation and pancreatic cancer. *Best Pract Res Clin Gastroenterol* **22**(1): 65-73

Moses AG, Maingay J, Sangster K, Fearon KC, Ross JA (2009) Pro-inflammatory cytokine release by peripheral blood mononuclear cells from patients with advanced pancreatic cancer: relationship to acute phase response and survival. *Oncol Rep* **21**(4): 1091-5

Mroczo B, Groblewska M, Gryko M, Kedra B, Szmitkowski M (2010) Diagnostic usefulness of serum interleukin 6 (IL-6) and C-reactive protein (CRP) in the differentiation between pancreatic cancer and chronic pancreatitis. *J Clin Lab Anal* **24**(4): 256-61

Pepys MB, Hirschfield GM (2003) C-reactive protein: a critical update. *J Clin Invest* **111**(12): 1805-12

Raimondi S, Lowenfels AB, Morselli-Labate AM, Maisonneuve P, Pezzilli R (2010) Pancreatic cancer in chronic pancreatitis; aetiology, incidence, and early detection. *Best Pract Res Clin Gastroenterol* **24**(3): 349-58

Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, Charrondière UR, Hémon B, Casagrande C, Vignat J, Overvad K, Tjønneland A, Clavel-Chapelon F, Thiébaud A, Wahrendorf J, Boeing H, Trichopoulos D, Trichopoulou A, Vineis P, Palli D, Bueno-De-Mesquita HB, Peeters PH, Lund E, Engeset D, González CA, Barricarte A, Berglund G, Hallmans G, Day NE, Key TJ, Kaaks R, Saracci R (2002) European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* **5**(6B): 1113-24

Talar-Wojnarowska R, Gasiorowska A, Smolarz B, Romanowicz-Makowska H, Kulig A, Malecka-Panas E (2009) Tumor necrosis factor alpha and interferon gamma genes polymorphisms and serum levels in pancreatic adenocarcinoma. *Neoplasma* **56**(1): 56-62

Trichopoulos D, Psaltopoulou T, Orfanos P, Trichopoulou A, Boffetta P (2006) Plasma C-reactive protein and risk of cancer: a prospective study from Greece. *Cancer Epidemiol Biomarkers Prev* **15**(2): 381-4


Vrieling A, Bueno-de-Mesquita HB, Boshuizen HC, Michaud DS, Severinsen MT, Overvad K, Olsen A, Tjønneland A, Clavel-Chapelon F, Boutron-Ruault MC, Kaaks R, Rohrmann S, Boeing H, Nothlings U, Trichopoulou A, Moutsiou E, Dilis V, Palli D, Krogh V, Panico S, Tumino R, Vineis P, van Gils CH, Peeters PH, Lund E, Gram IT, Rodriguez L, Agudo A, Larranaga N, Sanchez MJ, Navarro C, Barricarte A, Manjer J, Lindkvist B, Sund M, Ye W, Bingham S, Khaw KT, Roddam A, Key T, Boffetta P, Duell EJ, Jenab M, Gallo V, Riboli E (2010) Cigarette smoking, environmental tobacco smoke exposure and pancreatic cancer risk in the European Prospective Investigation into Cancer and Nutrition. *Int J Cancer* **126**(10): 2394-403

Whitcomb DC (2004) Inflammation and Cancer V. Chronic pancreatitis and pancreatic cancer. *Am J Physiol Gastrointest Liver Physiol* **287**(2): G315-9




### 1c) CRP

Subgroup	Ca / Co	OR (95% CI)	P int <sup>a</sup>
All	449 / 449	1.08 (0.99-1.18)	
BMI < median <sup>b</sup>	200 / 226	1.13 (1.00-1.28)	0.3
BMI ≥ median	250 / 226	1.01 (0.89-1.15)	
Non-diabetics	373 / 401	1.05 (0.95-1.15)	0.1
Diabetics <sup>c</sup>	59 / 34	1.17 (0.85-1.63)	
Never smoker	161 / 197	1.08 (0.94-1.23)	0.9
Former smoker	143 / 149	1.09 (0.93-1.29)	
Current smoker	141 / 101	1.04 (0.88-1.23)	
FUP ≤ 2yrs <sup>d</sup>	77 / 77	1.10 (0.90-1.35)	0.9
FUP > 2yrs	373 / 375	1.08 (0.98-1.18)	




### 1d) IL-6

Subgroup	Ca / Co	OR (95% CI)	P int <sup>a</sup>
All	424 / 424	1.09 (0.95-1.26)	
BMI < median <sup>b</sup>	193 / 216	1.19 (1.00-1.42)	0.4
BMI ≥ median	247 / 221	0.93 (0.76-1.13)	
Non-diabetics	364 / 388	1.06 (0.93-1.22)	0.02
Diabetics <sup>c</sup>	59 / 33	1.18 (0.71-1.96)	
Never smoker	155 / 193	1.03 (0.84-1.26)	0.2
Former smoker	144 / 142	1.16 (0.93-1.45)	
Current smoker	136 / 97	0.99 (0.77-1.28)	
FUP ≤ 2yrs <sup>d</sup>	74 / 74	1.09 (0.75-1.58)	0.9
FUP > 2yrs	366 / 363	1.09 (0.94-1.28)	




### 1a) sTNF-R1

Subgroup	Ca / Co	OR (95% CI)	P int <sup>a</sup>
All	390 / 390	1.39 (0.87-2.23)	
BMI < median <sup>b</sup>	191 / 209	1.14 (0.60-2.18)	0.3
BMI ≥ median	236 / 203	1.19 (0.68-2.08)	
Non-diabetics	354 / 366	1.08 (0.68-1.70)	0.003
Diabetics <sup>c</sup>	56 / 30	3.09 (0.84-11.36)	
Never smoker	155 / 184	1.07 (0.55-2.11)	0.05
Former smoker	133 / 131	1.10 (0.52-2.34)	
Current smoker	134 / 92	1.20 (0.54-2.68)	
FUP ≤ 2yrs <sup>d</sup>	73 / 69	0.81 (0.26-2.55)	0.3
FUP > 2yrs	354 / 343	1.55 (0.92-2.62)	



### 1b) sTNF-R2

Subgroup	Ca / Co	OR (95% CI)	P int <sup>a</sup>
All	414 / 414	1.55 (0.99-2.44)	
BMI < median <sup>b</sup>	179 / 204	0.89 (0.50-1.58)	0.04
BMI ≥ median	238 / 213	1.93 (1.11-3.37)	
Non-diabetics	349 / 375	1.12 (0.73-1.72)	0.001
Diabetics <sup>c</sup>	54 / 29	4.76 (1.11-20.37)	
Never smoker	148 / 187	0.92 (0.47-1.81)	0.001
Former smoker	133 / 136	1.40 (0.69-2.84)	
Current smoker	131 / 89	1.61 (0.77-3.37)	
FUP ≤ 2yrs <sup>d</sup>	73 / 73	0.97 (0.34-2.80)	0.3
FUP > 2yrs	344 / 344	1.72 (1.05-2.84)	



**Table 1.** Baseline characteristics of pancreatic cancer cases and matched controls

Variable	Cases (n=455)	Controls (n=455)	p-value <sup>a</sup>
Women, n (%)	235 (52)	235 (52)	matched
Age at recruitment [y], mean (range)	58 (30 – 76)	58 (30 – 76)	matched
Age at diagnosis [y], mean (range)	63 (37 – 82)	-	
Follow-up [y], mean (range)	5.3 (0 – 13)	-	
BMI [kg/m <sup>2</sup> ], mean ± SD			
Male	26.8 ± 3.6	26.7 ± 3.7	0.7
Female	26.5 ± 5.0	25.2 ± 4.3	0.002
Waist-hip ratio, mean ± SD			
Male	0.95 ± 0.06	0.95 ± 0.06	0.6
Female	0.82 ± 0.07	0.81 ± 0.06	0.09
Waist circumference [cm], mean ± SD			
Male	96.3 ± 9.9	96.7 ± 10.2	0.7
Female	84.4 ± 12.5	81.2 ± 10.7	0.001
Smoking status, n (%)			< 0.001
Never	162 (36)	198 (44)	
Former	145 (32)	151 (33)	
Current	143 (31)	101 (22)	
Unknown	5 (1)	5 (1)	
Alcohol intake at recruitment [g/d], mean ± SD			0.9
Male	21 ± 26	23 ± 31	
Female	9 ± 13	8 ± 11	
Fasting status, n (%)			matched
Fasting (≥ 6 hours)	118 (26)	113 (25)	
In between (3 - 6 hours)	78 (17)	78 (17)	
Non fasting (< 3 hours)	177 (39)	183 (40)	
Unknown	82 (18)	81 (18)	
Diabetes status, n (%)			
Self-reported diabetes at recruitment	33 (8)	19 (4)	0.05
Subjects HbA1c ≥ 6.5%	54 (12)	29 (6)	0.006
Self-reported diabetes or HbA1c ≥ 6.5%	59 (14)	34 (8)	0.01
Unknown	18 (4)	17 (4)	
CRP [mg/L], geometric mean (95% CI)			
Men	1.12 (0.97-1.29)	1.08 (0.94-1.25)	0.8
Women	1.24 (1.08-1.42)	0.97 (0.84-1.12)	0.02
IL-6 [pg/mL], geometric mean (95% CI)			
Men	1.79 (1.63-1.96)	1.69 (1.52-1.89)	0.6
Women	1.58 (1.43-1.74)	1.44 (1.31-1.59)	0.3
sTNF-R1 [ng/mL], geometric mean (95% CI)			
Men	1.33 (1.28-1.37)	1.36 (1.32-1.41)	0.3
Women	1.39 (1.34-1.44)	1.32 (1.28-1.36)	0.003
sTNF-R2 [ng/mL], geometric mean (95% CI)			
Men	2.31 (2.23-2.40)	2.28 (2.20-2.37)	0.5
Women	2.43 (2.35-2.51)	2.33 (2.26-2.40)	0.04

---

---

<sup>a</sup> *P* values for continuous variables were based on paired t tests; *p* values for categorical variables were based on generalized McNemar's tests

SD = standard deviation, CI = confidence interval

Note: matching factors were EPIC recruitment centre, sex, age at blood collection, date of blood donation, time of blood donation, fasting status, and use of hormones (in women)

**Table 2:** Correlation (95% CI) between inflammatory markers and selected covariates in control participants <sup>a</sup>

Covariate	CRP	IL-6	sTNF-R1	sTNF-R2
IL-6	0.44 (0.35 to 0.53)			
sTNF-R1	0.29 (0.18 to 0.39)	0.33 (0.22 to 0.42)		
sTNF-R2	0.27 (0.16 to 0.37)	0.23 (0.12 to 0.33)	0.65 (0.58 to 0.71)	
BMI	0.40 (0.30 to 0.49)	0.29 (0.18 to 0.39)	0.17 (0.06 to 0.28)	0.05 (-0.06 to 0.16)
Waist	0.32 (0.22 to 0.42)	0.31 (0.20 to 0.41)	0.21 (0.10 to 0.31)	0.10 (-0.01 to 0.21)
WHR	0.23 (0.13 to 0.34)	0.25 (0.14 to 0.35)	0.16 (0.04 to 0.26)	0.09 (-0.03 to 0.20)
HbA1c	0.16 (0.05 to 0.27)	0.09 (-0.02 to 0.20)	0.10 (-0.01 to 0.21)	0.01 (-0.10 to 0.12)
Diabetes <sup>b</sup>	1.37 (1.08 - 1.74)	1.49 (1.06 - 2.11)	1.41 (0.44 - 4.55)	1.30 (0.42 - 4.02)
Smoking <sup>c</sup>	1.30 (1.10 - 1.54)	1.36 (1.06 - 1.74)	1.60 (0.72 - 3.58)	0.97 (0.43 - 2.16)
Sex <sup>d</sup>	1.07 (0.95 - 1.21)	1.21 (1.02 - 1.44)	1.79 (0.97 - 3.31)	0.90 (0.50 - 1.62)
Age <sup>e</sup>	0.13 (0.03 to 0.23)	0.18 (0.08 to 0.28)	0.30 (0.20 to 0.39)	0.32 (0.22 to 0.41)

CRP denotes C-reactive protein, IL-6 interleukin 6, sTNF-R1 and sTNF-R2 soluble tumour necrosis factor receptor 1 and 2, BMI body mass index, waist waist circumference, WHR waist-to-hip ratio, HbA1c glycated haemoglobin, age age at recruitment.

<sup>a</sup> For continuous covariates Spearman's partial rank correlation coefficients were applied. For categorical covariates we used logistic regression. Both methods were performed in controls and adjusted for age, sex and EPIC recruitment centre if not stated otherwise.

<sup>b</sup> Diabetic (HbA1c  $\geq$  6.5% or self-reported diabetes at baseline) vs. non-diabetic participants

<sup>c</sup> Current vs. never smokers

<sup>d</sup> Men vs. women, adjusted for age and EPIC recruitment centre.

<sup>e</sup> Adjusted for sex and EPIC recruitment centre.

**Table 3.** Risk [OR (95% CI)] of pancreatic cancer by quartiles of CRP, IL-6, and sTNF receptors, all subjects combined and stratified by sex <sup>a</sup>

		Quartiles <sup>b</sup>				<i>P</i> trend <sup>c</sup>	OR for doubling in concentration
		1	2	3	4		
<b>CRP</b>	Quartile cut-offs [mg/L]	0.02 - 0.51	0.52 - 1.04	1.05 - 2.05	2.06 - 34.07		
	No. cases / controls (total 449 / 449)	88 / 112	112 / 112	130 / 113	119 / 112		
	Crude <sup>d</sup>	1.0	1.30 (0.88-1.94)	1.45 (1.00-2.10)	1.36 (0.92-2.01)	0.3	1.08 (0.99-1.18)
	Adjusted for smoking, BMI	1.0	1.25 (0.83-1.88)	1.20 (0.80-1.79)	1.02 (0.66-1.57)	0.6	1.01 (0.92-1.11)
	Men, crude	1.0	1.38 (0.76-2.52)	0.98 (0.56-1.70)	1.23 (0.68-2.21)	0.9	1.02 (0.90-1.15)
	Adjusted for smoking, BMI	1.0	1.39 (0.75-2.58)	0.93 (0.52-1.66)	1.09 (0.58-2.04)	0.7	1.00 (0.88-1.13)
	Women, crude	1.0	1.15 (0.67-1.97)	2.14 (1.27-3.59)	1.44 (0.85-2.47)	0.1	1.16 (1.02-1.31)
	Adjusted for smoking, BMI	1.0	1.19 (0.67-2.11)	1.65 (0.92-2.98)	0.99 (0.54-1.81)	0.6	1.02 (0.89-1.18)
<b>IL-6</b>	Quartile cut-offs [pg/ml]	0.16 - 0.94	0.95 - 1.57	1.58 - 2.65	2.66 - 9.66		
	No. cases / controls (total 424 / 424)	86 / 106	123 / 106	108 / 107	107 / 105		
	Crude <sup>d</sup>	1.0	1.45 (0.98-2.15)	1.28 (0.85-1.93)	1.30 (0.84-2.00)	0.6	1.09 (0.95-1.26)
	Adjusted for smoking, BMI	1.0	1.29 (0.86-1.94)	0.97 (0.62-1.51)	1.01 (0.64-1.61)	0.7	0.99 (0.85-1.16)
	Men, crude	1.0	2.02 (1.11-3.68)	1.73 (0.92-3.26)	1.36 (0.70-2.64)	0.9	1.07 (0.86-1.32)
	Adjusted for smoking, BMI	1.0	1.88 (1.00-3.51)	1.51 (0.75-3.04)	1.21 (0.60-2.45)	0.6	1.00 (0.80-1.25)
	Women, crude	1.0	1.10 (0.65-1.87)	1.01 (0.58-1.75)	1.29 (0.72-2.33)	0.4	1.12 (0.92-1.36)
	Adjusted for smoking, BMI	1.0	0.92 (0.52-1.62)	0.71 (0.38-1.33)	0.83 (0.43-1.60)	0.7	0.96 (0.77-1.19)
<b>sTNF-R1</b>	Quartile cut-offs [ng/ml]	0.75 - 1.13	1.14 - 1.31	1.32 - 1.58	1.59 - 2.95		
	No. cases / controls (total 390 / 390)	86 / 97	84 / 98	120 / 98	100 / 97		
	Crude <sup>d</sup>	1.0	0.97 (0.63-1.49)	1.41 (0.94-2.12)	1.23 (0.78-1.94)	0.2	1.39 (0.87-2.23)
	Adjusted for smoking, BMI	1.0	0.84 (0.54-1.32)	1.18 (0.77-1.82)	0.95 (0.58-1.55)	0.9	1.10 (0.66-1.81)
	Men, crude	1.0	0.72 (0.39-1.33)	0.81 (0.44-1.49)	0.71 (0.36-1.39)	0.4	0.67 (0.34-1.35)
	Adjusted for smoking, BMI	1.0	0.71 (0.38-1.35)	0.79 (0.42-1.49)	0.64 (0.31-1.29)	0.3	0.63 (0.30-1.32)
	Women, crude	1.0	1.23 (0.67-2.28)	2.25 (1.26-4.00)	1.97 (1.02-3.79)	0.02	2.74 (1.37-5.47)
	Adjusted for smoking, BMI	1.0	1.03 (0.53-1.99)	1.75 (0.93-3.27)	1.47 (0.72-3.02)	0.2	2.05 (0.97-4.34)
<b>sTNF-R2</b>	Quartile cut-offs [ng/ml]	0.83 - 1.95	1.96 - 2.31	2.32 - 2.68	2.69 - 4.82		
	No. cases / controls (total 414 / 414)	90 / 103	102 / 104	99 / 104	123 / 103		
	Crude <sup>d</sup>	1.0	1.17 (0.77-1.77)	1.18 (0.75-1.85)	1.52 (0.97-2.39)	0.07	1.55 (0.99-2.44)
	Adjusted for smoking, BMI	1.0	1.15 (0.74-1.77)	1.08 (0.68-1.72)	1.42 (0.89-2.27)	0.2	1.40 (0.88-2.23)
	Men, crude	1.0	1.06 (0.59-1.92)	0.98 (0.51-1.90)	1.20 (0.63-2.29)	0.6	1.24 (0.66-2.33)
	Adjusted for smoking, BMI	1.0	1.02 (0.55-1.88)	0.92 (0.46-1.81)	1.27 (0.65-2.46)	0.4	1.35 (0.69-2.61)
	Women, crude	1.0	1.28 (0.71-2.29)	1.40 (0.76-2.60)	1.92 (1.00-3.67)	0.05	1.95 (1.03-3.69)

	Quartiles <sup>b</sup>				<i>P</i> trend <sup>c</sup>	OR for doubling in concentration
	1	2	3	4		
Adjusted for smoking, BMI	1.0	1.22 (0.65-2.28)	1.17 (0.60-2.28)	1.72 (0.86-3.44)	0.1	1.60 (0.80-3.17)

CI = confidence interval, No. = number. CRP, IL-6, and sTNF receptor concentrations on continuous scales were log2 transformed. Smaller number of subjects due to missing laboratory values.

<sup>a</sup> Crude p-interaction over quartiles, for CRP = 0.03, IL-6 = 0.2, sTNF-R1 = 0.09, sTNF-R2 = 0.8. BMI and smoking adjusted p-interaction, for CRP = 0.03, IL-6 = 0.2, sTNF-R1 = 0.1, sTNF-R2 = 0.9

<sup>b</sup> Quartile cut points were based on the distribution of controls

<sup>c</sup> *P* trend test was based on median values of each quartile

<sup>d</sup> Logistic regression conditioned on matching factors (EPIC recruitment centre, sex, age at recruitment, date at entry in the cohort, time between blood sampling and last consumption of foods and drinks, hormone use). Adjusting variables in further model: smoking (former smokers adjusted for quitting smoking (< 10 or ≥ 10 years ago), current smokers adjusted for number of cigarettes (1-9, 10-19, or ≥ 20)), and BMI (continuous, [kg/m<sup>2</sup>]).

## Title and Caption of Figure 1

**Figure 1:** Crude relative risks [OR (95% CI)] of pancreatic cancer for a doubling in sTNF receptor concentrations, CRP, and IL-6, all and stratified by median BMI (26.2 for men, 24.6 for women), diabetes, smoking status, and length of follow-up ( $\leq 2$  vs.  $> 2$  yrs)

### Caption

NOTE: Stratified analysis using unconditional logistic regression were adjusted for matching factors (EPIC recruitment centre, sex, age at blood collection, date of blood donation, time of blood donation, fasting status, and use of hormones). Ca / Co = number of cases / controls. Size of squares is proportional to number of participants in the respective subgroup; squares represent ORs, with error bars indicating 95% CIs.

- <sup>a</sup> P for interaction was based on the Wald statistics, adjusted for matching factors.
- <sup>b</sup> Median BMI for male controls was 26.20 kg/m<sup>2</sup>, for female controls 24.61 kg/m<sup>2</sup>.
- <sup>c</sup> Diabetics included subjects with self-reported diabetes status at baseline and subjects with glycated haemoglobin (HbA1c) levels  $\geq 6.5\%$  or both.
- <sup>d</sup> FUP = follow-up time [years], using conditional logistic regression.