



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

**Functional characterization of the rod visual pigment of the echidna
(*Tachyglossus aculeatus*), a basal mammal**

Bickelmann, Constanze ; Morrow, James M ; Müller, Johannes ; Chang, Belinda S W

DOI: <https://doi.org/10.1017/S0952523812000223>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-75766>

Journal Article

Published Version

Originally published at:

Bickelmann, Constanze; Morrow, James M; Müller, Johannes; Chang, Belinda S W (2012). Functional characterization of the rod visual pigment of the echidna (*Tachyglossus aculeatus*), a basal mammal. *Visual Neuroscience*, 29(4-5):211-217.

DOI: <https://doi.org/10.1017/S0952523812000223>

Functional characterization of the rod visual pigment of the echidna (*Tachyglossus aculeatus*), a basal mammal

CONSTANZE BICKELMANN,^{1,*} JAMES M. MORROW,² JOHANNES MÜLLER,¹ AND BELINDA S.W. CHANG^{2,3,4}

¹Museum für Naturkunde – Leibniz-Institut für Evolutions- und Biodiversitätsforschung an der Humboldt-Universität zu Berlin, Berlin, Germany

²Department of Cell & Systems Biology, University of Toronto, Toronto, Ontario, Canada

³Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, Ontario, Canada

⁴Centre for the Analysis of Genome Evolution & Function, University of Toronto, Toronto, Ontario, Canada

*Present address: Paläontologisches Institut und Museum, Universität Zürich, Karl-Schmid-Strasse 4, 8006 Zurich, Switzerland

(RECEIVED March 7, 2012; ACCEPTED May 5, 2012; FIRST PUBLISHED ONLINE July 9, 2012)

Abstract

Monotremes are the most basal egg-laying mammals comprised of two extant genera, which are largely nocturnal. Visual pigments, the first step in the sensory transduction cascade in photoreceptors of the eye, have been examined in a variety of vertebrates, but little work has been done to study the rhodopsin of monotremes. We isolated the rhodopsin gene of the nocturnal short-beaked echidna (*Tachyglossus aculeatus*) and expressed and functionally characterized the protein *in vitro*. Three mutants were also expressed and characterized: N83D, an important site for spectral tuning and metarhodopsin kinetics, and two sites with amino acids unique to the echidna (T158A and F169A). The λ_{\max} of echidna rhodopsin (497.9 ± 1.1 nm) did not vary significantly in either T158A (498.0 ± 1.3 nm) or F169A (499.4 ± 0.1 nm) but was redshifted in N83D (503.8 ± 1.5 nm). Unlike other mammalian rhodopsins, echidna rhodopsin did react when exposed to hydroxylamine, although not as fast as cone opsins. The retinal release rate of light-activated echidna rhodopsin, as measured by fluorescence spectroscopy, had a half-life of 9.5 ± 2.6 min⁻¹, which is significantly shorter than that of bovine rhodopsin. The half-life of the N83D mutant was 5.1 ± 0.1 min⁻¹, even shorter than wild type. Our results show that with respect to hydroxylamine sensitivity and retinal release, the wild-type echidna rhodopsin displays major differences to all previously characterized mammalian rhodopsins and appears more similar to other nonmammalian vertebrate rhodopsins such as chicken and anole. However, our N83D mutagenesis results suggest that this site may mediate adaptation in the echidna to dim light environments, possibly *via* increased stability of light-activated intermediates. This study is the first characterization of a rhodopsin from a most basal mammal and indicates that there might be more functional variation in mammalian rhodopsins than previously assumed.

Keywords: Rhodopsin, Mammals, Monotremes, Retinal release rate, N83D mutagenesis

Introduction

The egg-laying monotremes are the most basal extant mammals and today comprise only five species: the duck-billed platypus (*Ornithorhynchus anatinus*; Shaw, 1799), the short-beaked echidna (*Tachyglossus aculeatus*; Shaw, 1792), and three species of long-beaked echidnas (*Zaglossus attenboroughi*; Flannery & Groves, 1998, *Z. bartoni*; Thomas, 1907, and *Z. brujini*; Peters & Doria, 1876). All living monotremes are nocturnal, homeothermic, and possess a low rate of reproduction (Dawson et al., 1979; Rismiller, 1999; Werneburg & Sánchez-Villagra, 2010). However, some diurnal activity has also been occasionally observed for echidnas (*T. aculeatus multiaculeatus*) (Rismiller & McKelvey, 2009). Overall, monotremes exhibit an

intriguing mosaic of reptilian and mammalian characters in terms of anatomy, physiology, and reproduction (Griffiths, 1989; Campbell & Reece, 2009). They are endemic to the Australia–New Guinea shelf (Dawson et al., 1979; Rismiller, 1999), with the short-beaked echidna being the most widely distributed extant monotreme and found both in Australia and in New Guinea (Griffiths, 1989; Nicol & Andersen, 2006). The origin of Monotremata presumably occurred sometime in the Late Triassic/Early Jurassic, a date supported by fossil as well as molecular data (Luo et al., 2002; Woodburne et al., 2003; Phillips et al., 2009). Based on fossil evidence, the divergence of modern platypus and echidna lineages occurred in the Early Cretaceous (Rowe et al., 2008).

Rhodopsin is the visual pigment responsible for dim light vision in the rod photoreceptors of vertebrates (Menon et al., 2001). While visual pigment (opsin) gene sequences have been isolated in a variety of vertebrates, including fish (Chinen et al., 2003), amphibians (Starace & Knox, 1998), reptiles including birds (Kawamura & Yokoyama, 1998), and mammals (Nathans & Hogness, 1983; Carvalho et al., 2006), functional studies of mammalian rhodopsins

Address correspondence and reprint requests to: Dr. Constanze Bickelmann, Paläontologisches Institut und Museum, Universität Zürich, Karl-Schmid-Strasse 4, 8006 Zurich, Switzerland. E-mail: constanze.bickelmann@pim.uzh.ch and Prof. Belinda S.W. Chang, Department of Cell & Systems Biology, University of Toronto, 25, Harbord St., Toronto, Ontario, Canada M5S 3G5. E-mail: belinda.chang@utoronto.ca

have been largely limited to bovine (Sakmar et al., 2002; Yan et al., 2002; Natochin et al., 2003) and human (Tam & Moritz, 2009; Pulagam & Palczewski, 2010), both diurnal therians. Recent evidence, however, suggests that female bovine may not follow a strictly diurnal activity pattern (Betteridge et al., 2010). Although gene sequences of echidna cone opsins and platypus visual pigments have been isolated (Davies et al., 2007; Wakefield et al., 2008), neither study included a thorough functional characterization of the expressed photopigments. Furthermore, there has been a notable lack of functional studies of nocturnal mammalian rhodopsins, which is surprising given that the molecular evolution of rhodopsin is thought to be linked to its photic environment (Zhao et al., 2009; Shen et al., 2010).

In this study, we isolate and functionally characterize the rhodopsin from the short-beaked echidna (*T. aculeatus*), a monotreme, which are the most basal living mammals, in order to investigate a number of functions known to differ among visual pigments, most notably between rod and cone opsins, including hydroxylamine stability (Kawamura & Yokoyama, 1998; Starace & Knox, 1998) and the rate of retinal release upon photoactivation (Farrens & Khorana, 1995; Yan et al., 2002). We perform site-directed mutagenesis in order to identify residues underlying functional differences found in the echidna rhodopsin in comparison with bovine rhodopsin, a largely diurnal therian mammal. Characterizing the rhodopsin of the echidna, one of the most basal of the living mammals, is crucial for understanding the evolution of mammalian rhodopsins.

Materials and methods

Rhodopsin cloning and site-directed mutagenesis

Blood samples from a female short-beaked echidna (*T. aculeatus*) were obtained from the Toronto Zoo (Toronto, Canada) and stored in Queen's Lysis buffer (Shaw et al., 2003). Genomic DNA was extracted from these blood samples using the DNeasy Blood and Tissue Kit (Qiagen, The Netherlands), and a Genome Walker library was synthesized according to the manufacturer's protocol (Universal GenomeWalker Kit; Clontech, Madison, WI). The echidna rhodopsin sequence was isolated from the Genome Walker library by polymerase chain reaction (PCR) using Genome Walker adapter primers (AP1 and AP2) combined with degenerate primers previously designed to target the platypus rhodopsin (Davies et al., 2007), as well as degenerate primers designed from other tetrapod rhodopsins (Table S1). The following cycling conditions were used: an initial 1-min denaturation at 95°C followed by 7 cycles of denaturation at 94°C for 25 s and primer annealing at 72°C for 3 min; another 32 cycles of denaturation at 94°C for 25 s and primer annealing at 67°C for 3 min; product extension was at 67°C for 7 min.

Site-directed mutagenesis primers were designed to generate the mutants N83D, T158A, and F169A (Table S2) using the following primers (Table S2). N83D was selected because it was previously shown to be involved in both spectral tuning (Fasick & Robsinson, 1998) and meta intermediate kinetics (Sugawara et al., 2010). Both T158 and F169 are amino acids unique to the echidna rhodopsin, with F169 being of particular interest, as it potentially interacts with the β -ionine ring of all-*trans*-retinal (Borhan et al., 2000; Palczewski et al., 2000). Amino acid numbering used in this study corresponds to bovine rhodopsin (Sakmar et al., 2002).

Phylogenetic analyses

The echidna rhodopsin gene sequence (GenBank accession: JX103830) was aligned to 12 other tetrapod rhodopsins downloaded

from the NCBI database (Table S3) and subjected to Bayesian phylogenetic analyses in MrBayes 3.2 (Ronquist & Huelsenbeck, 2003). The analysis was conducted using the GTR + I + G model, the best fitting model as determined by JModelTest (Posada, 2008), and was run for 1,000,000 generations sampling every 100 generations. Trees that were generated before stationary was reached were discarded (burn in = 1000), and the remaining trees were used to construct a consensus tree. Nodal support was assessed by posterior probability values ($\geq 95\%$ = statistical support).

Protein expression and purification

Once the gene sequence had been determined, the full coding sequence (minus the introns) of echidna rhodopsin was synthesized by GeneArt AG (Regensburg, Germany). The artificially synthesized sequence was then inserted into the p1D4-hrGFP II expression vector and thereby tagged with a nine amino acid sequence (TETSQVAPA) at the carboxy terminus to allow for immunoaffinity purification of expressed proteins from HEK293T cells (Morrow & Chang, 2010).

Expression vectors containing echidna rhodopsin and the mutants as well as bovine rhodopsin (8 μ g per 10-cm plate) were transiently transfected into cultured HEK293T cells using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) and harvested 48 h after transfection. Visual pigments were regenerated with 11-*cis*-retinal, generously provided by Dr. Rosalie Crouch (Medical University of South Carolina), then solubilized in 1% dodecylmaltoside, and purified with the 1D4 monoclonal antibody as previously described (Morrow & Chang, 2010; Morrow et al., 2011).

Spectroscopic assays

The ultraviolet-visible absorption spectra of purified visual pigments were recorded using a Cary 4000 double beam spectrophotometer (Agilent, Santa Clara, CA). Dark-light difference spectra were calculated by subtracting light-bleached absorbance spectra from respective dark spectra. Pigments were photoexcited with light from a fiber optic lamp for 60 s at 25°C. Reactivity to hydroxylamine was determined by incubating visual pigments in 50 mM hydroxylamine (Sigma-Aldrich, St. Louis, MO) at 25°C. Absorption spectra were taken every 3–5 min for 30 min and every 15 min for another 90 min. The rate of retinal release from light-activated rhodopsin was monitored using a Cary Eclipse fluorescence spectrophotometer (Agilent). This rate was measured by monitoring the increase in fluorescence due to the exit of retinal from the chromophore-binding pocket after photoactivation; a process that has previously been referred to as meta II decay (Farrens & Khorana, 1995). Excitation and emission wavelengths were set to 295 and 330 nm, respectively, with excitation and emission slit widths set to 1.5 and 10 nm. Fluorescence measurements were taken every 30 s for 45 min at 20°C. Data for retinal release assays was fit to a first-order exponential curve ($y = y_0 + a(1 - e^{-bx})$), with half-life values being calculated based on rate constant "b" ($t_{1/2} = \ln 2/b$).

Results and discussion

Echidna rhodopsin sequence analysis

Using a combination of degenerate primers designed from other mammalian rhodopsins, and Genome Walker adapter primers, we were able to isolate a full-length genomic sequence of echidna

rhodopsin. This sequence contained four introns (Table S4). The intron–exon boundaries are highly similar to the ones in another monotreme, the platypus (Davies et al., 2007). The translated amino acid echidna rhodopsin sequence contained many of the conserved motifs known to be important for visual pigment function (Sakmar et al., 2002), including a lysine residue (Lys 296) in the seventh transmembrane domain serving as the site of covalent attachment of the retinal chromophore, its counterion (Glu 113), a pair of cysteine residues known to form a structurally important disulphide bridge between helices 3 and 4 and another pair at the end of helix 8 known to be palmitoylated in bovine rhodopsin (Fig. 1). Phylogenetic analyses of the echidna rhodopsin coding sequence aligned with other tetrapod rhodopsin gene sequences placed the echidna as basal to all other mammalian sequences, as expected from its phylogenetic position within mammals (Fig. 2).

At highly conserved sites, the echidna rhodopsin bears the same residues shared with all other taxa (Table S3). Interestingly, at more variable sites, it shares a surprising number of residues in common with reptiles and amphibians, as opposed to therian mammals (Table S3). Shared residues between monotreme and reptile and amphibian rhodopsins include 7, 8, 13, 225, 346, and 348 (Table S3). Furthermore, serine and threonine residues at the C-terminus (349–353) are shared in common with amphibians and reptiles but were lost in placentals and marsupials (Table S3). These additional residues could be targets for phosphorylation, which in turn could affect the binding affinity of arrestin, a regulator of rhodopsin photochemistry (Sommer & Farrens, 2006). In contrast, the echidna sequence only shares four residues with therian mammals at more variable sites, where amino acids differ between Theria and nonmammals: 99, 100, 228, and 308 (Table S3). Substitutions unique to monotremes are found at residues 39, 169, and 344 (Table S3). Although site 169 is not very conserved among tetrapods, both echidna and platypus bear unique residues at this site (Table S3). At site 158, which is not highly conserved either, the echidna rhodopsin bears a unique substitution (Table S3).

A mosaic of derived and plesiomorphic characters in monotremes, as exemplified by our analysis of the echidna rhodopsin sequence, has also been reported from anatomic, genomic, physiological, and developmental studies (Bolk et al., 1934; Gresser & Noback, 1935; Griffiths, 1989; Young & Pettigrew, 1991; Warren et al., 2008; Werneburg & Sánchez-Villagra, 2010). This aspect of the echidna rhodopsin sequence would support the yet controversial Theria hypothesis that monotremes are sister to marsupials and placentals (Janke et al., 2002; Rowe et al., 2008). Moreover, the odd mosaic pattern in the echidna amino acid sequence might suggest unexpected structural and functional similarities with amphibian and reptilian rhodopsins, a hypothesis which would require further investigation.

Protein expression and spectroscopic assays

Echidna rhodopsin was successfully expressed *in vitro*, immunofinity purified, and regenerated with the 11-*cis*-retinal chromophore to produce a stable photopigment with an absorption maximum (λ_{\max}) at 497.9 ± 1.1 nm (Fig. 3). The λ_{\max} value of the platypus, the other extant monotreme, has been found to be 498 nm (Davies et al., 2007). Both are well within the characteristic range of λ_{\max} values for mammalian rhodopsins, which tend to be at around 500 nm (Menon et al., 2001; Bowmaker, 2008). Bovine rhodopsin, which was expressed alongside, showed a λ_{\max} of 500 nm, which falls within the published range (Oprian et al., 1987; Stavenga et al., 1993). We created three mutants to see if sites of interest cause any shift in properties known to differ between rod and cone opsins. Mutations of echidna-specific residues, T158A and F169A, were found to have λ_{\max} values similar to wild type of 498.0 ± 1.3 nm and 499.4 ± 0.1 nm, respectively. However, the N83D mutant caused a significant redshift in the absorbance spectrum, with a λ_{\max} of 503.8 ± 1.5 nm, compared to D83 wild-type echidna rhodopsin, which has its λ_{\max} at 497.9 ± 1.1 nm. This is consistent with previous studies that identified N83 as the cause of a blueshifted




Fig. 1. Two-dimensional structure of the echidna rhodopsin (modified from Sakmar et al., 2002). Residues shared with bovine rhodopsin are in light gray, whereas residues differing from bovine rhodopsin are highlighted in black. Note the insert at the C-terminus.




Fig. 2. Phylogenetic analysis of 13 tetrapod rhodopsin sequences, conducted in MrBayes 3.2. using the GTR + I + G model. Percentages indicate posterior probabilities. As expected, monotremes are situated basal to all other mammals.

λ_{\max} in the rhodopsins of some marine mammals (Fasick & Robinson, 2000). Both wild-type and mutant pigments as well as bovine rhodopsin converted to their biologically active meta II intermediates when activated with light, noted by a shift in λ_{\max} to approximately 380 nm, the characteristic absorbance maximum of metarhodopsin II (Fig. 3, Inset). This confirms that all synthesized visual pigments are able to properly activate in response to light.

The wild-type echidna rhodopsin and its mutants, together with bovine rhodopsin, were subjected to two spectroscopic assays shown in previous studies to differ between rod and cone visual pigments, hydroxylamine reactivity, and retinal release following photoactivation (Kawamura & Yokoyama, 1998; Starace & Knox, 1998; Chen et al., 2012). Both assays revealed significant differences between echidna and bovine rhodopsins and unexpected functional properties reminiscent of cone visual pigments. Monitoring hydroxylamine reactivity by absorption spectroscopy revealed that, unlike bovine




Fig. 3. Dark absorption spectra of the rhodopsin of the echidna (*T. aculeatus*). Note the λ_{\max} at 498 nm. Inset, difference spectra generated by taking an absorbance spectrum after bleaching with a fiber optic lamp for 30 s and subtracting it from the dark spectrum.

and other mammalian rhodopsins, wild-type echidna rhodopsin, along with T158A and F169A mutants, are in fact sensitive to hydroxylamine (Fig. 4). Moreover, fluorescence assays monitoring the decay of light-activated meta intermediates found a substantially faster rate of retinal release in echidna rhodopsin ($t_{1/2} = 9.5 \pm 2.6 \text{ min}^{-1}$) in comparison to bovine rhodopsin ($t_{1/2} = 13.5 \pm 1.3 \text{ min}^{-1}$; Fig. 5). The echidna N83D mutation resulted in even faster decay rates, decreasing the half-life to $5.1 \pm 0.1 \text{ min}^{-1}$ (Fig. 5), whereas mutations of echidna-specific substitutions (T158A, F169A) had little effect (data not shown).

Previous studies have shown that nonmammalian rhodopsins sometimes display more cone-like functional characteristics relative to bovine rhodopsin. For example, rhodopsin of the anole (*Anolis carolinensis*) is sensitive to hydroxylamine, yielding a complete reaction in around 8.5 h, whereas bovine rhodopsin is virtually unaffected for over 12 h (Kawamura & Yokoyama, 1998). Meanwhile, the decay rate of chicken metarhodopsin II is about twice as fast compared to both bovine and human rhodopsin (Imai et al., 1995; Lewis et al., 1997; Okada et al., 1994; Janz & Farrens, 2004).

Our study is the first to reveal such functional characteristics in a mammalian rhodopsin, with the echidna rhodopsin both reacting to hydroxylamine and having a faster rate of retinal release, a property connected to metarhodopsin decay rates, than bovine rhodopsin. It should also be noted that these differences among vertebrate rhodopsins are not as large as the more striking contrasts between rhodopsins and cone opsins, which are largely thought to be mediated by substitutions at two residues, Q122E and P189I (Imai et al., 1997, 2007; Kuwayama et al., 2002, 2005), which are invariant among vertebrate rhodopsins. However, our results contribute to the idea that intriguing functional differences exist within rhodopsins, not just between rhodopsins and cone opsins, and are worth investigating, especially in enigmatic and phylogenetically relevant species such as the echidna.




Fig. 4. Absorption spectra of echidna rhodopsin before and after 1 M hydroxylamine treatment. One mole of freshly prepared hydroxylamine diluted in phosphate buffered saline (PBS) was added at 25°C. Spectra were recorded with a scan rate of 400 nm/min, average and integration time of 0.1 and 0.12 s, respectively, data interval of 0.667 nm, and a slit width of 2 nm. Black curve shows dark absorption spectrum before hydroxylamine treatment. Light gray shows rhodopsin absorption after treatment for 2 h. Dark gray curve shows absorption after photoexcitement for 30 s with a fiber optic lamp. Inset, percentage of maximum absorbance over time for echidna wild type (light gray dots) and bovine rhodopsin (black dots) after hydroxylamine treatment. Note the decrease seen in echidna rhodopsin compared to the bovine.




Fig. 5. Fluorescence increase of retinal release following Schiff base hydrolysis due to photobleaching of rhodopsin. Excitation and emission wavelengths were 295 and 330 nm respectively, monitoring the decrease in fluorescent quenching of Trp265 by all-*trans*-retinal as the latter is released from the chromophore-binding pocket of rhodopsin. The $t_{1/2}$ values for bovine rhodopsin (black), echidna rhodopsin (light gray), and echidna rhodopsin N83D (dark gray) were 13.5, 9.5, and 5.1 min^{-1} , respectively. Data were collected for 45 min after photoactivation, at 30-s interval, with an integration time of 2 s.

A recent study has suggested that N83 may in fact be an adaptation to dim light environments, by increasing the rate of formation of the biologically active meta II rhodopsin intermediate, and thereby increasing photosensitivity in dim light in cichlid fishes and nocturnal bats (Sugawara et al., 2010). Echidna rhodopsin also has N83. When we mutated this residue to aspartic acid (D), the most common identity of site 83 in mammals, our results indicate a faster rate of retinal release compared to wild-type echidna rhodopsin. Although the precise mechanisms by which N83 affect meta intermediate kinetics remain unclear, previous studies have suggested that greater stability of one or more meta intermediates of light-activated echidna rhodopsin may allow for further transducin activation, sustaining signaling activity, and possibly even increasing the sensitivity of the photoreceptor cells (Heck et al., 2003; Lamb & Pugh, 2004; Imai et al., 2007). Increased photosensitivity could be advantageous, particularly for vision at low light levels. Therefore, our investigations of echidna rhodopsin function suggest that although its rhodopsin is sensitive to hydroxylamine and has a faster rate of retinal release upon light activation relative to other mammalian rhodopsins, it does possess a substitution, N83, which might be an adaptation to dim light environments.

In this study, we present the first detailed functional characterization of a rhodopsin from a most basal mammal, the echidna (*T. aculeatus*). Surprisingly, our results indicate that the rhodopsin of the echidna displays functional characteristics such as sensitivity to hydroxylamine and a faster metarhodopsin decay rate (as compared to bovine rhodopsin), which are more similar to nonmammalian rhodopsins, and, to a lesser extent, cone opsins. These results are intriguing and show that significant variation can occur even within mammalian rhodopsins.

Finally, although the rhodopsin of the echidna did not display overall functional characteristics in comparison with the more diurnal bovine rhodopsin that might be expected of a nocturnal pigment, our data do suggest nonetheless that N83 might be an adaptation to dim-light vision. In conclusion, the rhodopsin of the

echidna appears as enigmatic as the animal itself displaying unusual biochemical and functional characteristics, which differ from all other mammalian rhodopsins investigated, and may be due to the mosaic pattern of derived and plesiomorphic characters within its amino acid sequence.

Acknowledgments

We thank G. Crawshaw (Toronto) for the echidna blood samples. We are thankful to all members of the Chang Lab, especially to I. van Hazel (Toronto, Canada), as well as to M. Meixner, R. Schreiber, and A. Sonntag (all Berlin, Germany) for help in the lab. We thank E.C. Kirk (Austin, TX) for discussion on bovine activity patterns, as well as L.A. Tsuji, R.K. Schott, and the two anonymous reviewers for their constructive comments, which helped to improve an earlier version of the manuscript. This work was funded by the Deutsche Forschungsgemeinschaft (DFG Mu 1760/2-3, to C.B. and J.M.), a University Health Network (University of Toronto) Vision Science Research Program fellowship (J.M.M.), a National Sciences and Engineering Research Council Discovery grant (to B.S.W.C.), and a Government of Ontario Early Researcher Award (to B.S.W.C.).

References

- BETTERIDGE, K., COSTALL, D., BALLADUR, S., UPSDELL, M. & UMEMURA, K. (2010). Urine distribution and grazing behaviour of female sheep and cattle grazing a steep New Zealand hill pasture. *Animal Production Science* **50**, 624–629.
- BOLK, L., GÖPPERT, E., KALLIUS, E. & LUBOSCH, W. (1934). *Handbuch der vergleichenden Anatomie der Wirbeltiere*. Berlin, Germany: Urban & Schwarzenberg.
- BORHAN, B., SOUTO, M.L., IMAI, H., SHICHIDA, Y. & NAKANISHI, K. (2000). Movement of retinal along the visual transduction path. *Science* **288**, 2209–2212.
- BOWMAKER, J.K. (2008). Evolution of vertebrate visual pigments. *Vision Research* **48**, 2022–2041.
- CAMPBELL, N.A. & REECE, J.B. (2009). *Biologie*. München, Germany: Pearson Studium.
- CARVALHO, L.S., COWING, J.A., WILKIE, S.E., BOWMAKER, J.K. & HUNT, D.M. (2006). Shortwave visual sensitivity in tree and flying squirrels reflects changes in lifestyle. *Current Biology* **16**, R81–83.
- CHEN, M.H., KUEMMEL, C., BIRGE, R.R. & KNOX, B.E. (2012). Rapid retinal release from a cone visual pigment following photoactivation. *Biochemistry* **51**, 4117–4125.
- CHINEN, A., HAMAOKA, T., YAMADA, Y. & KAWAMURA, S. (2003). Gene duplication and spectral diversification of cone visual pigments of zebrafish. *Genetics* **163**, 663–675.
- DAVIES, W.L., CARVALHO, L.S., COWING, J.A., BEAZLEY, L.D., HUNT, D.M. & ARRESE, C.A. (2007). Visual pigments of the platypus: A novel route to mammalian colour vision. *Current Biology* **17**, R161–R163.
- DAWSON, T.J., GRANT, T.R. & FANNING, D. (1979). Standard metabolism of monotremes and the evolution of homeothermy. *Australian Journal of Zoology* **27**, 511–515.
- FARRENS, D.L. & KHORANA, H.G. (1995). Structure and function in rhodopsin. Measurement of the rate of metarhodopsin II decay by fluorescence spectroscopy. *The Journal of Biological Chemistry* **270**, 5073–5076.
- FASICK, J.I. & ROBINSON, P.R. (1998). Mechanism of spectral tuning in the dolphin visual pigments. *Biochemistry* **37**, 433–438.
- FASICK, J.I. & ROBINSON, P.R. (2000). Spectral-tuning mechanisms of marine mammal rhodopsins and correlations with foraging depth. *Visual Neuroscience* **17**, 781–788.
- FLANNERY, T.F. & GROVES, C.P. (1998). A revision of the genus *Zaglossus* (Monotremata, Tachyglossidae), with description of new species and subspecies. *Mammalia* **62**, 367–396.
- GRESSER, E.B. & NOBACK, C.V. (1935). The eye of the monotreme, Echidna hystrix. *Journal of Morphology* **58**, 279–284.
- GRIFFITHS, M. (1989). Tachyglossidae. In *Fauna of Australia. Mammalia*, ed. WALTON, D.W. & RICHARDSON, B.J. pp. 407–435. Canberra, Australia: Australian Capital Territory 1B.
- HECK, M., SCHADEL, S.A., MARETZKI, D., BARTL, F.J., RITTER, E., PALCZEWSKI, K. & HOFMANN, K.P. (2003). Signaling states of rhodopsin. Formation of the storage form, metarhodopsin III, from active metarhodopsin II. *The Journal of Biological Chemistry* **278**, 3162–3169.

- IMAI, H., IMAMOTO, Y., YOSHIZAWA, T. & SHICHIDA, Y. (1995). Difference in molecular properties between chicken green and rhodopsin as related to the functional difference between cone and rod photoreceptor cells. *Biochemistry* **34**, 10525–10531.
- IMAI, H., KEFALOV, V., SAKURAI, K., CHISAKA, O., UEDA, Y., ONISHI, A., MORIZUMI, T., FU, Y., ICHIKAWA, K., NAKATANI, K., HONDA, Y., CHEN, J., YAU, K.-W. & SHICHIDA, Y. (2007). Molecular properties of rhodopsin and rod function. *The Journal of Biological Chemistry* **282**, 6677–6684.
- IMAI, H., KOJIMA, D., OURA, T., TACHIBANAKI, S., TERAKITA, A. & SHICHIDA, Y. (1997). Single amino acid residue as a functional determinant of rod and cone visual pigments. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 2322–2326.
- JANKE, A., MAGNELL, O., WIECZOREK, G., WESTERMAN, M. & ARNASON, U. (2002). Phylogenetic analysis of 18S rRNA and the mitochondrial genomes of the wombat, *Vombatus ursinus*, and the spiny anteater, *Tachyglossus aculeatus*: Increased support for the Marsupionta hypothesis. *Journal of Molecular Evolution* **54**, 71–80.
- JANZ, J.M. & FARRENS, D.L. (2004). Role of the retinal hydrogen bond network in rhodopsin Schiff base stability and hydrolysis. *The Journal of Biological Chemistry* **279**, 55886–55894.
- KAWAMURA, S. & YOKOYAMA, S. (1998). Functional characterization of visual and nonvisual pigments of American chameleon (*Anolis carolinensis*). *Vision Research* **38**, 37–44.
- KUWAYAMA, S., IMAI, H., HIRANO, T., TERAKITA, A. & SHICHIDA, Y. (2002). Conserved proline residues at position 189 in cone visual pigments as a determinant of molecular properties different from rhodopsins. *Biochemistry* **41**, 15245–15252.
- KUWAYAMA, S., IMAI, H., MORIZUMI, T. & SHICHIDA, Y. (2005). Amino acid residues responsible for the meta-III decay rates in rod and cone visual pigments. *Biochemistry* **44**, 2208–2215.
- LAMB, T.D. & PUGH, E.N. Jr. (2004). Dark adaptation and the retinoid cycle of vision. *Progress in Retinal and Eye Research* **23**, 307–380.
- LEWIS, J.W., VAN KUIJK, F.J., CARRUTHERS, J.A. & KLIGER, D.S. (1997). Metarhodopsin III formation and decay kinetics: Comparison of bovine and human rhodopsin. *Vision Research* **37**, 1–8.
- LUO, Z.-X., KIELAN-JAWOROWSKA, Z. & CIFELLI, R.L. (2002). In quest for a phylogeny of Mesozoic mammals. *Acta Palaeontologica Polonica* **47**, 1–78.
- MENON, S.T., HAN, M. & SAKMAR, T.P. (2001). Rhodopsin: Structural basis of molecular physiology. *Physiological Reviews* **81**, 1659–1688.
- MORROW, J.M. & CHANG, B.S.W. (2010). The p1D4-hrGFP II expression vector: A tool for expressing and purifying visual pigments and other G protein-coupled receptors. *Plasmid* **64**, 162–169.
- MORROW, J.M., LAZIC, S. & CHANG, B.S.W. (2011). A novel rhodopsin-like gene expressed in zebrafish retina. *Visual Neuroscience* **28**, 325–335.
- NATHANS, J. & HOGNESS, D.S. (1983). Isolation, sequence analysis, and intron-exon arrangement of the gene encoding bovine rhodopsin. *Cell* **34**, 807–814.
- NATOCHIN, M., GASIMOV, K.G., MOUSSALF, M. & ARTEMYEV, N.O. (2003). Rhodopsin determinants for transducin activation: A gain-of-function approach. *The Journal of Biological Chemistry* **278**, 37574–37581.
- NICOL, S. & ANDERSEN, N.A. (2006). Body temperature as an indicator of egg-laying in the echidna, *Tachyglossus aculeatus*. *Journal of Thermal Biology* **31**, 483–490.
- OKADA, T., MATSUDA, T., KANDORI, H., FUKADA, Y., YOSHIZAWA, T. & SHICHIDA, Y. (1994). Circular dichroism of metaiodopsin II and its binding to transducin: A comparative study between meta II intermediates of iodopsin and rhodopsin. *Biochemistry* **33**, 4940–4946.
- OPRIAN, D.D., MOLDAI, R.S., KAUFMANN, R.J. & KHORANA, H.G. (1987). Expression of a synthetic bovine rhodopsin gene in monkey kidney cells. *Proceedings of the National Academy of Sciences of the United States of America* **84**, 8874–8878.
- PALCZEWSKI, K., KUMASAKA, T., HORI, T., BEHNKE, C.A., MOTOSHIMA, H., FOX, B.A., LE TRONG, I., TELLER, D.C., OKADA, T., STENKAMP, R.E., YAMAMOTO, M. & MIYANO, M. (2000). Crystal structure of rhodopsin: A G protein-coupled receptor. *Science* **289**, 739–745.
- PETERS, W. & DORIA, G. (1876). Diagnosi di alcune nuove specie di Marsupiali appartenenti Fauna papuana. *Annali del Museo Civico di Storia Naturale 'Giacomo Doria'* **7**: 541.
- PHILLIPS, M.J., BENNETT, T.H. & LEE, M.S.Y. (2009). Molecules, morphology, and ecology indicate a recent, amphibious ancestry for echidnas. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 17089–17094.
- POSADA, D. (2008). jModelTest: Phylogenetic model averaging. *Molecular Biology and Evolution* **25**, 1253–1256.
- PULAGAM, L.P. & PALCZEWSKI, K. (2010). Electrostatic compensation restores trafficking of the autosomal recessive retinitis pigmentosa E150K opsin mutant to the plasma membrane. *The Journal of Biological Chemistry* **285**, 29446–29456.
- RISMILLER, P. (1999). *The Echidna – Australia's Enigma*. Hong Kong: Hugh Lauter Levin Associates, Inc.
- RISMILLER, P. & MCKELVEY, M.W. (2009). Activity and behaviour of lactating echidnas (*Tachyglossus aculeatus multiaqueatus*) from hatching of egg to weaning of young. *Australian Journal of Zoology* **57**, 265–273.
- RONQUIST, F. & HUELSENBECK, J.P. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- ROWE, T., RICH, T.H., VICKERS-RICH, P., SPRINGER, M. & WOODBURNE, M.O. (2008). The oldest platypus and its bearing on divergence timing of the platypus and echidna clades. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 1238–1242.
- SAKMAR, T.P., MENON, S.T., MARIN, E.P. & AWAD, E.S. (2002). Rhodopsin: Insights from recent structural studies. *Annual Review of Biophysics and Biomolecular Structure* **31**, 443–484.
- SHAW, G. (1792). *Museum Leverianum, Containing Select Specimens from the Museum of the Late Sir Ashton Lever with Descriptions in Latin and English*. London: J. Parkinson.
- SHAW, G. (1799). The duck-billed platypus. In *The Naturalists' Miscellany*. London: F. P. Nodder.
- SHAW, C.N., WILSON, P.J. & WHITE, B.N. (2003). A reliable molecular method of gender determination for mammals. *Journal of Mammalogy* **84**, 123–128.
- SHEN, Y.-Y., LIU, J., IRWIN, D.M. & ZHANG, Y.-P. (2010). Parallel and convergent evolution of the dim-light vision gene RH1 in bats (Order: Chiroptera). *PloS ONE* **5**, e8838.
- SOMMER, M.E. & FARRENS, D.L. (2006). Arrestin can act as a regulator of rhodopsin photochemistry. *Vision Research* **46**, 4532–4546.
- STARACE, D.M. & KNOX, B.E. (1998). Cloning and expression of a *Xenopus* short wavelength cone pigment. *Experimental Eye Research* **67**, 209–220.
- STAVENGA, D.G., SMITS, R.P. & HOENDERS, B.J. (1993). Simple exponential functions describing the absorbance bands of visual pigment spectra. *Vision Research* **33**, 1011–1017.
- SUGAWARA, T., IMAI, H., NIKAI, M., IMAMOTO, Y. & OKADA, N. (2010). Vertebrate rhodopsin adaptation to dim light via rapid meta-II intermediate formation. *Molecular Biology and Evolution* **27**, 506–519.
- TAM, B.M. & MORITZ, O.L. (2009). The role of rhodopsin glycosylation in protein folding, trafficking, and light-sensitive retinal degeneration. *The Journal of Neuroscience* **29**, 15145–15154.
- THOMAS, M.O. (1907). A new *Acanthoglossus* from the island of Salawatti. *Annals and Magazine of Natural History London* **20**, 498–499.
- WAKEFIELD, M.J., ANDERSON, M., CHANG, E., WEI, K.-J., KAUL, R., GRAVES, J.A.M., GRÜTZNER, F. & DEEB, S.S. (2008). Cone visual pigments of monotremes: Filling the phylogenetic gap. *Visual Neuroscience* **25**, 257–264.
- WARREN, W.C., HILLIER, L.W., GRAVES, J.A.M., BIRNEY, E., PONTING, C.P., GRÜTZNER, F., BELOV, K., MILLER, W., CLARKE, L., CHINWALLA, A.T., YANG, S.-P., HEGER, A., LOCKE, D.P., MIETHKE, P., WATERS, P.D., VEYRINES, F., FULTON, L., FULTON, B., GRAVES, T., WALLIS, J., PUENTE, X.S., LÓPEZ-OTÍN, C., ORDÓÑEZ, G.R., EICHLER, E.E., CHEN, L., CHENG, Z., DEAKIN, J.E., ALSOP, A., THOMPSON, K., KIRBY, P., PAPPENFUSS, A.T., WAKEFIELD, M.J., OLENDER, T., LANCET, D., HUTTLER, G.A., SMIT, A.F.A., PASK, A., TEMPLE-SMITH, P., BATZER, M.A., WALKER, J.A., KONKEL, M.K., HARRIS, R.S., WHITTINGTON, C.M., WONG, E.S.W., GEMMELL, N.J., BUSCHIAZZO, E., VARGAS JENTZSCH, I.M., MERKEL, A., SCHMITZ, J., ZEMANN, A., CHURAKOV, G., KRIEGS, J.O., BROSIUS, J., MURCHISON, E.P., SACHIDANANDAM, R., SMITH, C., HANNON, G.J., TSEND-AYUSH, E., McMILLAN, D., ATTENBOROUGH, R., RENS, W., FERGUSON-SMITH, M., LEFÈVRE, C.M., SHARP, J.A., NICHOLAS, K.R., RAY, D.A., KUBE, M., REINHARDT, R., PRINGLE, T.H., TAYLOR, J., JONES, R.C., NIXON, B., DACHEUX, J.-L., NIWA, H., SEKITA, Y., HUANG, X., STARK, A., KHERADPOUR, P., KELLIS, M., FLICEK, P., CHEN, Y., WEBBER, C., HARDISON, R., NELSON, J., HALLSWORTH-PEPIN, K., DELEHAUNTY, K., MARKOVIC, C., MINX, P., FENG, Y., KREMITZKI, C., MITREVA, M., GLASSCOCK, J., WYLIE, T., WOHLDMANN, P., THIRU, P., NHAN, M.N., POHL, C.S., SMITH, S.M., HOU, S., RENFREE, M.B., MARDIS, E.R. & WILSON, R.K. (2008). Genome analysis of the platypus reveals unique signatures of evolution. *Nature* **455**, 175–256.
- WERNEBURG, I. & SÁNCHEZ-VILLAGRA, M.R. (2010). The early development of the echidna, *Tachyglossus aculeatus* (Mammalia:

- Monotremata), and patterns of mammalian development. *Acta Zoologica* **92**, 75–88.
- WOODBURNE, M.O., RICH, T.H. & SPRINGER, M.S. (2003). The evolution of tribospheny and the antiquity of mammalian clades. *Molecular Phylogenetics and Evolution* **28**, 360–385.
- YAN, E.C., KAZMI, M.A., DE, S., CHANG, B.S., SELBERT, C., MARIN, E.P., MATHIES, R.A. & SAKMAR, T.P. (2002). Function of extracellular loop 2 in rhodopsin: glutamic acid 181 modulates stability and absorption wavelength of metarhodopsin II. *Biochemistry* **41**, 3620–3627.
- YOUNG, H.M. & PETTIGREW, J.D. (1991). Cone photoreceptors lacking oil droplets in the retina of the echidna, *Tachyglossus aculeatus* (Monotremata). *Visual Neuroscience* **6**, 409–420.
- ZHAO, H., RU, B., TEELING, E.C., FAULKES, C.G., ZHANG, S. & ROSSITER, S.J. (2009). Rhodopsin molecular evolution in mammals inhabiting low light environments. *PLoS ONE* **4**, e8326.

Supplementary Data

Supplemental materials can be viewed in this issue of VNS by visiting <http://journals.cambridge.org/VNS>.

Table S1. Degenerate primers used for targeting the echidna rhodopsin by PCR based Genome Walking. The primers were designed from other tetrapod rhodopsins.

Table S2. Degenerate primers used in site-directed mutagenesis PCR.

Table S3. Tetrapod alignment including 13 rod opsin sequences used in Bayesian phylogenetic analyses. Accession numbers, obtained from NCBI: common frog (*Rana temporaria*): U59920.1; anole (*Anolis carolinensis*): L31503.1; chicken (*Gallus gallus*): NM_001030606.1; alligator (*Alligator mississippiensis*): U23802.1; platypus (*Ornithorhynchus anatinus*): NM_001082349.1; white-eared opossum (*Caluromys philander*): AY313946.1; fat-tailed dunnart (*Sminthopsis crassicaudatus*): AY159786.2; cat (*Felis felis*): NM_001009242.1; cattle (*Bos taurus*): NM_001014890.1; mouse (*Mus musculus*): NM_145383.1; human (*Homo sapiens*): BC112104.1; crab-eating macaque (*Macaca fascicularis*): S76579. Note the residues highlighted by a gray background, which indicate interesting sites and are discussed in the text.

Table S4. Full coding and non-coding sequence of the echidna rhodopsin. Bold numbers in gray indicate nucleotide positions of exon-intron boundaries.