



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Evidence for juvenile disc herniation in a *Homo erectus* boy skeleton

Häusler, Martin ; Schiess, Regula ; Böni, Thomas

DOI: <https://doi.org/10.1097/BRS.0b013e31827cd245>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-76396>

Journal Article

Accepted Version

Originally published at:

Häusler, Martin; Schiess, Regula; Böni, Thomas (2013). Evidence for juvenile disc herniation in a *Homo erectus* boy skeleton. *Spine*, 38:123-128.

DOI: <https://doi.org/10.1097/BRS.0b013e31827cd245>

Evidence for juvenile disc herniation in *Homo erectus* boy skeleton

Martin Haeusler, PhD, MD^{a,b,c,1}, Regula Schiess, MSc^b & Thomas Boeni KD, MD^{a,c}

^a *Centre for Evolutionary Medicine, Department of Anatomy, University of Zuerich, Winterthurerstrasse 190, 8057 Zuerich, Switzerland*

^b *Anthropological Institute and Museum, University of Zuerich, Winterthurerstrasse 190, 8057 Zuerich, Switzerland*

^c *Orthopaedische Universitaetsklinik Balgrist, Forchstrasse 340, 8008 Zuerich, Switzerland*

¹ Corresponding author:

Centre for Evolutionary Medicine, Department of Anatomy, University of Zuerich,
Winterthurerstrasse 190, 8057 Zuerich, Switzerland

Tel. +41 44 635 55 26, Fax: ++41 44 635 57 99, email: mfh@aim.uzh.ch

Conflicts of Interest: none

ACKNOWLEDGEMENTS

We thank Emma Mbua, Meave Leakey, and the staff of the National Museums of Kenya, Nairobi, as well as the Kenyan Ministry of Education, Science and Technology for access to the Nariokotome boy skeleton. Traveling expenses have been supported by the A.H. Schultz Foundation.

ABSTRACT

Study Design: An analysis and differential diagnosis of bony alterations in the lower lumbar vertebrae of a *Homo erectus* boy skeleton.

Objective: To analyze low back problems during early human evolution.

Summary of Background Data: Back problems in modern humans are often attributed to our upright, bipedal locomotion that is thought to place huge mechanical stresses on the vertebral column. However, little is known of this situation during the course of human evolution.

Methods: We analyzed the lower lumbar spine of the most complete early hominid skeleton, the 1.5 million year old *Homo erectus* boy KNM-WT 15000 from Nariokotome, Kenya, who died at an age of approximately 8 years. We use bony alterations as indirect evidence for disc disease in the absence of soft tissue.

Results: We describe an extensive osteophytic anterior curved remodeling of the left superior articular process of L5 and formation of a new joint at the underside of the left pedicle of L4. This indicates collisional facet joint subluxation, most likely as the result of juvenile traumatic disc herniation.

Conclusions: This indirect evidence of possible juvenile disc herniation in a *Homo erectus* boy skeleton represents the earliest known case of this typical human ailment that is intricately linked to upright bipedalism. The extensive bony remodeling of the articular processes of L4 and L5 suggests that the disc herniation occurred several months before his death. Disabling backache and recurrent sciatica might have at least temporarily restricted his daily activities, which indicates advanced social care and nursing in early *Homo*. We hypothesize that the early *Homo* intervertebral discs were more vulnerable to injury compared to modern humans due to a relatively small vertebral cross-sectional area.

KEY POINTS

- Facet joint subluxation and formation of a new joint between the tip of the superior articular facet of L5 and the underside of the pedicle of L4 probably indicates juvenile disc herniation in a 1.5 million year old *Homo erectus* boy skeleton.
- Disc herniation therefore seems to be a very ancient disease in human prehistory and probably intimately linked to the evolution of bipedal locomotion.
- The extensive bony alterations in the L4-5 facet joint indicates that this *Homo erectus* boy survived a substantial period of time with this potentially disabling condition, which implies some form of advanced social care and nursing at that time.
- The evolution of an increased vertebral cross-sectional area in modern humans might have made our intervertebral discs less vulnerable to injury than in our ancestors.

INTRODUCTION

Disorders of the lower back are the foremost musculoskeletal problem of modern humans. Thus, 60 to 80% of all people are affected by low back pain at some point in their life.¹ They constitute an increasing public health issue in Western societies with immense socioeconomic costs.² However, despite enormous advances in medicine there is no compelling evidence that the prevalence of musculoskeletal disorders of the lower back has changed in the last decades nor is significantly different throughout the world.^{3,4} Nevertheless, a few risk factors have been causally linked to spinal degeneration and low back pain, including genetic predisposition⁵⁻⁹, age and repetitive heavy physical activities, particularly activities that involve bending and twisting.^{7,10-12}

Presumably the most important origin of mechanical strain on our vertebral column stems from our upright, bipedal posture and locomotion, which led some authors to suggest that the true origin of back problems is to be sought in our evolutionary history.^{13,14} In fact, spinal pathologies are remarkably uncommon in our closest relatives, the quadrupedal great apes.¹⁵ On the other hand, a study of bipedal rats suggests that prolonged upright posture induces degenerative changes in intervertebral discs of the rat lumbar spine.¹⁶ Furthermore, a study of monozygotic twins showed that heredity just explains 32% of the variance in disc degeneration at the L4-S1 levels, age an additional 7%, and physical loading exposure an additional 2%.¹⁷ The 59% of the variance that remained unexplained might be attributed to lifetime exposure associated with everyday activity and upright posture and locomotion.¹⁸

However, little is known of the adaptations and failures of the spinal column during the course of human evolution, as vertebrae have rarely survive in the fossil record. Only a handful of fossils preserve more than a few vertebrae. The most complete fossil spinal column

belongs to a juvenile male *Homo erectus* skeleton from Nariokotome, Kenya.^{19,20} His lower lumbar spine presents an unusual pathology^{21,22} that we will analyse here.

MATERIAL AND METHODS

The KNM-WT 15000 *Homo erectus* skeleton was found in 1.53 million year old lake sediments at Nariokotome at the western shore of Lake Turkana, Kenya.^{19,23} It is the best preserved early hominid skeleton discovered to date and most of our knowledge about the anatomy and biology of early *Homo* is based on this exceptionally complete skeleton. It only lacks the first six cervical and two thoracic vertebrae, both radii, and the majority of the hand and foot bones. Pelvis and skull robusticity suggest that it is from a male individual. The long bones imply a stature of 157 cm,²⁴ and an analysis of tooth microanatomy indicates an age at death of approximately eight years.^{25,26} Epiphyseal closure pattern and skeletal development are, however, comparable to 13.5 – 15-year-old modern humans, which entails an ape-like, rapid growth velocity in *H. erectus*.^{27,28} Nevertheless, body shape and proportion of the Nariokotome boy were already very human-like, and the lumbar lordosis was well developed.²⁹⁻³¹ Recently discovered additional vertebral and rib material shows that his lumbar region was composed of five rather than six vertebrae as previously thought.^{20,32-34} His spinal biomechanics, posture and mode of locomotion were therefore probably very similar to modern humans.

RESULTS

A visual analysis of the spinal column of Nariokotome boy discloses striking asymmetries in shape and length of the articular processes of the last two lumbar vertebrae as

the only obvious pathology of the axial skeleton (Fig. 1). The left superior facet of L5 is markedly shortened. Its tip is bent anteriorly and articulates with a nearthrosis at the inferior side of the left pedicle of L4 (Fig. 2). This extra joint is discernable in the form of a prominent eburnated knob, suggesting intra vitam bone remodeling rather than post-mortem deformation as interpreted previously.¹⁹

DIFFERENTIAL DIAGNOSIS

An equivalent condition with a nearthrosis between the superior articular process and the underside of the superjacent pedicle is typical of advanced stages of disc space narrowing.³⁵⁻³⁷ Here, the facet joints of the affected vertebrae may sub-luxate and finally collide, resulting in anterior curved osteophytic remodeling of the tips of the superior articular facets.³⁶ The affliction is often unilateral as in the Nariokotome boy. We attribute the marked asymmetry in length of the other facet joints of the last two lumbar vertebrae to the juvenile age of the Nariokotome boy and the remaining potential for growth, which allowed for a compensative lengthening of the articular processes to offset the obliquity of the vertebral bodies of L4 and L5.

The differential diagnosis of narrowed intervertebral disc space in adolescents includes trauma, degeneration, congenital anomalies and infection. There is no evidence for the latter in the vertebral column of KNM-WT 15000 because in discitis, irregularities of the vertebral bodies would be expected rather than an exclusive involvement of the posterior elements of the vertebrae. However, in accordance to the juvenile age of the specimen, the cartilaginous superior and inferior vertebral growth plates are not preserved.

Several congenital anomalies are associated with juvenile disc herniation, including block and hemivertebrae, spondylolisthesis, asymmetrical sacralisation and lumbalisation, a six-element long lumbar vertebral column, spina bifida and scoliosis.³⁸⁻⁴⁰ None of these anomalies are present in the spine of the Nariokotome boy save for a spina bifida occulta affecting the last three sacral vertebrae, which can be considered a normal variation without clinical relevance, or it might be solely attributable to his juvenile age.²² A previous report on the axial skeleton of Nariokotome boy described distortions of the vertebrae and ribs and interpreted them as indicating severe congenital scoliosis.⁴¹ However, a rearrangements of his ribs based on the new vertebral and rib material revealed a perfectly symmetric rib cage that challenges the diagnosis of a structural scoliosis.^{20,42}

Degeneration and trauma are more likely factors contributing to disc space narrowing. Incipient degenerative processes are often already detectable in intervertebral discs of adolescents.^{6-9,17,18,38,39,43-47} A sudden impact or a fall, particularly with torsion of the flexed spine, could also cause an annulus tear or an endplate fracture. These injuries may induce accelerated disc degeneration, although this has only been shown experimentally.⁴⁸⁻⁵⁰ The massive unilateral osteophytic bone remodeling of the L4–5 facet joint of the *Homo erectus* youth might, however, also suggest a traumatic disc herniation, perhaps with a concomitant traumatic disruption of the left facet joint capsule. In fact, in contrast to adults between 40 and 75% of juvenile disc herniations are traumatic in origin or related to sports injury, rather than degenerative.^{38,39,51-55} The level of the presumed disc herniation of the Nariokotome boy at L4–5 also perfectly fits the frequency distribution of disc herniations in modern human children with 97% occurring at L4–5 and L5–S1.⁵⁵

Facet joint tropism of 16° , which can be observed at the L4–5 level of the Nariokotome boy vertebral column, might be another possible morphological risk factor for disc lesions (Fig. 3). Particularly in adolescent patients this asymmetric sagittal angulation of the facet joints has been related to increased intervertebral shear forces, although its role is controversially discussed.⁵⁶⁻⁵⁸

DISCUSSION

If our interpretation of the telescoping subluxation and formation of a new joint between the two last lumbar vertebrae of the 1.5 million year old Nariokotome *Homo erectus* boy is correct, it provides evidence for the earliest disc lesion in human prehistory. Although there are reports of naturally occurring disc extrusions in certain dog breeds^{59,60} and cats,^{61,62} traumatic disc rupture in a penguin,⁶³ and of disc degeneration in baboons,⁶⁴ macaques⁶⁵ and sand rats,^{66,67} lumbar disc disease is otherwise restricted to humans. This emphasizes the importance of biomechanical factors in its etiology as well as its relation to the evolution of upright bipedalism and lumbar lordosis. *Homo erectus* is the earliest human ancestor with a truly modern body shape and spinal curvature.^{20,29-31} Thus, it would not be unexpected if he already suffered from this typical human disease.

On the other hand, this is, to our knowledge, the first study that deduces traumatic disc lesion from skeletal remains without preserved soft tissue. This is owed to a number of unique circumstances. The nearthrosis between the tip of the superior articular process of L5 and the underside of the pedicle of L4 suggests that concurrent to the disc lesion the Nariokotome boy probably experienced a substantial trauma that may also have disrupted the facet joint capsula. Further, the immature age of the Nariokotome boy and the remaining potential for

growth facilitated bony remodeling of the contiguous articular processes secondary to asymmetrical disc space narrowing and a compensatory scoliosis. Moreover, the extensive curved bony remodeling of the left facet joint L4/5 indicates that the Nariokotome boy survived a considerable period of time with this potentially disabling condition. Patients with similar findings of bony impingement are reported to have a history of between at least six months and several years of recurrent low back pain and sciatica.^{35,68,69}

Although back symptoms only weakly correlate with the amount of pathologic changes it can be hypothesized that the Nariokotome boy was temporarily restricted in walking, bending and other daily activities. In addition, pediatric patients with lumbar disk hernias tend to be more disabled than adults.^{39,52,54} This contributes to evidence for advanced social care in other early *Homo* fossils.^{70,71}

Lumbar disc herniation is common in modern humans with a prevalence of about 2-5% and a peak incidence at age 40 to 45.⁷²⁻⁷⁴ However, in adolescents disc herniation is rare and exceptional in children before puberty.⁷⁵ In fact, so far only ten cases have been reported in the worldwide literature in children younger than ten years.^{52,75} Yet, skeletal rather than chronological age seems to be critical. Although the age at death of the Nariokotome boy is estimated to approximately eight years based on dental microanatomy, epiphyseal fusion pattern rather is comparable to 13 – 15-year-old modern humans.²⁵⁻²⁸ This is in good agreement with the known vulnerable phase for disc lesions during the period of accelerated growth.⁵⁴

An important characteristic of the *Homo erectus* spinal column is a relatively small cross-sectional area of the vertebral bodies compared to those of average modern humans.²⁹ Because stress is directly proportional to the force over the loaded area, a small cross-section

makes vertebrae and discs more susceptible to injury. This might not only account for the disc herniation in the Nariokotome boy, but also for surprisingly frequent Scheuermann's disease in earlier australopithecines.^{76,77} We might therefore speculate that our spinal column has been shaped by a long process of natural selection to become less vulnerable than that of our ancestors. Increased vertebral cross-sectional area might indeed be considered one of the most important vertebral adaptations to habitual bipedal locomotion. The evolutionary history of modern humans should therefore not be blamed for the widespread back problems. Rather, our results corroborate the hypothesis that the vertebral column of modern humans is an optimized compromise between mobility and stability.⁷⁸

REFERENCES

1. Andersson GBJ. Epidemiological features of chronic low-back pain. *Lancet* 1999;354:581-5.
2. Katz JN. Lumbar disc disorders and low-back pain: Socioeconomic factors and consequences. *J Bone Joint Surg Am* 2006;88A:21-4.
3. Waddell G. Low back pain: A twentieth century health care enigma. *Spine* 1996;21:2820-5.
4. Louw QA, Morris LD, Grimmer-Somers K. The prevalence of low back pain in Africa: a systematic review. *BMC Musculoskelet Disord* 2007;8:105.
5. Videman T, Sarna S, Battié MC, et al. The long-term effects of physical loading and exercise lifestyles on back-related symptoms, disability, and spinal pathology among men. *Spine* 1995;20:699-709.

6. Sambrook PN, MacGregor AJ, Spector TD. Genetic influences on cervical and lumbar disc degeneration: a magnetic resonance imaging study in twins. *Arthritis Rheum* 1999;42:366-72.
7. Zhang YG, Sun Z, Zhang Z, et al. Risk factors for lumbar intervertebral disc herniation in Chinese population: a case-control study. *Spine* 2009;34:E918-22.
8. Varlotta GP, Brown MD, Kelsey JL, et al. Familial predisposition for herniation of a lumbar disc in patients who are less than twenty-one years old. *J Bone Joint Surg Am* 1991;73:124-8.
9. Matsui H, Terahata N, Tsuji H, et al. Familial predisposition and clustering for juvenile lumbar disc herniation. *Spine* 1992;17:1323-8.
10. Hoogendoorn WE, van Poppel MN, Bongers PM, et al. Physical load during work and leisure time as risk factors for back pain. *Scand J Work Environ Health* 1999;25:387-403.
11. Suri P, Hunter DJ, Jouve C, et al. Inciting events associated with lumbar disc herniation. *Spine J* 2010;10:388-95.
12. Sørensen IG, Jacobsen P, Gyntelberg F, et al. Occupational and other predictors of herniated lumbar disc disease-a 33-year follow-up in the Copenhagen male study. *Spine* 2011;36:1541-6.
13. Krogman WM. The scars of human evolution. *Sci Am* 1951;185:54-7.
14. Anderson R. Human evolution, low back pain, and dual-level control. In Trevathan WR, Smith EO, McKenna JJ eds. *Evolutionary Medicine*. Oxford: Oxford University Press, 1999:333-49.
15. Jurmain RD. Degenerative joint disease in African great apes: an evolutionary perspective. *J Hum Evol* 2000;39:185-203.

16. Liang QQ, Zhou Q, Zhang M, et al. Prolonged upright posture induces degenerative changes in intervertebral discs in rat lumbar spine. *Spine* 2008;33:2052-8.
17. Battié MC, Videman T, Gibbons LE, et al. 1995 Volvo Award in clinical sciences. Determinants of lumbar disc degeneration. A study relating lifetime exposures and magnetic resonance imaging findings in identical twins. *Spine* 1995;20:2601-12.
18. Battié MC, Videman T, Parent E. Lumbar disc degeneration: epidemiology and genetic influences. *Spine* 2004;29:2679-90.
19. Walker A, Leakey R eds. *The Nariokotome Homo erectus skeleton*. Berlin: Springer, 1993.
20. Haeusler M, Schiess R, Boeni T. New vertebral and rib material point to modern bauplan of the Nariokotome *Homo erectus* skeleton. *J Hum Evol* 2011;61:575-82.
21. Haeusler M. Traumatic spinal injury in the KNM-WT 15000 *Homo erectus* skeleton. *Am J Phys Anthropol* 2012;Suppl. 54:157-8.
22. Schiess R, Haeusler M, Langenegger E. Wie pathologisch ist die Wirbelsäule des Nariokotome Boys KNM-WT 15'000 (*Homo erectus*)? *Bulletin der Schweizerischen Gesellschaft für Anthropologie* 2006;12:13-22.
23. Brown F, Harris J, Leakey R, et al. Early *Homo erectus* skeleton from west Lake Turkana, Kenya. *Nature* 1985;316:788-92.
24. Ruff C. Body size prediction from juvenile skeletal remains. *Am J Phys Anthropol* 2007;133:698-716.
25. Dean MC, Leakey MG, Reid DJ, et al. Growth processes in teeth distinguish modern humans from *Homo erectus* and earlier hominins. *Nature* 2001;414:628-31.

26. Zihlman A, Bolter D, Boesch C. Wild chimpanzee dentition and its implications for assessing life history in immature hominin fossils. *Proc Natl Acad Sci USA* 2004;101:10541-3.
27. Smith BH. The physiological age of KNM-WT 15000. In Walker A, Leakey R eds. *The Nariokotome Homo erectus skeleton*. Berlin: Springer, 1993:195-220.
28. Tardieu C. Short adolescence in early hominids: infantile and adolescent growth of the human femur. *Am J Phys Anthropol* 1998;107:163-78.
29. Latimer B, Ward CV. The thoracic and lumbar vertebrae. In Walker A, Leakey R eds. *The Nariokotome Homo erectus Skeleton*. Berlin: Springer, 1993:266-93.
30. Haeusler M, McHenry HM. Body proportions of *Homo habilis* reviewed. *J Hum Evol* 2004;46:433-65.
31. Haeusler M, McHenry HM. Evolutionary reversals of limb proportions in early hominids? Evidence from KNM-ER 3735 (*Homo habilis*). *J Hum Evol* 2007;53:383-405.
32. Haeusler M, Martelli S, Boeni T. Vertebrae numbers of the early hominid lumbar spine. *J Hum Evol* 2002;43:621-43.
33. Haeusler M, Schiess R, Boeni T. Modern or distinct axial bauplan in early hominins? A reply to Williams (2012). *J Hum Evol* 2012;63:557-9.
34. Williams SA. Modern or distinct axial bauplan in early hominins? Comments on Haeusler et al. (2011). *J Hum Evol* 2012;63:552-6.
35. Hadley LA. Anatomico-roentgenographic studies of the posterior spinal articulations. *Am J Roentgenol Radium Ther Nucl Med* 1961;86:270-6.

36. Jinkins JR. Acquired degenerative changes of the intervertebral segments at and suprajacent to the lumbosacral junction. A radioanatomic analysis of the nondiscal structures of the spinal column and perispinal soft tissues. *Eur J Radiol* 2004;50:134-58.
37. Crock HV. Normal and pathological anatomy of the lumbar spinal nerve root canals. *J Bone Joint Surg Br* 1981;63B:487-90.
38. Børgeesen SE, Vang PS. Herniation of the lumbar intervertebral disk in children and adolescents. *Acta Orthop Scand* 1974;45:540-9.
39. Rugtveit A. Juvenile lumbar disc herniations. *Acta Orthop Scand* 1966;37:348-56.
40. Parisini P, Di Silvestre M, Greggi T, et al. Lumbar disc excision in children and adolescents. *Spine* 2001;26:1997-2000.
41. Latimer B, Ohman JC. Axial dysplasia in *Homo erectus*. *J Hum Evol* 2001;40:A12.
42. Schiess R, Haeusler M, Langenegger E. How pathological is the Nariokotome boy KNM-WT 15000 (*Homo erectus*)? *Am. J. Phys. Anthropol. Suppl.* 2006;42:159.
43. Miller JA, Schmatz C, Schultz AB. Lumbar disc degeneration: correlation with age, sex, and spine level in 600 autopsy specimens. *Spine* 1988;13:173-8.
44. Battié MC, Videman T, Kaprio J, et al. The Twin Spine Study: contributions to a changing view of disc degeneration. *Spine J* 2009;9:47-59.
45. Patel AA, Spiker WR, Daubs M, et al. Evidence for an inherited predisposition to lumbar disc disease. *J Bone Joint Surg Am* 2011;93:225-9.
46. Kumar R, Kumar V, Das NK, et al. Adolescent lumbar disc disease: findings and outcome. *Childs Nerv Syst* 2007;23:1295-9.
47. Powell MC, Wilson M, Szypryt P, et al. Prevalence of lumbar disc degeneration observed by magnetic resonance in symptomless women. *Lancet* 1986;2:1366-7.

48. Hancock MJ, Battie MC, Videman T, et al. The Role of Back Injury or Trauma in Lumbar Disc Degeneration An Exposure-Discordant Twin Study. *Spine* 2010;35:1925-9.
49. Osti OL, Vernon-Roberts B, Fraser RD. 1990 Volvo Award in experimental studies. Anulus tears and intervertebral disc degeneration. An experimental study using an animal model. *Spine* 1990;15:762-7.
50. Hadjipavlou AG, Tzermiadianos MN, Bogduk N, et al. The pathophysiology of disc degeneration: a critical review. *J Bone Joint Surg Br* 2008;90:1261-70.
51. Kurihara A, Kataoka O. Lumbar disc herniation in children and adolescents. A review of 70 operated cases and their minimum 5-year follow-up studies. *Spine* 1980;5:443-51.
52. Fakouri B, Nnadi C, Boszczyk B, et al. When is the appropriate time for surgical intervention of the herniated lumbar disc in the adolescent? *Journal of Clinical Neuroscience* 2009;16:1153-6.
53. Bradbury N, Wilson LF, Mulholland RC. Adolescent disc protrusions. A long-term follow-up of surgery compared to chymopapain. *Spine* 1996;21:372-7.
54. Slotkin JR, Mislow JM, Day AL, et al. Pediatric disk disease. *Neurosurg Clin N Am* 2007;18:659-67.
55. Pietilä TA, Stendel R, Kombos T, et al. Lumbar disc herniation in patients up to 25 years of age. *Neurol Med Chir (Tokyo)* 2001;41:340-4.
56. Cyron BM, Hutton WC. Articular tropism and stability of the lumbar spine. *Spine* 1980;5:168-72.
57. Ishihara H, Matsui H, Osada R, et al. Facet joint asymmetry as a radiologic feature of lumbar intervertebral disc herniation in children and adolescents. *Spine* 1997;22:2001-4.

58. Lee DY, Ahn Y, Lee SH. The influence of facet tropism on herniation of the lumbar disc in adolescents and adults. *J Bone Joint Surg Br* 2006;88:520-3.
59. Goggin JE, Li AS, Franti CE. Canine Intervertebral Disk Disease - Characterization by Age, Sex, Breed, and Anatomic Site of Involvement. *Am J Vet Res* 1970;31:1687-&.
60. Levine JM, Levine GJ, Kerwin SC, et al. Association between various physical factors and acute thoracolumbar intervertebral disk extrusion or protrusion in Dachshunds. *Javma-Journal of the American Veterinary Medical Association* 2006;229:370-5.
61. Muñana KR, Olby NJ, Sharp NJ, et al. Intervertebral disk disease in 10 cats. *J Am Anim Hosp Assoc* 2001;37:384-9.
62. Kathmann I, Cizinauskas S, Rytz U, et al. Spontaneous Lumbar Intervertebral Disc Protrusion in Cats: Literature Review and Case Presentations. *Journal of Feline Medicine and Surgery* 2000;2:207-12.
63. Emerson CL, Eurell JAC, Brown MD, et al. Ruptured intervertebral disc in a juvenile king penguin (*Aptenodytes patagonica*). 1990.
64. Lauerman WC, Platenberg RC, Cain JE, et al. Age-Related Disk Degeneration - Preliminary-Report of a Naturally-Occurring Baboon Model. *J Spinal Disord* 1992;5:170-4.
65. Nuckley DJ, Kramer PA, Del Rosario A, et al. Intervertebral disc degeneration in a naturally occurring primate model: Radiographic and biomechanical evidence. *J Orthop Res* 2008;26:1283-8.
66. Moskowitz RW, Ziv I, Denko CW, et al. Spondylosis in Sand Rats a Model of Intervertebral Disc Degeneration and Hyperostosis. *J Orthop Res* 1990;8:401-11.
67. Silberberg R, Aufdermaur M, Adler JH. Degeneration of the Intervertebral Disks and Spondylosis in Aging Sand Rats. *Arch Pathol Lab Med* 1979;103:231-5.

68. Hadley LA. Apophyseal subluxation. *J Bone Joint Surg Am* 1936;53:428-33.
69. Hadley LA. Intervertebral joint subluxation, bony impingement and foramen encroachment with nerve root changes. *Am J Roentgenol Radium Ther Nucl Med* 1951;65:377-402.
70. Lordkipanidze D, Vekua A, Ferring R, et al. The earliest toothless hominin skull. *Nature* 2005;434:717 - 8.
71. Bonmatí A, Gómez-Olivencia A, Arsuaga JL, et al. Middle Pleistocene lower back and pelvis from an aged human individual from the Sima de los Huesos site, Spain. *Proc Natl Acad Sci USA* 2010;107:18386-91.
72. Heliövaara M. Risk factors for low back pain and sciatica. *Ann Med* 1989;21:257-64.
73. Dammers R, Koehler PJ. Lumbar disc herniation: level increases with age. *Surg Neurol* 2002;58:209-12; discussion 12-3.
74. Kaila-Kangas L, Leino-Arjas P, Karppinen J, et al. History of physical work exposures and clinically diagnosed sciatica among working and nonworking Finns aged 30 to 64. *Spine* 2009;34:964-9.
75. Dang L, Liu Z. A review of current treatment for lumbar disc herniation in children and adolescents. *Eur Spine J* 2010;19:205-14.
76. Cook DC, Buikstra JE, DeRousseau CJ, et al. Vertebral pathology in the Afar australopithecines. *Am J Phys Anthropol* 1983;60:83-101.
77. Haeusler M, Schiess R, Boeni T. Evolutionary adaptations of the hominid vertebral column. 2nd Annual Meeting of the European Society for the study of Human Evolution, Bordeaux, France, September 21-22, 2012, 2012.
78. Putz RLV, Müller-Gerbl M. The vertebral column - a phylogenetic failure? A theory explaining the function and vulnerability of the human spine. *Clin Anat* 1996;9:205-12.

FIGURES




Figure 1. Facet joint subluxation of the two last lumbar vertebrae of the *Homo erectus* boy KNM-WT 15000. A schematic representation of the preserved vertebrae is shown on the left, photographs of the original fossil vertebrae L4 and L5 in inferior and superior view, respectively (middle), and in dorsal view (right). The broken off left lateral process of L4 is graphically reconstructed in semitransparent colours. Note the collisional curved remodelling of the left superior articular facet of L5 (simple arrows) and the nearthrosis it has built at the underside of L4 (double arrow).




Figure 2. A, Digital models of articulated vertebrae L4 and L5 of KNM-WT 15000, in dorsal view. The models have been generated from high-resolution 3D surface scans of casts of the original vertebrae (Polygon Technology, Darmstadt, Germany). The extra joint (nearthrosis, arrow) and the asymmetric articular processes imply an oblique position of the vertebrae relative to each other. B, Section through the nearthrosis (see A), left lateral view. Cropped parts of the vertebrae are hatched. Compare the remodelled superior articular facet of L5 (arrow) to the normal shape of a superior articular facet in L4. Scale bar 3 cm.




Figure 3. Section through the superior articular processes of a digital model of vertebra L5. Cropped parts are hatched. The model has been generated from high-resolution 3D surface scans of a cast of the original vertebra (Polygon Technology, Darmstadt, Germany). Scale bar 3 cm.