



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Methylphenidate enhances neural stem cell differentiation

Bartl, Jasmin ; Mori, Takatoshi ; Riederer, Peter ; Ozawa, Hiroki ; Grünblatt, Edna

Abstract: Background: The psychostimulant methylphenidate (MPH) is the first choice of drug treatment in Attention-Deficit/Hyperactivity Disorder (ADHD). Since therapy often begins at a time when the brain is still developing and the long-term consequences of MPH are still not fully clarified, we examined the influences of an acute treatment with MPH on the differentiation and proliferation of murine neural stem cells (mNSC). Findings and conclusion We found that MPH enhanced neuronal differentiation and inhibited neural proliferation.

DOI: <https://doi.org/10.1186/2049-9256-1-5>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://www.zora.uzh.ch/78277>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 2.0 Generic (CC BY 2.0) License.

Originally published at:

Bartl, Jasmin; Mori, Takatoshi; Riederer, Peter; Ozawa, Hiroki; Grünblatt, Edna (2013). Methylphenidate enhances neural stem cell differentiation. *Journal of Molecular Psychiatry*, 1:5.

DOI: <https://doi.org/10.1186/2049-9256-1-5>

SHORT REPORT

Open Access

Methylphenidate enhances neural stem cell differentiation

Jasmin Bartl^{1*}, Takatoshi Mori², Peter Riederer³, Hiroki Ozawa² and Edna Grünblatt^{1,3,4}

Abstract

Background: The psychostimulant methylphenidate (MPH) is the first choice of drug treatment in Attention-Deficit/Hyperactivity Disorder (ADHD). Since therapy often begins at a time when the brain is still developing and the long-term consequences of MPH are still not fully clarified, we examined the influences of an acute treatment with MPH on the differentiation and proliferation of murine neural stem cells (mNSC).

Findings and conclusion: We found that MPH enhanced neuronal differentiation and inhibited neural proliferation.

Keywords: Methylphenidate, Neural stem cells, Neuronal maturation, Cell proliferation

Findings

Background

ADHD is one of the most frequent psychiatric disorders in children and adolescents, with up to 5% affected worldwide and similar prevalence rates throughout different cultural settings [1,2]. It is characterized by developmentally inappropriate levels of inattention, hyperactivity, and/or impulsivity. These core symptoms lead to impairment in home, school, and peer contexts. Stimulants, particularly MPH, are part of the first-line treatment therapies for ADHD [3] and in recent years, the number of prescriptions has increased nearly tenfold [4]. Mechanistically, MPH is a high-affinity inhibitor of the dopamine transporter and a middle-affinity inhibitor to the norepinephrine transporter [5]. However, given the extent of the prescribed use of MPH [6] exposure to MPH during the early stages of brain development raises some concern for public health due to possible adverse long-term effects such as neurogenesis, neuronal development, or receptor density. It is not clear whether such changes occur; if they do, whether they are related to medication or to the condition itself; if they are caused by medication, whether they have functional significance; and whether any changes are helpful or harmful to mental development. Still, the full mechanism of action of MPH has not yet been elucidated. A recent study by Lee and colleagues (2012) investigated *in vivo* the effect of

chronically treatment of MPH on cell proliferation and neuronal differentiation in adolescent mice brain tissue [7]. They could demonstrate that 10 mg/kg MPH treatment for 28 days enhances cell proliferation as well as neuroblast differentiation in contrast to Lagace and colleagues (2006), who detected an inhibition of survival of adult-generated neurons in the temporal hippocampus of adolescent rats after 16 days treatment with 2 mg/kg of MPH [8]. In concern to the contrasting results and experimental designs of *in vivo* studies, we investigated the *in vitro* effect of an acute treatment with MPH using murine neural stem cells (mNSC) originating from hippocampal tissue of embryonic mice E15. NSC are neurosphere-forming cells and can serve as a model for basic neurodevelopmental processes as well as a potential source of neurodegenerative disease [9]. They are clonogenic, self-renewing, and multipotent cells with plasticity to proliferate and to differentiate into all cell types of the central nervous system (CNS) such as glia/astrocytes, oligodendrocytes, and neurons [10]. Such a model enables the study of neuronal development, differentiation, and neuronal cell death mechanisms *in vitro*.

Material and methods

Murine neural stem cell (mNSC) sphere culturing

mNSCs were derived from the hippocampus tissue of albino mouse (Charles River, Japan) embryos on embryonic day 15. Cells were cultured in a medial hormone mix (MHM) with 10 µg/ml epidermal (EGF; Sigma aldrich, Japan, Cat-No: SRP3196) and 10 µg/ml basic fibroblast growth factors (bFGF; Sigma aldrich, Japan, Cat-No: F5392)

* Correspondence: jasmin.bartl@kjpzdzh.ch

¹Hospital of Child and Adolescent Psychiatry, University of Zurich, Winterthurerstr. 180, Room L84/86, 8057, Zurich, Switzerland
Full list of author information is available at the end of the article

for six days at 37°C and 5% CO₂. After six days, the grown NSC spheres were harvested and prepared for differentiation and/or proliferation studies (for methods, see below).

Differentiation study

mNSC spheres were collected and plated on a 30 µg/ml poly-l-lysine (Sigma aldrich, Japan, Cat-No: P1274) and 20 µg/ml laminine (Invitrogen, Japan, Cat-No: 23017-015) coated 8-well glass cover-slip and incubated for another 4 days. Each well had a final concentration of 22.8x10⁴ cells/ml cultured in MHM containing 1% fetal bovine serum (FBS) and without EGF and bFGF. The cells were treated directly after seeding to the cover-slip using different concentration (0 nM, 1 nM, 10 nM, 100 nM) of methylphenidate (MPH; Sigma aldrich, Japan, Cat-No: M2892). On the last day, cells were fixed with 4% paraformaldehyde (Fluka, Japan, Cat-No: 76240) for 20 minutes at room temperature (RT) and stained with different antibodies (see "Immunocytochemistry").

BrdU incorporation

The proliferation of NSC was identified by in vitro labeling with the thymidine analog 5-bromo-2-desoxyuridine (BrdU; Wako, Japan, Cat-No: 023-15563). NSC spheres were collected and plated on coated 8-well glass cover-slip and incubated for 24 h. Each well had a final concentration of 22.8x10⁴ cells/ml cultured in MHM, containing EGF and bFGF and without FBS. The cells were treated with different concentration of MPH (0 nM, 1 nM, 10 nM, 100 nM). After 24 h, BrdU was added to each well to reach


a final concentration of 10 µM and incubated for 4 h at RT. Before BrdU immunostaining, DNA was denatured and the nucleus-membrane was broken by treating cells with 2M HCL for 35 min at RT; afterward, cells were rinsed with 1% phosphate buffer saline (PBS), followed by treatment with sodium borate (pH=8.5) for 10 min at RT for the neutralization of HCL. The prepared cells were stained against BrdU (see "Immunocytochemistry").


Immunocytochemistry

Fixed and prepared cells were stained for differentiation studies with rabbit monoclonal antibody against glial fibrillary acidic protein (GFAP; 1:200, Sigma aldrich, USA, Cat-No: C4546) and/or with mouse monoclonal antibody against β-tubulin III (Tuj 1; 1:300, Sigma aldrich, USA, Cat-No: T3952) overnight at 4°C. For proliferation studies, cells were stained with rat monoclonal antibody against BrdU (1:100, Accurate Chemical & Scientific, Japan, Cat-No: OBT0030) overnight at 4°C. In both studies, fixed cells were also stained with Hoechst 33258 (1:100; Invitrogen, Cat-No: H3569) for 15 min at RT to visualize the cell nuclei. After overnight incubation, the primary antibodies were visualized with secondary antibodies against rabbit, mouse, or rat conjugated to the following fluorochromes: Alexa Fluor[®] -488 and Alexa Fluor[®] -555 (Life Technologies, Japan).

Statistical analysis

The stainings were analyzed for the cell count of astrocytes, immature neurons and proliferated cells in comparison to the total number of cells using the Mann-






Whitney (U-Rang) Test in the StatView software program (Stat View 5.0. software, SAS Institute Inc. Cary, NC, USA). A p-value < 0.05 was set as significant. There were at least five independent experiments, and four wells per slide were analyzed. In each case, the total amount of Hoechst stained cells/well were manually counted, and afterward, either BrdU positive, GFAP positive, or Tuj 1 positive cells were counted and the percentage of positive cells to the total amount was calculated. In the proliferation and

differentiation studies a comparison of the treated mNSC to the control mNSC (MPH untreated) was done.

Results

After 1 nM MPH treatment, the neuronal differentiation was significantly enhanced, and 53% more immature neurons could be detected compared to untreated mNSCs (Figure 1). After 10 nM of MPH, the enhancement of neuronal differentiation was over 80%, but the highest



dose of MPH (100 nM) showed a reduction of 39% (Figure 1). In contrast, all tested concentration (1-100 nM) of MPH inhibited the proliferation of mNSC (Figure 2). For neuronal maturation, it is important that the neural stem cell stops proliferating and starts the outgrowth of neuritis to develop into full neurons, which seems to be enhanced by the concentration of 1 nM and 10 nM of used MPH. In contrast, MPH treatment at the given dose range did not significantly affect the development of astrocytes in comparison to untreated mNSC (Figure 3).

Availability of supporting data

Discussion

In this work, we presented the effects of low doses (1-100 nM) MPH treatment at mNSC proliferation and neuronal maturation. In 2010 we could already demonstrate, that especially low doses of MPH have the strongest impact on gene expression and cell proliferation in neuronal cell line culture [11]. Our results suggest that MPH seems to support neuronal maturation in a specific range and enhances the neuronal outcome. This hypothesis fits the fact that children with ADHD have atypical or typical but delayed maturation of the prefrontal cortex [12]. Structural imaging in ADHD provides evidence for a global maturational delay based on reduced gray and white matter volume and cortical thickness in ADHD relative to controls through childhood and adolescence [13,14]. MPH seems to balance this retardation of neuronal development, but the mechanism of action is still not known and needs further investigation [15]. However, few studies have addressed the structural correlates of psychostimulant treatment. Castellanos et al. (2002) found that prior treatment with psychostimulants in children with ADHD was associated at study entry with greater white matter lobar volumes relative to stimulant-naïve children with ADHD and volumes that lie closer to the range of their typically developing counterparts, suggesting a neuroprotective effect [16]. An independent study of 30 children with ADHD examining the regions implicated in the pathogenesis of the disorder, similarly, found that treatment with psychostimulants was associated with a more normative volume of the caudate and anterior cingulate cortex [17]. Additionally, the group of Antonello Bonci (2010) recently showed that MPH can influence neuronal plasticity in the amygdala and, thus, can improve learning performance [18]. Within the inherent limitations of a preliminary study, we find great influences of MPH on neuronal maturation, which may prove activity-dependent neuronal plasticity. However, one has to take into account that this *in vitro* experiment did not investigate any MPH influences on neurogenesis, which can be another possible mechanism of action of MPH.

Abbreviations

ADHD: Attention-Deficit/Hyperactivity Disorder; bFGF: basic fibroblast growth factor; BrdU: 5-bromo-2-deoxyuridine; CNS: Central nervous system;

CO₂: Carbon dioxide; EGF: Epidermal growth factor; FBS: Fetal bovine serum; GFAP: Glial fibrillary acidic protein; HCL: Hydrochloric acid; MHM: Medial hormone mix; mNSC: Murine neural stem cell; MPH: Methylphenidate; nM: Nano mol; RT: Room temperature; Tuj1: Beta tubulin III.

Competing interests

The authors declare no competing financial or non-financial interests.

Authors' contributions

JB prepared NSCs, carried out the experiments, performed statistical analyses and wrote the manuscript. TM helped design the experiments and supervised JB in material and methods. PR designed the experiments and conceived the project. HO contributed reagents and supervised JB in Nagasaki, Japan. EG conceived the project, designed experiments, helped write the manuscript and reviewed the statistical analyses. All authors read and approved the final manuscript.

Acknowledgments

Jasmin Bartl was supported by the "Deutscher Akademischer Austausch Dienst (DAAD)" with a travel grant to Japan. The DAAD had no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Author details

¹Hospital of Child and Adolescent Psychiatry, University of Zurich, Winterthurerstr. 180, Room L84/86, 8057, Zurich, Switzerland. ²Division of Neuropsychiatry, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki, Japan. ³Department of Psychiatry, Psychosomatics and Psychotherapy, University Hospital of Wuerzburg, Wuerzburg, Germany. ⁴Neuroscience Center Zurich, University of Zurich and ETH Zurich, Switzerland.

Received: 23 November 2012 Accepted: 18 January 2013

Published: 23 April 2013

References

1. Biederman J, Faraone SV: Attention-deficit hyperactivity disorder. *Lancet* 2005, **366**(9481):237-248.
2. Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA: The worldwide prevalence of ADHD: a systematic review and meta-regression analysis. *Am J Psychiatry* 2007, **164**(6):942-948.
3. Gunther T, Herpertz-Dahlmann B, Konrad K: Sex differences in attentional performance and their modulation by methylphenidate in children with attention-deficit/hyperactivity disorder. *J Child Adolesc Psychopharmacol* 2010, **20**(3):179-186.
4. Schubert I, Koster I, Lehmkuhl G: The changing prevalence of attention-deficit/hyperactivity disorder and methylphenidate prescriptions: a study of data from a random sample of insureds of the AOK Health Insurance Company in the German State of Hesse, 2000-2007. *Dtsch Arztebl Int* 2010, **107**(36):615-621.
5. Han DD, Gu HH: Comparison of the monoamine transporters from human and mouse in their sensitivities to psychostimulant drugs. *BMC Pharmacol* 2006, **6**:6.
6. Bogle KE, Smith BH: Illicit methylphenidate use: a review of prevalence, availability, pharmacology, and consequences. *Curr Drug Abuse Rev* 2009, **2**(2):157-176.
7. Lee TH, Lee CH, Kim IH, Yan BC, Park JH, Kwon SH, Park OK, Ahn JH, Cho JH, Won MH, Kim SK: Effects of ADHD therapeutic agents, methylphenidate and atomoxetine, on hippocampal neurogenesis in the adolescent mouse dentate gyrus. *Neurosci Lett* 2012, **524**(2):84-88.
8. Lagace DC, Yee JK, Bolanos CA, Eisch AJ: Juvenile administration of methylphenidate attenuates adult hippocampal neurogenesis. *Biol Psychiatry* 2006, **60**(10):1121-1130.
9. Tropepe V, Sibilia M, Ciruna BG, Rossant J, Wagner EF, van der Kooy D: Distinct neural stem cells proliferate in response to EGF and FGF in the developing mouse telencephalon. *Dev Biol* 1999, **208**(1):166-188.
10. Onorati M, Camnasio S, Binetti M, Jung CB, Moretti A, Cattaneo E: Neurotropic self-renewing neural stem (NS) cells derived from mouse induced pluripotent stem (iPS) cells. *Mol Cell Neurosci* 2010, **43**(3):287-295.

11. Bartl J, Link P, Schlosser C, Gerlach M, Schmitt A, Walitza S, Riederer P, Grunblatt E: **Effects of methylphenidate: the cellular point of view.** *Atten Defic Hyperact Disord* 2010, **2**(4):225–232.
12. Vaidya CJ: **Neurodevelopmental abnormalities in ADHD.** *Curr Top Behav Neurosci* 2012, **9**:49–66.
13. Seidman LJ, Valera EM, Makris N: **Structural brain imaging of attention-deficit/hyperactivity disorder.** *Biol Psychiatry* 2005, **57**(11):1263–1272.
14. Mostofsky SH, Rimrod SL, Schafer JG, Boyce A, Goldberg MC, Pekar JJ, Denckla MB: **Atypical motor and sensory cortex activation in attention-deficit/hyperactivity disorder: a functional magnetic resonance imaging study of simple sequential finger tapping.** *Biol Psychiatry* 2006, **59**(1):48–56.
15. Rubia K, Halari R, Christakou A, Taylor E: **Impulsiveness as a timing disturbance: neurocognitive abnormalities in attention-deficit hyperactivity disorder during temporal processes and normalization with methylphenidate.** *Philos Trans R Soc Lond B Biol Sci* 2009, **364**(1525):1919–1931.
16. Castellanos FX, Lee PP, Sharp W, Jeffries NO, Greenstein DK, Clasen LS, Blumenthal JD, James RS, Ebens CL, Walter JM, Zijdenbos A, Evans AC, Giedd JN, Rapoport JL: **Developmental trajectories of brain volume abnormalities in children and adolescents with attention-deficit /hyperactivity disorder.** *JAMA* 2002, **288**(14):1740–1748.
17. Semrud-Clikeman M, Pliszka SR, Lancaster J, Liotti M: **Volumetric MRI differences in treatment-naive vs chronically treated children with ADHD.** *Neurology* 2006, **67**(6):1023–1027.
18. Tye KM, Tye LD, Cone JJ, Hekkelman EF, Janak PH, Bonci A: **Methylphenidate facilitates learning-induced amygdala plasticity.** *Nat Neurosci* 2010, **13**(4):475–481.

doi:10.1186/2049-9256-1-5

Cite this article as: Bartl et al.: Methylphenidate enhances neural stem cell differentiation. *Journal of Molecular Psychiatry* 2013 1:5.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

