Year: 2013

Pure bipolar plasma vaporization of the prostate: The Zürich experience

Kranzbühler, Benedikt ; Wettstein, Marian Severin ; Fankhauser, Christian Daniel ; Grossmann, Nico Christian ; Poyet, Cédric ; Largo, Remo ; Fischer, Boris ; Zimmermann, Matthias ; Sulser, Tullio ; Müller, Alexander ; Hermanns, Thomas

Abstract: Introduction and objectives: Bipolar plasma vaporization (BPV) has been introduced as an alternative to transurethral resection of the prostate (TURP). Promising short-term results but inferior mid-term results compared to TURP have been reported following first-generation bipolar electrovaporization. Outcome data following second-generation BPV is still scarce. The aim of this investigation was to evaluate the intra- and post-operative outcome of contemporary BPV in a center with long-standing expertise on laser vaporization of the prostate. Methods: A consecutive series of 83 patients undergoing BPV in a tertiary referral center was prospectively evaluated. The investigated outcome parameters included the maximum flow rate (Q_{max}), postvoid residual volume, IPSS / QoL and PSA tests. Follow-up investigations took place after six weeks, six months and twelve months. Wilcoxon signed-rank test was used to compare pre- and post-treatment parameters. Results: The median (range) pre-operative prostate volume was 41ml (17-111ml). The pre-operative IPSS, QoL, Q_{max} and residual volume were 16 (2-35), 4 (0-6), 10.1ml/sec (3-29.3ml/sec) and 87ml (0-1000ml), respectively. One third of the patients were undergoing platelet aggregation inhibition. No intra-operative complications occurred. Post-operatively, 13 patients (15.7%) had to be re-catheterized. Three patients (3.6%) had clot retention and 28 patients (34%) reported any grade of dysuria. After six weeks, all outcome parameters improved significantly and remained improved over the twelve months observation period (IPSS: 3 (0-2); QoL: 1 (0-4); Q_{max}: 17.2ml/s (3.2-56ml/s); residual volume 11ml (0-190ml)). The PSA-reduction was 60% at study conclusion. Three patients (3.6%) developed a urethral stricture and four patients (4.8%) bladder neck sclerosis. Re-resections were not necessary. Conclusions: Contemporary BPV is a safe and efficacious treatment option even for patients undergoing platelet aggregation inhibition. Early urinary retention and temporary dysuria seem to be specific side effects of the treatment. Bleeding complications are rare. Long-term follow-up is needed to confirm these promising short-term results.

DOI: <https://doi.org/10.1089/end.2013.0335>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-79321>

Journal Article

Accepted Version

Originally published at:

Kranzbühler, Benedikt; Wettstein, Marian Severin; Fankhauser, Christian Daniel; Grossmann, Nico Christian; Poyet, Cédric; Largo, Remo; Fischer, Boris; Zimmermann, Matthias; Sulser, Tullio; Müller, Alexander; Hermanns, Thomas (2013). Pure bipolar plasma vaporization of the prostate: The Zürich experience. *Journal of Endourology*, 27(10):1261-1266.

DOI: <https://doi.org/10.1089/end.2013.0335>

Pure bipolar plasma vaporization of the prostate: The Zürich experience

Benedikt Kranzbühler (benedikt.kranzbuehler@usz.ch)

Marian Severin Wettstein (marianseverin.wettstein@usz.ch)

Christian D. Fankhauser (christian.fankhauser@usz.ch)

Nico Grossmann (nico.grossmann@usz.ch)

Oliver Gross (oliver.gross@usz.ch)

Cédric Poyet (cedric.poyet@usz.ch)

Remo Largo (remo.largo@usz.ch)

Boris Fischer (boris.fischer@usz.ch)

Matthias Zimmermann (matthias.zimmermann@usz.ch)

Tullio Sulser (tullio.sulser@usz.ch)

Alexander Müller (alex.mueller@usz.ch)

Thomas Hermanns* (thomas.hermanns@usz.ch)

Department of Urology, University Hospital Zürich, University of Zürich, Switzerland

*Corresponding author:

Department of Urology

University Hospital Zürich

Frauenklinikstr. 10

8091 Zürich

Switzerland

Phone: +41-44-2551111

Fax: +41-44-2554566

Running title: Bipolar plasma vaporization of the prostate

Introduction and objectives: Bipolar plasma vaporization (BPV) has been introduced as an alternative to transurethral resection of the prostate (TURP). Promising short-term results but inferior mid-term results compared to TURP have been reported following first-generation bipolar electro-vaporization. Outcome data following second-generation BPV is still scarce. The aim of this investigation was to evaluate the intra- and post-operative outcome of contemporary BPV in a center with long-standing expertise on laser vaporization of the prostate.

Methods: A consecutive series of 83 patients undergoing BPV in a tertiary referral center was prospectively evaluated. The investigated outcome parameters included the maximum flow rate (Qmax), postvoid residual volume, IPSS / Qol and PSA tests. Follow-up investigations took place after six weeks, six months and twelve months. Wilcoxon signed-rank test was used to compare pre- and post-treatment parameters.

Results: The median (range) pre-operative prostate volume was 41ml (17-111ml). The pre-operative IPSS, Qol, Qmax and residual volume were 16 (2-35), 4 (0-6), 10.1ml/sec (3-29.3ml/sec) and 87ml (0-1000ml), respectively. One third of the patients were undergoing platelet aggregation inhibition. No intra-operative complications occurred. Post-operatively, 13 patients (15.7%) had to be re-catheterized. Three patients (3.6%) had clot retention and 28 patients (34%) reported any grade of dysuria. After six weeks, all outcome parameters improved significantly and remained improved over the twelve months observation period (IPSS: 3 (0-2); Qol: 1 (0-4); Qmax: 17.2ml/s (3.2-56ml/s); residual volume 11ml (0-190ml)). The PSA-reduction was 60% at study conclusion. Three patients (3.6%) developed a urethral stricture and four patients (4.8%) bladder neck sclerosis. Re-resections were not necessary.

Conclusions: Contemporary BPV is a safe and efficacious treatment option even for patients undergoing platelet aggregation inhibition. Early urinary retention and temporary dysuria seem to be specific side effects of the treatment. Bleeding complications are rare. Long-term follow-up is needed to confirm these promising short-term results.

Introduction

Monopolar transurethral resection of the prostate (TURP) has been shown to be an efficacious and durable treatment option for patients with lower urinary tract symptoms caused by prostatic enlargement¹⁻³. It is considered the surgical reference standard for patients with a prostate volume of less than 80ml⁴. However, despite many technical refinements over the last decades it is still associated with a relevant rate of complications mainly caused by bleeding and the so-called TUR-syndrome^{2, 5, 6}. The morbidity of TURP and the growing number of patients with significant cardiovascular co-morbidities or with an imperative indication for platelet aggregation inhibition (PAI) or anti-coagulation prompted the development of novel, minimally invasive alternatives. Bipolar TURP and 532nm laser vaporization are two safe and efficacious minimally invasive treatment options. They both are characterized by improved hemostatic properties and the possibility to perform intra-operative irrigation with isotonic saline. A lower rate of bleeding complications compared to conventional TURP and excellent clinical outcomes have been reported for these procedures⁷⁻⁹. Prostate vaporization using the 532nm laser is even safe in patients undergoing PAI or anti-coagulation¹⁰.

Bipolar plasma vaporization (BPV) of the prostate has been introduced to further improve the hemostatic properties of bipolar TURP. Contemporary BPV using a hemispherical vaporization electrode combines the advantages of the bipolar technique (affordable equipment, saline irrigation) with the benefits of the 532nm laser technique (tissue vaporization, reduced bleeding, excellent overview and short learning curve). However, outcome data following contemporary BPV of the prostate is still scarce. Initial reports are very promising with low intra- and peri-operative morbidity and functional short-term results comparable or even superior to TURP¹¹⁻¹³. The aim of the present investigation was to evaluate the intra-, peri- and postoperative outcome of the initial BPV series of our center, which has a long-standing expertise on prostate vaporization using the 532nm laser.

Patients and methods

A consecutive series of patients undergoing BPV for prostatic bladder outlet obstruction in a tertiary referral centre between August 2009 and November 2011 was prospectively evaluated. Approval for this observational study was obtained from the local ethics committee and all enrolled patients provided written informed consent.

The attending urologist generally made the indication for BPV after unsuccessful medical treatment or if the patient had refused medical therapy for his symptomatic prostatic bladder outlet obstruction. BPV was offered to all patients without PAI or anticoagulation but also to those undergoing PAI.

Pre-operatively, patients were asked to complete the International Prostate Symptom Score (IPSS) and quality of life (QoL) questionnaire. Transrectal ultrasound of the prostate, uroflowmetry and postvoid residual volume measurement were performed. The pre-operative blood work included a prostate specific antigen (PSA) test. Patients with a PSA value >4 ng/ml or a suspicious digital rectal examination underwent BPV only if a pre-operative prostate biopsy showed no malignant results. Urinalysis and a urine culture were also routinely performed.

All patients with a normal pre-operative urine status received intravenous single-shot antibiotic prophylaxis with Trimethoprim-Sulfamethoxazole 160/80mg 30-60 minutes prior to the operation. In patients with significant leukocyturia, antibiotic treatment was initiated pre-operatively and continued for at least five days after catheter removal.

The procedure was carried out under either general or spinal anesthesia. A SurgMaster UES-40 generator and the hemispherical bipolar HF-vaporization electrode were used in combination with a 24-F OES continuous-flow Iglesias resectoscope (all Olympus Winter & Ibe GmbH, Hamburg, Germany). The power output used for tissue vaporization and tissue coagulation was 290-320W and 150-170W, respectively. The operation was performed using

the vaporization mode and coagulation was only used selectively for localized bleeding. All procedures were carried out under continuous low-pressure irrigation using an automated irrigation-suction pump (Endo Fluid Management System Urology, Future Medical System SA, Genève, Switzerland) and pre-warmed isotonic saline (37°C).

The operation was carried out analogous to a conventional TURP. After insertion of the cystoscope and careful inspection of the prostate and bladder, the ureteral orifices were identified. Vaporization was initiated at the bladder neck to ablate the median lobe of the prostate. The procedure was continued at the lateral lobes. Finally, the anterior part of the prostate and the apical / para-collicular region were vaporized until the appearance of a TURP-like cavity. Extensive coagulation of the cavity at the end of the operation was generally avoided. In patients with a prostate volume of less than 30ml an additional bladder neck incision was regularly performed. At the end of the procedures a 20-F three-way irrigation catheter was inserted and continuous irrigation with isotonic saline was initiated. The operative time (from insertion of the cystoscope to insertion of the catheter), the total amount of irrigation fluid and all intra-operative complications were recorded.

Generally, the catheter was removed after three days. In the case of persisting hematuria, catheter-removal was postponed. After removal of the catheter uroflowmetry and postvoid-residual volume measurements were done. Peri-operative complications, the duration of catheterization as well as re-catheterizations were recorded.

The patients were regularly seen in follow-up after six weeks, six months and twelve months in the outpatient clinic. At each follow-up visit uroflowmetry, residual volume measurements and a PSA test were performed. Additionally, the IPSS and Qol questionnaire was completed. Furthermore, the patients were asked to report symptoms of dysuria.

All data are presented as median and range. Statistical analysis was done using IBM SPSS Statistics software 20.0 (IBM, Armonk, NY, USA). The pre-operative variables were

compared to post-operative variables using the Wilcoxon signed-rank test. All p-values <0.05 were considered statistically significant.

Results

A total of 83 patients were included in this study. The pre-operative patients characteristics are displayed in Table 1. Twenty-six patients (31.3 %) were operated under ongoing PAI with acetylsalicylic acid. One patient had dual PAI with additional clopidrogel medication. In all six patients with oral anticoagulation the coumarin medication was peri-operatively replaced by low molecular weight heparin at therapeutical doses. Heparin was stopped 24 hours before the operation and re-initiated with sub-therapeutical doses after the procedure. It was then gradually increased and replaced by coumarin if no macrohematuria occurred.

All patients underwent the pre-operative, initial post-operative and six weeks assessment. A total of 75 (90%) and 71 (86%) patients were available for the evaluation after six and twelve months, respectively.

The intra- and peri-operative results are shown in Table 2. BPV was performed by a total of five surgeons. Three senior surgeons did 67 procedures (81%) and two senior residents did 16 supervised procedures (19%) in form of a teaching operation. Pure BPV was successfully completed in 82 patients (98.8%). Conversion to conventional bipolar TURP was necessary in one patient with a prostate volume of 110ml due to intra-operative diffuse hemorrhage, which resulted in poor visibility. The patient was not undergoing PAI but was under antibiotic therapy for an asymptomatic urinary tract infection. Major intra-operative complications did not occur and intra- or peri-operative blood transfusions were not necessary in any of the patients.

A total of 32 patients (36%) had a positive pre-operative urine culture. Nine of these patients had an unsuspecting urine status. Post-operatively, a positive urine culture was found in nine

patients who subsequently were treated with antibiotics. Only one of these patients had an indwelling catheter pre-operatively. None of the patients developed symptoms of a systemic urinary tract infection or an urosepsis.

Four of the thirteen patients who needed post-operative re-catheterization had an indwelling catheter pre-operatively. Three patients were discharged with a catheter of which two were successfully removed after thirteen days. In one patient with persistent urinary retention a suprapubic cystostomy was performed for permanent solution. Peri-operatively, one patient developed gross hematuria with clot retention. Following cystoscopic clot evacuation transurethral coagulation of the ablation cavity was undertaken. One patient developed a small bowel volvulus associated with an adhesive strangulation three days after the operation. After laparotomy a segment resection and entero-enterostomy was performed. Seven days later an anastomotic leak was detected and a revision of the entero-enteostomy was necessary. The patient was discharged 36 days after BPV.

Table 3 summarizes the post-operative outcome parameters. The maximum flow rate (Q_{max}) and the residual volume were already investigated after catheter removal. The Q_{max} increased non-significantly to 14ml/s ($p=0.06$), whereas the reduction of the residual volume was already statistically significant at this early post-operative assessment ($p<0.001$). After six weeks all functional outcome parameters improved significantly compared to the baseline values (Wilcoxon $p<0.001$ (IPPS, Qol, residual volume) and Wilcoxon $p=0.006$ (Q_{max})). A further significant improvement was detectable after six months for the IPSS and Qol and after twelve month for the IPSS. All parameters remained significantly improved over the entire twelve months observation period (Table 3). The PSA value was already significantly reduced after six weeks ($p<0.001$). After twelve months a 60% reduction of the initial PSA value was detectable (Table 3).

Table 4 illustrates further post-operative results. Of the 15 patients who were found to have a positive urine culture six weeks after surgery, six had an indwelling catheter and ten a positive urine culture pre-operatively. All patients were treated with oral antibiotics. One patient was diagnosed with a pyelonephritis one month after the operation. The urine was found to be positive for mycobacterium tuberculosis. An initial four-drug regimen was followed by a three-drug treatment for nine months. Subsequently, mycobacteria were not detectable anymore. Four patients were diagnosed with a urinary tract infection between the six weeks and six months follow-up. Afterwards, urinary tract infections were not diagnosed anymore. Two patients (2.4%) presented with delayed hematuria and clot retention after four and six weeks, respectively. Both patients were hospitalized for cystoscopic clot evacuation and subsequent coagulation. One of these patients was undergoing dual PAI medication and therefore, laser coagulation of the ablation cavity was performed. In three symptomatic patients (4%) a *de novo* urethral stricture was diagnosed and treated by visual internal urethrotomy after six and eight months and by a simple dilatation procedure after eleven months, respectively. All of the four patients who developed a symptomatic bladder neck sclerosis were treated by transurethral bladder neck incision after 2 months (n=1) four months (n=2) and seven months (n=1), respectively. Only one of the patients had an initial prostate volume of less than 30ml and none of the patients had a urinary tract infection pre- or post-operatively.

Discussion

This prospective investigation of intra-, peri- and post-operative outcomes following pure contemporary BPV of the prostate revealed that the procedure is safe, has a low morbidity and results in significant improvements of all investigated outcome parameters. No major intra-operative complications occurred and conversion to conventional bipolar transurethral

resection was necessary in only one patient with a large prostate. Post-operative bleeding complications, which required re-interventions but no transfusions, occurred in three patients of whom one was undergoing dual PAI. Interestingly, two of these complications occurred with a delay of four to six weeks after the operation. Urinary tract infections were relatively frequent pre-operatively but also in the early post-operative period. One third of the patients reported dysuria after six weeks. After 6 months urinary symptoms were generally rare. Re-operations due to persistent obstructive tissue were not necessary but seven patients (8%) were treated for symptomatic urethral strictures (n=3) or bladder neck sclerosis (n=4).

Kaplan and Te introduced electro-vaporization of the prostate using a rollerball electrode in 1995¹⁴. Despite promising functional results, this monopolar technique was soon abandoned due to higher rates of irritative voiding symptoms and stress urinary incontinence compared to conventional TURP^{11, 15}. In 2001 Botto and colleagues reported their initial results of a bipolar vaporization technique using a special bar electrode¹⁶. Subsequently several studies revealed a lower intra- and perioperative morbidity and functional short-term results comparable to conventional TURP¹⁷⁻²⁰. However, mid-term results have been reported to be inferior compared to TURP^{7, 21}.

In 2008 a novel bipolar device using an ergonomic, hemispherical electrode and a different energy setting was launched²². To date, only limited data is available for this contemporary BPV technique. In 2010 Reich and colleagues reported their initial clinical experience of a very small series of 30 patients with a follow-up of six months¹¹. Four different surgeons from two centers did the operations. Their short-term functional results were comparable to the results of the present study. Bleeding complications did not occur and one patient with a large initial prostate volume required a re-operation four weeks after the initial procedure.

Robert and colleagues reported a French multicenter observational study of 106 consecutive patients with a short-term follow-up of three months¹³. Patients were recruited from eight

different centers. A maximum of three surgeons per center performed the operations resulting in an average of only seven operations per surgeon. Improvements of the IPSS and bother score were comparable to the improvements observed in the present study. However a high failure rate of eight percent after three months indicates that tissue ablation might have been insufficient in a relevant proportion of patients in this study. Additionally, three major complications (urethral necrosis with urinary fistulas (n=2) and significant bladder necrosis (n=1)) occurred in three consecutive patients from one center. These complications were considered to be associated with material failure. The case of a small bowel volvulus requiring a segment resection represents a serious complication in our series. It is likely that the development of the volvulus is somehow related to the initial surgery. However, we do not believe that this complication is specifically related to the BPV procedure.

The same group from France reported a *post-hoc* retrospective comparison of 54 patients under oral anticoagulation from their initial study and 57 patients undergoing TURP under oral anticoagulation²³. Comparable functional results after three months but significantly less bleeding complications after BPV were seen. It is noteworthy that most of the patients were not under oral anticoagulation but under PAI. Furthermore, PAI was stopped in the majority of the patients and oral anticoagulation was replaced by heparin in all patients prior to surgery. In the present investigation, 30% of the patients were under ongoing PAI. Intra-operative bleeding complications did not occur in these patients. However, dual PAI and urinary tract infections might increase the risk of intra- and perioperative bleeding complications.

Two randomized-trials comparing BPV with monopolar and bipolar TURP were conducted by Geavlete and colleagues^{12, 24}. In their first study they reported six-months results of 155 patients undergoing either BPV (n=75) or monopolar TURP (n=80) done by a single surgeon²⁴. The operative time, catheterization period and hospital stay were significantly

shorter in the BPV arm. Furthermore, all relevant outcome parameters were in favor of BPV and the complication rate was significantly lower following BPV compared to TURP. The mean operative time was 35.1min and the mean prostate volume decreased from 56.2ml to 16.8ml after six months. Compared to the present investigation, the Qmax in their study was slightly higher and the re-catheterization rate and dysuria rate lower. Urethral strictures and re-operations were not reported.

The second trial by the same group indicated to evaluate long-term results of a randomized trial of 510 patients who underwent BPV (n=170), monopolar TURP (n=170) and bipolar TURP (n=170)¹². However, with a follow-up of only 18-months, we think that the results of this trial represent rather short-term than long-term results. This large-scale single center trial revealed a clear benefit for BPV in terms of complications and clinical outcome. The intra-operative complication rate was significantly lower in the BPV arm compared to both, the monopolar and the bipolar TURP arm. Post-operative irritative voiding symptoms and urethral strictures were comparable between the three groups. However, a significantly lower re-catheterization rate, a lower rate of bladder neck sclerosis and fewer re-treatments were detectable following BPV. Furthermore, improvements of IPSS, Qol and Qmax were significantly better in the BPV arm. It remains unclear why the bipolar TURP arm and the additional 95 and 90 patients in the BPV and monopolar TURP arm, respectively, were not reported in the initial series which was published only 12 months earlier. Long-term follow-up will show if these excellent results are durable.

In the present study the duration of catheterization was significantly longer compared to the other studies^{11, 13, 24}. This difference is not related to a higher rate of bleeding complications but is mainly due to local reimbursement policies in Switzerland. The present investigation also revealed a lower Qmax compared to the studies by Geavlete and colleagues^{12, 24}. The improvement of the Qmax in the present investigation is comparable to what has been

reported in the studies by Reich and Robert, respectively^{11, 13}. However, it seems also to be significantly lower than the improvement reported following monopolar TURP^{7, 24}. All other investigated functional outcome parameters are comparable to the results reported after BPV and monopolar TURP. The studies of Geavlete and colleagues also showed a lower complication rate compared to the present investigation. We found more urinary tract infections, re-catheterizations and higher rates of dysuria and bladder neck sclerosis. Dysuria, early post-operative urinary retention and urinary tract infections have been reported to be frequent after 532nm laser vaporization of the prostate⁷. The higher rate of these complications in the present study might indicate that these complications are associated with prostate vaporization in general. Given the wide range of the reported rate of bladder neck sclerosis following BPV among different studies (0.6% to 7.2%), the rate in the present study appears to be rather average^{12, 25}. Furthermore, it seems to be comparable to what has been reported for monopolar and bipolar TURP^{26, 27}. In our collective, re-treatments were not necessary during the twelve months observation period. This is lower compared to published BPV and TURP data. A longer follow-up has to confirm this promising result.

The generally excellent coagulation properties of the bipolar vaporization device result in an excellent overview during the procedure and, in our opinion, make this treatment modality particularly appealing for teaching operations. However, we also realized that the procedure has to be performed in a speedy manner in order to achieve sufficient tissue ablation in a reasonable time. The experience of the surgeon is likely to influence the speed and efficiency of tissue vaporization during BPV. In the study by Geavlete, a single surgeon did all procedures. This might explain the excellent ablation rate of approximately 40ml in 35 minutes and the above-average clinical outcome²⁴. The high retreatment rate in the French study might reflect the insufficient tissue ablation of surgeons who were rather inexperienced using this technique¹³. All senior surgeons in the present study were experienced in prostate

vaporizations using the 532nm laser. This experience might improve the vaporization efficiency particularly in the learning phase of the BPV procedure. The PSA reduction of 60% and the lack of necessity for re-operations substantiate this assumption.

A drawback of the present investigation is the lack of a control group, which limits the ability to compare the results with those of other techniques. However, given the sparsely available data of contemporary BPV, our study adds important information to the published evidence. Furthermore, we did not design and power this study for subgroup analyses, which might allow identifying predictors of clinical outcome following BPV. A larger prospective investigation would be required to formally identify these predictors and our results can be helpful in designing such a study. The long-standing experience with prostate vaporization techniques in our center, the longer follow-up period, the larger number of patients and the lower number of surgeons who performed the procedure compared to most of the other BPV studies are specific advantages of our investigation.

Conclusion

Contemporary BPV is a safe and efficacious treatment option even for patients undergoing PAI. BPV results in significant improvements of all relevant outcome parameters over a twelve months period. Bleeding complications are rare. Early urinary retention and temporary dysuria seem to be specific side effects of the treatment. A multi-institutional study with long-term follow-up is needed to confirm the promising short-term results and to further evaluate the durability of the procedure.

Acknowledgements

The authors acknowledge Ms. Alexandra Veloudios for the excellent organization of the patient care during this study.

Author disclosure statement

The authors declare that no competing financial interests exist.

References:

1. Madersbacher S, Lackner J, Brossner C, et al. Reoperation, myocardial infarction and mortality after transurethral and open prostatectomy: a nation-wide, long-term analysis of 23,123 cases. *Eur Urol*. 2005 Apr;47(4):499-504.
2. Reich O, Gratzke C, Bachmann A, et al. Morbidity, mortality and early outcome of transurethral resection of the prostate: a prospective multicenter evaluation of 10,654 patients. *The Journal of Urology*. 2008 Jul;180(1):246-9.
3. Roos NP, Wennberg JE, Malenka DJ, et al. Mortality and reoperation after open and transurethral resection of the prostate for benign prostatic hyperplasia. *N Engl J Med*. 1989 Apr 27;320(17):1120-4.
4. Oelke M, Bachmann A, Descazeaud A, et al. EAU Guidelines on the Treatment and Follow-up of Non-neurogenic Male Lower Urinary Tract Symptoms Including Benign Prostatic Obstruction. *Eur Urol*. 2013 Mar 13
5. Hahn RG. Fluid absorption in endoscopic surgery. *British Journal of Anaesthesia*. 2006 Jan;96(1):8-20.
6. Mebust WK, Holtgrewe HL, Cockett AT, Peters PC. Transurethral prostatectomy: immediate and postoperative complications. Cooperative study of 13 participating

- institutions evaluating 3,885 patients. *J Urol*, 141: 243-247, 1989. *The Journal of Urology*. 2002 Jan;167(1):5-9.
7. Ahyai SA, Gilling P, Kaplan SA, et al. Meta-analysis of functional outcomes and complications following transurethral procedures for lower urinary tract symptoms resulting from benign prostatic enlargement. *Eur Urol*. 2010 Sep;58(3):384-97.
 8. Mamoulakis C, Ubbink DT, de la Rosette JJ. Bipolar versus monopolar transurethral resection of the prostate: a systematic review and meta-analysis of randomized controlled trials. *Eur Urol*. 2009 Nov;56(5):798-809.
 9. Teng J, Zhang D, Li Y, et al. Photoselective vaporization with the green light laser vs transurethral resection of the prostate for treating benign prostate hyperplasia: a systematic review and meta-analysis. *BJU Int*. 2013 Feb;111(2):312-23.
 10. Ruszat R, Wyler S, Forster T, et al. Safety and effectiveness of photoselective vaporization of the prostate (PVP) in patients on ongoing oral anticoagulation. *Eur Urol*. 2007 Apr;51(4):1031-8;
 11. Reich O, Schlenker B, Gratzke C, et al. Plasma vaporisation of the prostate: initial clinical results. *Eur Urol*. 2010 Apr;57(4):693-7.
 12. Geavlete B, Georgescu D, Multescu R, Stanescu F, Jecu M, Geavlete P. Bipolar plasma vaporization vs monopolar and bipolar TURP-A prospective, randomized, long-term comparison. *Urology*. 2011 Oct;78(4):930-5.
 13. Robert G, Descazeaud A, Delongchamps NB, et al. Transurethral plasma vaporization of the prostate: 3-month functional outcome and complications. *BJU Int*. 2012 Aug;110(4):555-60.
 14. Kaplan SA, Te AE. A comparative study of transurethral resection of the prostate using a modified electro-vaporizing loop and transurethral laser vaporization of the prostate. *The Journal of Urology*. 1995 Nov;154(5):1785-90.

15. Gallucci M, Puppo P, Perachino M, et al. Transurethral electrovaporization of the prostate vs. transurethral resection. Results of a multicentric, randomized clinical study on 150 patients. *Eur Urol*. 1998;33(4):359-64.
16. Botto H, Lebre T, Barre P, Orsoni JL, Herve JM, Lugagne PM. Electrovaporization of the prostate with the Gyrus device. *Journal of Endourology*. 2001 Apr;15(3):313-6.
17. Dincel C, Samli MM, Guler C, Demirbas M, Karalar M. Plasma kinetic vaporization of the prostate: clinical evaluation of a new technique. *Journal of Endourology*. 2004 Apr;18(3):293-8.
18. Dunsmuir WD, McFarlane JP, Tan A, et al. Gyrus bipolar electrovaporization vs transurethral resection of the prostate: a randomized prospective single-blind trial with 1 y follow-up. *Prostate Cancer and Prostatic Diseases*. 2003;6(2):182-6.
19. Hon NH, Brathwaite D, Hussain Z, et al. A prospective, randomized trial comparing conventional transurethral prostate resection with PlasmaKinetic vaporization of the prostate: physiological changes, early complications and long-term followup. *The Journal of Urology*. 2006 Jul;176(1):205-9.
20. Karaman MI, Kaya C, Ozturk M, Gurdal M, Kirecci S, Pirincci N. Comparison of transurethral vaporization using PlasmaKinetic energy and transurethral resection of prostate: 1-year follow-up. *Journal of Endourology*. 2005 Jul-Aug;19(6):734-7.
21. Kaya C, Ilktac A, Gokmen E, Ozturk M, Karaman IM. The long-term results of transurethral vaporization of the prostate using plasmakinetic energy. *BJU Int*. 2007 Apr;99(4):845-8.
22. Bucuras V, Bardan R. Bipolar vaporization of the prostate: is it ready for the primetime? *Therapeutic Advances in Urology*. 2011 Dec;3(6):257-61.
23. Delongchamps NB, Robert G, de la Taille A, et al. Surgical management of BPH in patients on oral anticoagulation: transurethral bipolar plasma vaporization in saline

- versus transurethral monopolar resection of the prostate. Can J Urol. 2011 Dec;18(6):6007-12.
24. Geavlete B, Multescu R, Dragutescu M, Jecu M, Georgescu D, Geavlete P. Transurethral resection (TUR) in saline plasma vaporization of the prostate vs standard TUR of the prostate: 'the better choice' in benign prostatic hyperplasia? BJU Int. 2010 Dec;106(11):1695-9.
 25. Højgaard M, Menne S, Thomsen F, FAhrenkrug L, Schou J. Photoselective Vaporization of the Prostate vs. Mushroom treatment: Clinical outcome in two Comparable Populations. Eur Urol Suppl. 2011;10(2):152-3.
 26. Mayer EK, Kroeze SG, Chopra S, Bottle A, Patel A. Examining the 'gold standard': a comparative critical analysis of three consecutive decades of monopolar transurethral resection of the prostate (TURP) outcomes. BJU Int. 2012 Dec;110(11):1595-601.
 27. Mamoulakis C, Schulze M, Skolarikos A, et al. Midterm results from an international multicentre randomised controlled trial comparing bipolar with monopolar transurethral resection of the prostate. Eur Urol. 2013 Apr;63(4):667-76.

Abbreviations Used

BPV	= Bipolar plasma vaporization
IPSS	= International Prostate Symptom Score
PAI	= Platelet aggregation inhibition
PSA	= Prostate specific antigen
Qmax	= Maximum urinary flow rate
Qol	= Quality of life
TURP	= Transurethral resection of the prostate

Table 1 - Baseline patient characteristics

Number of patients	83
Age (y)	67 (48 – 89)
Prostate volume (ml)	41 (17 – 111)
PSA (ng/ml)	2.69 (0.26 – 22.5)
IPSS	16 (2 – 35)
QoL	4 (0 – 6)
Qmax (ml/s)	10.1 (3 – 29.3)
Residual volume (ml)	87 (0 – 1000)
Indwelling catheter (n)	20 (24.1%)
ASA score	2 (1-3)
1	8 (9.6 %)
2	50 (60.2%)
3	25 (30.1%)
Coagulation modifiers (n)	26 (31.3%)
Acetylsalicylic acid (n)	26 (31.3%)
Clopidrogel (n)	1 (1.2%)
Coumarine (n)	0 (0%)
Dual therapy (n)	1 (1.2%)
Positive urine culture (n)	32 (38.5%)
Indwelling catheter (n)	18 (21.6%)
No catheter (n)	14 (16.8%)
Data presented as median (range) or number (percent). PSA = prostate-specific antigen, IPSS = International Prostate Symptom score, QoL = Quality of life, Qmax = Maximum flow rate, ASA score = American Society of Anaesthesiology score.	

Table 2: Intra- and peri-operative results

Operative time (min)	80 (34-145)
Irrigation volume (l)	22.5 (12-39)
General anesthesia (n)	60 (72.2%)
Spinal anesthesia (n)	23 (27.8%)
Blood transfusion (n)	0 (0%)
Duration of catheterization (d)	3 (2-13)
Duration of hospitalization (d)	4 (2-11)
Re-catheterization (n)	13 (15.6%)
Urinary retention (n)	10 (12%)
High residual volume + UTI (n)	2 (2.4%)
Clot retention (n)	1 (1.2%)
Data presented as median (range) or number (percent). UTI = urinary tract infection	

Table 3: Baseline and post-operative outcome parameters

Parameter	Baseline	Post-operative	6 weeks	6 months	12 months
IPSS	16 (2 – 35)	N/A	8* (0 – 29)	3* [◦] (0 – 24)	3* [◦] (0 – 20)
QoL	4 (0 – 6)	N/A	2* (0 – 6)	1* [◦] (0 – 4)	1* (0 – 4)
Qmax (ml/s)	10.1 (3 – 29.3)	14.0 (3.8 – 47)	16.7* (2.3 – 52)	16.7* (5 – 53.8)	17.2* (3.3 – 56)
Residual volume (ml)	87 (0 – 1000)	20* (0 – 900)	10* (0 – 437)	6.5* (0 – 500)	11* (0 – 190)
PSA (ng/ml)	2.69 (0.26 – 22.5)	N/A	1.36* (0.04 – 9.56)	1.27* (0.15 – 6.98)	1.11* (0.08 – 7.18)
Data presented as median (range) IPSS = International Prostate Symptom score, QoL = Quality of life, Qmax = Maximum flow rate, PSA = prostate-specific antigen, N/A = not applicable * indicates a statistically significant improvement compared to the baseline value (Wilcoxon signed-rank test) ◦ indicates a statistically significant improvement compared to the preceding assessment (Wilcoxon signed-rank test)					

Table 4: Post-operative results

Urinary tract infection	
≤ 6 weeks (n)	15 (18%)
6 weeks – 6 months (n)	4 (4.8%)
6 months – 12 months (n)	0 (0%)
Delayed clot retention (≤ 6 weeks; n)	2 (2.4%)
Dysuria	
6 weeks (n)	28 (34%)
6 months (n)	2 (2.4%)
12 months (n)	3 (3.6%)
Urethral stricture (n)	3 (3.6%)
Bladder neck sclerosis (n)	4 (4.8%)
Re-resection (n)	0 (0%)
Data presented as median (range) or number (percent).	