



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
Main Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2014

---

## **Computed tomography of the abdomen of calves during the first 105 days of life: III. Urinary tract and adrenal glands**

Braun, Ueli ; Schnetzler, C ; Augsburg, Heinz ; Bettschart-Wolfensberger, Regula ; Ohlerth, Stefanie

**Abstract:** Computed tomographic (CT) findings of the urinary tract and adrenal glands of five healthy male calves in the first 105 days of life were compared with corresponding cadaver slices. The structures seen on CT images were identified using the corresponding cadaver slices. CT produced exact images of the kidneys, urinary bladder, urethra and adrenal glands, but reliable images of the ureters were only obtained near the renal hilus. There was excellent agreement between the structures on the CT images and the tissue slices. The structure and vessels of the kidneys, the origin of the ureters, the location, size and content of the urinary bladder and the course of the urethra in the pelvis and penis were evident on images. The size and volume of the kidneys and the length and width of the adrenal glands increased significantly during the study, but the ureteral and urethral diameters changed little.

DOI: [https://doi.org/10.1024/0036-7281/a\\_000583](https://doi.org/10.1024/0036-7281/a_000583)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-98724>

Journal Article

Accepted Version

Originally published at:

Braun, Ueli; Schnetzler, C; Augsburg, Heinz; Bettschart-Wolfensberger, Regula; Ohlerth, Stefanie (2014). Computed tomography of the abdomen of calves during the first 105 days of life: III. Urinary tract and adrenal glands. *Schweizer Archiv für Tierheilkunde*, 156(5):237-247.

DOI: [https://doi.org/10.1024/0036-7281/a\\_000583](https://doi.org/10.1024/0036-7281/a_000583)

1 **Computed tomography of the abdomen of calves during the first 105 days of life: III. Urinary**  
2 **tract and adrenal glands**

3  
4 U. Braun<sup>1</sup>, C. Schnetzler<sup>1</sup>, H. Augsburger<sup>2</sup>, R. Bettschart<sup>3</sup>, S. Ohlerth<sup>4</sup>

5  
6 <sup>1</sup>Department of Farm Animals, <sup>2</sup>Institute of Veterinary Anatomy, <sup>3</sup>Division of Anaesthesiology and  
7 <sup>4</sup>Division of Diagnostic Imaging, Vetsuisse Faculty, University of Zurich

8  
9 **Summary**

10 Computed tomographic (CT) findings of the urinary tract and adrenal glands of five healthy male  
11 calves in the first 105 days of life were compared with corresponding cadaver slices. The structures  
12 seen on CT images were identified using the corresponding cadaver slices. CT produced exact  
13 images of the kidneys, urinary bladder, urethra and adrenal glands, but reliable images of the ureters  
14 were only obtained near the renal hilus. There was excellent agreement between the structures on  
15 the CT images and the tissue slices. The structure and vessels of the kidneys, the origin of the  
16 ureters, the location, size and content of the urinary bladder and the course of the urethra in the  
17 pelvis and penis were evident on images. The size and volume of the kidneys and the length and  
18 width of the adrenal glands increased significantly during the study, but the ureteral and urethral  
19 diameters changed little.

20  
21 **Keywords:** computed tomography, cattle, calf, urinary tract, kidney, urinary bladder, ureter, urethra,  
22 adrenal glands

23  
24 **Computertomographie des Abdomens von Kälbern vom ersten bis zum 105. Lebenstag: III.**  
25 **Harnapparat und Nebennieren**

26 In der vorliegenden Arbeit werden die computertomographischen (CT) Befunde an Harnapparat und  
27 Nebennieren von 5 gesunden männlichen Kälbern von der Geburt bis zum Alter von 105 Tagen  
28 beschrieben und mit denjenigen der postmortalen Untersuchung verglichen. Nieren, Harnblase,  
29 Urethra und Nebennieren konnten computertomographisch exakt dargestellt werden, während die  
30 Ureteren nur im Bereich des Hilus renalis zuverlässig identifiziert werden konnten. Die  
31 Topographie der Organe bzw. der Strukturen und der anatomischen Schnittpräparate stimmten sehr  
32 gut überein. An den Nieren waren die Struktur und die Blutgefäße, an den Ureteren deren

33 Anfangsteil, an der Harnblase Lage, Grösse und Inhalt und an der Urethra der Verlauf im Becken  
34 und Penis zu sehen. Die Ausdehnung und das Volumen der Nieren nahmen von der ersten bis zur  
35 sechsten Untersuchung signifikant zu. Das Gleiche gilt für die Länge und Breite der Nebennieren,  
36 während sich die Durchmesser von Ureteren und Urethra kaum veränderten.

37

38 Schlüsselwörter: Computertomographie, Rind, Kalb, Harnapparat, Niere, Harnblase, Ureter,  
39 Urethra, Nebennieren

40

## 41 **Introduction**

42 Computed tomography (CT) is an established imaging modality for the examination of the urinary  
43 tract in cats and dogs. It has also been used to study the urinary tract in adult goats by comparing  
44 transverse, dorsal and sagittal CT images with corresponding anatomical cadaver slices (Irmer,  
45 2010; Braun et al., 2011c). Diagnostic procedures in calves with suspected urinary tract disorders  
46 include clinical examination, urinalysis, haematological analyses, radiography, cystoscopy (in nanny  
47 goats) and ultrasonography (Weisser, 2000). Computed tomography is indispensable for the  
48 diagnosis of urinary tract disorders in humans (Dobry and Danuser, 2009) and small animals.  
49 Because there have been no CT studies of the urinary tract in calves, the goal of this investigation  
50 was to document CT findings of the urinary tract in five healthy male calves in the first 105 days of  
51 life and to compare the findings with corresponding anatomical cadaver slices.

52

## 53 **Animals, Material and Methods**

54 See communication I.

55

### 56 **Kidneys**

57 Depending on which plane provided the best images, the maximum length of the kidneys was  
58 determined in the sagittal or dorsal plane by measuring the distance between the cranial and caudal  
59 poles. The same plane was used to measure the maximum width of the kidneys, defined as the  
60 distance between the hilus and the lateral border. Multiple transverse sections were used to calculate  
61 the kidney volume (in cm<sup>3</sup>) and to determine the parenchymal density in a 4-cm<sup>2</sup> area between the  
62 hilus and the lateral border (in HU) as described in detail (Schnetzler, 2012).

63

### 64 **Ureters**

65 The diameter of both ureters was measured near the kidney in the plane that provided the best  
66 images of the structures.

67

68 Urinary bladder

69 The distance between the apex of the bladder and the pelvic brim was measured in the sagittal  
70 plane. The density of the urine was determined in a 5-cm<sup>2</sup> area 5 mm from the bladder wall.

71

72 Urethra

73 The urethral diameter was measured in the transverse plane.

74

75 Adrenal glands

76 The maximum length and width of the adrenal glands were measured in the plane that provided the  
77 best images of these organs.

78

79 **Results**

80 All structures in the CT images could be accurately identified based on transverse, sagittal and  
81 horizontal anatomical cadaver sections (Fig. 1, 2). Visual comparison of the CT images with the  
82 corresponding cadaver slices was made in the transverse plane at each vertebra from the 6th thoracic  
83 vertebra to the middle part of the sacrum (Schnetzler, 2012).

84

85 Kidneys

86 Both kidneys were easily identified in all calves based on the unique pattern afforded by the  
87 reniculi. Contrast enhancement rendered the kidneys hyperdense relative to the surrounding tissue.

88 The density of the renal cortex was greater than that of the medulla, and the renal pelvis was  
89 hypodense.

90

91 Left kidney

92 The left kidney extended from the 3rd lumbar vertebra to the sacrum but could only be identified  
93 consistently at the level of the 5th lumbar vertebra (Fig. 3). In all calves, it was seen from the 4th to  
94 the 6th lumbar vertebra at the first examination and from the 5th and 6th lumbar vertebra in the last  
95 examination. During the study period, it became displaced by the expanding dorsal sac of the rumen  
96 from the left side toward the median or sometimes into the right hemiabdomen (Fig. 4). In the dorsal

97 and sagittal planes, the left renal artery was seen running from the aorta to the kidney and in the  
98 right paramedian region, the left renal vein was observed running from the kidney to the caudal vena  
99 cava (Fig. 5, 6). Both blood vessels appeared as straight structures and were only a few centimetres  
100 in length. They had a cranioventral course to the renal hilus until the age of three weeks, when it  
101 changed to a craniodorsal course relative to the left kidney. The latter change was the result of  
102 displacement of the kidney to the right paramedian region by the rumen accompanied by a rotation  
103 of the renal hilus. The ureter was seen exiting from the hilar region. The left kidney underwent  
104 significant growth during the study period; it was 10.8 cm long and 6.0 cm wide at the first  
105 examination and 14.6 cm long and 7.1 cm wide at the last examination ( $P < 0.05$ ) (Tab. 1). During  
106 the same time period, the mean volume increased from 149.4 to 307.5 cm<sup>3</sup> ( $P < 0.05$ ). The mean  
107 parenchymal density decreased significantly from 42.8 to 27.2 HU during the same period.

108

#### 109 Right kidney

110 The right kidney was imaged from the 13th thoracic to the 6th lumbar vertebra (Fig. 7) but could  
111 only be seen consistently at the level of the 3rd lumbar vertebra. It was seen in all calves at the level  
112 of the 3rd and 4th lumbar vertebrae at the first examination and from the 2nd thoracic to 3rd lumbar  
113 vertebra at the last examination. It was located retroperitoneally, embedded in perirenal fat, always  
114 in the right hemiabdomen cranial to the left kidney and in contact with the liver in the renal  
115 depression (Fig. 8 - 10). It appeared oval in all planes. As for the left kidney, the right renal artery  
116 and vein were seen medially. The hilus and ureter exiting from the left kidney were directed  
117 ventromedially. The length, width, volume and density of the right and left kidneys were similar  
118 (Tab. 2).

119

#### 120 Ureters

121 The renal attachment of the ureter at the hilus was always seen in the right kidney (Fig. 8, 9) and in  
122 over 80 % of examinations of the left kidney. Both ureters ran caudally in a dorsomedial direction  
123 but were only occasionally identified; their site of entry into the urinary bladder was also not usually  
124 seen. The left and right ureters had similar diameters, which measured 0.3 cm on average at the first  
125 examination and 0.5 and 0.7 cm, respectively, at the last examination (Tab. 1, 2).

126

#### 127 Urinary bladder

128 The urinary bladder was identified as an oval to circular structure at the level of the sacrum on the  
129 pelvic floor beneath the uterus at all examinations (Fig. 11). Depending on the degree of fill it  
130 reached beyond the pelvic brim into the abdomen.

131

#### 132 Urethra

133 The urethra was seen exiting from the neck of the urinary bladder in all three planes. It ran caudally  
134 below the rectum to the ischiadic arc, where it turned back cranioventrally and continued along the  
135 penis to the tip of the penis in the prepuce (Fig. 12, 13). The mean urethral diameter was 1.5 cm at  
136 the first examination and 1.7 cm at the last examination (Tab. 3).

137

#### 138 Adrenal glands

139 The adrenal glands were best seen in the transverse plane and appeared as oval to bean-shaped  
140 structures. The left adrenal gland was located at the level of the cranial pole of the left kidney in the  
141 immediate vicinity of the renal vessels (Fig. 6 A). It was exactly in the median and ventral to the  
142 aorta and the caudal vena cava. The right adrenal gland was located medial to the hilus of the right  
143 kidney and immediately cranial to or between the renal vein and artery (Fig. 14, 15). From the first  
144 to the last examination, the mean length of the left adrenal gland increased from 1.01 to 1.53 cm and  
145 the width from 0.66 to 0.91 cm (Tab. 4); similar increases were recorded for the right adrenal gland.

146

#### 147 **Discussion**

148 The kidneys were readily identified in calves at all ages because of their unmistakable shape, the  
149 unique structure of the reniculi and the strong contrast enhancement. Differentiation of  
150 neighbouring organs was also aided by the surrounding hypodense perirenal fat. The size and  
151 volume of the kidneys of newborn calves were the same as those of adult goats (Braun et al.,  
152 2011c), but increased considerably by the age of 105 days. The length, width and volume of the left  
153 kidney of goats were on average 9.3 cm, 5.0 cm and 154.9 cm<sup>3</sup>, respectively (Braun et al., 2011c),  
154 compared with 14.6 cm, 7.1. cm and 307.5 cm<sup>3</sup> in 105-day-old calves. There was considerable  
155 individual variation in the density of the renal parenchyma, but there were no differences between  
156 the kidneys of individual calves. This is in agreement with the results of CT of human kidneys, in  
157 which the parenchyma appeared relatively homogeneous and had a density of 30 to 50 HU (Sagel et  
158 al., 2002). The renal density of the calves at the last examination (27.2 HU left, 30.1 HU right) was  
159 similar to that of adult goats (18.5 to 39.5 HU). A difference in density between paired kidneys of

160 more than 5 HU indicates urolithiasis in human medicine (Goldman et al., 2004). Computed  
161 tomography is the diagnostic technique of choice in people with suspected pyelonephritis (Craig et  
162 al., 2008), kidney tumours (Dobry and Danuser, 2009; Griffin et al., 2009) and kidney trauma (Lee  
163 et al., 2007), and is indispensable for the examination of patients with renal colic attributable to  
164 urolithiasis (Jindal and Ramchandani, 2007). It is likely that CT will improve the diagnosis of  
165 urinary tract diseases in calves.

166 The ureters were seen at the hilus of the right kidney in all scans and at the hilus of the left kidney in  
167 80 % of scans, but their course to and junction with the urinary bladder were only occasionally seen,  
168 in contrast to CT scans in goats (Braun et al., 2011c). We assume that dilated or otherwise diseased  
169 ureters would also be visible on CT scans in calves.

170 In contrast to the ureters, the urinary bladder and urethra were always seen at the level of the  
171 sacrum. Omphaloarteritis, omphalourachitis and various other urachal abnormalities are common  
172 problems in calves and can be diagnosed quite reliably using ultrasonography (Lischer and Steiner,  
173 1993, 1994; Lischer et al., 1994). Computed tomography should further improve images of  
174 umbilical and urachal disease processes and facilitate the planning of the surgical treatment. While  
175 ultrasonographic imaging of the urinary bladder is straightforward, imaging of the urethra of bull  
176 calves is much more difficult and reports are scant (Weisser, 2000); the penile urethra was only  
177 vaguely seen during urination. Concrement in the urethra is readily seen on radiographs and should  
178 also be visible on CT images.

179 Unlike the umbilical vein, which could be seen on CT scans until the age of three weeks  
180 (communication I), the umbilical artery was never identified caudal to the umbilicus, which was in  
181 agreement with ultrasonographic findings (Lischer, 1991; Watson et al., 1994).

182 The adrenal glands were reliably seen on CT scans, although their measurements varied greatly.  
183 Measurements of smaller structures are less precise than those of larger ones (Voorhout, 1990), but  
184 also depend on the position of the organ relative to the tomographic plane (Voorhout, 1990). The  
185 adrenal glands did not grow significantly during the study period and their size at the age of 105  
186 days corresponded to measurements obtained in adult goats and dogs (Assheuer and Sager, 1997;  
187 Braun et al., 2011c). In the calf, adrenal disease is rare and CT examination of the adrenal glands is  
188 of little clinical importance. This is in contrast to the goat, in which adrenal cortical adenoma  
189 (Smith and Sherman, 2009), medullary pheochromocytoma (De Gritz, 1997) and adrenal  
190 hypocortisolism (Swart et al., 1996; Engelbrecht et al., 2000) have been reported, although adrenal  
191 disease is generally considered to be rare.

192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223

## **Conclusions**

This study has shown that CT is very useful for the examination of the urogenital tract of calves. The transverse, horizontal and sagittal cadaver sections were essential for identification of the structures on CT images. This study provides reference values for CT examination of calves with suspected diseases of the urinary tract and adrenal glands.

## **References**

- Ahvenjärvi, S., Skiba, B. Huhtanen, P.:* Effect of heterogenous digesta chemical composition on the accuracy of measurements of fiber flow in dairy cows. *J. Anim. Sci.* 2001, 79: 1611-1620.
- Assheuer, J., Sager, M.:* MRI and CT Atlas of the Dog. Blackwell Science, Berlin 347-421, 1997.
- Becker, M., Heun, C., Tsousis, G., Bollwein, H.:* Application of computed tomography for the evaluation of obstetrically relevant measurements in German Holstein-Friesian calves. *Theriogenology* 2011, 75: 1052-1056.
- Braun, U., Marmier, O.:* Ultrasonographic examination of the small intestine of cows. *Vet. Rec.* 1995, 136: 239-244.
- Braun, U., Amrein, E.:* Ultrasonographic examination of the caecum and the proximal and spiral ansa of the colon of cattle. *Vet. Rec.* 2001, 149: 45-48.
- Braun, U., Blessing, S.:* Ultrasonographic examination of the omasum in 30 healthy cows. *Vet. Rec.* 2006, 159: 812-815.
- Braun, U.:* Ultrasonography of the gastrointestinal tract in cattle. *Vet. Clin. North Am. (Food Anim. Pract.)* 2009, 25: 567-590.
- Braun, U., Irmer, M., Augsburg, H., Jud, R., Ohlerth, S.:* Computed tomography of the abdomen in Saanen goats: I. Reticulum, rumen and omasum. *Schweiz. Arch. Tierheilk.* 2011a, 153: 307-313.



224 *Braun, U., Irmer, M., Augsburg, H., Müller, U., Jud, R., Ohlerth, S.:* Computed tomography of  
225 the abdomen in Saanen goats: II. Liver, spleen, abomasum, and intestine. *Schweiz. Arch. Tierheilk.*  
226 2011b, 153: 314-320.  
227

228 *Braun, U., Irmer, M., Augsburg, H., Ohlerth, S.:* Computed tomography of the abdomen in  
229 Saanen goats: III. Kidneys, ureters and urinary bladder. *Schweiz. Arch. Tierheilk.* 2011c, 153: 321-  
230 329.  
231

232 *Braun, U., Gautschi, A.:* Ultrasonography of the reticulum, rumen, omasum and abomasum in 10  
233 calves before, during and after ingestion of milk. *Schweiz. Arch. Tierheilk.* 2012, 154: 287-297.  
234

235 *Braun, U., Gautschi, A., Tschuor, A., Hässig, M.:* Ultrasonography of the reticulum, rumen,  
236 omasum and abomasum before, during and after ingestion of hay and grass silage in 10 calves. *Res.*  
237 *Vet. Sci.* 2012, 93: 1407-1412.  
238

239 *Braun, U., Schweizer, A., Trösch, L.:* Ultrasonography of the rumen of dairy cows. *BMC Vet. Res.*  
240 2013, 9:44.  
241

242 *Breiman, R. S., Beck, J. W., Korobkin, M., Glenny, R., Akwari, O. E., Heaston, D. K., Moore, A. V.,*  
243 *Ram, P. C.:* Volume determinations using computed tomography. *Am. J. Roentgenol.* 1982, 138:  
244 329-333.  
245

246 *Buczinski, S., Duval, J., D'Anjou, M. A., Francoz, D., Fecteau, G.:* Portocaval shunt in a calf:  
247 Clinical, pathologic, and ultrasonographic findings. *Can. Vet. J.* 2007, 48: 407-410.  
248

249 *Craig, W. D., Wagner, B. J., Travis, M. D.:* Pyelonephritis: radiologic-pathologic review.  
250 *Radiographics* 2008, 28: 255-277.  
251

252 *De Gritz, B. G.:* Hereditary caprine pheochromocytoma. *Zbl. Vetmed. A* 1997, 44: 313-316.  
253

254 *Dirksen, G.:* Krankheiten von Haube und Pansen bei Milchkalb und Jungrind. In: Innere Medizin  
255 und Chirurgie des Rindes. Hrsg. G. Dirksen, H.-D. Gründer, M. Stöber. Parey Buchverlag, Berlin,  
256 2002, 455-469.

257

258 *Dobry, E., Danuser, H.:* Bildgebung der Nieren und Harnwege. Ther. Umsch. 2009, 66: 39-42.

259

260 *Engelbrecht, Y., Herselman, T., Louw, A., Swart, P.:* Investigation of the primary cause of  
261 hypoadrenocorticism in South African angora goats (*Capra aegagrus*): a comparison with Boer goats  
262 (*Capra hircus*) and Merino sheep (*Ovis aries*). J. Anim. Sci. 2000, 78: 371-379.

263

264 *Fike, J. R., Drury, E. M., Zook, B. C., Davis, D. O., Thompson, J. E., Chaney, E., Bradley, E. W.:*  
265 Canine anatomy as assessed by computerized tomography. Am. J. Vet. Res. 1980, 41: 1823-1832.

266

267 *Flöck, M.:* Ultraschalldiagnostik von Entzündungen der Nabelstrukturen, persistierendem Urachus  
268 und Umbilikalhernie beim Kalb. Berl. Münch. Tierärztl. Wschr. 2003, 116: 2-11.

269

270 *Gautschi, A.:* Sonographische Untersuchungen an Haube, Pansen, Psalter und Labmagen von 30  
271 Kälbern. Dissertation, Universität Zürich, 2010.

272

273 *Goldman, S. M., Faintuch, S., Ajzen, S. A., Christofalo, D. M., Araújo, M. P., Ortiz, V., Srougi, M.,*  
274 *Kenney, P. J., Szejnfeld, J.:* Diagnostic value of attenuation measurements of the kidney on  
275 unenhanced helical CT of obstructive ureterolithiasis. Am. J. Roentgenol. 2004, 182: 1251-1254.

276

277 *Gore, R. M., Balthazar, E. J., Ghahremani, G. G., Miller, F. H.:* CT features of ulcerative colitis  
278 and Crohn's disease. Am. J. Roentgenol. 1996, 167: 3-15.

279

280 *Griffin, N., Grant, L. A., Bharwani, N., Sohaib, S. A.:* Computed tomography in metastatic renal cell  
281 carcinoma. Semin. Ultrasound CT MRI 2009, 30: 359-366.

282

283 *Heidemann, A., Grunert, E.:* Ultraschalldiagnostik als Entscheidungshilfe für das weitere Vorgehen  
284 bei Nabelentzündungen des neugeborenen Kalbes. Prakt. Tierarzt 1995, 76: 742-746.

285

286 *Hummel, J., K.-H. Südekum, D. Bayer, S. Ortmann, W. J. Streich, J.-M. Hatt, M. Clauss:* Physical  
287 characteristics of reticuloruminal contents of oxen in relation to forage type and time after feeding.  
288 *J. Anim. Physiol. Anim. Nutr.* 2009, 93: 209-220.  
289

290 *Irmer, M.:* Computertomographische Untersuchung des Abdomens bei 30 Ziegen. Dissertation,  
291 Universität Zürich, 2010.  
292

293 *James, S., Balfe, D. M., Lee, J. K. T., Picus, D.:* Small-bowel disease: Categorization by CT  
294 examination. *Am. J. Roentgenol.* 1987, 148: 863-868.  
295

296 *Jindal, G., Ramchandani, P.:* Acute flank pain secondary to urolithiasis: radiologic evaluation and  
297 alternate diagnoses. *Radiol. Clin. North Am.* 2007, 45: 395-410, vii.  
298

299 *Kayaalp, C., Arda, K., Oto, A., Oran, M.:* Liver volume measurement by spiral CT: an in vitro  
300 study. *Clin. Imaging* 2002, 26: 122-124.  
301

302 *König, H. E., Sautet, J., Liebich, H.-G.:* Anatomie der Haussäugetiere. Schattauer Verlag, Stuttgart,  
303 2005.  
304

305 *Kovács, P. L., Südekum, K. H., Stangassinger, M.:* Effects of intake level of a mixed diet on  
306 chewing activity and on particle size of ruminated boli, ruminal digesta fractions and faeces of  
307 steers. *Reprod. Nutr. Dev.* 1997, 37: 517-528.  
308

309 *Krüger, S. S.:* Sonographische Untersuchungen an Haube, Pansen, Psalter, Labmagen, Milz und  
310 Leber von Kälbern von der Geburt bis zum Alter von 100 Tagen. Dissertation, Universität Zürich,  
311 2012.  
312

313 *Lee, Y. J., Oh, S. N., Rha, S. E., Byun, J. Y.:* Renal trauma. *Radiol. Clin. North Am.* 2007, 45: 581-  
314 592, ix.  
315

316 *Lee, K., Yamada, K., Tsuneda, R., Kishimoto, M., Shimizu, J., Kobayashi, Y., Furuoka, H., Matsui,*  
317 *T., Sasaki, N., Ishii, M., Inokuma, H., Iwasaki, T., Miyake, Y.:* Clinical experience of using  
318 multidetector-row CT for the diagnosis of disorders in cattle. *Vet. Rec.* 2009, 165: 559-562.

319

320 *Lee, K., Ikeo, S., Kishimoto, M., Shimizu, J., Iwasaki, T., Miyake, Y., Yamada, K.:* Use of a saline  
321 chaser to reduce the dose of contrast material for dynamic CT in cattle. *Vet. Rec.* 2010, 167: 170-  
322 173.

323

324 *Lischer, C.:* Sonographische Darstellung der Nabelinvolution beim Kalb. Dissertation, Universität  
325 Zürich, 1991.

326

327 *Lischer, C. J., Steiner, A.:* Ultrasonography of umbilical structures in calves. Part I:  
328 Ultrasonographic description of umbilical involution in clinically healthy calves. *Schweiz. Arch.*  
329 *Tierheilk.* 1993, 135: 221-230.

330

331 *Lischer, C. J., Steiner, A.:* Ultrasonography of the umbilicus in calves. Part 2: Ultrasonography,  
332 diagnosis and treatment of umbilical diseases. *Schweiz. Arch. Tierheilk.* 1994, 136: 227-241.

333

334 *Lischer, C. J., Iselin, U., Steiner, A.:* Ultrasonographic diagnosis of urachal cyst in three calves. *J.*  
335 *Am. Vet. Med. Assoc.* 1994, 204: 1801-1804.

336

337 *Lubbers, B. V., Apley, M. D., Coetzee, J. F., Mosier, D. A., Biller, D. S., Mason, D. E., Henao-*  
338 *Guerrero, P. N.:* Use of computed tomography to evaluate pathologic changes in the lungs of calves  
339 with experimentally induced respiratory tract disease. *Am. J. Vet. Res.* 2007, 68: 1259-1264.

340

341 *Mortele, K. J., McTavish, J., Ros, P. R.:* Current techniques of computed tomography, helical CT,  
342 multidetector CT, and 3D reconstruction. *Clin. Liver Dis.* 2002, 6: 29-52.

343

344 *Nickel, R., Schummer, A., Seiferle, E.:* Eingeweide. In: *Lehrbuch der Anatomie der Haustiere*, Vol.  
345 2. Eds. J. Frewein, H. Gasse, R. Leiser, H. Roos, H. Thomé, B. Vollmerhaus, H. Waibl. Parey  
346 Buchverlag, Stuttgart, 2004, 103-434.

347

348 *Noda, T., Todani, T., Watanabe, Y., Yamamoto, S.:* Liver volume in children measured by computed  
349 tomography. *Pediatr. Radiol.* 1997, 27: 250-252.

350

351 *Nuss, K., Schnetzler, C., Hagen, R., Schwarz, A., Kircher, P.:* Klinische Anwendung der  
352 Computertomographie beim Rind. *Tierärztl. Prax.* 2011, 39 (G): 317-324.  
353

354 *Radostits, O. M., Gay, C. C., Hinchcliff, K. W., Constable, P. D.:* *Veterinary Medicine. A Textbook*  
355 *of the Diseases of Cattle, Horses, Sheep, Pigs, and Goats.* Philadelphia, Saunders Elsevier, 2007.  
356

357 *Rivero, M. A., Vázquez, J. M., Gil, F., Ramírez, J. A., Vilar, J. M., De Miguel, A., Arencibia, A.:*  
358 *CT-soft tissue window of the cranial abdomen in clinically normal dogs: An anatomical description*  
359 *using macroscopic cross-sections with vascular injection.* *Anat. Histol. Embryol.* 2009, 38: 18-22.  
360

361 *Sagel, S. S., Stanley, R. J., Levitt, R. G., Geisse, G.:* *Computed tomography of the kidney.* 1977. *J.*  
362 *Urol.* 2002, 167: 458-468.  
363

364 *Salomon, F.-V., Geyer, H., Gille, U.:* *Verdauungsapparat.* In: *Anatomie für die Tiermedizin.* Enke  
365 *Verlag, Stuttgart, 2008, 235-323.*  
366

367 *Samii, V. F., Biller, D. S., Koblik, P. D.:* *Normal cross-sectional anatomy of the feline thorax and*  
368 *abdomen: comparison of computed tomography and cadaver anatomy.* *Vet. Radiol. Ultrasound*  
369 *1998, 39: 504-511.*  
370

371 *Schnetzler, C.:* *Computertomographische Untersuchung des Abdomens von Kälbern von der Geburt*  
372 *bis zum Alter von 105 Tagen.* Dissertation, Universität Zürich, 2012.  
373

374 *Schnorr, B., Kressin, M.:* *Entwicklung der Verdauungsorgane.* In: *Embryologie der Haustiere.* Enke  
375 *Verlag, Stuttgart, 2011, 161-185.*  
376

377 *Steiner, A., Lischer, C. J., Oertle, C.:* *Marsupialization of umbilical vein abscesses with*  
378 *involvement of the liver in 13 calves.* *Vet. Surg.* 1993, 22: 184-189.  
379

380 *Stieger, S. M., Zwingenberger, A., Pollard, R. E., Kyles, A. E., Wisner, E. R.:* *Hepatic volume*  
381 *estimation using quantitative computed tomography in dogs with portosystemic shunts.* *Vet. Radiol.*  
382 *Ultrasound* 2007, 48: 409-413.

383

384 *Smith M. C., Sherman, D. M.:* Goat Medicine. Wiley-Blackwell, 2009.

385

386 *Swart, P., Engelbrecht, Y., Herselmann, T.:* An investigation of hypo-adrenocorticism in Angora  
387 goats. *Endocr. Res.* 1996, 22: 563-565.

388

389 *Teixeira, M., Gil, F., Vazquez, J. M., Cardoso, L., Arencibia, A., Ramirez-Zarzosa, G., Agut, A.:*  
390 Helical computed tomographic anatomy of the canine abdomen. *Vet. J.* 2007, 174: 133-138.

391

392 *Thrall, D. E.:* Textbook of Veterinary Diagnostic Radiology. W. B. Saunders, Philadelphia, 2007,  
393 625-802.

394

395 *Tschuor, A., Clauss, M.:* Investigations on the stratification of forestomach contents in ruminants:  
396 an ultrasonographic approach. *Eur. J. Wildl. Res.* 2008, 54: 627-633.

397

398 *Urata, K., Kawasaki, S., Matsunami, H., Hashikura, Y., Ikegami, T., Ishizone, S., Momose, Y.,*  
399 *Komiyama, A., Makuuchi, M.:* Calculation of child and adult standard liver volume for liver  
400 transplantation. *Hepatology* 1995, 21: 1317-1321.

401

402 *Voorhout, G.:* X-ray-computed tomography, nephrotomography, and ultrasonography of the adrenal  
403 glands of healthy dogs. *Am. J. Vet. Res.* 1990, 51: 625-631.

404

405 *Watson, E., Mahaffey, M. B., Crowell, W., Selcer, B. A., Morris, D. D., Seginak, L.:*  
406 Ultrasonography of the umbilical structures in clinically normal calves. *Am. J. Vet. Res.* 1994, 55:  
407 773-780.

408

409 *Weisser, P.:* Sonographische Befunde am Harnapparat des gesunden Kalbes im Alter von 1 bis 6  
410 Monaten. Dissertation, Universität München, 2000.

411

## 412 **Correspondence**

413 Ueli Braun, Departement für Nutztiere, Winterthurerstrasse 260, CH-8057 Zürich, E-Mail:

414 [ubraun@vetclinics.uzh.ch](mailto:ubraun@vetclinics.uzh.ch); Fax: ++41 44 63 58 904

415 **Legend to figures**

416 Figure 1: Comparison of a transverse CT image in a soft-tissue window setting (A) and the  
417 corresponding anatomical slice (B) at the level of the 3rd lumbar vertebra in a 104-day-old Holstein-  
418 Friesian bull calf. 1 Right kidney, 2 Renal sinus, 3 Renal vein, 4 Right ureter, 5 Liver, 6 Rumen, 7  
419 Abomasum, 8 Small intestine, 9 Large intestine, 10 Caudal vena cava, L Left, R Right.

420

421 Figure 2: Comparison of a transverse CT image in a soft-window setting (A) and the corresponding  
422 anatomical slice (B) at the level of the 6th lumbar vertebra in a 104-day-old Holstein-Friesian bull  
423 calf. 1 Left kidney, 2 Left ureter, 3 Caudal vena cava, 4 Aorta, 5 Rumen, 6 Small intestine, 7 Large  
424 intestine, L Left, R Right.

425

426 Figure 3: Visibility of the left kidney on transverse CT images in five Holstein-Friesian bull calves.  
427 The images were taken at different lumbar vertebrae and the sacrum during six CT scans in the first  
428 104 days of life. The different shades of red indicate the number of calves in which the kidney was  
429 visible at the respective levels. 20 %, 40 %, 60 %, 80 % and 100 %, visible in 1, 2, 3, 4 and in all  
430 calves, respectively.

431

432 Figure 4: Transverse CT images of the abdomen at the level of the 1st lumbar vertebra in a one-day-  
433 old (A) and at the level of the 6th lumbar vertebra in a 104-day-old Holstein-Friesian bull calf (B). 1  
434 Left kidney, 2 Right kidney, 3 Aorta, 4 Left ureter, 5 Rumen, 6 Abomasum, 7 Intestines, L Left, R  
435 Right.

436

437 Figure 5: Dorsal CT images of the abdomen at the level of the aorta in a 21-day-old (A) and 103-  
438 day-old Holstein-Friesian bull calf (B). 1 Left kidney, 2 Right kidney, 3 Aorta, 4 Left renal vein, 5  
439 Left renal artery, 6 Liver, 7 Spleen, 8 Rumen, 9 Intestines, L Left, R Right.

440

441 Figure 6: Sagittal CT images of the abdomen at the level of the left adrenal gland-in a two-day-old  
442 (A) and at the level of the caudal vena cava in a 103-day-old Holstein-Friesian bull calf (B). 1 Left  
443 kidney, 2 Right kidney, 3 Caudal vena cava, 4 Liver, 5 Left renal vein, 6 Left adrenal gland, 7  
444 Urinary bladder, 8 Abomasum, Cr Cranial, Cd Caudal.

445

446 Figure 7: Visibility of the right kidney on transverse CT images in five Holstein-Friesian bull  
447 calves. The images were taken at different thoracic and lumbar vertebrae during six CT scans in the  
448 first 105 days of life. The different shades of red indicate the number of calves in which the right  
449 kidney was visible at the respective levels. 20 %, 40 %, 60 %, 80 % and 100 %, visible in 1, 2, 3, 4  
450 and in all calves, respectively.

451

452 Figure 8: Transverse CT image of the abdomen at the level of the 4th lumbar vertebra in a one-day-  
453 old Holstein-Friesian bull calf. 1 Right kidney, 2 Right ureter, 3 Caudal vena cava, 4 Aorta, 5 Left  
454 kidney, 6 Liver, 7 Abomasum, L Left, R Right.

455

456 Figure 9: Dorsal CT image at the level of the aorta in a 104-day-old calf. 1 Right kidney, 2 Aorta, 3  
457 Right renal vein, 4 Left kidney, 5 Urinary bladder, 6 Liver, L Left, R Right.

458

459 Figure 10: Sagittal CT image at the level of the right hip joint in a 104-day-old Holstein-Friesian  
460 bull calf. 1 Right kidney, 2 Right ureter, 3 Caudal vena cava, 4 Liver, 5 Reticulum, 6 Abomasum, 7  
461 Omasum, 8 Rumen, Cr Cranial, Cd Caudal.

462

463 Figure 11: Transverse CT image at the level of the sacrum in a one-day-old Holstein-Friesian bull  
464 calf. 1 Urinary bladder, 2 Right ureter, 3 Left ureter, 4 Rectum, L Left, R Right.

465

466 Figure 12: Transverse CT image at the level of the pelvis in a two-day-old Holstein-Friesian bull  
467 calf. 1 Penis with urethra, 2 Testes, 3 Rectum, L Left, R Right.

468

469 Figure 13: Sagittal CT image at the level of the aorta in a 40-day-old Holstein-Friesian bull calf. 1  
470 Urethra, 2 Urinary bladder, 3 Testis, 4 Left kidney, 5 Aorta, Cr Cranial, Cd Caudal.

471

472 Figure 14: Transverse CT image at the level of the 2nd lumbar vertebra in a 104-day-old Holstein-  
473 Friesian bull calf. 1 Right adrenal gland, 2 Caudal vena cava, 3 Right renal vein, 4 Aorta, 5 Right  
474 kidney, 6 Liver, 7 Abomasum, L Left, R Right.

475



476 Figure 15: Dorsal CT image at the level of the aorta in a one-day-old Holstein-Friesian bull calf. 1  
477 Right adrenal gland, 2 Caudal vena cava, 3 Right renal vein, 4 Aorta, 5 Right kidney, 6 Liver, 7 Left  
478 kidney, 8 Spleen, 9 Rumen, L Left, R Right.

479 *Table 1: CT measurements of the left kidney and left ureter in five Holstein-Friesian bull calves during the first 105 days of life (mean ± sd, range).*

Variable	Examination					
	1	2	3	4	5	6
Length (cm)	10.8 ± 1.05 (9.2 – 11.8)	11.0 ± 1.72 <sup>a</sup> (9.0 – 12.6)	12.2 ± 1.57 <sup>a</sup> (10.4 – 13.5)	13.2 ± 1.11 <sup>b,§</sup> (11.6 – 14.0)	14.0 ± 1.13 <sup>b</sup> (12.7 – 15.5)	14.6 ± 1.09 (12.8 – 15.7)
Width (cm)	6.0 ± 0.65 (5.0 – 6.7)	5.9 ± 1.19 <sup>a</sup> (4.8 – 7.8)	7.6 ± 0.33 <sup>a,§</sup> (7.1 – 8.0)	7.6 ± 0.57 (7.0 – 8.5)	7.0 ± 0.93 (6.0 – 8.3)	7.1 ± 0.91 (6.3 – 8.4)
Volume (cm <sup>3</sup> )	149.4 ± 23.64 <sup>a</sup> (116.5 – 177.2)	185.4 ± 29.51 <sup>a,b,§</sup> (142.0 – 224.4)	248.3 ± 26.63 <sup>b,c</sup> (217.4 – 283.7)	310.3 ± 28.5 <sup>c</sup> (278.4 – 348.9)	287.9 ± 31.97 <sup>d</sup> (248.7 – 332.9)	307.5 ± 32.67 <sup>d</sup> (272.8 – 358.0)
Parenchymal density (HU)	42.8 ± 3.15 (39.1 – 46.6)	44.6 ± 5.94 (34.9 – 49.2)	38.2 ± 4.77 (33.7 – 44.4)	38.8 ± 7.80 (29.7 – 50.3)	30.9 ± 11.7 <sup>§</sup> (14.0 – 46.4)	27.2 ± 11.81 (16.6 – 42.9)
Ureteral diameter (cm)	0.3 ± 0.08 (0.2 – 0.4)	0.3 ± 0.07 (0.2 – 0.4)	0.3 ± 0.07 <sup>o,a</sup> (0.3 – 0.4)	0.3 ± 0.13 <sup>a</sup> (0.2 – 0.6)	0.5 ± 0.34 <sup>o</sup> (0.3 – 1.0)	0.5 ± 0.12 <sup>o,§</sup> (0.4 – 0.7)

480

481 <sup>o</sup> Measured in 4 of 5 calves

482 <sup>a, b, c, d</sup> Within rows measurements with identical indices are different (P < 0.05)

483 <sup>§</sup> First significant difference compared with examination 1 (P < 0.05)

484

485 *Table 2: CT measurements of the right kidney and left ureter in five Holstein-Friesian bull calves during the first 105 days of life (mean  $\pm$  sd, range).*

Variable	Examination					
	1	2	3	4	5	6
Length (cm)	11.8 $\pm$ 1.03 (10.2 – 12.7)	11.9 $\pm$ 1.51 <sup>a</sup> (9.5 – 13.3)	12.9 $\pm$ 1.52 <sup>a,b,§</sup> (10.4 – 14.2)	14.3 $\pm$ 1.17 <sup>b,c</sup> (12.5 – 15.7)	15.0 $\pm$ 1.32 <sup>c</sup> (13.0 – 16.7)	15.7 $\pm$ 1.30 (13.7 – 17.0)
Width (cm)	6.4 $\pm$ 0.82 (5.4 – 7.2)	6.5 $\pm$ 0.77 <sup>a</sup> (5.5 – 7.5)	7.1 $\pm$ 1.10 <sup>a</sup> (5.3 – 8.3)	8.0 $\pm$ 0.74 <sup>§</sup> (7.3 – 8.8)	7.7 $\pm$ 0.52 (7.2 – 8.6)	8.0 $\pm$ 0.77 (7.2 – 9.2)
Volume (cm <sup>3</sup> )	147.5 $\pm$ 20.82 <sup>a</sup> (125.6 – 182.0)	179.1 $\pm$ 26.55 <sup>a,b,§</sup> (143.0 – 202.4)	238.0 $\pm$ 19.08 <sup>b,c</sup> (213.3 – 262.1)	292.4 $\pm$ 17.00 <sup>c</sup> (272.7 – 317.7)	270.9 $\pm$ 41.77 <sup>d</sup> (225.0 – 339.2)	288.7 $\pm$ 43.22 <sup>d</sup> (260.1 – 365.3)
Parenchymal density (HU)	45.2 $\pm$ 1.05 (44.6 – 47.1)	44.4 $\pm$ 5.53 (36.4 – 51.2)	35.9 $\pm$ 6.73 <sup>§</sup> (29.7 – 44.0)	34.6 $\pm$ 8.6 (23.6 – 46.2)	30.5 $\pm$ 3.50 (26.8 – 34.5)	30.1 $\pm$ 11.64 (21.5 – 48.6)
Ureteral diameter (cm)	0.3 $\pm$ 0.07 (0.2 – 0.4)	0.3 $\pm$ 0.05 (0.3 – 0.4)	0.4 $\pm$ 0.20 (0.3 – 0.7)	0.4 $\pm$ 0.24 (0.3 – 0.9)	0.6 $\pm$ 0.20 <sup>§</sup> (0.4 – 0.8)	0.7 $\pm$ 0.06 (0.6 – 0.8)

486

487 <sup>a, b, c, d</sup> Within rows measurements with identical indices are different (P < 0.05)

488 <sup>§</sup> First significant difference compared with examination 1 (P < 0.05)

489 *Table 3: CT measurements of the urinary bladder and urethra in five Holstein-Friesian bull calves during the first 105 days of life (mean ± sd,*  
 490 *median, range).*

Variable	Examination					
	1	2	3	4	5	6
Distance between apex of bladder and pelvic brim	6.5 ± 3.52 (4.0 – 12.6)	6.1 ± 1.97 (2.9 – 7.8)	7.3 ± 2.33 (5.0 – 10.5)	7.9 ± 4.0 (2.9 – 12.8)	6.4 ± 1.97* (4.3 – 8.2)	4.4 ± 0.35 <sup>#,§</sup> (4.2 – 4.7)
Urinary density (HU)	10.6 (7.4 – 39.2)	9.5 (5.6 – 15.6)	10.8 (-2.7 – 18.8)	7.5 (1.2 – 37.7)	20.0 (-3.0 – 30.2)	27.5 <sup>#</sup> (21.4 – 33.5)
Urethral diameter (cm)	1.5 ± 0.33 (1.1 – 1.9)	1.5 ± 0.27 (1.2 – 1.9)	1.4 ± 0.27 (1.1 – 1.8)	1.6 ± 0.16 (1.4 – 1.8)	1.5 ± 0.24 (1.1 – 1.7)	1.7 ± 0.24 <sup>°</sup> (1.5 – 2.0)

491

492 <sup>°</sup> Measured in 4 of 5 calves

493 <sup>\*</sup> Measured in 3 of 5 calves

494 <sup>#</sup> Measured in 2 of 5 calves

495 *Table 4: CT measurements of the adrenal glands in five Holstein-Friesian bull calves during the first 105 days of life (mean ± sd, range, all*  
 496 *measurements in cm).*

Variable	Untersuchung					
	1	2	3	4	5	6
Length of right gland	1.27 ± 0.23 (0.89 – 1.45)	1.46 ± 0.46 (0.96 – 2.06)	1.47 ± 0.32 (1.18 – 1.95)	1.65 ± 0.26 <sup>§</sup> (1.30 – 1.96)	1.69 ± 0.38 <sup>°</sup> (1.26 – 2.04)	1.78 ± 0.31 (1.38 – 2.13)
Width of right gland	0.77 ± 0.17 0.53 – 0.98	0.87 ± 0.34 0.58 – 1.44	0.85 ± 0.22 (0.49 – 1.10)	0.85 ± 0.22 (0.51 – 1.06)	0.89 ± 0.17 <sup>§</sup> (0.71 – 1.10)	0.92 ± 0.21 (0.68 – 1.18)
Length of left gland	1.01 ± 0.14 (0.88 – 1.23)	0.99 ± 0.28 (0.68 – 1.32)	1.33 ± 0.23 <sup>§</sup> (0.96 – 1.58)	1.42 ± 0.64 (0.90 – 2.46)	1.28 ± 0.22 <sup>°,*</sup> (0.98 – 1.53)	1.53 ± 0.21 (1.39 – 1.90)
Width of left gland	0.66 ± 0.06 (0.59 – 0.75)	0.55 ± 0.13 <sup>a</sup> (0.36 – 0.69)	0.76 ± 0.17 <sup>a</sup> (0.55 – 1.02)	0.94 ± 0.33 (0.52 – 1.28)	0.74 ± 0.22 (0.51 – 0.98)	0.91 ± 0.15 <sup>§</sup> (0.77 – 1.13)

497

498 <sup>°</sup> Measured in 4 of 5 calves

499 <sup>a</sup> Measurements are different (P < 0.05)

500 <sup>§</sup> First significant difference compared with examination 1 (P < 0.05)

501 <sup>\*</sup> Significant difference compared with right adrenal gland (P < 0.05)